

Erratum to: Immunogenicity and antigenic relationships among spike proteins of porcine epidemic diarrhea virus subtypes G1 and G2

Xiaobo Wang¹ · Jianfei Chen¹ · Da Shi¹ · Hongyan Shi¹ · Xin Zhang¹ ·
Jing Yuan¹ · Shibo Jiang² · Li Feng¹

Published online: 3 February 2016
© Springer-Verlag Wien 2016

Erratum to: Arch Virol DOI 10.1007/s00705-015-2694-6

In the original publication, Table 2 was overlooked by the author during proof checking. The correct version of Table 2 is given below:

Table 2 The luminescence intensity of fluorescence of anti-S polyclonal antibodies against different types of virus

	Dilution of anti-CV777 S polyclonal antisera						
	1600	3200	6400	12800	25600	51200	102400
CV777	+++	+++	++	++	+	+	-
LNCT2	+++	+++	++	++	+	-	-
Vero E6	-	-	-	-	-	-	-
	Dilution of anti-LNCT2 S polyclonal antisera						
	1600	3200	6400	12800	25600	51200	102400
LNCT2	+++	+++	++	++	+	+	-
CV777	+++	+++	++	++	+	-	-
Vero E6	-	-	-	-	-	-	-

The luminescence intensity was divided into four degrees: (++++) strong, (++) moderate, (+) weak and (-) negative

Also, the text in page 7 should read as below:

The average titer of anti-SLNCT2 PAbs neutralizing CV777 reached 1:230, while the titer of neutralizing LNCT2 was 1:387.

The online version of the original article can be found under doi:10.1007/s00705-015-2694-6.

✉ Li Feng
fl@hvri.ac.cn

¹ Division of Swine Infectious Diseases, State Key Laboratory of Veterinary Biotechnology, Harbin Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Harbin 150001, People's Republic of China

² Key Laboratory of Medical Molecular Virology of Ministries of Education and Health, Shanghai Public Health Clinical Center and Shanghai Medical College, Fudan University, Shanghai 200032, People's Republic of China