



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Associations between mortality from COVID-19 in two Italian regions and outdoor air pollution as assessed through tropospheric nitrogen dioxide



Tommaso Filippini^a, Kenneth J. Rothman^{b,c}, Silvia Cocchio^d, Elena Narne^e, Domenico Mantoan^f, Mario Saia^e, Alessia Goffi^g, Fabrizio Ferrari^g, Giuseppe Maffei^g, Nicola Orsini^h, Vincenzo Baldo^d, Marco Vinceti^{a,c,*}

^a Environmental, Genetic and Nutritional Epidemiology Research Center (CREAGEN), Section of Public Health, Department of Biomedical, Metabolic and Neural Sciences, University of Modena and Reggio Emilia, Modena, Italy

^b RTI Health Solutions, Research Triangle Park, NC, USA

^c Department of Epidemiology, Boston University School of Public Health, Boston, MA, USA

^d Department of Cardiac Thoracic Vascular Sciences and Public Health, University of Padua, Italy

^e Azienda Zero of Veneto Region, Padua, Italy

^f Department for Health and Social Affairs of Veneto Region, Venice, Italy


^g TerrAria, Milan, Italy

^h Department of Global Public Health, Karolinska Institutet, Stockholm, Sweden

HIGHLIGHTS

- Air pollution may increase susceptibility to SARS-CoV-2 infection.
- We examined COVID-19 mortality in the two regions in Northern Italy.
- Satellite-detected tropospheric nitrogen dioxide was used to assess air pollution.
- We found a positive association between NO₂ levels and COVID-19 mortality rates.

GRAPHICAL ABSTRACT



ARTICLE INFO

Article history:

Received 22 September 2020

Received in revised form 24 October 2020

Accepted 25 October 2020

Available online 4 November 2020

Editor: Damia Barcelo

Keywords:

Covid-19
Coronavirus
Air pollution
Nitrogen dioxide
Mortality
Public health

ABSTRACT

After the appearance of COVID-19 in China last December 2019, Italy was the first European country to be severely affected by the outbreak. The first diagnosis in Italy was on February 20, 2020, followed by the establishment of a light and a tight lockdown on February 23 and on March 8, 2020, respectively. The virus spread rapidly, particularly in the North of the country in the 'Padan Plain' area, known as one of the most polluted regions in Europe. Air pollution has been recently hypothesized to enhance the clinical severity of SARS-CoV-2 infection, acting through adverse effects on immunity, induction of respiratory and other chronic disease, upregulation of viral receptor ACE-2, and possible pathogen transportation as a virus carrier. We investigated the association between air pollution and subsequent COVID-19 mortality rates within two Italian regions (Veneto and Emilia-Romagna). We estimated ground-level nitrogen dioxide through its tropospheric levels using data available from the Sentinel-5P satellites of the European Space Agency Copernicus Earth Observation Programme before the lockdown. We then examined COVID-19 mortality rates in relation to the nitrogen dioxide levels at three 14-day lag points after the lockdown, namely March 8, 22 and April 5, 2020. Using a multivariable negative binomial regression model, we found an association between nitrogen dioxide and COVID-19 mortality. Although ecological data provide only weak evidence, these findings indicate an association between air pollution levels and COVID-19 severity.

© 2020 Elsevier B.V. All rights reserved.

* Corresponding author at: Dipartimento di Scienze Biomediche, Metaboliche e Neuroscienze – UNIMORE, Via Campi 287, 41125 Modena, Italy.
E-mail address: marco.vinceti@unimore.it (M. Vinceti).

1. Introduction

After the first indigenous Italian case of COVID-19 diagnosed on February 20, 2020 in a small municipality of Lombardy region, Codogno, the SARS-CoV-2 outbreak rapidly swept across all of Italy, resulting in more than 240,000 confirmed positive cases and almost 35,000 deaths by mid-July (CPD, 2020; Docea et al., 2020). Public health measures to stem the spread included tracking positive cases and related contacts and quarantining them, inviting the population to use face masks, and encouraging social distancing and hand washing. In addition, Italian authorities swiftly adopted two major interventions, the two national lockdowns to limit the spread of the virus, following two local lockdown adopted in the Codogno area and in the small municipality of Vò, Veneto region (Lavezzo et al., 2020). The first national lockdown was instituted on February 23, 2020, with a set of decrees in the most affected regions (i.e. Lombardy, Veneto and Emilia-Romagna). The second was on March 8, 2020 in the most affected areas and immediately extended the following day to all of Italy (Gabutti et al., 2020; Ministry of Health, 2020b). Only the second lockdown was seen to be effective in slowing the infection, with a reversal of the epidemic curve beginning 9 days after its implementation (Vinceti et al., 2020).

The quick spread of SARS-CoV-2 infection first in China, then Italy and the remainder of the world, prompted researchers to study factors associated with higher susceptibility to the infection or to disease severity. These factors included medical history, life-style factors, and environmental risk factors such as climate indicators and air pollutants. Some studies suggested a possible relation of virus outbreak with high levels of environmental air pollution (Barcelo, 2020; Bashir et al., 2020; Copat et al., 2020; Filippini et al., 2020; Li et al., 2020; Ogen, 2020; Pequeno et al., 2020; Setti et al., 2020b; Tsatsakis et al., 2020; Wu et al., 2020; Zhu et al., 2020). Putative mechanisms included enhanced susceptibility to respiratory viral infections through damage and inflammation of lung cells, immune dysregulation, and hyper-activation of inflammatory cytokines and chemokines (Chen et al., 2010; Comunian et al., 2020; Conticini et al., 2020; Peng et al., 2020), up-regulation of viral receptor ACE-2 (Borro et al., 2020; Tung et al., 2020), or air pathogen transportation by particulate matter suitable as virus carrier (Comunian et al., 2020; Manoj et al., 2020).

A link between air pollution and COVID-19 was also suggested by the observation that the SARS-CoV-2 outbreak first spread in Northern Italy in the Po basin area ('Po Valley' or 'Padan Plain'), a flat area that is the most heavily industrialized and polluted area of the country, and one of the most polluted of Europe (EEA, 2019). Tropospheric levels of nitrogen dioxide (NO₂) in early April of 2020 were positively associated with prevalence rates of COVID-19 within the three most affected Italian regions (Lombardy, Veneto and Emilia-Romagna) (Filippini et al., 2020), with the association becoming stronger at levels above 130 µmol/m².

In addition, high nitrogen dioxide levels have been associated with increased mortality for all causes, cardiovascular and respiratory mortality (Brunekreef et al., 2009; Eum et al., 2019; Hoek et al., 2013), and also pneumonia in older adults (Eum et al., 2019). A recent spatial analysis reported that up to 78% of COVID-19 deaths occurred in the five European areas located in Italy and Spain that also had the highest nitrogen dioxide levels, thus indicating a possible contribution to fatality by long-term exposure to nitrogen dioxide (Ogen, 2020). In addition, a recent study from China showed a positive association between short-term exposure to air pollutants (i.e. particulate matter, sulfur dioxide, carbon monoxide, ozone and nitrogen dioxide) and newly diagnosed COVID-19 confirmed cases (Zhu et al., 2020). Higher COVID-19 mortality was also correlated with relatively poor air quality in several other countries (Pansini and Fornacca, 2020). However, results from some recent studies have been inconsistent, with some reporting a null (Bontempi, 2020b; Briz-Redon et al., 2020) or inverse association (Zoran et al., 2020a) between air pollution and SARS-CoV-2 infection incidence. Therefore, this putative correlation appears to be worthy of further investigation. In addition, infection rates may not entirely correlate with the most severe clinical outcomes

of COVID-19, such as hospital intensive-care admission or mortality. Furthermore, infection rates also depend on the period of infection, the individual viral load, age and sex of the infected person, presence of comorbidities, blood type, and the efficiency of the health care system (Bontempi et al., 2020; Latz et al., 2020; Liu et al., 2020).

In this study, we aimed to investigate if the relation previously reported between air pollution assessed through satellite-detected tropospheric nitrogen dioxide and the early phases of the SARS-CoV-2 outbreak was associated with a higher COVID-19 severity, using available mortality data from two of the previously investigated three regions in Northern Italy.

2. Methods

2.1. Study area

We studied the population of two Northern Italy regions, Veneto and Emilia-Romagna, for which we were able to obtain the daily number of COVID-19 deaths by province of diagnosis. Such data were available in the period February 23–April 15, 2020 for Veneto region from the database of Veneto Region 'COVID-19 working group', and February 23–April 22, 2020 for Emilia-Romagna region from the database of the 'Public Health Authority' of Emilia-Romagna Region. Mortality data from the third region, Lombardy, were not available at a provincial level. For Veneto and Emilia-Romagna, we computed daily province-specific mortality rates, based on provincial population data retrieved from the Italian National Institute of Statistics website (ISTAT, 2020a).

2.2. Environmental air pollution assessment

We based our air pollution assessment in the study population on daily information tropospheric nitrogen dioxide levels, as available from the Copernicus Earth Observation Programme Open Access Hub-Sentinel-5P mission (Filippini et al., 2020). In particular, we focused on the geolocated tropospheric column of nitrogen dioxide with a spatial resolution of around 7 × 7 km² reported by ESA Sentinel-5P made publicly available for all of Europe from the European Space Agency (ESA, 2020). Using both Near Real Time (NRT) processing (typically 3 h from sensing) and Offline (OFFL) processing (5 days after sensing time) (Eskes et al., 2019), we collected all satellite images with quality index greater than 0.5 related to the study area of interest for each day of the analysis period. Finally, we computed population-weighted spatial average of nitrogen dioxide values (in µmol/m²) for each province. We also validated satellite data through measured ground-level nitrogen dioxide concentration by ARPA (Environmental Protection Regional Agency) monitoring stations (Filippini et al., 2020). When nitrogen dioxide values could not be considered reliable for one or more days, i.e. when satellite coverage in those days decreased below 30% of the spatial units (provinces) because of cloud-cover or surface snow or ice that affected satellite image reliability (Eskes et al., 2019), we imputed the missing data with a regression measured ground level concentrations from values averaged from the two-three days before and after. Imputation was needed for an average of two days over the entire study period.

To control for potential environmental confounders, we retrieved temperature data publicly available from the ERAS model reanalysis of the European Centre for Medium-Range Weather Forecasts website (ECMWF, 2020), and relative humidity that has been calculated using environmental temperature and dewpoint temperature according to a method described by Lawrence (Lawrence, 2005). As additional potential confounders, we considered the presence of international airports for which traffic flow exceeded 100,000 passengers in January 2020 (ASSAEROPORTI, 2020). Based on this criterion, we included in our model the airports of Venice, Verona and Treviso (Veneto region) and Bologna (Emilia-Romagna region). Finally, we accounted for population mobility using information on the mobile telephone daily movements available through the Call Detail Records (CDR) of anonymous data of

the SIM cards processed and made available by Teralytics for the period February 1–March 27, 2020 (Polzer, 2020), as explained in detail elsewhere (Vinceti et al., 2020).

2.3. Data analysis

To assess the average amount of air pollution preceding the onset of the SARS-CoV-2 outbreak, we modeled the time-series of daily nitrogen dioxide tropospheric levels from February 1 to 24 using a linear regression taking into account heteroskedasticity and autocorrelation up to 7 days using a Newey–West estimator (Newey and West, 1987). We also modeled these values after the lockdowns. We then examined the relation between nitrogen dioxide levels before February 24, using the estimated nitrogen dioxide levels on February 12, which is the midpoint of the period before the lockdown, and COVID-19 mortality rate at several time points: 14 days (corresponding to the period between outbreak onset and the establishment of the tight lockdown on March 8), 28 and 42 days, corresponding to March 8, March 22, and April 5. We modeled COVID-19 mortality rates using multivariable negative binomial regression analysis. Nitrogen dioxide was modeled with restricted cubic splines with three knots at fixed percentile of its distribution (10th, 50th and 90th percentiles) (Harrell, 2001) to assess the shape of the relation, adjusting for possible confounders. We considered province-specific

information on other possible confounding factors abstracted from the National Institute of Statistic website (ISTAT, 2020a), including population density and aging index in 2019, and percentage of population commuting daily for work or school, percentage of single-member families, and percentage of dwellings with one resident at 2011 census. In the multivariable analysis, we adjusted for population density, aging index, people mobility measured from telephone movements before the lockdown, temperature ($^{\circ}\text{C}$) and relative humidity in the three subsequent periods, and airport presence. In subsequent models, we alternatively added population percentage commuting outside the municipality of residence on a daily basis, percentage of families including only one member, or percentage of dwellings occupied by only one resident.

3. Results

Fig. 1 shows tropospheric nitrogen dioxide levels that we estimated for the study area before the onset of the outbreak and the establishment of the lockdowns. Provinces characterized by the highest levels above $130\ \mu\text{mol}/\text{m}^2$ were Verona ($181\ \mu\text{mol}/\text{m}^2$), Padua ($147\ \mu\text{mol}/\text{m}^2$) and Vicenza ($141\ \mu\text{mol}/\text{m}^2$) in Veneto, and Piacenza ($140\ \mu\text{mol}/\text{m}^2$) in Emilia-Romagna. Table 1 reports number of COVID-19 deaths, mortality rate (per 100,000 people), pollutant levels on the three selected time points (March 8, March 22, and April 5) in Veneto




Fig. 1. Average NO₂ tropospheric levels in the study area before the lockdown of February 23, 2020, based on data from the Sentinel-5P mission from Copernicus Earth Observation Programme of the European Space Agency and the European Union.

Table 1
Cumulative number of total COVID-19 deaths, mortality rate (per 100,000 people) and nitrogen dioxide (NO₂) levels (μmol/m²), before the lockdown, on March 8, March 22, and April 5 in Veneto and Emilia-Romagna regions.

	Population at Jan 1 2019	Death cases			Mortality rate (per 100,000)			NO ₂ levels (μmol/m ²)			
		Mar 8	Mar 22	Apr 5	Mar 8	Mar 22	Apr 5	Before Feb 24		After Feb 24	
										Feb 24–Mar 8	Mar 8–Mar 22
Veneto	4,905,854	97	546	908	1.98	11.13	18.51	136	94	75	50
Belluno (BL)	202,950	2	19	37	0.99	9.36	18.23	50	56	31	29
Padua (PD)	937,908	16	93	174	1.71	9.92	18.55	147	88	84	55
Rovigo (RO)	234,937	0	9	16	0.00	3.83	6.81	118	68	56	42
Treviso (TV)	887,806	37	125	173	4.17	14.08	19.49	108	96	59	42
Venice (VE)	853,338	30	84	130	3.52	9.84	15.23	126	94	66	46
Verona (VR)	926,497	8	142	257	0.86	15.33	27.74	181	114	97	59
Vicenza (VI)	862,418	4	74	121	0.46	8.58	14.03	141	97	84	54
Emilia-Romagna	4,459,477	99	1060	2287	2.22	23.77	51.28	109	81	66	38
Bologna (BO)	1,014,619	2	78	243	0.20	7.69	23.95	109	86	71	37
Ferrara (FE)	345,691	1	11	45	0.29	3.18	13.02	104	71	55	39
Forlì-Cesena (FC)	394,627	0	12	46	0.00	3.04	11.66	80	61	51	29
Modena (MO)	705,393	2	89	264	0.28	12.62	37.43	117	95	70	42
Parma (PR)	451,631	41	301	529	9.08	66.65	117.13	121	87	72	42
Piacenza (PC)	287,152	42	388	673	14.62	135.12	234.37	140	113	90	58
Ravenna (RA)	389,456	0	9	38	0.00	2.31	9.76	95	67	60	31
Reggio Emilia (RE)	531,891	10	103	289	1.88	19.36	54.33	118	83	68	43
Rimini (RN)	339,017	1	69	160	0.29	20.35	47.120	90	53	40	23

and Emilia-Romagna regions. Fourteen days after the tight lockdown (March 8), mortality rates in Veneto and Emilia-Romagna were almost the same (1.98 and 2.22 per 100,000, respectively). In the subsequent periods, provinces reporting higher mortality rates were Verona (15.33 and 27.74 per 100,000 on March 22 and April 5, respectively) in Veneto, and Piacenza (135.12 and 234.37 per 100,000 on March 22 and April 5, respectively) and Parma (66.65 and 117.13 per 100,000 on March 22 and April 5, respectively) in Emilia-Romagna (Table 1).

Using three 14-day lag points after the lockdown (on March 8, March 22, and April 5), we fit a negative binomial regression between nitrogen dioxide levels before the lockdown and the cumulative mortality rates at these time points. Although the precision of the model was low, our data were compatible with an increase in COVID-19 related mortality rates at the three time points for levels of nitrogen dioxide above 100 μmol/m², particularly 28 and 42 days after the lockdown (Fig. 2). The association between mortality and air pollution exhibited a non-linear shape particularly in the last period of analysis, with a rapidly increasing rate at the highest levels of nitrogen dioxide. Further adjustment for population percentage commuting outside the municipality of residence, percentage of families including only one member, or percentage of dwellings occupied by only one resident did not change the results meaningfully (Supplemental Figs. S1–S3).

4. Discussion

Our data support a positive, non-linear association between high nitrogen dioxide tropospheric levels and subsequent COVID-19 mortality rates at different time periods in 16 Northern Italy provinces severely hit by the epidemic, which provide evidence supporting an effect of health pollution in increasing the case-fatality ratio of the disease. The association was stronger when we used cumulative mortality in the longest period as an outcome. These results are similar to those yielded by a previous study of ours carried out in Emilia-Romagna, Veneto and Lombardy and using daily SARS-CoV-2 new infections (Filippini et al., 2020). In that study, the association between nitrogen dioxide levels and SARS-CoV-2 infection incidence was not linear, with risk starting to increase at around 130 μmol/m²; at values above that, the steepness of the two slopes were comparable. The evidence for an association between air pollution and mortality starting at slightly lower nitrogen dioxide levels in the present study could stem from an association between air pollution and severe COVID-19, mediated by a high chronic disease

prevalence existing already at relatively low levels of nitrogen dioxide exposure. Evidence is clear that a higher prevalence of comorbidities may lead to a more clinically severe form of COVID-19 and to a higher mortality from the disease (Onder et al., 2020; Reilev et al., 2020; Ssentongo et al., 2020).

Some studies indicate a gradual association between average air pollution levels and mortality from COVID-19, rather than a threshold effect. A nation-wide study carried out using data on over 3000 US counties reported that an increase of 1 μg/m³ in particular matter (PM_{2.5}) was associated with a 15% increase in mortality from COVID-19 (Wu et al., 2020). Similarly, an Italian study assessing air pollution exposure using PM_{2.5} concentrations in Northern Italy municipalities found a positive association with mortality related to COVID-19, a 1 μg/m³ increase being associated with a 9% increase in COVID-19 mortality (Coker et al., 2020). In addition, two Spanish studies found a higher incidence and mortality of COVID-19 in urban and industrial areas compared with rural areas in Tarragona (Marques et al., 2020), and an association between COVID-19 spread (including mortality) and nitrogen dioxide exposure in Catalonia (Saez et al., 2020). On the other hand, inconsistent results were reported from a study assessing the correlation between urban air pollution in the Milan metropolitan area and number of COVID-19 deaths, which found a negative correlation with air pollution assessed through surface levels of both PM_{2.5} and PM₁₀ (Zoran et al., 2020b), and ground levels of nitrogen dioxide (Zoran et al., 2020a), but a positive one using ground levels of ozone, possibly due to the enhanced formation of outdoor airborne secondary aerosols (Zoran et al., 2020a). Similarly, a study of seven Spain cities found little association between outdoor air pollutant levels as assessed through traffic monitoring stations and COVID-19 transmission (Briz-Redon et al., 2020).

An association between outdoor air pollution and incidence, mortality and severe clinical course of COVID-19 is not unexpected and may be mediated by different factors and mechanisms. Long-term exposure to air pollutant levels has been linked to chronic lung inflammation, thus possibly promoting an increased severity of COVID-19 syndrome (Shi et al., 2020). Air pollutants have also been shown to alter the immunological status and thus increasing individual susceptibility to infectious diseases (Ibironke et al., 2019; Marchini et al., 2020; Rivas-Santiago et al., 2015; Williams et al., 2011). As specifically regards traffic-generated nitrogen dioxide, it has been shown to be associated with increased risk of rhinitis in adult population (Cesaroni et al., 2008; de




Fig. 2. Negative binomial regression analysis between nitrogen dioxide (NO_2) tropospheric levels ($\mu\text{mol}/\text{m}^3$) before the spread of the outbreak and COVID-19 mortality rate (deaths per 100,000) in the three periods after the lockdown dates (A: February 24–March 8; B: February 24–March 22; C: February 24–April 5). Adjusted COVID-19 mortality rate (solid line) with 95% confidence interval (dash lines) was estimated with a multivariable model, taking into account population density, aging index, people mobility measured from telephone movements before the lockdown, temperature ($^{\circ}\text{C}$) and relative humidity (%) in the three subsequent periods, and airport presence. Shaded circles are weighted on number of deaths corresponding to the adjusted mortality rates at each time point.

Marco et al., 2002) and other upper respiratory illnesses (Chen et al., 2007), as well as to a higher risk of asthma, weakened lung function (Bowatte et al., 2017; Cai et al., 2017; de Marco et al., 2002; McCreanor et al., 2007), and exacerbation of chronic obstructive pulmonary disease (Ghanbari Ghazikali et al., 2016; Pfeiffer et al., 2019). The geographic area investigated in the present study is also known to be one of the most polluted in Europe (EEA, 2019) and its high levels of

air pollutants have been previously associated with higher mortality and hospitalization for both cardiovascular and respiratory diseases (Carugno et al., 2016; Fattore et al., 2011). A study carried out in the Bergamo area, which had one of the highest excess of mortality rates (+571% in March 2020) due to COVID-19 in Northern Italy (ISTAT, 2020b), found SARS-CoV-2 RNA in particulate matter samples, supporting the hypothesis that air pollutants may enhance virus spread (Setti et al., 2020a). Similarly, another study carried out using data of 55 Northern Italy provinces found some evidence for an accelerated diffusion of SARS-CoV-2 in cities with higher number of days exceeding the limit set for PM_{10} of $50 \mu\text{g}/\text{m}^3$ (Coccia, 2020).

Our study has important limitations. Because this study is based upon aggregated rather than individual-level data, the study could be affected by ecologic bias. We could not take into account individual risk factors that may be associated with higher susceptibility to virus infection, including factors such as working in health care delivery, residence in nursing homes, as well as individual contact with infected people or travel to China or other infected countries (Bontempi, 2020a; Bontempi et al., 2020; Domingo et al., 2020). In addition, the two investigated regions may slightly differ in term of health care and public health measures provided by the Local Health Authorities during the pandemic, though the presence of a National Health System and guidelines for health care workers should have attenuated this issue (Ministry of Health, 2020a). Also, the study size provided only limited precision of the effect estimates, as reflected in the broad confidence intervals for our dose-response curve. Nonetheless, study findings still appear to support a positive, non-linear association between air pollution measures and increased COVID-19 mortality.

Funding

This study was supported by grant 'UNIMORE FAR 2019 Interdisciplinare Linea FCRMO - Fondazione Cassa di Risparmio di Modena' and grant 'UNIMORE FAR IMPULSO 2020' (No. 494/2020).

CRedit authorship contribution statement

MV and TF designed the original study, and with KJR analyzed and interpreted the data, and wrote the manuscript. NO designed and carried out data analysis with TF. SC, EN, DM, MS and VB collected and validated population data. AG, FF and GM downloaded and processed environmental data. All authors read and approved the final manuscript.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Dr. Paola Angelini from the Public Health Authority of the Emilia-Romagna Region and the COVID-19 working group of Veneto Region for providing data necessary for this study.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2020.143355>.

References

- ASSAEROPORTI, 2020. January 2020 Statistics. Associazione Italiana Gestori di Aeroporti http://assaerporti.com/statistiche_202001/ (Accessed: June 12, 2020).
- Barcelo, D., 2020. An environmental and health perspective for COVID-19 outbreak: meteorology and air quality influence, sewage epidemiology indicator, hospitals

- disinfection, drug therapies and recommendations. *J. Environ. Chem. Eng.* 8, 104006. <https://doi.org/10.1016/j.jece.2020.104006>.
- Bashir, M.F., Ma, B., Bilal, Komal, B., Bashir, M.A., Tan, D., et al., 2020. Correlation between climate indicators and COVID-19 pandemic in New York, USA. *Sci. Total Environ.* 728, 138835. <https://doi.org/10.1016/j.scitotenv.2020.138835>.
- Bontempi, E., 2020a. Commercial exchanges instead of air pollution as possible origin of COVID-19 initial diffusion phase in Italy: more efforts are necessary to address interdisciplinary research. *Environ. Res.* 188, 109775. <https://doi.org/10.1016/j.envres.2020.109775>.
- Bontempi, E., 2020b. First data analysis about possible COVID-19 virus airborne diffusion due to air particulate matter (PM): the case of Lombardy (Italy). *Environ. Res.* 186, 109639. <https://doi.org/10.1016/j.envres.2020.109639>.
- Bontempi, E., Vergalli, S., Squazzoni, F., 2020. Understanding COVID-19 diffusion requires an interdisciplinary, multi-dimensional approach. *Environ. Res.* 188, 109814. <https://doi.org/10.1016/j.envres.2020.109814>.
- Borro M, Di Girolamo P, Gentile G, De Luca O, Preissner R, Marcolongo A, et al. Evidence-based considerations exploring relations between SARS-CoV-2 pandemic and air pollution: involvement of PM2.5-mediated up-regulation of the viral receptor ACE-2. *Int. J. Environ. Res. Public Health* 2020; 17: 5573. doi: <https://doi.org/10.3390/ijerph17155573>.
- Bowatte, G., Erbas, B., Lodge, C.J., Knibbs, L.D., Gurrin, L.C., Marks, G.B., et al., 2017. Traffic-related air pollution exposure over a 5-year period is associated with increased risk of asthma and poor lung function in middle age. *Eur. Respir. J.* 50, 1602357. <https://doi.org/10.1183/13993003.02357-2016>.
- Briz-Redon, A., Belenguier-Sapina, C., Serrano-Aroca, A., 2020. A city-level analysis of air pollution, climate and COVID-19 early spread during the Spanish lockdown. *medRxiv* <https://doi.org/10.1101/2020.08.09.20171041>.
- Brunekreef, B., Beelen, R., Hoek, G., Schouten, L., Bausch-Goldbohm, S., Fischer, P., et al., 2009. Effects of long-term exposure to traffic-related air pollution on respiratory and cardiovascular mortality in the Netherlands: the NLCS-AIR study. *Res. Rep. Health Eff. Inst.* 5–71 (discussion 73–89).
- Cai, Y., Zijlema, W.L., Doiron, D., Blangiardo, M., Burton, P.R., Fortier, I., et al., 2017. Ambient air pollution, traffic noise and adult asthma prevalence: a BioSHare approach. *Eur. Respir. J.* 49, 1502127. <https://doi.org/10.1183/13993003.02127-2015>.
- Carugno, M., Consonni, D., Randi, G., Catelan, D., Grisotto, L., Bertazzi, P.A., et al., 2016. Air pollution exposure, cause-specific deaths and hospitalizations in a highly polluted Italian region. *Environ. Res.* 147, 415–424. <https://doi.org/10.1016/j.envres.2016.03.003>.
- Cesaroni, G., Badaloni, C., Porta, D., Forastiere, F., Perucci, C.A., 2008. Comparison between various indices of exposure to traffic-related air pollution and their impact on respiratory health in adults. *Occup. Environ. Med.* 65, 683–690. <https://doi.org/10.1136/oem.2007.037846>.
- Chen, T.-M., Kuschner, W.G., Gokhale, J., Shofer, S., 2007. Outdoor air pollution: nitrogen dioxide, sulfur dioxide, and carbon monoxide health effects. *Am J Med Sci* 333, 249–256. <https://doi.org/10.1097/MAJ.0b013e31803b900f>.
- Chen, P.S., Tsai, F.T., Lin, C.K., Yang, C.Y., Chan, C.C., Young, C.Y., et al., 2010. Ambient influenza and avian influenza virus during dust storm days and background days. *Environ. Health Perspect.* 118, 1211–1216. <https://doi.org/10.1289/ehp.0901782>.
- Coccia, M., 2020. Factors determining the diffusion of COVID-19 and suggested strategy to prevent future accelerated viral infectivity similar to COVID. *Sci. Total Environ.* 729, 138474. <https://doi.org/10.1016/j.scitotenv.2020.138474>.
- Coker, E., Cavalli, L., Fabrizi, E., Guastella, G., Lippo, E., Parisi, M.L., et al., 2020. *The Effects of Air Pollution on COVID-19 Related Mortality in Northern Italy. Nota di Lavoro. Fondazione Eni Enrico Mattei, Milan, Italy.*
- Comunian, S., Dongo, D., Milani, C., Palestini, P., 2020. Air pollution and COVID-19: the role of particulate matter in the spread and increase of COVID-19's morbidity and mortality. *Int. J. Environ. Res. Public Health* 17, 4487. <https://doi.org/10.3390/ijerph17124487>.
- Coticini, E., Frediani, B., Caro, D., 2020. Can atmospheric pollution be considered a cofactor in extremely high level of SARS-CoV-2 lethality in Northern Italy? *Environ. Pollut.* 261, 114465. <https://doi.org/10.1016/j.envpol.2020.114465>.
- Copat, C., Cristaldi, A., Fiore, M., Grasso, A., Zucarello, P., Signorelli, S.S., et al., 2020. The role of air pollution (PM and NO2) in COVID-19 spread and lethality: a systematic review. *Environ. Res.* 191, 110129. <https://doi.org/10.1016/j.envres.2020.110129>.
- CPD, 2020. Italian Civil Protection Department: COVID-19 data. <https://github.com/pcm-dpc/COVID-19>, (Accessed: June 12, 2020).
- de Marco, R., Poli, A., Ferrari, M., Accordini, S., Giammanco, G., Bugiani, M., et al., 2002. The impact of climate and traffic-related NO2 on the prevalence of asthma and allergic rhinitis in Italy. *Clin. Exp. Allergy* 32, 1405–1412. <https://doi.org/10.1046/j.1365-2745.2002.01466.x>.
- Docea, A.O., Tsatsakis, A., Albulescu, D., Cristea, O., Zlatian, O., Vinceti, M., et al., 2020. A new threat from an old enemy: reemergence of coronavirus (review). *Int. J. Mol. Med.* 45, 1631–1643. <https://doi.org/10.3892/ijmm.2020.4555>.
- Domingo, J.L., Marques, M., Rovira, J., 2020. Influence of airborne transmission of SARS-CoV-2 on COVID-19 pandemic. *A review. Environ. Res.* 188, 109861. <https://doi.org/10.1016/j.envres.2020.109861>.
- ECMWF, 2020. ERA5. <https://www.ecmwf.int/en/forecasts/datasets/reanalysis-datasets/era5>, (Accessed: June 12, 2020).
- EEA, 2019. In: European Environment Agency (Ed.), EEA Report No 10/2019 - Air Quality in Europe 2019. EEA, Copenhagen, Denmark <https://www.eea.europa.eu/publications/air-quality-in-europe-2019> (Accessed: June 12, 2020).
- ESA, 2020. Sentinel-5P Data. European Space Agency https://www.esa.int/Applications/Observing_the_Earth/Copernicus/Sentinel-5P, (Accessed: June 12, 2020).
- Eskes, H., van Geffen, J., Boersma, F., Eichmann, K.-U., Apituley, A., Pedergana, M., et al., 2019. Sentinel-5 Precursor/TROPOMI Level 2 Product User Manual Nitrogen dioxide. RNM Institute <https://sentinel.esa.int/documents/247904/2474726/Sentinel-5P-Level-2-Product-User-Manual-Nitrogen-Dioxide>, (Accessed: 12 June 2020).
- Eum, K.D., Kazemparkouhi, F., Wang, B., Manjourides, J., Pun, V., Pavlu, V., et al., 2019. Long-term NO2 exposures and cause-specific mortality in American older adults. *Environ. Int.* 124, 10–15. <https://doi.org/10.1016/j.envint.2018.12.060>.
- Fattore, E., Paiano, V., Borgini, A., Tittarelli, A., Bertoldi, M., Crosignani, P., et al., 2011. Human health risk in relation to air quality in two municipalities in an industrialized area of Northern Italy. *Environ. Res.* 111, 1321–1327. <https://doi.org/10.1016/j.envres.2011.06.012>.
- Filippini, T., Rothman, K.J., Goffi, A., Ferrari, F., Maffei, G., Orsini, N., et al., 2020. Satellite-detected tropospheric nitrogen dioxide and spread of SARS-CoV-2 infection in Northern Italy. *Sci. Total Environ.* 739, 140278. <https://doi.org/10.1016/j.scitotenv.2020.140278>.
- Gabutti, G., d'Anchera, E., Sandri, F., Savio, M., Stefanati, A., 2020. Coronavirus: update related to the current outbreak of COVID-19. *Infect. Dis. Ther.* 9, 1–13. <https://doi.org/10.1007/s40121-020-00295-5>.
- Ghanbari Ghazikali, M., Heibati, B., Naddafi, K., Kloog, I., Oliveri Conti, G., Polosa, R., et al., 2016. Evaluation of chronic obstructive pulmonary disease (COPD) attributed to atmospheric O3, NO2, and SO2 using Air Q Model (1911–2012 year). *Environ. Res.* 144, 99–105. <https://doi.org/10.1016/j.envres.2015.10.030>.
- Harrell, F.E., 2001. *Regression Modeling Strategies: With Applications to Linear Models, Logistic and Ordinal Regression, and Survival Analysis.* Springer-Verlag, New York.
- Hoek, G., Krishnan, R.M., Beelen, R., Peters, A., Ostro, B., Brunekreef, B., et al., 2013. Long-term air pollution exposure and cardio-respiratory mortality: a review. *Environ. Health* 12, 43. <https://doi.org/10.1186/1476-069X-12-43>.
- Ibironke, O., Carranza, C., Sarkar, S., Torres, M., Choi, H.T., Nwoko, J., et al., 2019. Urban air pollution particulates suppress human T-cell responses to *Mycobacterium tuberculosis*. *Int. J. Environ. Res. Public Health* 16, 4112. <https://doi.org/10.3390/ijerph16214112>.
- ISTAT, 2020a. Data from the Italian National Institute of Statistic. <https://www.istat.it/it/popolazione-e-famiglie> (Accessed: June 12, 2020).
- ISTAT, 2020b. Impatto dell'epidemia COVID-19 sulla mortalità totale della popolazione residente primo quadrimestre 2020. https://www.epicentro.iss.it/coronavirus/pdf/Rapp_ISTAT_ISS_3Giugno.pdf. (Accessed 4 June 2020).
- Latz, C.A., DeCarlo, C., Boitano, L., Png, C.Y.M., Patell, R., Conrad, M.F., et al., 2020. Blood type and outcomes in patients with COVID-19. *Ann. Hematol.* 99, 2113–2118. <https://doi.org/10.1007/s00277-020-04169-1>.
- Lavezzo, E., Franchin, E., Ciavarella, C., Cuomo-Dannenburg, G., Barzon, L., Del Vecchio, C., et al., 2020. Suppression of a SARS-CoV-2 outbreak in the Italian municipality of Vo'. *Nature* 584, 425–429. <https://doi.org/10.1038/s41586-020-2488-1>.
- Lawrence, M.G., 2005. The relationship between relative humidity and the dewpoint temperature in moist air: a simple conversion and applications. *Bull. Am. Meteorol. Soc.* 86, 225–234. <https://doi.org/10.1175/BAMS-86-2-225>.
- Li, H., Xu, X.L., Dai, D.W., Huang, Z.Y., Ma, Z., Guan, Y.J., 2020. Air pollution and temperature are associated with increased COVID-19 incidence: a time series study. *Int. J. Infect. Dis.* 97, 278–282. <https://doi.org/10.1016/j.ijid.2020.05.076>.
- Liu, T., Liang, W., Zhong, H., He, J., Chen, Z., He, G., et al., 2020. Risk factors associated with COVID-19 infection: a retrospective cohort study based on contacts tracing. *Emerg. Microbes Infect.* 9, 1546–1553. <https://doi.org/10.1080/22221751.2020.1787799>.
- Manoj, M.G., Satheesh Kumar, M.K., Valsaraj, K.T., Sivan, C., Vijayan, S.K., 2020. Potential link between compromised air quality and transmission of the novel corona virus (SARS-CoV-2) in affected areas. *Environ. Res.* 190, 110001. <https://doi.org/10.1016/j.envres.2020.110001>.
- Marchini, T., Zirikli, A., Wolf, D., 2020. Pathogenic role of air pollution particulate matter in cardiometabolic disease: evidence from mice and humans. *Antioxid. Redox Signal.* 33, 263–279. <https://doi.org/10.1089/ars.2020.8096>.
- Marques, M., Rovira, J., Nadal, M., Domingo, J.L., 2020. Effects of air pollution on the potential transmission and mortality of COVID-19: a preliminary case-study in Tarragona Province (Catalonia, Spain). *Environ. Res.* 192, 110315. <https://doi.org/10.1016/j.envres.2020.110315>.
- McCreanor, J., Cullinan, P., Nieuwenhuijsen, M.J., Stewart-Evans, J., Malliarou, E., Jarup, L., et al., 2007. Respiratory effects of exposure to diesel traffic in persons with asthma. *N. Engl. J. Med.* 357, 2348–2358. <https://doi.org/10.1056/NEJMoA071535>.
- Ministry of Health, 2020a. Novel coronavirus. <http://www.salute.gov.it/portale/nuovocoronavirus/dettaglioContenutiNuovoCoronavirus.jsp?lingua=italiano&id=5373&area=nuovoCoronavirus&menu=vuoto> (Accessed: September 10, 2020).
- Ministry of Health, 2020b. Official Italian statistics on SARS-CoV-2. March 8, 2020. http://www.salute.gov.it/imgs/C_17_pagineAree_5351_0_file.pdf (Accessed: June 12, 2020).
- Newey, W.K., West, K.D., 1987. A simple, positive semi-definite, heteroskedasticity and autocorrelation consistent covariance matrix. *Econometrica* 55, 703–708. <https://doi.org/10.2307/1913610>.
- Ogen, Y., 2020. Assessing nitrogen dioxide (NO2) levels as a contributing factor to coronavirus (COVID-19) fatality. *Sci. Total Environ.* 726, 138605. <https://doi.org/10.1016/j.scitotenv.2020.138605>.
- Onder, G., Rezza, G., Brusaferro, S., 2020. Case-fatality rate and characteristics of patients dying in relation to COVID-19 in Italy. *JAMA* 323, 1775–1776. <https://doi.org/10.1001/jama.2020.4683>.
- Pansini, R., Fornacca, D., 2020. Higher virulence of COVID-19 in the air-polluted regions of eight severely affected countries. *medRxiv* <https://doi.org/10.1101/2020.04.30.20086496>.
- Peng, L., Zhao, X., Tao, Y., Mi, S., Huang, J., Zhang, Q., 2020. The effects of air pollution and meteorological factors on measles cases in Lanzhou, China. *Environ. Sci. Pollut. Res.* 27, 13524–13533. <https://doi.org/10.1007/s11356-020-07903-4>.
- Pequeno, P., Mendel, B., Rosa, C., Bosholn, M., Souza, J.L., Baccaro, F., et al., 2020. Air transportation, population density and temperature predict the spread of COVID-19 in Brazil. *PeerJ* 8, e9322. <https://doi.org/10.7717/peerj.9322>.

- Pfeffer, P.E., Donaldson, G.C., Mackay, A.J., Wedzicha, J.A., 2019. Increased chronic obstructive pulmonary disease exacerbations of likely viral etiology follow elevated ambient nitrogen oxides. *Am. J. Respir. Crit. Care Med.* 199, 581–591. <https://doi.org/10.1164/rccm.201712-2506OC>.
- Polzer, G., 2020. La mappa della nostra era glaciale: così il coronavirus ha congelato l'Italia. *La Repubblica*, Italy. <https://lab.gedidigital.it/repubblica/2020/cronaca/coronavirus-mappa-italia-impatto-sulla-mobilita/?ref=RHPPLF-BH-I252606083-C8-P3-S1.8-T1>. (Accessed 12 June 2020).
- Reilev, M., Kristensen, K.B., Pottogard, A., Lund, L.C., Hallas, J., Ernst, M.T., et al., 2020. Characteristics and predictors of hospitalization and death in the first 11 122 cases with a positive RT-PCR test for SARS-CoV-2 in Denmark: a nationwide cohort. *Int J Epidemiol.* dyaa140 <https://doi.org/10.1093/ije/dyaa140>.
- Rivas-Santiago, C.E., Sarkar, S., Cantarella, P., Osornio-Vargas, A., Quintana-Belmares, R., Meng, Q., et al., 2015. Air pollution particulate matter alters antimicrobial respiratory epithelium innate immunity. *Infect. Immun.* 83, 2507–2517. <https://doi.org/10.1128/IAI.03018-14>.
- Saez, M., Tobias, A., Barcelo, M.A., 2020. Effects of long-term exposure to air pollutants on the spatial spread of COVID-19 in Catalonia, Spain. *Environ. Res.* 191, 110177. <https://doi.org/10.1016/j.envres.2020.110177>.
- Setti, L., Passarini, F., De Gennaro, G., Barbieri, P., Perrone, M.G., Borelli, M., et al., 2020a. SARS-Cov-2RNA found on particulate matter of Bergamo in Northern Italy: first evidence. *Environ. Res.* 188, 109754. <https://doi.org/10.1016/j.envres.2020.109754>.
- Setti, L., Passarini, F., De Gennaro, G., Barbieri, P., Licen, S., Perrone, M.G., et al., 2020b. Potential role of particulate matter in the spreading of COVID-19 in Northern Italy: first observational study based on initial epidemic diffusion. *BMJ Open* 10, e039338. <https://doi.org/10.1136/bmjopen-2020-039338>.
- Shi, Y., Wang, Y., Shao, C., Huang, J., Gan, J., Huang, X., et al., 2020. COVID-19 infection: the perspectives on immune responses. *Cell Death Differ.* 27, 1451–1454. <https://doi.org/10.1038/s41418-020-0530-3>.
- Ssentongo, P., Ssentongo, A.E., Heilbrunn, E.S., Ba, D.M., Chinchilli, V.M., 2020. Association of cardiovascular disease and 10 other pre-existing comorbidities with COVID-19 mortality: a systematic review and meta-analysis. *PLoS One* 15, e0238215. <https://doi.org/10.1371/journal.pone.0238215>.
- Tsatsakis, A., Petrakis, D., Nikolouzakis, T.K., Docea, A.O., Calina, D., Vinceti, M., et al., 2020. COVID-19, an opportunity to reevaluate the correlation between long-term effects of anthropogenic pollutants on viral epidemic/pandemic events and prevalence. *Food Chem. Toxicol.* 141, 111418. <https://doi.org/10.1016/j.fct.2020.111418>.
- Tung, N.T., Cheng, P.C., Chi, K.H., Hsiao, T.C., Jones, T., BeruBe, K., et al., 2020. Particulate matter and SARS-CoV-2: a possible model of COVID-19 transmission. *Sci. Total Environ.* 750, 141532. <https://doi.org/10.1016/j.scitotenv.2020.141532>.
- Vinceti, M., Filippini, T., Rothman, K.J., Ferrari, F., Goffi, A., Maffei, G., et al., 2020. Lock-down timing and efficacy in controlling COVID-19 using mobile phone tracking. *EClinicalMedicine* 25, 100457. <https://doi.org/10.1016/j.eclinm.2020.100457>.
- Williams, L., Ulrich, C.M., Larson, T., Wener, M.H., Wood, B., Chen-Levy, Z., et al., 2011. Fine particulate matter (PM_{2.5}) air pollution and immune status among women in the Seattle area. *Arch. Environ. Occup. Health* 66, 155–165. <https://doi.org/10.1080/19338244.2010.539636>.
- Wu, X., Nethery, R.C., Sabath, B.M., Braun, D., Dominici, F., 2020. Exposure to air pollution and COVID-19 mortality in the United States: a nationwide cross-sectional study. *medRxiv* <https://doi.org/10.1101/2020.04.05.20054502>.
- Zhu, Y., Xie, J., Huang, F., Cao, L., 2020. Association between short-term exposure to air pollution and COVID-19 infection: evidence from China. *Sci. Total Environ.* 727, 138704. <https://doi.org/10.1016/j.scitotenv.2020.138704>.
- Zoran, M.A., Savastru, R.S., Savastru, D.M., Tautan, M.N., 2020a. Assessing the relationship between ground levels of ozone (O₃) and nitrogen dioxide (NO₂) with coronavirus (COVID-19) in Milan, Italy. *Sci. Total Environ.* 740, 140005. <https://doi.org/10.1016/j.scitotenv.2020.140005>.
- Zoran, M.A., Savastru, R.S., Savastru, D.M., Tautan, M.N., 2020b. Assessing the relationship between surface levels of PM_{2.5} and PM₁₀ particulate matter impact on COVID-19 in Milan, Italy. *Sci. Total Environ.* 738, 139825. <https://doi.org/10.1016/j.scitotenv.2020.139825>.