

Published in final edited form as:

*Neuropharmacology*. 2018 April 01; 132: 58–70. doi:10.1016/j.neuropharm.2017.06.029.

## Relevance of tissue specific subunit expression in channelopathies

Hartwig Seitter, Alexandra Koschak\*

University of Innsbruck, Institute of Pharmacy, Pharmacology and Toxicology, Center for Chemistry and Biomedicine, Innrain 80-82/III, 6020 Innsbruck, Austria

### Abstract

Channelopathies are a diverse group of human disorders that are caused by mutations in genes coding for ion channels or channel-regulating proteins. Several dozen channelopathies have been identified that involve both non-excitabile cells as well as electrically active tissues like brain, skeletal and smooth muscle or the heart. In this review, we start out from the general question which ion channel genes are expressed tissue-selectively. We mined the human gene expression database Human Protein Atlas (HPA) for tissue-enriched ion channel genes and found 85 genes belonging to the ion channel families. Most of these genes were enriched in brain, testis and muscle and a complete list of the enriched ion channel genes is provided. We further focused on the tissue distribution of voltage-gated calcium channel (VGCC) genes including different brain areas and the retina based on the human gene expression from the FANTOM5 dataset. The expression data is complemented by an overview of the tissue-dependent aspects of L-type calcium channel (LTCC) function, dysfunction and pharmacology, as well as of their splice variants. Finally, we focus on the pathology of tissue-restricted LTCC channelopathies and their treatment options.

This article is part of the Special Issue entitled ‘Channelopathies.’

### Keywords

Tissue expression; Calcium channel; Channelopathy; Retina; Congenital stationary night blindness

## 1 Introduction

The physiology and functionality of cell types and tissues is based on the underlying gene expression diversity within the different body parts of a living being. The question arises which of these genes are ubiquitously expressed and which genes and combinations endow organs with their unique tissue identity. In channelopathies, i.e. human pathologies caused by dysfunction of ion channels, the tissue-dependence can profoundly affect disease manifestation and treatment options.

---

\*Corresponding author. alexandra.koschak@uibk.ac.at (A. Koschak).

Voltage-gated  $\text{Ca}^{2+}$  channels (VGCCs) are interesting case examples of ion channels that have very selective or very widespread tissue distribution. VGCCs generate  $\text{Ca}^{2+}$  signals for many important physiological functions such as neurotransmission, muscle contraction, hormone secretion as well as regulation of gene expression (Catterall, 2011). Like other ion channels they form part of signaling complexes together with molecules, such as receptors, kinases, phosphatases and calmodulin (Catterall, 2011). Most VGCCs exist in a hetero-oligomeric complex of several subunits with the  $\alpha 1$ -subunit forming the  $\text{Ca}^{2+}$ -selective channel pore. Their biophysical and pharmacological properties are determined by their  $\alpha 1$ -subunit. Based on structural and pharmacological similarities of their  $\alpha 1$ -subunits, VGCCs can be grouped into two major families: LTCCs ( $\text{Ca}_v1$  family; Cav1.1-Cav1.4) and non-LTCCs ( $\text{Ca}_v2$ , Cav2.1-Cav2.3;  $\text{Ca}_v3$  families, Cav3.1-Cav3.3). The ten different  $\alpha 1$ -subunit isoforms associate with different  $\beta$ -subunits ( $\beta 1-4$ ) and  $\alpha 2\delta$  subunits ( $\alpha 2\delta 1-4$ ) and in some cases also  $\gamma$ -subunits. For a detailed overview we refer to a recent review by Zamponi and colleagues (Zamponi et al., 2015).

LTCC channelopathies are particularly interesting because some show obvious signs of tissue-selective phenotypes, whereas others are caused by less focally expressed channel genes that nevertheless primarily affect certain tissues. As for example skeletal muscle diseases like malignant hyperthermia or hypokalemic periodic paralysis (Jurkat-Rott et al., 2002) are caused by mutations in the very tissue-selectively expressed *CACNA1S* gene that encodes for Cav1.1 LTCCs. Also the *CACNA1F* gene that translates into Cav1.4 LTCCs exhibits very selective expression in the retina and therefore shows a tissue-specific disease manifestation (Zeitze et al., 2015). In contrast the two other LTCC members, Cav1.2 and Cav1.3 do not show a similar restriction of expression localization. Of note, some of their associated channelopathies (e.g. Timothy syndrome in Cav1.2) also highlight aspects of splice variant expression differences and the importance of the cellular environment for a phenotypic exhibition.

As ‘channel diggers’ we would highly benefit from a better overview of the tissue-selectivity of gene expression and/or even cell specific expression to unearth their functional mysteries.

## 2 Tissue specificity in voltage-gated ion channel genes according to large-scale screenings

Details on ion channel expression in single cell types are slowly accumulating. However it is also worthwhile to look at the overall ion channel repertoires on the tissue level from human transcriptome and proteome screenings. Recently large-scale screenings of human gene expression in tissues have been undertaken by several consortiums that allow for a broader view on gene expression localization in the human body (for an overview (Uhlen et al., 2016)). Here we made use of two datasets, the Human Protein Atlas ([www.proteinatlas.org](http://www.proteinatlas.org); (Uhlen et al., 2015)) and FANTOM5 ([fantom.gsc.riken.jp/5/](http://fantom.gsc.riken.jp/5/); (Lizio et al., 2015)). Within the framework of the Human Protein Atlas (HPA), Illumina-sequencing of RNA (RNA-seq) was performed on 32 human tissues, capturing the mRNA levels of transcripts within each tissue. Although mRNA level to protein level ratios can vary between different gene products depending on e.g. translation rates and protein half-life (Eden et al., 2011), overall the data

from RNA-seq is in good agreement with corresponding protein amounts (Wilhelm et al., 2014). The genes contained in the HPA are categorized according to their tissue-selective expression. Of the putative 19628 protein coding genes, 7367 genes were classified as expressed in all tissues, while 7835 genes showed elevated expression in subsets of tissues (3328 were mixed, 1098 not detected). The most selectively expressed subclass among the elevated genes are so-called tissue-enriched genes (defined by an expression value in the highest-expressing tissue that is at least five-fold of the second highest expression value; tissue specificity score  $TS \geq 5$ ).

From the HPA list of 2547 tissue-enriched genes we pulled out the 85 genes that belong to the ion channel families (according to the IUPHAR ion channel list, (Southan et al., 2016)) to obtain an overview of which ion channels are expressed most selectively (Table 1) and could thus give rise to a tissue-specific ion channel function in health and disease. The majority of tissue-enriched ion channel genes were found in a low number of tissues. As expected, the four tissues with the most ion channel genes comprised electrically active tissues like cerebral cortex, skeletal muscle and heart, but also testis contained a number of selectively expressed ion channel genes (Fig. 1). Yet, the fraction of channel genes was different for these tissues. While testis contained the largest number of tissue-enriched genes (Fig. 1A), only few of those genes code for ion channels (Fig. 1B). Cerebral cortex, skeletal muscle and heart included a much higher fraction of ion channels that was comparable within these three tissues. Cerebral cortex displayed the second highest number of tissue-enriched genes, therefore most of the ion channels were indeed found there. In Fig. 1C the ion channel genes selectively expressed in cerebral cortex, testis, skeletal muscle and heart are summarized and sorted according to their expression strengths within the given tissue. The numbers refer to Table 1 to facilitate retrieval of additional information.

The dataset can be looked at in detail from different angles: which genes are found in a given tissue or which tissues express a certain gene? For the former we concentrated on VGCC expression in three different tissue subsets: brain regions, muscle tissues and lymphoid tissues (Fig. 2). It is evident that there is only partial selectivity of expression of individual VGCC subunits, e.g.  $Ca_v1.1$  and  $Ca_v1.4$  LTCCs which show a very restricted expression in skeletal muscle and retinal tissue, respectively (Fig. 2A).  $Ca_v2$  channels and  $\gamma$  subunits showed an overall CNS-focused expression pattern. The most striking differences in the FANTOM5 brain samples were seen with respect to  $Ca_v2.1$  which was highly expressed in the cerebellum and  $Ca_v2.3$  that was most prominent in caudate (Fig. 2B). In general, the human data compared well with the data from mouse brain samples; e. g. evidence from quantitative RT-PCR analyses in hippocampal and cerebellar tissue found similar relationships in  $Ca_v2.1$ ,  $Ca_v2.3$ ,  $\beta_4$ ,  $\alpha_2\delta-1$  and  $\alpha_2\delta-2$  as in the FANTOM5 data, albeit e.g.  $Ca_v1.3$  and  $\beta_2$  showed a different ratio (Schlick et al., 2010).

In the muscle tissues a clear-cut segregation in the use of LTCCs and  $\beta$ -subunits was apparent. In skeletal muscle:  $Ca_v1.1$  and  $\beta_1$ ; in heart and smooth muscle:  $Ca_v1.2$  and  $\beta_2$  while all muscle tissues are alike in their expression of  $\alpha_2\delta-1$  as the main  $\alpha_2\delta$  subunit (Fig. 2C). Smooth muscle showed additionally high expression levels of  $Ca_v3.2$  as has previously been demonstrated ((Harraz et al., 2015a; Harraz et al., 2015b), for review (Kuo et al., 2014). Other findings suggest that smooth muscle cells of the CNS vasculature also express

Cav3.1 (Fernandez et al., 2015), rat retina arterioles) and Cav3.3 (Harraz et al., 2015b), human cerebral arteries). The  $\gamma_1$  subunit was only found in skeletal muscle but with a striking abundance (see inset in Fig. 2C for comparison of skeletal muscle VGCC subunit expression strengths). Overall this expression map recapitulated well what is known about muscle tissue  $\text{Ca}^{2+}$  channel repertoires (Catterall et al., 2005).

LTCCs have been reported repeatedly also in cells of the lymphoid lineage, mainly in different types of T-cells (for an overview, see (Davenport et al., 2015; Badou et al., 2013)). In the FANTOM5 lymphoid tissues the only LTCCs with prominent expression were Cav1.2 and Cav1.3 (Fig. 2D). Both have been described previously (Cabral et al., 2010; Robert et al., 2014). Cav1.1 (Matza et al., 2016) and Cav1.4 (Robert et al., 2014) (Omilusik et al., 2011; McRory et al., 2004) have also been reported in immune cells/ tissues but were not detected in the FANTOM5 lymphoid tissues (with exception of a low amount of Cav1.1 in the tonsil tissue). It is unclear if this could be attributed to the activation state of the immune cells because Cav1.4 expression was reported in naive T-cells as well as upon activation (Omilusik et al., 2011), while a novel variant of Cav1.1 (lacking exon 29 like in embryonic skeletal muscle but with alternative N-terminal exons; Table 2 and see also below) was found in T-cells after T-cell receptor stimulation (Matza et al., 2016). Among auxiliary subunits, all  $\beta$  subunits as well as  $\alpha_2\delta$ -2 have been described in immune cells (Robert et al., 2014; Jha et al., 2015; Cabral et al., 2010), which conforms to FANTOM5 lymphoid tissue data.

Moreover we compared expression data of the human retina with mouse retina quantitative RT-PCR data from Knoflach et al. (2013) (Fig. 2E). To this end we normalized all expression values to the highest expressed gene,  $\alpha_2\delta$ -4. While the quantitative expression values differed, the qualitative relationships between different VGCC subunits within each species were conserved. In both mouse and human retinas Cav1.4 was the most abundantly expressed pore-forming  $\alpha$ 1 subunit, CaV2.1 the most abundant CaV2 member and  $\beta_2$  was the main  $\beta$  subunit. The outcome for Cav1.4 and  $\beta_2$  is compelling confirming the suitability of mouse models for investigations in human Cav1.4 related channelopathies (see also below; (Zeitz et al., 2015)). The major difference between the two datasets lies in the high expression values for Cav1.2 and Cav1.3 in the human retina data. Most likely this can be explained by abundant RNA from retinal pigment epithelium (RPE) in the samples used for FANTOM5, which is evident in the high expression values of RPE-specific genes (e.g. PMEL, RGR, RPE65) in the retina data set (not shown). In fact, both Cav1.2 and Cav1.3 have been found prominently in RPE (Wimmers et al., 2008; Muller et al., 2014).

Fig. 3 takes a look at the FANTOM5 expression data from the other angle: Which tissues express a given VGCC gene. It is immediately obvious that Cav1.1 and Cav1.4 have very restricted expression mostly in a single tissue, while Cav1.2 and Cav1.3 were found in a large number of different tissues. All three CaV2 channel isoforms were detected almost exclusively in different parts of the brain, where also  $\beta_4$ ,  $\alpha_2\delta$ -3 and Cav3.3, to a smaller degree, Cav3.1 were more prominently found. Most of the  $\gamma$  subunits were also mainly localized in the CNS. Other tissue-elevated expression profiles included  $\beta_1$ ,  $\gamma_1$ ,  $\gamma_6$  (skeletal muscle),  $\alpha_2\delta$ -2 (lung) and  $\alpha_2\delta$ -4 (retina).

While FANTOM5 data is largely ignorant of alternative splicing (except 5' transcription start sites), the HPA data set can also be used to check for the expression of (Ensembl-annotated) splice variants. As an example the two Cav1.1 splice variants, one 'full-length' and Cav1.1<sub>29</sub> (Ensembl identifiers ENST00000362061, ENST00000367338; see also chapter 4) have a slightly different expression distribution. While in skeletal muscle samples the 'full-length' Cav1.1 is ~5-fold more abundant than Cav1.1<sub>29</sub>, it seems to dominate in esophagus.

Some caveats of these databases have to be kept in mind: on one hand only tissues that have been sampled can be compared, on the other hand tissues can be heterogeneous and therefore have a very different channel set in different parts of the organ. This was obvious from the Cav1.4 expression in the HPA data where it is deemed non-specific because there is no retina sample included. Moreover contributions from retina and retinal pigment epithelium could not be distinguished in the FANTOM 5 dataset because presumably these two tissues were not separated during the sample preparation. In general, both the subunit composition of multimeric channels as well as other interactions can affect the severity of channel dysfunctions. Co-expression in a tissue or even a celltype does not yet prove a functional interaction, but these catalogues can help to identify common candidates for protein interaction networks or even new players, including also non-channel proteins.

### 3 Tissue specific expression of L-type Ca<sup>2+</sup>channels

We found a great heterogeneity in the tissue expression of LTCC isoforms. This diversity is expected to impact also on different functional roles of LTCCs in the different tissues. Indeed their physiological importance reaches from excitation-contraction coupling in muscle, to stimulus secretion coupling in sensory and endocrine cells, cardiac pace-making, as well as neuronal firing to learning and memory (Catterall et al., 2005). Cav1.2 and Cav1.3 channels are both expressed in heart, brain, endocrine cells, and auditory and vestibular receptor cells (for review see (Zamponi et al., 2015)). In some neurons, adrenal chromaffin cells, sinoatrial node and atrial cardiomyocytes those isoforms are often even expressed in the same cell type (Olson et al., 2005; Chan et al., 2007; Dragicevic et al., 2014; Marcantoni et al.; Mangoni et al., 2003). Their differential contribution to cellular processes and their physiological functions can therefore only be distinguished in Cav1.2- and Cav1.3-deficient animals (for reviews on animal models see (Striessnig and Koschak, 2008; Hofmann et al., 2014). Cav1.1 and Cav1.4  $\alpha$ 1 transcripts were not found at significant levels in the brain (Figs. 2B and 3; Sinnegger-Brauns et al., 2009). Expression in a limited subset of (brain) neurons or - with respect to Cav1.4 mRNA expression - contamination with pineal gland tissue where Cav1.4 has been reported to be expressed (McRory et al., 2004; Yu et al., 2015a) can't however be entirely excluded. Of note, immunohistochemical analyses that previously reported expression of Cav1.1 in retinal tissue (Specht et al., 2009; Michalakis et al., 2014; Tummala et al., 2014) were carried out with a Cav1.1-directed antibody that was recently discovered to cross-react with GPR179 (Hasan et al., 2016). Additional western blot and mass spectro-metric analyses of Cav1.1 immunoprecipitated proteins also failed to detect any matching Cav1.1 peptides, however a novel splice variant with exons 1–16 skipped and two novel exons termed 16a and 16b has been found by RNAseq (Hasan et al., 2016).

Being described as an atypical prototypical VGCC Cav1.1 is expressed solely in skeletal muscle where it serves as voltage sensor for excitation-contraction coupling and contributes excitation-coupled  $\text{Ca}^{2+}$  entry (for review see (Bannister and Beam, 2013)). In skeletal muscle Cav1.1 is located within the junctional membranes of the t-tubule system where it physically interacts with ryanodine-sensitive release channels in the sarcoplasmic reticulum in order to trigger rapid  $\text{Ca}^{2+}$  release and mediated muscle contraction. Cav1.4 expression however is primarily restricted to the retina. Cav1.4 immunoreactivity has been first described in the outer and inner plexiform layers of chicken retina (Firth et al., 2001). Its expression at photoreceptor release sites located in close vicinity to the typical horseshoe-shaped ribbon synapses has been confirmed in many studies on rodent and zebrafish retina (Mansergh et al., 2005; Specht et al., 2009; Liu et al., 2013; Knoflach et al., 2013; Michalakis et al., 2014; Regus-Leidig et al., 2014; Jia et al., 2014; An et al., 2015; Grabner et al., 2015). Cav1.4 expression in retinal bipolar cells has also been suggested (Berntson et al., 2003; Busquet et al., 2010; Morgans et al., 2005) but functional assessment of the channel subunit composition is lacking. The LTCC identity of these neurons is therefore still ambiguous. As already mentioned above, Cav1.4 was further detected outside the retina in mouse, including tissue of the immune system and the pineal gland (McRory et al., 2004; Kotturi and Jefferies, 2005; Yu et al., 2015a).

#### 4 L-type $\text{Ca}^{2+}$ channel diversity by alternative splicing and its consequences

As the reader can take from Table 2, literature about the post-transcriptional mechanism of alternative splicing in VGCCs is rapidly emerging as is the information about the functional heterogeneity encoded by those alternatively spliced gene products. Alternative splicing is important in all four  $\alpha 1$  isoforms of LTCC. For example one Cav1.1 variant that differs in the length of the IVS3-S4 linker due to skipping of exon 29 (Cav1.1<sub>29</sub>, see also Table 2) was abundantly expressed in mouse and human myotubes (Tuluc et al., 2009) and has also been reported in mouse spleen (Matza et al., 2016). Of note, muscle weakness in the pathogenesis of human myotonic dystrophy type 1 (DM1) and type 2 (DM2) has been related to the aberrant Cav1.1 splicing and altered gating properties in the resulting Cav1.1<sub>29</sub> channels (Tang et al., 2012; Santoro et al., 2014).

A rodent splice variant of Cav1.2 that contained the mutually exclusive exons 21 and 22 (Cav1.2<sub>e21+22</sub>) resulted in nonconducting Cav1.2 channels. In heterologously transfected HEK-293 cells Cav1.2<sub>e21+22</sub> in turn reduced the number of cell surface expressed wild type Cav1.2 channels by enhanced ubiquitination and subsequent proteasomal degradation, that consequently also decreased Cav1.2 mediated  $\text{Ca}^{2+}$  influx (Hu et al., 2016). This variant was highly expressed in the neonatal heart but no expression in brain was reported. Under stressful conditions also adult mouse and human hearts showed increased abundance indicating that this splice variant may even contribute to heart failure (Hu et al., 2016). Another study showed that in a rat myocardial infarction model the expression of a number of alternatively spliced exons were changed. Upon ligation of a descending artery a novel Cav1.2<sub>9\*/delta33</sub> channel (inclusion of exon 9\* and deletion of exon 33) was generated in the scar region that exhibited hyperpolarized shifts of the voltage-dependence of activation



and inactivation in the heterologous expression system (Liao et al., 2009). Intriguingly, in Cav1.2 the binding site for DHPs - which specifically block LTCCs – is encoded by the exclusive exons 8 and 8a (see also Table 2). Cav1.2 channels containing exon 8 are expressed in smooth muscle and display a 10-fold greater affinity for DHP blockers than exon 8a containing channels which comprise the cardiac form (Welling et al., 1997). However, in human heart tissue samples - taken from heart transplant recipients - a subset of tissues expressed predominantly the exon 8 containing smooth muscle isoform, rather than the cardiac isoform (Welling et al., 1997; Wang et al., 2006). This finding suggested that in affected individuals therapeutic DHP treatment could cause strong unwanted effects in the heart. This is a clear example highlighting that interindividual variability in alternative splicing can severely modulate pharmacological effects through distinct tissue expression and therefore cause harm to patients.

Furthermore, also drug action on Cav1.3 channels depends on the effects of alternative splicing. An intramolecular interaction between a distal and a proximal regulatory domain – forming an intrinsic C-terminal modulatory mechanism, short CTM - within the Cav1.3 C-terminus is a major determinant of their voltage- and  $\text{Ca}^{2+}$ -dependent gating kinetics (Striessnig, 2007). Splicing in the C-terminus of Cav1.3  $\alpha 1$  subunits gives rise to fundamentally different channels with either ‘long’ or ‘short’ C-termini. Thereby removal of the CTM domains by alternative splicing generates the ‘short’ Cav1.3 (Cav1.3<sub>42A</sub>, Cav1.3<sub>43S</sub>) channels. These splice isoforms activated at a more negative voltage range and exhibited different voltage- and  $\text{Ca}^{2+}$ -dependent inactivation properties of these channels (Bock et al., 2011; Singh et al., 2008; Tan et al., 2012). At this point it is important to know that the action of DHPs depends on the inactivated channel state (Bean et al., 1986; Hamilton et al., 1987; Koschak et al., 2001; Berjukow and Hering, 2001) such that DHPs possess a higher affinity for the inactivated channel conformation. The amount of channel block is therefore expected to change with the availability of inactivated channel states. Of note, the expression of both short and long variants showed a certain tissue specificity. The long variant seemed to predominate in the heart whereas expression of both variants were about equally abundant in the brain (Bock et al., 2011). Together, this suggests that DHP drugs might act differently on Cav1.3 channel variants expressed in different tissues to fine-tune channel function (Huang et al., 2014; Ortner et al., 2017). Also in Cav1.4, the most abundantly expressed LTCC in the retina, alternative splicing occurs. In a transcript scanning approach of human retina samples, 19 Cav1.4 splice variants were identified (Tan et al., 2012). One of those also resulted in a truncation of the channels’ C-terminus (Ca<sub>v</sub>1.4 ex43\*). Similar to Cav1.3 channels, such structural change is also expected to severely affect the voltage- and  $\text{Ca}^{2+}$ -dependent gating properties of Cav1.4 channels (Singh et al., 2006; Wahl-Schott et al., 2006). Indeed, the loss of CTM in the splice variant Ca<sub>v</sub>1.4 ex43\* resulted in robust  $\text{Ca}^{2+}$ -dependent inactivation and a hyperpolarized activation range (Haeseleer et al., 2016). Such gating changes might be even pathological for visual signaling (Singh et al., 2006; Wahl-Schott et al., 2006). Though a recent heterologous expression study provided evidence that coexpression of CaBP4 - a member of a family of  $\text{Ca}^{2+}$ -binding proteins (CaBPs) related to CaM and expressed in retinal photoreceptors (Haeseleer et al., 2004) – could at least in part compensate for the expected Cav1.4 gating changes by binding to the remaining C-terminal domain of the Ca<sub>v</sub>1.4 C-terminus (Haeseleer et al.,

2016). To which extent these naturally occurring alternative splice variants indeed add to the properties of native Cav1.4 currents (von Gersdorff and Matthews, 1996; Rabl and Thoreson, 2002) or whether the abundance of particular splice variants is changed under pathophysiological conditions - as seen for all other LTCCs (Table 2) - is not yet resolved. Details on these functionally different Cav1.4 splice variants in other tissues (lymphocytes or pineal gland) are yet to be elucidated.

## 5 Manifestation of Cav1.4-related channelopathies and treatment challenges

Tissue expression of a certain protein often correlates with histological and functional features seen in animal models and humans as well as the clinical features in case of a dysfunctional protein. Among the LTCC family Cav1.4 is one of the finest examples.

The role of Cav1.4 is well supported by the fact that mutations in the *CACNA1F* gene can cause several forms of human retinal diseases (OMIM #300071, #300476, #300600, for review see (Zeitz et al., 2015)). The majority of Cav1.4 mutations were identified in congenital stationary night blindness type 2 (CSNB2) patients who are presenting with typical symptoms like low visual acuity, myopia, nystagmus, strabismus, photophobia and variable levels of night blindness (Bech-Hansen et al., 1998a,b). CSNB2 mainly involves males due to the X-linked nature of the disease but recent reports substantiated that heterozygote females can also be affected (Hope et al., 2005; Michalakis et al., 2014). Both loss of channel function and increased channel activity can lead to alterations in photoreceptor synapse formation. This has been demonstrated in Cav1.4 deficient mice (Cav1.4<sup>-/-</sup> (Morgans et al., 2001; Mansergh et al., 2005; Raven et al., 2008; Zabouri and Haverkamp, 2013; Specht et al., 2009)) and also in Cav1.4 mutant mice. The latter contain a point mutation at position Ile756Thr (corresponding to Ile745Thr in the patients) that induced enhanced channel activity obvious from a strong hyperpolarizing shift in the voltage-dependence of activation and slowing of the inactivation properties (Cav1.4-IT; (Hemara-Wahanui et al., 2005)). A large body of work done by many labs highlighted that Cav1.4 channels are central for the proper formation and maturation of photoreceptor synapses (Morgans et al., 2001; Mansergh et al., 2005; Chang et al., 2006; Bayley and Morgans, 2007; Raven et al., 2008; Zabouri and Haverkamp, 2013; Specht et al., 2009; Liu et al., 2013; Regus- Leidig et al., 2014; Michalakis et al., 2014).

Functional data from CSNB2 patients still implied an incomplete defect of the ON and OFF bipolar cells or their synapses in the rod and cone visual pathways (Miyake et al., 1986; Miyake, 2002). Indeed signal transmission was not completely blocked but some reduced activity remained in Cav1.4 mutant retinas (Chang et al., 2006; Knoflach et al., 2015). In the Cav1.4-IT mouse model the left shift in the voltage-dependence of channel activation was hypothesized to reduce the dynamic range of photoreceptor activity. In line with this hypothesis only a fraction of Cav1.4-IT ganglion cells responded to light stimulation in multielectrode array recordings from whole-mounted retinas, and spatial responses as well as changes in their contrast sensitivity profile were decreased in these mutants (Knoflach et al., 2015). Of note, strongly reduced visual acuity in fact turned out to be a major complaint



in CSNB2 patients (Bijveld et al., 2013) and patients carrying the respective Cav1.4 mutation also showed pronounced vision loss (up to 80% (Hope et al., 2005)).

When we consider treatment of retinal Cav1.4-related channelopathies we face several challenges. The very restricted expression of Cav1.4 channels in retinal tissue should be a clear advantage for Cav1.4 targeted therapy. But still, in systemic pharmacotherapy due to the expression of Cav1.4 in immune cells or pineal gland tissue (see above) the potential effects on the immune system or the sleep/wake cycle would have to be controlled for, even though no obvious deficits in circadian rhythm or immune system have been documented so far in CSNB2 patients. The main problem however lies in the specificity of available drugs. Currently all LTCC blockers that are in clinical use, as for example the DHP nifedipine or other DHPs (Striessnig et al., 2015), are considered non-selective because Cav1.2, Cav1.3 and Cav1.4 channels only slightly differ in their DHP sensitivity (Koschak et al., 2001; Xu and Lipscombe, 2001; Koschak et al., 2003). Due to their high binding affinity to Cav1.2 channels DHPs mainly have cardiovascular effects ( $Ca_v1.2$  expression in cardiovascular tissue, see Figs. 2C and 3; (Moosmang et al., 2003; Zhang et al., 2006)). For this reason, DHPs are not applicable in CSNB2 patients – and also for treatment of other CNS disorders – considering the severe side effects of the necessary therapeutic doses (for review see (Zamponi et al., 2015)). When considering restoration of channel function either by pharmacological chaperoning in the retina (<http://clinicaltrials.gov/ct2/show/study/NCT01233609>) or gene-based retinal therapies.

(Sengillo et al., 2017), treatments likely would have to start early during retinal development because of the crucial role of Cav1.4 for proper synapse formation already mentioned above.

## 6 Gene therapy of Cav1.4 channelopathies

Treatment e.g. of skeletal muscle by gene therapy is considered in humans (for review (Saada et al., 2016; Sommese et al., 2017)) and has been demonstrated in animal models (Gruntman et al., 2013; Qiao et al., 2011) (Wang et al., 2014) even without the use of viral vectors (Lu et al., 2003), though the necessary systemic approach to treat all muscles in the body bears certain risks given by the widespread distribution of muscle tissues. The focal expression of Cav1.4 in the eye on the other hand provides a number of advantages which are briefly outlined in the following.

The eye is not only easily accessible but it is also a small sized and confined organ. Thus, low amounts of viral vectors are sufficient and there is little risk of a systemic spread of the vector. For this reason and for reasons explained below, the eye has been targeted in gene therapy (Sengillo et al., 2017; Scholl et al., 2016). In particular, intravitreal injections of vectors are easy to perform and constitute a routine procedure in eye hospitals without the need for inpatient admission. Intravitreal injections can reach large areas of the retina (Da Costa et al., 2016), a clear advantage over subretinal delivery, where a needle has to penetrate the retina to inject vectors between retina and choroid. Of note, adeno-associated viruses (AAV) that effectively target photoreceptors even from intravitreal injections (e.g. AAV.7m8) have been developed and applied successfully in gene therapy in mice (Dalkara et al., 2013). Alternatively, the functionality of selective promoters for expression in rods

(Koch et al., 2012) or (S-)cones (Michalakis et al., 2010) without cell-type specific tropism has been demonstrated. Targeting other cell types is also feasible due to the comparatively well-known cell types in the retina and the knowledge about cell-type specific genes (Siegert et al., 2012) and promoters (Ivanova et al., 2010). In the case of Cav1.4 it might be necessary to treat not only photoreceptors but also bipolar cells that express Cav1.4 (Berntson et al., 2003), so a mixed approach of different vectors/tropisms or promoters could be required. Targeting Cav1.4 might still be challenging due to the size of the *CACNA1F* gene (OMIM 300100) and the limited packaging capacity of AAVs. However, split-intein-mediated protein transsplicing methods have already been used successfully to manipulate other LTCCs (Subramanyam et al., 2013).

Viral vector-based gene therapy benefits from the immune-privileged status and the specific location of the retina in the eye. The blood-retina barrier provides an advantage in the protection of treated cells from the patients' immune responses by limiting potential inflammatory reactions (Bennett, 2003). It has however been observed that intravitreal injections can trigger an immune response not observed by subretinal injections (Li et al., 2008). Still, the enclosed structure of the eye in conjunction with the blood-retina barrier provides a spatial restriction of gene therapeutic treatments almost unrivalled by other organs. These features are relevant for all genetic interventions, in particular in CRISPR/Cas9 applications (Hung et al., 2016; Latella et al., 2016; Bakondi et al., 2016; Yanik et al., 2017; Ruan et al., 2017) where off-target effects might not be fully excluded (Fu et al., 2014). Together, the eye/retina provides a high level of containment with limited immune response, making applications comparatively safe. Indeed several clinical trials of gene therapy for eye diseases in patients have been undertaken in recent years (Bainbridge et al., 2008; Maguire et al., 2008; Cideciyan et al., 2008). Cav1.4 mutations would therefore be amenable for gene therapeutic treatments in the foreseeable future.

## Funding

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 674901 to AK, and was supported by the Austrian Science Fund (FWF P-26881-B23, P-29359 to AK), the Center of Molecular Biosciences Innsbruck and the University of Innsbruck.

## References

- An J, Zhang L, Jiao B, Lu F, Xia F, Yu Z, Zhang Z. *Cacna1f* gene decreased contractility of skeletal muscle in rat model with congenital stationary night blindness. *Gene*. 2015; 562:210–219. [PubMed: 25748727]
- Badou A, Jha MK, Matza D, Flavell RA. Emerging roles of L-type voltagegated and other calcium channels in T lymphocytes. *Front Immunol*. 2013; 4:243. [PubMed: 24009608]
- Bainbridge JW, Smith AJ, Barker SS, Robbie S, Henderson R, Balaggan K, Viswanathan A, Holder GE, Stockman A, Tyler N, Petersen-Jones S, et al. Effect of gene therapy on visual function in Leber's congenital amaurosis. *N Engl J Med*. 2008; 358:2231–2239. [PubMed: 18441371]
- Bakondi B, Lv W, Lu B, Jones MK, Tsai Y, Kim KJ, Levy R, Akhtar AA, Breunig JJ, Svendsen CN, Wang S. In Vivo CRISPR/Cas9 gene editing Corrects retinal dystrophy in the S334ter-3 rat model of autosomal dominant retinitis pigmentosa. *Mol Ther*. 2016; 24:556–563. [PubMed: 26666451]
- Bannister RA, Beam KG. Ca(V)1.1: the atypical prototypical voltage-gated Ca(2)(+) channel. *Biochim Biophys Acta*. 2013; 1828:1587–1597. [PubMed: 22982493]

- Bayley PR, Morgans CW. Rod bipolar cells and horizontal cells form displaced synaptic contacts with rods in the outer nuclear layer of the nob2 retina. *J Comp Neurol.* 2007; 500:286–298. [PubMed: 17111373]
- Bean BP, Sturek M, Puga A, Hermsmeyer K. Calcium channels in muscle cells isolated from rat mesenteric arteries: modulation by dihydropyridine drugs. *Circ Res.* 1986; 59:229–235. [PubMed: 2427250]
- Bech-Hansen NT, Boycott KM, Gratton KJ, Ross DA, Field LL, Pearce WG. Localization of a gene for incomplete X-linked congenital stationary night blindness to the interval between DXS6849 and DXS8023 in Xp11.23. *Hum Genet.* 1998a; 103:124–130. [PubMed: 9760193]
- Bech-Hansen NT, Naylor MJ, Maybaum TA, Pearce WG, Koop B, Fishman GA, Mets M, Musarella MA, Boycott KM. Loss-of-function mutations in a calcium-channel alpha1-subunit gene in Xp11.23 cause incomplete X-linked congenital stationary night blindness. *Nat Genet.* 1998b; 19:264–26. [PubMed: 9662400]
- Bennett J. Immune response following intraocular delivery of recombinant viral vectors. *Gene Ther.* 2003; 10:977–982. [PubMed: 12756418]
- Berjukow S, Hering S. Voltage-dependent acceleration of Ca(v)1.2 channel current decay by (+)- and (-)-isradipine. *Br J Pharmacol.* 2001; 133:959–966. [PubMed: 11487504]
- Berntson A, Taylor WR, Morgans CW. Molecular identity, synaptic localization, and physiology of calcium channels in retinal bipolar cells. *J Neurosci Res.* 2003; 71:146–151. [PubMed: 12478624]
- Bijveld MM, Florijn RJ, Bergen AA, van den Born LI, Kamermans M, Prick L, Riemsdag FC, van Schooneveld MJ, Kappers AM, van Genderen MM. Genotype and phenotype of 101 Dutch patients with congenital stationary night blindness. *Ophthalmology.* 2013; 120:2072–208. [PubMed: 23714322]
- Bock G, Gebhart M, Scharinger A, Jangsangthong W, Busquet P, Poggiani C, Sartori S, Mangoni ME, Sinnegger-Brauns MJ, Herzig S, Striessnig J, et al. Functional properties of a newly identified C-terminal splice variant of Cav1.3 L-type Ca<sup>2+</sup> channels. *J Biol Chem.* 2011; 286:42736–42748. [PubMed: 21998310]
- Busquet P, Nguyen NK, Schmid E, Tanimoto N, Seeliger MW, Ben-Yosef T, Mizuno F, Akopian A, Striessnig J, Singewald N. CaV1.3 L-type Ca<sup>2+</sup> channels modulate depression-like behaviour in mice independent of deaf phenotype. *Int J Neuropsychopharmacol.* 2010; 13:499–513. [PubMed: 19664321]
- Cabral MD, Paulet PE, Robert V, Gomes B, Renoud ML, Savignac M, Leclerc C, Moreau M, Lair D, Langelot M, Magnan A, et al. Knocking down Cav1 calcium channels implicated in Th2 cell activation prevents experimental asthma. *Am J Respir Crit Care Med.* 2010; 181:1310–1317. [PubMed: 20167851]
- Catterall WA. Voltage-gated calcium channels. *Cold Spring Harb Perspect Biol.* 2011; 3:a003947. [PubMed: 21746798]
- Catterall WA, Perez-Reyes E, Snutch TP, Striessnig J. International Union of Pharmacology. XLVIII. Nomenclature and structure-function relationships of voltage-gated calcium channels. *Pharmacol Rev.* 2005; 57:411–425. [PubMed: 16382099]
- Chan CS, Guzman JN, Ilijic E, Mercer JN, Rick C, Tkatch T, Meredith GE, Surmeier DJ. ‘Rejuvenation’ protects neurons in mouse models of Parkinson’s disease. *Nature.* 2007; 447:1081–1086. [PubMed: 17558391]
- Chang B, Heckenlively JR, Bayley PR, Brecha NC, Davisson MT, Hawes NL, Hirano AA, Hurd RE, Ikeda A, Johnson BA, McCall MA, et al. The nob2 mouse, a null mutation in *Cacna1f*: anatomical and functional abnormalities in the outer retina and their consequences on ganglion cell visual responses. *Vis Neurosci.* 2006; 23:11–24. [PubMed: 16597347]
- Cideciyan AV, Aleman TS, Boye SL, Schwartz SB, Kaushal S, Roman AJ, Pang JJ, Sumaroka A, Windsor EA, Wilson JM, Flotte TR, et al. Human gene therapy for RPE65 isomerase deficiency activates the retinoid cycle of vision but with slow rod kinetics. *Proc Natl Acad Sci U S A.* 2008; 105:15112–15117. [PubMed: 18809924]
- Da Costa R, Roger C, Segelken J, Barben M, Grimm C, Neidhardt J. A novel method combining vitreous aspiration and intravitreal AAV2/8 injection results in retina-wide transduction in adult mice. *Invest Ophthalmol Vis Sci.* 2016; 57:5326–5334. [PubMed: 27784063]

- Dalkara D, Byrne LC, Klimczak RR, Visel M, Yin L, Merigan WH, Flannery JG, Schaffer DV. In vivo-directed evolution of a new adeno-associated virus for therapeutic outer retinal gene delivery from the vitreous. *Sci Transl Med.* 2013; 5:189ra176.
- Davenport B, Li Y, Heizer JW, Schmitz C, Perraud AL. Signature channels of excitability no more: L-type channels in immune cells. *Front Immunol.* 2015; 6:375. [PubMed: 26257741]
- Dragicevic E, Poetschke C, Duda J, Schlaudraff F, Lammel S, Schiemann J, Fauler M, Hetzel A, Watanabe M, Lujan R, Malenka RC, et al. Cav1.3 channels control D2-autoreceptor responses via NCS-1 in substantia nigra dopamine neurons. *Brain.* 2014; 137:2287–2302. [PubMed: 24934288]
- Eden E, Geva-Zatorsky N, Issaeva I, Cohen A, Dekel E, Danon T, Cohen L, Mayo A, Alon U. Proteome half-life dynamics in living human cells. *Science.* 2011; 331:764–768. [PubMed: 21233346]
- Fernandez JA, McGahon MK, McGeown JG, Curtis TM. CaV3.1 T-Type Ca<sup>2+</sup> channels contribute to myogenic signaling in rat retinal arterioles. *Invest Ophthalmol Vis Sci.* 2015; 56:5125–5132. [PubMed: 26241400]
- Firth SI, Morgan IG, Boelen MK, Morgans CW. Localization of voltagesensitive L-type calcium channels in the chicken retina. *Clin Exp Ophthalmol.* 2001; 29:183–187. [PubMed: 11446465]
- Fu Y, Sander JD, Reyon D, Cascio VM, Joung JK. Improving CRISPR-Cas nuclease specificity using truncated guide RNAs. *Nat Biotechnol.* 2014; 32:279–284. [PubMed: 24463574]
- Grabner CP, Gandini MA, Rehak R, Le Y, Zamponi GW, Schmitz F. RIM1/2-Mediated facilitation of Cav1.4 channel opening is required for Ca<sup>2+</sup>-stimulated release in mouse rod photoreceptors. *J Neurosci.* 2015; 35:13133–13147. [PubMed: 26400943]
- Gruntman AM, Bish LT, Mueller C, Sweeney HL, Flotte TR, Gao G. Gene transfer in skeletal and cardiac muscle using recombinant adeno-associated virus. *Curr Protoc Microbiol.* 2013; doi: 10.1002/9780471729259.mc14d03s28
- Haeseleer F, Imanishi Y, Maeda T, Possin DE, Maeda A, Lee A, Rieke F, Palczewski K. Essential role of Ca<sup>2+</sup>-binding protein 4, a Cav1.4 channel regulator, in photoreceptor synaptic function. *Nat Neurosci.* 2004; 7:1079–1087. [PubMed: 15452577]
- Haeseleer F, Williams B, Lee A. Characterization of C-terminal splice variants of Cav1.4 Ca<sup>2+</sup> channels in human retina. *J Biol Chem.* 2016; 291:15663–15673. [PubMed: 27226626]
- Hamilton SL, Yatani A, Brush K, Schwartz A, Brown AM. A comparison between the binding and electrophysiological effects of dihydropyridines on cardiac membranes. *Mol Pharmacol.* 1987; 31:221–231. [PubMed: 2436031]
- Harraz OF, Brett SE, Zechariah A, Romero M, Puglisi JL, Wilson SM, Welsh DG. Genetic ablation of CaV3.2 channels enhances the arterial myogenic response by modulating the RyR-BKCa axis. *Arterioscler Thromb Vasc Biol.* 2015a; 35:1843–1851. [PubMed: 26069238]
- Harraz OF, Visser F, Brett SE, Goldman D, Zechariah A, Hashad AM, Menon BK, Watson T, Starreveld Y, Welsh DG. CaV1.2/CaV3.x channels mediate divergent vasomotor responses in human cerebral arteries. *J Gen Physiol.* 2015b; 145:405–418. [PubMed: 25918359]
- Hasan N, Ray TA, Gregg RG. CACNA1S expression in mouse retina: novel isoforms and antibody cross-reactivity with GPR179. *Vis Neurosci.* 2016; 33:E009. [PubMed: 27471951]
- Hemara-Wahanui A, Berjukow S, Hope CI, Dearden PK, Wu SB, Wilson-Wheeler J, Sharp DM, Landon-Treweek P, Clover GM, Hoda JC, Striessnig J, et al. A CAC-NA1F mutation identified in an X-linked retinal disorder shifts the voltage dependence of Cav1.4 channel activation. *Proc Natl Acad Sci U S A.* 2005; 102:7553–7558. [PubMed: 15897456]
- Hofmann F, Flockerzi V, Kahl S, Wegener JW. L-type CaV1.2 calcium channels: from in vitro findings to in vivo function. *Physiol Rev.* 2014; 94:303–326. [PubMed: 24382889]
- Hope CI, Sharp DM, Hemara-Wahanui A, Sissingh JI, Landon P, Mitchell EA, Maw MA, Clover GM. Clinical manifestations of a unique X-linked retinal disorder in a large New Zealand family with a novel mutation in CAC-NA1F, the gene responsible for CSNB2. *Clin Exp Ophthalmol.* 2005; 33:129–136. [PubMed: 15807819]
- Hu Z, Wang JW, Yu D, Soon JL, de Kleijn DP, Foo R, Liao P, Colecraft HM, Soong TW. Aberrant splicing promotes proteasomal degradation of L-type CaV1.2 calcium channels by Competitive binding for CaVbeta subunits in cardiac hypertrophy. *Sci Rep.* 2016; 6:35247. [PubMed: 27731386]

- Huang H, Ng CY, Yu D, Zhai J, Lam Y, Soong TW. Modest Ca<sub>v</sub>1.342-selective inhibition by compound 8 is beta-subunit dependent. *Nat Commun.* 2014; 5:4481. [PubMed: 25057870]
- Hung SS, Chrysostomou V, Li F, Lim JK, Wang JH, Powell JE, Tu L, Daniszewski M, Lo C, Wong RC, Crowston JG, et al. AAV-mediated CRISPR/Cas gene editing of retinal cells in vivo. *Invest Ophthalmol Vis Sci.* 2016; 57:3470–3476. [PubMed: 27367513]
- Ivanova E, Hwang GS, Pan ZH. Characterization of transgenic mouse lines expressing Cre recombinase in the retina. *Neuroscience.* 2010; 165:233–243. [PubMed: 19837136]
- Jha A, Singh AK, Weissgerber P, Freichel M, Flockerzi V, Flavell RA, Jha MK. Essential roles for Cavbeta2 and Cav1 channels in thymocyte development and T cell homeostasis. *Sci Signal.* 2015; 8:ra103. [PubMed: 26486172]
- Jia S, Muto A, Orisme W, Henson HE, Parupalli C, Ju B, Baier H, Taylor MR. Zebrafish *Cacna1fa* is required for cone photoreceptor function and synaptic ribbon formation. *Hum Mol Genet.* 2014; 23:2981–2994. [PubMed: 24419318]
- Jurkat-Rott K, Lerche H, Lehmann-Horn F. Skeletal muscle channelopathies. *J Neurol.* 2002; 249:1493–1502. [PubMed: 12420087]
- Knoflach D, Kerov V, Sartori SB, Obermair GJ, Schmuckermair C, Liu X, Sothilingam V, Garrido MG, Baker SA, Glosmann M, Schicker K, et al. Cav1.4 IT mouse as model for vision impairment in human congenital stationary night blindness type 2. *Channels (Austin).* 2013; 7:503–513. [PubMed: 24051672]
- Knoflach D, Schicker K, Glosmann M, Koschak A. Gain-of-function nature of Cav1.4 L-type calcium channels alters firing properties of mouse retinal ganglion cells. *Channels (Austin).* 2015; 9:298–306. [PubMed: 26274509]
- Koch S, Sothilingam V, Garcia Garrido M, Tanimoto N, Becirovic E, Koch F, Seide C, Beck SC, Seeliger MW, Biel M, Muhlfriedel R, et al. Gene therapy restores vision and delays degeneration in the CNGB1(-/-) mouse model of retinitis pigmentosa. *Hum Mol Genet.* 2012; 21:4486–4496. [PubMed: 22802073]
- Koschak A, Reimer D, Huber I, Grabner M, Glosmann H, Engel J, Striessnig J.  $\alpha$ 1D(Ca<sub>v</sub>1.3) subunits can form L-type Ca<sub>2+</sub> channels activating at negative voltages. *J Biol Chem.* 2001; 276:22100–22106. [PubMed: 11285265]
- Koschak A, Reimer D, Walter D, Hoda JC, Heinzle T, Grabner M, Striessnig J. Cav1.4 $\alpha$ 1 subunits can form slowly inactivating dihydropyridine-sensitive L-type Ca<sup>2+</sup> channels lacking Ca<sup>2+</sup>-dependent inactivation. *J Neurosci.* 2003; 23:6041–6049. [PubMed: 12853422]
- Kotturi MF, Jefferies WA. Molecular characterization of L-type calcium channel splice variants expressed in human T lymphocytes. *Mol Immunol.* 2005; 42:1461–1474. [PubMed: 15899519]
- Kuo IY, Howitt L, Sandow SL, McFarlane A, Hansen PB, Hill CE. Role of T-type channels in vasomotor function: team player or chameleon? *Pflugers Arch.* 2014; 466:767–779. [PubMed: 24482062]
- Latella MC, Di Salvo MT, Cocchiarella F, Benati D, Grisendi G, Comitato A, Marigo V, Recchia A. In vivo editing of the human mutant rhodopsin gene by electroporation of plasmid-based CRISPR/Cas9 in the mouse retina. *Mol Ther Nucleic Acids.* 2016; 5:e389. [PubMed: 27874856]
- Li Q, Miller R, Han PY, Pang J, Dinculescu A, Chiodo V, Hauswirth WW. Intraocular route of AAV2 vector administration defines humoral immune response and therapeutic potential. *Mol Vis.* 2008; 14:1760–1769. [PubMed: 18836574]
- Liao P, Li G, Yu de J, Yong TF, Wang JJ, Wang J, Soong TW. Molecular alteration of Ca(v)1.2 calcium channel in chronic myocardial infarction. *Pflugers Arch.* 2009; 458:701–711. [PubMed: 19263075]
- Liu X, Kerov V, Haeseleer F, Majumder A, Artemyev N, Baker SA, Lee A. Dysregulation of Ca(v)1.4 channels disrupts the maturation of photoreceptor synaptic ribbons in congenital stationary night blindness type 2. *Channels (Austin).* 2013; 7:514–523. [PubMed: 24064553]
- Lizio M, Harshbarger J, Shimoji H, Severin J, Kasukawa T, Sahin S, Abugessaisa I, Fukuda S, Hori F, Ishikawa-Kato S, Mungall CJ, et al. Gateways to the FANTOM5 promoter level mammalian expression atlas. *Genome Biol.* 2015; 16:22. [PubMed: 25723102]
- Liu QL, Bou-Gharios G, Partridge TA. Non-viral gene delivery in skeletal muscle: a protein factory. *Gene Ther.* 2003; 10:131–142. [PubMed: 12571642]




- Maguire AM, Simonelli F, Pierce EA, Pugh EN Jr, Mingozzi F, Bennicelli J, Banfi S, Marshall KA, Testa F, Surace EM, et al. Safety and efficacy of gene transfer for Leber's congenital amaurosis. *N Engl J Med.* 2008; 358:2240–2248. [PubMed: 18441370]
- Mangoni ME, Couette B, Bourinet E, Platzer J, Reimer D, Striessnig J, Nargeot J. Functional role of L-type Cav1.3 Ca<sup>2+</sup> channels in cardiac pacemaker activity. *Proc Natl Acad Sci U S A.* 2003; 100:5543–5548. [PubMed: 12700358]
- Mansergh F, Orton NC, Vessey JP, Lalonde MR, Stell WK, Tremblay F, Barnes S, Rancourt DE, Bech-Hansen NT. Mutation of the calcium channel gene *Cacnal1f* disrupts calcium signaling, synaptic transmission and cellular organization in mouse retina. *Hum Mol Genet.* 2005; 14:3035–3046. [PubMed: 16155113]
- Marcantoni A, Vandael DH, Mahapatra S, Carabelli V, Sinnegger-Brauns MJ, Striessnig J, Carbone E. Loss of Cav1.3 channels reveals the critical role of L-type and BK channel coupling in pacemaking mouse adrenal chromaffin cells. *J Neurosci.* 30:491–504.
- Matza D, Badou A, Klemic KG, Stein J, Govindarajulu U, Nadler MJ, Kinet JP, Peled A, Shapira OM, Kaczmarek LK, Flavell RA. T cell receptor mediated calcium entry requires alternatively spliced Cav1.1 channels. *PLoS One.* 2016; 11:e0147379. [PubMed: 26815481]
- McRory JE, Hamid J, Doering CJ, Garcia E, Parker R, Hamming K, Chen L, Hildebrand M, Beedle AM, Feldcamp L, Zamponi GW, et al. The *CACNA1F* gene encodes an L-type calcium channel with unique biophysical properties and tissue distribution. *J Neurosci.* 2004; 24:1707–1718. [PubMed: 14973233]
- Michalakis S, Muhlfriedel R, Tanimoto N, Krishnamoorthy V, Koch S, Fischer MD, Becirovic E, Bai L, Huber G, Beck SC, Fahl E, et al. Restoration of cone vision in the *CNGA3*<sup>-/-</sup> mouse model of congenital complete lack of cone photoreceptor function. *Mol Ther.* 2010; 18:2057–2063. [PubMed: 20628362]
- Michalakis S, Shaltiel L, Sothilingam V, Koch S, Schludi V, Krause S, Zeitz C, Audo I, Lancelot ME, Hamel C, Meunier I, et al. Mosaic synaptopathy and functional defects in Cav1.4 heterozygous mice and human carriers of *CSNB2*. *Hum Mol Genet.* 2014; 23:1538–1550. [PubMed: 24163243]
- Miyake Y. Establishment of the concept of new clinical entities—complete and incomplete form of congenital stationary night blindness. *Nippon Ganka Gakkai Zasshi.* 2002; 106:737–755. [PubMed: 12610835]
- Miyake Y, Yagasaki K, Horiguchi M, Kawase Y, Kanda T. Congenital stationary night blindness with negative electroretinogram. A new classification. *Arch Ophthalmol.* 1986; 104:1013–1020. [PubMed: 3488053]
- Moosmang S, Schulla V, Welling A, Feil R, Feil S, Wegener JW, Hofmann F, Klugbauer N. Dominant role of smooth muscle L-type calcium channel Cav1.2 for blood pressure regulation. *EMBO J.* 2003; 22:6027–6034. [PubMed: 14609949]
- Morgans CW, Bayley PR, Oesch NW, Ren G, Akileswaran L, Taylor WR. Photoreceptor calcium channels: insight from night blindness. *Vis Neurosci.* 2005; 22:561–568. [PubMed: 16332266]
- Morgans CW, Gaughwin P, Maleszka R. Expression of the alpha1F calcium channel subunit by photoreceptors in the rat retina. *Mol Vis.* 2001; 7:202–209. [PubMed: 11526344]
- Muller C, Mas Gomez N, Ruth P, Strauss O. CaV1.3 L-type channels, maxiK Ca(2+)-dependent K(+) channels and bestrophin-1 regulate rhythmic photoreceptor outer segment phagocytosis by retinal pigment epithelial cells. *Cell Signal.* 2014; 26:968–978. [PubMed: 24407175]
- Olson PA, Tkatch T, Hernandez-Lopez S, Ulrich S, Ilijic E, Mugnaini E, Zhang H, Bezprozvany I, Surmeier DJ. G-protein-coupled receptor modulation of striatal Cav1.3 L-type Ca<sup>2+</sup> channels is dependent on a Shankbinding domain. *J Neurosci.* 2005; 25:1050–1062. [PubMed: 15689540]
- Omilusik K, Priatel JJ, Chen X, Wang YT, Xu H, Choi KB, Gopaul R, McIntyre-Smith A, Teh HS, Tan R, Bech-Hansen NT, et al. The Ca(v)1.4 calcium channel is a critical regulator of T cell receptor signaling and naive T cell homeostasis. *Immunity.* 2011; 35:349–360. [PubMed: 21835646]
- Ortner NJ, Bock G, Dougalis A, Kharitonova M, Duda J, Hess S, Tuluc P, Pomberger T, Stefanova N, Pitterl F, Ciossek T, et al. Lower affinity of isradipine for L-type Ca<sup>2+</sup> channels during substantia nigra dopamine neuron-like activity: implications for neuroprotection in Parkinson's disease. *J Neurosci.* 2017 Jun 7. doi: 10.1523/JNEUROSCI.2946-16.2017




- Qiao C, Koo T, Li J, Xiao X, Dickson JG. Gene therapy in skeletal muscle mediated by adeno-associated virus vectors. *Methods Mol Biol.* 2011; 807:119–140. [PubMed: 22034028]
- Rabl K, Thoreson WB. Calcium-dependent inactivation and depletion of synaptic cleft calcium ions combine to regulate rod calcium currents under physiological conditions. *Eur J Neurosci.* 2002; 16:2070–2077. [PubMed: 12473074]
- Raven MA, Orton NC, Nassar H, Williams GA, Stell WK, Jacobs GH, Bech-Hansen NT, Reese BE. Early afferent signaling in the outer plexiform layer regulates development of horizontal cell morphology. *J Comp Neurol.* 2008; 506:745–758. [PubMed: 18076080]
- Regus-Leidig H, Atorf J, Feigenspan A, Kremers J, Maw MA, Brandstatter JH. Photoreceptor degeneration in two mouse models for congenital stationary night blindness type 2. *PLoS One.* 2014; 9:e86769. [PubMed: 24466230]
- Robert V, Triffaux E, Paulet PE, Guery JC, Pelletier L, Savignac M. Protein kinase C-dependent activation of CaV1.2 channels selectively controls human TH2-lymphocyte functions. *J Allergy Clin Immunol.* 2014; 133:1175–1183. [PubMed: 24365142]
- Ruan GX, Barry E, Yu D, Lukason M, Cheng SH, Scaria A. CRISPR/Cas9-Mediated genome editing as a therapeutic approach for Leber congenital amaurosis 10. *Mol Ther.* 2017; 25:331–341. [PubMed: 28109959]
- Saada YB, Dib C, Lipinski M, Vassetzky YS. Genome- and cell-based strategies in therapy of muscular dystrophies. *Biochem (Mosc).* 2016; 81:678–690.
- Santoro M, Piacentini R, Masciullo M, Bianchi ML, Modoni A, Podda MV, Ricci E, Silvestri G, Grassi C. Alternative splicing alterations of Ca<sup>2+</sup> handling genes are associated with Ca<sup>2+</sup> signal dysregulation in myotonic dystrophy type 1 (DM1) and type 2 (DM2) myotubes. *Neuropathol Appl Neurobiol.* 2014; 40:464–476. [PubMed: 23888875]
- Schlick B, Flucher BE, Obermair GJ. Voltage-activated calcium channel expression profiles in mouse brain and cultured hippocampal neurons. *Neuroscience.* 2010; 167:786–798. [PubMed: 20188150]
- Scholl HP, Strauss RW, Singh MS, Dalkara D, Roska B, Picaud S, Sahel JA. Emerging therapies for inherited retinal degeneration. *Sci Transl Med.* 2016; 8:368rv366.
- Sengillo JD, Justus S, Cabral T, Tsang SH. Correction of monogenic and common retinal disorders with gene therapy. *Genes (Basel).* 2017; 8
- Siegert S, Cabuy E, Scherf BG, Kohler H, Panda S, Le YZ, Fehling HJ, Gaidatzis D, Stadler MB, Roska B. Transcriptional code and disease map for adult retinal cell types. *Nat Neurosci.* 2012; 15(487-495):S481–S482.
- Singh A, Gebhart M, Fritsch R, Sinnegger-Brauns MJ, Poggiani C, Hoda JC, Engel J, Romanin C, Striessnig J, Koschak A. Modulation of voltage- and Ca<sup>2+</sup>-dependent gating of CaV1.3 L-type calcium channels by alternative splicing of a C-terminal regulatory domain. *J Biol Chem.* 2008 Jul 25; 283(30):20733–20744. [PubMed: 18482979]
- Singh A, Hamedinger D, Hoda JC, Gebhart M, Koschak A, Romanin C, Striessnig J. C-terminal modulator controls Ca<sup>2+</sup>-dependent gating of Ca<sup>v</sup>1.4 L-type Ca<sup>2+</sup> channels. *Nat Neurosci.* 2006; 9:1108–1116. [PubMed: 16921373]
- Sinnegger-Brauns MJ, Huber IG, Koschak A, Wild C, Obermair GJ, Einzinger U, Hoda JC, Sartori SB, Striessnig J. Expression and 1,4-dihydropyridine-binding properties of brain L-type calcium channel isoforms. *Mol Pharmacol.* 2009; 75:407–414. [PubMed: 19029287]
- Sommese L, Zullo A, Schiano C, Mancini FP, Napoli C. Possible muscle repair in the human cardiovascular system. *Stem Cell Rev.* 2017 Apr; 13(2):170–191.
- Southan C, Sharman JL, Benson HE, Faccenda E, Pawson AJ, Alexander SP, Buneman OP, Davenport AP, McGrath JC, Peters JA, Spedding M, et al. The IUPHAR/BPS Guide to PHARMACOLOGY in 2016: towards curated quantitative interactions between 1300 protein targets and 6000 ligands. *Nucleic Acids Res.* 2016; 44:D1054–D1068. [PubMed: 26464438]
- Specht D, Wu SB, Turner P, Dearden P, Koentgen F, Wolfrum U, Maw M, Brandstatter JH, tom Dieck S. Effects of presynaptic mutations on a postsynaptic Cacna1s calcium channel colocalized with mGluR6 at mouse photoreceptor ribbon synapses. *Invest Ophthalmol Vis Sci.* 2009; 50:505–515. [PubMed: 18952919]
- Striessnig J. C-terminal tailoring of L-type calcium channel function. *J Physiol.* 2007; 585:643–644. [PubMed: 18084049]

- Striessnig J, Koschak A. Exploring the function and pharmacotherapeutic potential of voltage-gated Ca<sup>2+</sup> channels with gene knockout models. *Channels (Austin)*. 2008; 2:233–251. [PubMed: 18719397]
- Striessnig J, Ortner NJ, Pinggera A. Pharmacology of L-type calcium channels: novel drugs for old targets? *Curr Mol Pharmacol*. 2015; 8:110–122. [PubMed: 25966690]
- Subramanyam P, Chang DD, Fang K, Xie W, Marks AR, Colecraft HM. Manipulating L-type calcium channels in cardiomyocytes using split-intein protein transsplicing. *Proc Natl Acad Sci U S A*. 2013; 110:15461–15466. [PubMed: 24003157]
- Tan GM, Yu D, Wang J, Soong TW. Alternative splicing at C terminus of Ca(V)<sub>1.4</sub> calcium channel modulates calcium-dependent inactivation, activation potential, and current density. *J Biol Chem*. 2012; 287:832–847. [PubMed: 22069316]
- Tang ZZ, Yarotsky V, Wei L, Sobczak K, Nakamori M, Eichinger K, Moxley RT, Dirksen RT, Thornton CA. Muscle weakness in myotonic dystrophy associated with misregulated splicing and altered gating of Ca(V)<sub>1.1</sub> calcium channel. *Hum Mol Genet*. 2012; 21:1312–1324. [PubMed: 22140091]
- Tuluc P, Molenda N, Schlick B, Obermair GJ, Flucher BE, Jurkat-Rott K. A CaV<sub>1.1</sub> Ca<sup>2+</sup> channel splice variant with high conductance and voltagesensitivity alters EC coupling in developing skeletal muscle. *Biophys J*. 2009; 96:35–44. [PubMed: 19134469]
- Tummala SR, Neinstein A, Fina ME, Dhingra A, Vardi N. Localization of Cacna1s to ON bipolar dendritic tips requires mGluR6-related cascade elements. *Invest Ophthalmol Vis Sci*. 2014; 55:1483–1492. [PubMed: 24519419]
- Uhlen M, Fagerberg L, Hallstrom BM, Lindskog C, Oksvold P, Mardinoglu A, Sivertsson A, Kampf C, Sjostedt E, Asplund A, Olsson I, et al. Proteomics. Tissue-based map of the human proteome. *Science*. 2015; 347:1260419. [PubMed: 25613900]
- Uhlen M, Hallstrom BM, Lindskog C, Mardinoglu A, Ponten F, Nielsen J. Transcriptomics resources of human tissues and organs. *Mol Syst Biol*. 2016; 12:862. [PubMed: 27044256]
- von Gersdorff H, Matthews G. Calcium-dependent inactivation of calcium current in synaptic terminals of retinal bipolar neurons. *J Neurosci*. 1996; 16:115–122. [PubMed: 8613777]
- Wahl-Schott C, Baumann L, Cuny H, Eckert C, Griessmeier K, Biel M. Switching off calcium-dependent inactivation in L-type calcium channels by an autoinhibitory domain. *Proc Natl Acad Sci U S A*. 2006; 103:15657–15662. [PubMed: 17028172]
- Wang D, Papp AC, Binkley PF, Johnson JA, Sadee W. Highly variable mRNA expression and splicing of L-type voltage-dependent calcium channel alpha subunit 1C in human heart tissues. *Pharmacogenet Genomics*. 2006; 16:735–745. [PubMed: 17001293]
- Wang D, Zhong L, Nahid MA, Gao G. The potential of adeno-associated viral vectors for gene delivery to muscle tissue. *Expert Opin Drug Deliv*. 2014; 11:345–364. [PubMed: 24386892]
- Welling A, Ludwig A, Zimmer S, Klugbauer N, Flockerzi V, Hofmann F. Alternatively spliced IS6 segments of the alpha 1C gene determine the tissuespecific dihydropyridine sensitivity of cardiac and vascular smooth muscle L-type Ca<sup>2+</sup> channels. *Circ Res*. 1997; 81:526–532. [PubMed: 9314833]
- Wilhelm M, Schlegl J, Hahne H, Gholami AM, Lieberenz M, Savitski MM, Ziegler E, Butzmann L, Gessulat S, Marx H, Mathieson T, et al. Mass-spectrometry-based draft of the human proteome. *Nature*. 2014; 509:582–587. [PubMed: 24870543]
- Wimmers S, Coepplius L, Rosenthal R, Strauss O. Expression profile of voltage-dependent Ca<sup>2+</sup> channel subunits in the human retinal pigment epithelium. *Graefes Arch Clin Exp Ophthalmol*. 2008; 246:685–692. [PubMed: 18270725]
- Xu W, Lipscombe D. Neuronal Cav1.3α1 L-type channels activate at relatively hyperpolarized membrane potentials and are incompletely inhibited by dihydropyridines. *J Neurosci*. 2001; 21:5944–5951. [PubMed: 11487617]
- Yanik M, Muller B, Song F, Gall J, Wagner F, Wende W, Lorenz B, Stieger K. In vivo genome editing as a potential treatment strategy for inherited retinal dystrophies. *Prog Retin Eye Res*. 2017; 56:1–18. [PubMed: 27623223]
- Yu H, Seo JB, Jung SR, Koh DS, Hille B. Noradrenaline upregulates T-type calcium channels in rat pinealocytes. *J Physiol*. 2015a; 593:887–904. [PubMed: 25504572]

- Yu NY, Hallstrom BM, Fagerberg L, Ponten F, Kawaji H, Carninci P, Forrest AR, Fantom C, Hayashizaki Y, Uhlen M, Daub CO. Complementing tissue characterization by integrating transcriptome profiling from the Human Protein Atlas and from the FANTOM5 consortium. *Nucleic Acids Res.* 2015b; 43:6787–6798. [PubMed: 26117540]
- Zabouri N, Haverkamp S. Calcium channel-dependent molecular maturation of photoreceptor synapses. *PLoS One.* 2013; 8:e63853. [PubMed: 23675510]
- Zamponi GW, Striessnig J, Koschak A, Dolphin AC. The physiology, pathology, and pharmacology of voltage-gated calcium channels and their future therapeutic potential. *Pharmacol Rev.* 2015; 67:821–870. [PubMed: 26362469]
- Zeitc C, Robson AG, Audo I. Congenital stationary night blindness: an analysis and update of genotype-phenotype correlations and pathogenic mechanisms. *Prog Retin Eye Res.* 2015; 45:58–110. [PubMed: 25307992]
- Zhang J, Berra-Romani R, Sinnegger-Brauns MJ, Striessnig J, Blaustein MP, Matteson DR. Role of Cav1.2 L-type Ca<sup>2+</sup> channels in vascular tone: effects of nifedipine and Mg<sup>2+</sup>. *Am J Physiol Heart Circ Physiol.* 2007 Jan; 292(1):H415–H425. [PubMed: 16980345]




**Fig. 1. Localization of all tissue-enriched genes and of the tissue-enriched channel genes.** (A) The largest number of tissue-enriched genes were found in testis (brown) and cerebral cortex (green) samples (tissue-enriched genes are defined by at least five-fold higher expression in one tissue compared to the tissue with the second highest expression). Only tissues with  $\geq 1\%$  of the total number of tissue-enriched genes are labelled. (B) Within each tissue, a different fraction of tissue-enriched genes belong to the ion channel families. While the electrically active tissues (cerebral cortex, skeletal (dark blue) and heart muscle (light blue)) have a higher-than-average percentage of channel genes, in testis only a low fraction of the tissue-enriched genes are channels. (C) Most tissue-enriched channels are found in cerebral cortex (green), testis (brown) and skeletal muscle (dark blue). Genes are listed in order of their expression strength (highest expression (TPM) on top). The numbering refers to Table 1. All data was taken from the Human Protein Atlas (retrieved from [www.proteinatlas.org](http://www.proteinatlas.org) on 11.01.2017).



**Fig. 2. VGCC gene expression in different human tissues and mouse retina.**

(A) Heatmap of VGCC gene expression strengths across tissues. Darker hues indicate higher expression (TgPM = tags per million). Because the HPA dataset contained only a part of the brain (cerebral cortex) and no retinal samples, we chose FANTOM5 data for this comparison which were shown to highly correlate with the HPA (Yu et al., 2015b). Note that Ca<sub>v</sub>3.3 and γ<sub>6</sub> data was not available from FANTOM5 and using another database for the comparison of absolute values was not possible. (B) Expression of VGCCs in different brain regions. Note that “Brain” is a whole-brain sample. (C) Expression of VGCCs in muscle tissue. The inset shows the skeletal muscle calcium channel subunit encoding genes. Note the high abundance of γ<sub>1</sub>. (D) Expression of VGCCs in lymphoid tissues (E) Comparison of expression of VGCC in mouse and human retina. Data were normalized to α<sub>v</sub>δ-4. FANTOM5 data was retrieved through Human Protein Atlas website [www.proteinatlas.org](http://www.proteinatlas.org) on 11.01.2017. VGCC nomenclature according to Catterall et al. (2005).





**Fig. 3. Distribution of VGCC gene expression in human tissues.**

Tissues names are given where the expression exceeds 5% of the total expression in all tissues (except for  $\beta_2$ : retina: 4.9%, heart 4.6%). The colour code is the same throughout: brain tissues (green hues), muscle tissues (blue hues), other tissues are indicated by different colours. Less expressed tissues are combined as ‘others’ (grey). Some of the VGCC genes show highly abundant expression (>50% of total):  $Ca_v1.1$ ,  $Ca_v1.4$ ,  $Ca_v2.1$ ,  $Ca_v2.3$ ,  $\alpha_2\delta-4$ ,  $\gamma_1$ ,  $\gamma_2$ ,  $\gamma_4$ ,  $\gamma_6$ ,  $\gamma_8$ . Others are particularly expressed in several brain areas (together >50% of total):  $Ca_v2.2$ ,  $Ca_v3.1$ ,  $\beta_4$ ,  $\alpha_2\delta-3$ ,  $\gamma_3$ ,  $\gamma_7$ . Some are expressed preferentially (>33% of total):  $\beta_1$ ,  $\alpha_2\delta-2$ ,  $\gamma_5$ . The remaining genes showed widespread expression:  $Ca_v1.2$ ,  $Ca_v1.3$ ,  $Ca_v3.2$ ,  $\beta_2$ ,  $\beta_3$ ,  $\alpha_2\delta-1$ . According to the tissue specificity score (TS) calculation of the Human Protein Atlas (highest expression/2nd highest expression), the tissue-enriched



VGCC are (TS score):  $\text{Ca}_V1.4$  ( $\infty$ ),  $\text{Ca}_V1.1$  (22.8),  $\gamma_1$  (21.0),  $\gamma_6$  (9.0),  $\text{Ca}_V2.1$  (7.6) and  $\alpha_2\delta-4$  (6.2). Note that  $\text{Ca}_V3.3$  and  $\gamma_6$  expression data stems from the GTEx dataset (no FANTOM5 data available on HPA website). FANTOM5 data was retrieved through [www.proteinatlas.org](http://www.proteinatlas.org) on 11.01.2017, GTEx on 12.06.2017).

**Table 1**  
**List of tissue-enriched channel genes in human tissues.**

We pulled out all ion channel genes (according to IUPHAR list of ion channel family members) from the dataset of tissue-enriched genes found in the Human Protein Atlas (HPA). Genes in the HPA tissue-enriched gene list have the highest selectivity of expression (expression in one tissue at least five-fold higher than all other tissues). In total, we found 85 channel genes, numbered from highest tissue specificity to lowest. Black squares indicate the tissue in which the gene is enriched. Abbreviations: a.k.a., also known as; TS, tissue specificity score; TPM, transcripts per million (expression strength in the indicated tissue). Note: TPM = 1 is considered as “expressed”. Voltage-gated Ca<sup>2+</sup> channel subunits and directly associated channels are in **bold**, auxiliary/modulatory/regulatory subunits in *italics*. Human Protein Atlas version 16, Release date: 04.12.2016; Data retrieved on 10.01.2017.

#	Gene	a.k.a	adrenal gland	cerebral cortex	epididymis	esophagus	fallopian tube	heart muscle	kidney	lung	pancreas	parathyroid gland	prostate
1	GRIN1	NMDAR1											
2	<i>CATSPERD</i>	<i>TMEM146</i>											
3	KCNJ1	Kir1.1											
4	KCNJ9	Kir3.3											
5	<i>CACNG7</i>												
6	GJA8	CX50											
7	AQP12A	AQP12											
8	CATSPER4												
9	GABRA1												
10	KCNV2	Kv8.2											
11	GABRA5												

#	Gene	a.k.a	adrenal gland	cerebral cortex	epididymis	esophagus	fallopian tube	heart muscle	kidney	lung	pancreas	parathyroid gland	prostate
12	<i>CATSCSPER4</i>												
13	KCNA7	Kv1.7											
14	CACNA1S	Cav1.1											
15	<i>CACNG8</i>												
16	KCNF1	Kv5.1											
17	CATSPER1	CATSPER											
18	<b>RYR1</b>												
19	<i>CACNG3</i>												
20	CHRNE	ACHRE											
21	CATSPER3	CACRC											
22	<i>CACNG1</i>	<i>CACNLG</i>											
23	GRIN2B	NMDAR2B											
24	KCNU1	KCa5.1											
25	<i>KCNG2</i>												
26	KCNV1	Kv8.1											

#	Gene	a.k.a	adrenal gland	cerebral cortex	epididymis	esophagus	fallopian tube	heart muscle	kidney	lung	pancreas	parathyroid gland	prostate
27	SCN2A	Nav1.2											
28	CHRNA2												
29	CHRNA2	ACHRG											
30	SCN5A	Nav1.5											
31	CHRNA10												
32	CLCN1	CLC1											
33	GABRG1												
34	KCNK12	K2p12.1											
35	GABRB1												
36	CHRNA2												
37	CHRNA2	ACHRD											
38	KCNQ3	Kv7.3											
39	CACNG2	<i>stargazi</i>											
40	GABRA4												
41	KCNH1	Kv10.1											

#	Gene	a.k.a	adrenal gland	cerebral cortex	epididymis	esophagus	fallopian tube	heart muscle	kidney	lung	pancreas	parathyroid gland	prostate
42	KCNK4	K2p4.1											
43	GABRB2												
44	KCNC2	Kv3.2											
45	<i>KCNRG</i>												
46	PKD2L2	TRPP5											
47	SCN3B												
48	<i>c g 2</i>												
49	<i>DCEPG</i>												
50	ASIC1												
51	GABRD												
52	GRIA4	GluA4											
53	GRID2	GluD2											
54	KCNN1	KCa2.1											
55	GJB2	CX26											
56	<b>RYR2</b>												
57	GABRG2												

#	Gene	a.k.a	adrenal gland	cerebral cortex	epididymis	esophagus	fallopian tube	heart muscle	kidney	lung	pancreas	parathyroid gland	prostate
58	ASIC4												
59	GRIA2	GluA2											
60	KCNC1	Kv3.1											
61	TRPM8												
62	CACNA1E	Cav2.3											
63	CHRNA3												
64	GABRA3												
65	HCN1												
66	KCNJ6	Kir3.2											
67	KCNK7	K2p7.1											
68	KCNQ2	Kv7.2											
69	TRPM7												
70	ANO3	TMEM16C											
71	BSND												



#	Gene	a.k.a	adrenal gland	cerebral cortex	epididymis	esophagus	fallopian tube	heart muscle	kidney	lung	pancreas	parathyroid gland	prostate
72	CACNA2D2												
73	CHRN3												
74	GABRA2												
75	GABRR1												
76	GRIA1	GluA1											
77	GRIK3	GluK3											
78	KCNA1	Kv1.1											
79	KCNH7	Kv11.3											
80	SCN3A	Nav1.3											
81	SCN8A	Nav1.6											
82	GJB4	CX30.3											
83	GRIN2A	NMDAR2A											
84	KCNA2	Kv1.2											
85	SCN2B												

#	Gene	a.k.a	adrenal gland	cerebral cortex	epididymis	esophagus	fallopian tube	heart muscle	kidney	lung	pancreas	parathyroid gland	prostate

**Table 2**  
**Expression of VGCC  $\alpha 1$  subunit splice variants.**

Both tissue source and species are indicated. Existing channelopathies or specific interactions with drugs are indicated in bold (disease) or italic (pharmacology). Abbreviation: n.i., not indicated.

<b><math>\alpha 1</math> subunit</b>	<b>Splice variant</b>	<b>Tissue expression</b>	<b>Species</b>	<b>Disease pharmacology</b>	<b>Reference PMID</b>
Cav1.1	exon 29	spleen (CD4 T cells)	mouse		26815481
	exon 29	muscle biopsy	human	<b>Myotonic dystrophy type 1 and type 2</b>	23888875
	exon 29	muscle biopsy	human	<b>Myotonic dystrophy type 1 and type 2</b>	22140091
	exon 29	muscle, primary myotubes, C2C12 line	Human mouse		19134469
Cav1.2	Cav1.2 <sub>e21+22</sub>	heart	mouse	<b>Cardiac hypertrophy, heart failure</b>	27731386
	Cav1.2 <sub>33L</sub>	heart (neonatal, adult)	mouse		25694430
	1896G>A	n.i.	human	<b>Brugada syndrome</b>	24321233
	Cavh2 <sub>73</sub>	cardiovascular tissue	human		23926129
		brain, A7r5 cells	rat		
	Cav1.2 <sub>9*</sub>	aorta cDNA library	human		21998324
	exons 8, 8a	cortex	mouse	<b>Timothy Syndrome</b>	21282112
	exons 8/8a, 9, 33	cardiac and smooth muscle	n.i.	<i>Diltiazem sensitivity</i>	20649567
	exon 9*	cerebral artery myocytes	rabbit		19717733
	exons 9*, 33	embryonic cortex	mouse		19564422
	exons 9*, 33	heart	rat	<b>Myocardial infarction rat model</b>	19263075
	exons 1a, 1, 8a, 8, 9*, 21, 22, 31, 32, 32-6nt, 33	heart	rat	<b>Hypertensive rat model</b>	18070605
	exons 8/8a, 9, 33	cardiac and smooth muscle		<i>Nifedipine sensitivity</i>	17916557
	exon 9*	cartilage, bone, fat, liver, kidney, aorta, bladder, heart tissues, CNS	human		15916803
	exon 9*	smooth muscle	rat		15381693
	exons 21, 22	brain	rat	<i>Isradipine sensitivity</i>	7737988
	a1C-a	heart	rat	<i>Nisoldipine sensitivity</i>	9314833
	a1C-b	aorta			
	exons 8, 8a, 31, 32, 40–43	n.i.	human	<i>Isradipine sensitivity</i>	9607315
	Cav1.3	exons 42S, 42L, 43S, 43L	whole-brain, brain regions, inner ear,	human	
		sinoatrial node	mouse		21998310
C-terminus		testis, heart, pancreas, kidney, lung	hamster		21093409
exon 43S		whole-brain, brain regions, heart, eye	human		21998310
			mouse		
	exons 42, 42S 43S, 41, 44, 48S	whole-brain, brain regions, spinal cord	mouse		21998309

<b><math>\alpha</math>1 subunit</b>	<b>Splice variant</b>	<b>Tissue expression</b>	<b>Species</b>	<b>Disease pharmacology</b>	<b>Reference PMID</b>
	exons 42, 42A	whole brain, brain regions	human		18482979
	Cav1.3 <sub>b</sub>	cortical neuron culture	rat		17287512
	Cav1.3 <sub>IQ</sub>	cochlea inner hair cell	rat		17050708
	Cav1.3 <sub>a</sub>	striatal medium spiny neurons	mouse		15689540
	Cav1.3 <sub>b</sub>				
	Cav1.3 <sub>1a</sub>				
	Cav1.3 <sub>1b</sub>				
	Cav1.2 <sup>-/-</sup> cardiomyocyte	mouse		12900400	
	Cav1.3 <sub>short</sub>	neuroendocrine cells, whole-brain, pituitary GH3 cells	rat		11514547
	exons 31, 32	ventricular myocytes	human	<b>Heart failure</b>	10888251
	IIIS2	cochlea	chicken		9405709
	26 ins I <sub>-II</sub>				
	10 ins IVS2-S3				
	IVS3-IVS4	whole-brain, heart, lung, aorta	rat		9405178
	rCACN4A, 4B, C-terminus, I-II loop, IVS3-S4	RINm5F cell line	rat		7760845
Cav1.4	exon 45,	retina	human		27226626
	exon 47,		monkey		
	exon p45,47		mouse		
	transcript scanning	retina Marathon <sup>®</sup> -Ready cDNA	human		22069316
	CaV1.4a, b	Jurkat T cells, peripheral blood T	human		15899519
Cav2.1	exons 31 –34, 37	lymphocytes, activated T cells			24344901
	C-terminus	neurosecretory cells			21550405
	exons 46, 47	n.i.		<b>Spinocerebellar ataxia type 6</b>	20129625
	splicing mutations	blood samples	human	<b>Episodic ataxia type 2</b>	19242091
	exon 47	n.i.	n.i.	<b>Familial hemiplegic migraine type 1</b>	18390553
	synprint	neuroendocrine cells	rat		18835329
	exon1	whole-brain, cultured cells			17291689
	exon 2 exon 47 ins	Purkinje and granule cells	human patient	<b>Spinocerebellar ataxia type 6</b>	16278278
	exons 37a, 37b	whole-brain, cerebellum	mouse		10321243
			rat		21150296
	exons 28, 34, 47	cerebellum, Purkinje neurons	rat		24369063
	$\alpha$ 1A-a, $\alpha$ 1A-b	brain regions	rat		22836269
Cav2.2	exons 18a, 24a, 31a, 37a, 37b	whole-brain, cerebellum dorsal root ganglion	mouse		19125229
	exons 37a, 37b	n.i.	mouse	<i>Analgesic treatment</i>	
	exon 37a	dorsal root ganglia	mouse		
	exon 18a ins	hippocampus,	rat	<b>Chronic inflammatory pain</b>	

<b><math>\alpha</math>1 subunit</b>	<b>Splice variant</b>	<b>Tissue expression</b>	<b>Species</b>	<b>Disease pharmacology</b>	<b>Reference PMID</b>
	exon 1	spinal cord			
	exons 18a, 24a, 31a, 37a, 37b	dorsal root ganglia	rat		14715140
	exon 1	fetal and adult brain	human		11756491
	exon 2				
	II–III loop	superior cervical and dorsal root ganglia, spinal cord, brain regions	rat		10864934
	synprint II–III loop	ventral mesencephalon	rat		10501220
	IIIS3-S4 IVS3-S4	whole-brain, superior cervical ganglia	rat		9010213
	$\alpha$ 1E <sub>L</sub>	retina, brain regions	rat		9582423
Cav3.1	transcript scanning	whole-brain, thalamus, cortex, cerebellum, hippocampus	mouse	<b>Absence epilepsy (lethargic, tottering)</b>	19480703
	exon 14 exon26				
	exons 25, 26	dorsal thalamus	rat		17707654
	exons 14, 25, 26, 34, 35, 38B	infant brain	human		10548410
Cav3.2	$\pm$ exon 25	heart tissues		<b>Hypertension-associated cardiac hypertrophy</b>	20699644
Cav3.3	exons 33, 34	brain regions, pineal gland	rat		12297319