



Circulating Metabolites and the Development of Type 2 Diabetes in Chinese Adults

Diabetes Care 2022;45:477–480 | <https://doi.org/10.2337/dc21-1415>

Fiona Bragg,^{1,2} Christiana Kartsonaki,^{1,2}
Yu Guo,³ Michael Holmes,^{1,2}
Huaidong Du,^{1,2} Canqing Yu,^{4,5} Pei Pei,⁶
Ling Yang,^{1,2} Donghui Jin,⁷
Yiping Chen,^{1,2} Dan Schmidt,¹
Daniel Avery,¹ Jun Lv,^{4,5} Junshi Chen,⁸
Robert Clarke,^{1,2} Michael Hill,¹
Liming Li,^{4,5} Iona Millwood,^{1,2} and
Zhengming Chen^{1,2}

OBJECTIVE

To assess prospective associations of circulating metabolites with the risk of type 2 diabetes (T2D) among Chinese adults.

RESEARCH DESIGN AND METHODS

A case-cohort study within the 8-year prospective China Kadoorie Biobank comprised 882 participants with incident T2D and 789 subcohort participants. Nuclear magnetic resonance metabolomic profiling quantified 225 metabolites in stored baseline plasma samples. Cox regression related individual metabolites with T2D risk, adjusting for potential confounders and fasting time.

RESULTS

After correction for multiple testing, 163 metabolites were significantly associated with the risk of T2D ($P < 0.05$). There were strong positive associations of VLDL particle size, the ratio of apolipoprotein B to apolipoprotein A-1, branched-chain amino acids, glucose, and triglycerides with T2D, and inverse associations of HDL-cholesterol, HDL particle size, and relative n-3 and saturated fatty acid concentrations.

CONCLUSIONS

In Chinese adults, metabolites across diverse pathways were independently associated with T2D risk, providing valuable etiological insights and potential to improve T2D risk prediction.

Understanding of type 2 diabetes (T2D) etiological pathways is fundamental to disease prevention and development of new therapies. Comprehensive profiling of circulating metabolites in diverse populations is essential for improved understanding of the molecular basis of T2D. With the largest T2D population globally (1), and with marked differences in genetic, environmental, and lifestyle factors when compared with more widely studied Western populations, large-scale investigation of T2D-associated metabolomic profiles in the Chinese population is needed.

RESEARCH DESIGN AND METHODS

The China Kadoorie Biobank (CKB) includes 512,715 adults aged 30–79 years, who were recruited from 10 areas of China in 2004–2008 (2). The baseline survey collected detailed information on medical history, sociodemographic and lifestyle factors, and physical measurements. Venous blood samples were collected (with time since last food recorded) for immediate on-site measurement of random plasma

¹Clinical Trial Service Unit & Epidemiological Studies Unit (CTSU), Nuffield Department of Population Health, University of Oxford, Oxford, U.K.

²Medical Research Council Population Health Research Unit, Nuffield Department of Population Health, University of Oxford, Oxford, U.K.

³Fuwai Hospital Chinese Academy of Medical Sciences, National Center for Cardiovascular Diseases, Beijing, China

⁴Department of Epidemiology & Biostatistics, School of Public Health, Peking University Health Science Center, Beijing, China

⁵Peking University Center for Public Health and Epidemic Preparedness & Response, Beijing, China

⁶Chinese Academy of Medical Sciences, Beijing, China

⁷Hunan Centre for Disease Control and Prevention, Changsha, Hunan, China

⁸China National Center for Food Safety Risk Assessment, Beijing, China

Corresponding author: Zhengming Chen, zhengming.chen@ndph.ox.ac.uk

Received 6 July 2021 and accepted 9 November 2021

F.B. and C.K. are joint first authors. L.L., I.M., and Z.C. are joint senior authors.

© 2022 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at <https://www.diabetesjournals.org/journals/pages/license>.

glucose (RPG) concentrations (SureStep Plus system; LifeScan, Milpitas, CA) and those with a RPG ≥ 140 mg/dL and < 200 mg/dL were invited for fasting glucose testing the following day. Remaining blood samples were retained for long-term storage. Participants were followed up through ongoing linkage to death and disease (including diabetes) registries and to the national health insurance system, providing data on ICD-10-coded morbidity and mortality. Local, national, and international ethics approval was obtained. All participants provided written informed consent.

We selected, using simple random sampling, 900 from 7,721 participants with incident T2D (ICD-10 E11) recorded by 1 January 2017 (median 8.6 [interquartile range 5.2] years follow-up) and without self-reported or newly diagnosed diabetes (3) at baseline, and a subcohort of 905 participants from 31,443 randomly selected genotyped participants. The main analyses included 882 participants with T2D and a subcohort of 789 (including 26 with diabetes) after excluding participants with inadequate plasma samples or with case status mismatch or diabetes at baseline.

Nuclear magnetic resonance (NMR)-based profiling quantified 225 metabolites (directly measured or ratios of these) in baseline plasma samples (4). Cox proportional hazards models, using the Prentice pseudopartial likelihood (5) and correcting for multiple testing (6), were fitted to estimate T2D hazard ratios (HRs) for metabolites, adjusting for sociodemographic and lifestyle factors, fasting time, adiposity, and family history of diabetes. Metabolites were examined as categorical (split at quartiles) and continuous (per 1-SD increment) variables. Additional analyses further adjusted for RPG.

RESULTS

At recruitment, participants with T2D were, on average, older than subcohort participants (55.1 [SD 9.6] vs. 51.9 [10.6] years), had higher adiposity levels (BMI 25.7 [3.6] vs. 23.9 [3.6] kg/m²; waist circumference 85 [10] vs. 80 [10] cm), and were more likely to have a family history of T2D (10% vs. 8%). Use of lipid-lowering medication was rare ($< 1\%$). Overall, 163 of the 225 quantified metabolite measures were independently associated with

T2D ($P < 0.05$), with continuous, largely linear, relationships. Following additional adjustment for RPG, 147 significant associations remained.

There were strong positive associations of apolipoprotein B-to-apolipoprotein A-1 ratio (HR 1.79 per 1-SD higher) and triglyceride concentrations (1.78) with T2D, and a weaker positive association of VLDL-cholesterol concentration (1.27) (Figure 1). Higher HDL-cholesterol concentrations were associated with lower T2D risk (0.48). Each 1-SD increment in mean VLDL and HDL particle sizes was associated with 74% higher and 57% lower risks of T2D, respectively.

Leucine and isoleucine were both strongly positively associated with T2D ($\sim 80\%$ higher risk per 1-SD), as was valine (HR 2.05). There were weaker positive associations of aromatic amino acids (phenylalanine 1.37; tyrosine 1.21), alanine (1.59) and glutamine (1.22). Higher relative concentrations of total n-3 fatty acids were associated with lower T2D risk (0.72), with a particularly strong inverse association (0.46) of docosahexaenoic acid. In contrast, each 1-SD increment in total n-6 fatty acids was associated with 17% higher T2D risk. There were positive and inverse associations, respectively, of monounsaturated (1.30) and saturated (0.62) fatty acids. Lactate, acetoacetate, and 3-hydroxybutyrate were modestly positively associated with T2D (1.49, 1.31, and 1.21, respectively), but there was no association of glycoprotein acetyls.

With the exception of moderately stronger associations for lipoprotein measures among younger participants, no marked differences in the associations of metabolites were observed by age, sex, or urban/rural residence, and associations remained largely unchanged after exclusion of the first 2 years of follow-up.

CONCLUSIONS

This is the largest prospective investigation in China of the associations of diverse circulating metabolites with T2D risk. Among relatively lean Chinese adults, large numbers of metabolites, across varied pathways, were found to be independently associated with T2D. Furthermore, the majority of these associations persisted after adjustment for glycemia.

The lipid and lipoprotein profile associated with higher T2D risk in the current

study, including higher concentrations of VLDL-cholesterol and triglycerides, lower concentrations of HDL-cholesterol, and smaller mean HDL and larger mean VLDL particle sizes, broadly replicates previous findings (7). However, the current associations are moderately more extreme than those previously observed; for example, in a cohort of $\sim 12,000$ young adults in Finland using the same NMR metabolomics platform (7). The focus of the metabolomics platform used on lipid and lipoprotein measures, and the correlation between these, partly explains the large number of significant associations observed. However, metabolites associated with T2D were diverse and included branched-chain amino acids—leucine, isoleucine, and valine—the associations of which were among the strongest observed. This is consistent with previous investigations (8,9) and with genetic association studies suggesting higher branched-chain amino acid concentrations result from insulin resistance and, in turn, cause T2D (10,11). The observed lipid and lipoprotein profile is also consistent with the importance of insulin resistance as a precursor of T2D (12), even in comparatively lean populations such as in CKB.

The higher risks of T2D with higher concentrations of alanine, phenylalanine, and tyrosine reflect well-established associations (7–9). We also observed higher T2D risk among participants with higher glutamine concentrations, in contrast to other study findings showing null (9) or even inverse (7) associations. Further larger studies are needed to clarify this relationship. Although consistent with previous findings in predominantly European population studies (13), the notably lower risks of T2D among individuals with lower relative total n-3 fatty acid and docosahexaenoic acid concentrations represent the first large-scale evidence of such associations in a Chinese population.

Our study has several strengths, including large sample size, prospective design, investigation of metabolite-T2D associations in a comparatively understudied population, and use of a validated NMR platform (14). Moreover, the infrequent use of lipid-lowering medications in CKB reduces treatment-associated biases.

The study also has limitations. Incident T2D comprised diagnosed cases




Figure 1—Associations of circulating metabolites with risk of incident T2D (n = 882), adjusted for fasting time. Adjusted for age (numeric), sex, study area (10 areas), education (6 categories), fasting time (numeric), smoking (ever regular vs. other), alcohol drinking (ever regular vs. other), physical activity (metabolic equivalent of task hours/day, numeric), dietary factors (frequency of consumption of meat, fish, fresh fruit, dairy products; 4 times/week or more vs. other), family history of diabetes (any first-degree relative vs. none), BMI (numeric), and waist circumference (numeric). Squares represent the HR per 1-SD higher metabolite. Horizontal lines represent the corresponding 95% CI. Fatty acid ratios represent ratios of individual to total fatty acids. *P ≤ 0.05, **P ≤ 0.01, ***P ≤ 0.001 after adjustment for multiple testing using Benjamini-Hochberg correction.

only, and resulting misclassification would likely underestimate metabolite-associated risks. A similar effect may have resulted from reliance on single metabolite measures, precluding adjustment for intraindividual variation. Furthermore, the targeted NMR platform used is not comprehensive. However, the strong focus on lipids and lipoproteins is clearly relevant to cardiometabolic disorders such as T2D. Finally, use of nonfasting blood samples would be expected to increase interindividual variation in metabolite concentrations. However, all analyses adjusted for fasting time, and additional adjustment for dietary variables did not appreciably alter risk estimates.

In summary, our study shows that, among relatively lean Chinese adults, a diverse range of circulating metabolites is independently associated with T2D risk. These findings highlight the value of NMR metabolomic profiling for advancing understanding of molecular derangements associated with T2D, with potential translational relevance for enhanced T2D prediction and, ultimately, prevention. Moreover, they may inform selection of metabolites for investigation in future studies to establish causality (e.g., Mendelian randomization

studies), facilitating discovery of novel therapeutic targets.

Acknowledgments. The chief acknowledgment is to the participants, the project staff, and the China National Centre for Disease Control and Prevention and its regional offices for access to death and disease registries. The Chinese National Health Insurance scheme provides electronic linkage to all hospital admission data.

Funding. The baseline survey and the first resurvey were supported by a research grant from the Kadoorie Charitable Foundation in Hong Kong. The long-term continuation of the project is supported by program grants from the UK Wellcome Trust (088158/Z/09/Z, 104085/Z/14/Z, 202922/Z/16/Z, 212946/Z/18/Z), the Chinese Ministry of Science and Technology (2011BAI09B01, 2012–2014), the Chinese National Natural Science Foundation (91843302), and the National Key Research and Development Program of China (2016YFC0900500, 2016YFC0900501, 2016YFC0900504, 2016YFC1303904). The British Heart Foundation (CH/1996001/9454), UK Medical Research Council (MC_UU_00017/1, MC_UU_12026/2, MC_U137686851), and Cancer Research UK (C16077/A29186, C500/A16896) provide core funding to the Oxford Clinical Trial Service Unit and Epidemiological Studies Unit for China Kadoorie Biobank.

Duality of Interest. No potential conflicts of interest relevant to this article were reported.

Author Contributions. F.B. drafted the manuscript. F.B., C.K., Y.G., M.Ho., H.D., C.Y., P.P., L.Y., D.J., Y.C., D.S., D.A., J.L., J.C., R.C., M.Hi., L.L.,

I.M., and Z.C. contributed to critical revision of the manuscript for important intellectual content. F.B., C.K., Y.G., C.Y., P.P., L.Y., D.J., Y.C., J.L., J.C., M.Hi., L.L., and Z.C. contributed to acquisition, analysis, or interpretation of data. F.B., C.K., J.C., L.L., and Z.C. contributed to concept and design. Y.G., C.Y., P.P., D.J., D.S., D.A., J.L., J.C., M.Hi., and L.L., provided administrative, technical, or material support. Y.G., C.Y., J.L., J.C., L.L., I.M., and Z.C. obtained funding. F.B., C.K. and Z.C. are the guarantors of this work and, as such, had full access to all the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.

References

- International Diabetes Federation. Diabetes Atlas. 9th ed. Brussels, International Diabetes Federation, 2019
- Chen Z, Chen J, Collins R, et al.; China Kadoorie Biobank (CKB) Collaborative Group. China Kadoorie Biobank of 0.5 million people: survey methods, baseline characteristics and long-term follow-up. *Int J Epidemiol* 2011;40:1652–1666
- Bragg F, Li L, Smith M, et al.; China Kadoorie Biobank Collaborative Group. Associations of blood glucose and prevalent diabetes with risk of cardiovascular disease in 500 000 adult Chinese: the China Kadoorie Biobank. *Diabet Med* 2014;31:540–551
- Würtz P, Kangas AJ, Soininen P, Lawlor DA, Davey Smith G, Ala-Korpela M. Quantitative serum nuclear magnetic resonance metabolomics in large-scale epidemiology: a primer on -omic technologies. *Am J Epidemiol* 2017;186:1084–1096
- Prentice RL. A case-cohort design for epidemiologic cohort studies and disease prevention trials. *Biometrika* 1986;73:1–11

6. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J R Stat Soc Series B Stat Methodol* 1995;57:289–300
7. Ahola-Olli AV, Mustelin L, Kalimeri M, et al. Circulating metabolites and the risk of type 2 diabetes: a prospective study of 11,896 young adults from four Finnish cohorts. *Diabetologia* 2019;62:2298–2309
8. Guasch-Ferré M, Hruby A, Toledo E, et al. Metabolomics in prediabetes and diabetes: a systematic review and meta-analysis. *Diabetes Care* 2016;39:833–846
9. Qiu G, Zheng Y, Wang H, et al. Plasma metabolomics identified novel metabolites associated with risk of type 2 diabetes in two prospective cohorts of Chinese adults. *Int J Epidemiol* 2016;45:1507–1516
10. Lotta LA, Scott RA, Sharp SJ, et al. Genetic predisposition to an impaired metabolism of the branched-chain amino acids and risk of type 2 diabetes: a Mendelian randomisation analysis. *PLoS Med* 2016;13:e1002179
11. Mahendran Y, Jonsson A, Have CT, et al. Genetic evidence of a causal effect of insulin resistance on branched-chain amino acid levels. *Diabetologia* 2017;60:873–878
12. Garvey WT, Kwon S, Zheng D, et al. Effects of insulin resistance and type 2 diabetes on lipoprotein subclass particle size and concentration determined by nuclear magnetic resonance. *Diabetes* 2003;52:453–462
13. Qian F, Ardisson Korat AV, Imamura F, et al.; Fatty Acids and Outcomes Research Consortium (FORCE). n-3 Fatty acid biomarkers and incident type 2 diabetes: an individual participant-level pooling project of 20 prospective cohort studies. *Diabetes Care* 2021;44:1133–1142
14. Holmes MV, Millwood IY, Kartsonaki C, et al.; China Kadoorie Biobank Collaborative Group. Lipids, lipoproteins, and metabolites and risk of myocardial infarction and stroke. *J Am Coll Cardiol* 2018;71:620–632