



Correction to: Public Health Approaches to Type 2 Diabetes Prevention: the US National Diabetes Prevention Program and Beyond

Stephanie M. Gruss¹ · Kunthea Nhim¹ · Edward Gregg² · Miriam Bell¹ · Elizabeth Luman¹ · Ann Albright¹

Published online: 27 June 2020

© Springer Science+Business Media, LLC, part of Springer Nature 2020

Correction to: Current Diabetes Reports (2019) 19: 78
<https://doi.org/10.1007/s11892-019-1200-z>

The original version of this review article unfortunately contained a mistake in the first paragraph of “Measures and Methods” section. Reference citation “[40]” should be changed to “[37]”.

The sentence should read: “Average weight loss and physical activity minutes per week were calculated among participants who attended ≥ 3 sessions in the first 6 months and whose time from first session attended to last session attended was ≥ 9 months. CDC considers this threshold to be the minimum dose to begin seeing lifestyle and weight change that can impact type2 diabetes [37].”

Publisher’s Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

The online version of the original article can be found at <https://doi.org/10.1007/s11892-019-1200-z>

✉ Stephanie M. Gruss
inf6@cdc.gov

Kunthea Nhim
knhim@cdc.gov

Edward Gregg
e.gregg@imperial.ac.uk

Miriam Bell
omw5@cdc.gov

Elizabeth Luman
ecl7@cdc.gov

Ann Albright
aga6@cdc.gov

¹ Division of Diabetes Translation, Centers for Disease Control and Prevention, 4770 Buford Hwy., Mailstop F75, Atlanta, GA 30341, USA

² Department of Epidemiology and Biostatistics, School of Public Health, Imperial College London, London, UK