



OPEN

Mindfulness and Behavior Change

Zev Schuman-Olivier, MD, Marcelo Trombka, MD, David A. Lovas, MD, Judson A. Brewer, MD, PhD, David R. Vago, PhD, Richa Gawande, PhD, Julie P. Dunne, PhD, RN, PMHNP-BC, Sara W. Lazar, PhD, Eric B. Loucks, PhD, and Carl Fulwiler, MD, PhD

Abstract: Initiating and maintaining behavior change is key to the prevention and treatment of most preventable chronic medical and psychiatric illnesses. The cultivation of mindfulness, involving acceptance and nonjudgment of present-moment experience, often results in transformative health behavior change. Neural systems involved in motivation and learning have an important role to play. A theoretical model of mindfulness that integrates these mechanisms with the cognitive, emotional, and self-related processes commonly described, while applying an integrated model to health behavior change, is needed. This integrative review (1) defines mindfulness and describes the mindfulness-based intervention movement, (2) synthesizes the neuroscience of mindfulness and integrates motivation and learning mechanisms within a mindful self-regulation model for understanding the complex effects of mindfulness on behavior change, and (3) synthesizes current clinical research evaluating the effects of mindfulness-based interventions targeting health behaviors relevant to psychiatric care. The review provides insight into the limitations of current research and proposes potential mechanisms to be tested in future research and targeted in clinical practice to enhance the impact of mindfulness on behavior change.

Keywords: health behavior, mental disorders, mindfulness, motivation, neuroscience, self-management, self-regulation

INTRODUCTION

Health behavior refers to any behavior that affects physical or mental health or quality of life.¹ Unhealthy behaviors such as tobacco smoking, alcohol and substance use, excessive eating, and

From the Department of Psychiatry, Harvard Medical School (Drs. Schuman-Olivier, Trombka, Gawande, Lazar, and Fulwiler); Cambridge Health Alliance, Center for Mindfulness and Compassion, Cambridge, MA (Drs. Schuman-Olivier, Trombka, Gawande, Dunne, and Fulwiler); Department of Psychiatry, Hospital de Clinicas de Porto Alegre, Porto Alegre, Rio Grande do Sul, Brazil (Dr. Trombka); Department of Psychiatry, School of Medicine, Dalhousie University (Dr. Lovas); Brown Mindfulness Center (Drs. Brewer and Loucks), School of Public Health (Dr. Loucks), and Department of Psychiatry (Dr. Brewer), Brown University; Osher Center for Integrative Medicine, Department of Physical Medicine and Rehabilitation, Vanderbilt University Medical Center, Nashville, TN (Dr. Vago); Boston College School of Nursing (Dr. Dunne); Department of Psychiatry, Massachusetts General Hospital, Boston, MA (Dr. Lazar); Department of Psychiatry, School of Medicine, University of Massachusetts (Dr. Fulwiler).

Supported by National Institutes of Health (NIH) Helping to End Addiction Long-Term (HEAL) Initiative, award nos. R21AT010125, R33AT010125 administered by the National Center for Complementary and Integrative Health (Dr. Schuman-Olivier); Brazil Ministry of Education CAPES scholarship no. 88887.363065/2019-00 (Dr. Trombka); NIH Science of Behavior Change Common Fund Program, award nos. UH2AT009145 and UH3AT009145 administered by the National Center for Complementary and Integrative Health (Dr. Loucks).

Original manuscript received 1 April 2020; revised manuscript received 22 June 2020, accepted for publication subject to revision 21 July 2020; revised manuscript received 10 August 2020.

Correspondence: Zev Schuman-Olivier, MD, Director, Center for Mindfulness and Compassion, 1035 Cambridge St., Suite 21A, Cambridge, MA 02141. Email: zschuman@cha.harvard.edu

Copyright © 2020 The Author(s). Published by Wolters Kluwer Health, Inc. on behalf of the President and Fellows of Harvard College. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

DOI: 10.1097/HRP.0000000000000277

nonadherence to medical regimens account for a substantial proportion of global disease morbidity and mortality, and for 40%–50% of the risk for early death in the United States.^{2–6} All-cause mortality is inversely associated with healthy lifestyle behaviors, and changes in unhealthy behaviors can lead to improved physical and mental health outcomes.⁷ Despite widespread awareness, it remains exceptionally difficult to initiate and maintain health behavior change.²

Health behavior theories are used to understand and predict health behaviors,^{8–16} and several behavior-change interventions have become popular.^{17,18} A growing body of evidence suggests that mindfulness-based interventions (MBIs) are effective in reducing harmful health behaviors,¹⁹ catalyzing chronic disease self-management and health behavior change,²⁰ and improving physical and mental health outcomes.^{21–23} This article provides a narrative review with three main aims: (1) to define mindfulness and describe the evolving context and content for the MBI movement, (2) to describe an updated “mindful self-regulation” model that integrates motivation and learning mechanisms essential for behavior change and is grounded in emerging neuroscientific evidence, and (3) to synthesize current clinical research on MBIs targeting health behaviors relevant to psychiatry.

MINDFULNESS AND MBIs

Mindfulness is commonly defined as the awareness that arises when paying attention to the present moment nonjudgmentally.²⁴ In 1881, the English scholar Rhys Davids translated the word *mindfulness* from the Pali word *sati* found in Buddhist texts, which meant “memory, recollection, calling-to-mind, being-aware-of, certain specified facts”²⁵ but which has also been described as “lucid awareness”²⁶ or “bare attention.”²⁷

The terms *mindfulness* and *meditation* are increasingly conflated. General integrative practices (e.g., visualization, yoga) are often described as “mindfulness” in public discourse, diluting the word’s meaning. Importantly, not all meditation is mindfulness and not all mindfulness is meditation. Meditation is a practice that self-regulates the body and mind by engaging a specific attentional set.²⁸ In *mindfulness meditation* (MM), the practice is to pay attention to present-moment experience with an orientation of curiosity, openness, acceptance, nonreactivity, and nonjudgment.^{29,30}

Several core types of MM are taught in MBIs, along an attentional continuum that ranges from *focused attention*, which involves directing and sustaining attention on an object, disengaging from distractors (e.g., mind wandering), and returning attention to the object, to *open monitoring*, which has no explicit focus but cultivates metacognitive monitoring with a nonreactive awareness of the flow of cognition, emotions, and sensations.³¹ MBIs also use other evocative meditations to cultivate specific emotions (e.g., compassion, loving-kindness) that support MM practice.^{32,33} MM may produce relaxation in the body (i.e., the *relaxation response*, a voluntary, wakeful, hypometabolic state of parasympathetic dominance),³⁴ but relaxation is not necessarily the objective. Rather, MM is an active and intentional practice of cultivating awareness of present-moment experience that may include strong emotions, difficult thoughts, or unpleasant sensations.³⁵ MM cultivates both awareness and equanimity, an even-minded mental state or dispositional tendency toward all experiences/objects, regardless of their affective valence (pleasant/unpleasant/neutral).³³ Several valuable models describe potential psychological and neurobiological mechanisms through which MM could exert its salutary effects.^{30,36–45}

Persistent tensions have emerged throughout the MBI movement between (1) adaptation/openness to change versus fidelity/quality/safety, (2) drives for innovation/novelty versus ownership/tradition, and (3) non-attachment to labels versus valuing lineage with respect for teachers. To avoid confusion and to remain inclusive and neutral to these natural tensions, we describe a brief history of the movement—providing context for, and meaning to, various definitions.

Jon Kabat-Zinn began developing the first MBI, *mindfulness-based stress reduction* (MBSR), in 1977, integrating Buddhist insight (Pali: Vipassana) meditation,^{46,47} other contemplative practices (e.g., Zen, yoga),^{47,48} and modern psychological theories about stress and stress coping for application in health care.^{49–52} This work spawned a new secular, mainstream pedagogy for MM training, with a package of practices (i.e., body scan, mindful breathing, sitting meditation, informal mindfulness) bundled in an eight-week program that was designed to cultivate mindful attention in daily life, thereby helping patients to cope with stress, pain, and other chronic conditions. The first adaptation of MBSR, *mindfulness-based cognitive therapy* for depression,^{53–56} constituted, along with MBSR, the first generation of MBIs.⁵⁷ More than a decade later, these approaches would come to be called *mindfulness-*

based programs (MBPs), which have in common the “systematic and sustained training in formal and informal MM practices (for *both* teacher and participants).”^{57,58} Crane and colleagues⁵⁷ distinguish MBPs from general MBIs, which were adapted from MBSR and mindfulness-based cognitive therapy; although these approaches also focus on cultivating MM practice, they have not always followed the same teacher-training process or other MBP standards. Together MBIs and MBPs focus on MM practice and differ from *mindfulness-informed interventions*,⁵⁹ or “third-wave” interventions⁶⁰ (e.g., *dialectical behavior therapy* [DBT], *acceptance and commitment therapy* [ACT]),^{61,62} which feature mindfulness as a component within a larger suite of techniques and mechanisms of change without an explicit focus on MM practice.^{57,59} Research publications on MBIs began increasing exponentially in the 2000s, in parallel with the proliferation of MBIs for specific conditions, and by 2020, MBIs have emerged for many chronic mental and physical illnesses.^{21,63,64} Academic mindfulness centers also emerged as dissemination nodes for MBPs,⁶⁵ encouraging standards for formal teacher training and teacher competency assessments to ensure program integrity.⁶⁶

This review will focus primarily on the areas with the strongest evidence relating to mindfulness and behavior change. We will describe and expand existing models of *mindful self-regulation* based on neurobiological mechanisms of mindfulness, motivation, and learning.^{38,67,68} Then we will review data from meta-analyses and well-designed randomized, controlled trials (RCTs) of MBPs affecting health behaviors. Given the more heterogeneous nature of related interventions (e.g., MBIs, DBT, ACT, *mindful self-compassion*, *integrative mind-body therapy*),^{61–63,69,70} we will limit references to these interventions except when MBP research is scarce. It should be noted, however, that DBT and ACT have made a substantial contribution to the understanding of behavior change, with strong meta-analytic evidence for a range of psychiatric disorders—which has contributed to a broader reconceptualization of the mechanisms of behavior change brought about by third-wave cognitive and behavioral therapies.^{60,71}

MINDFUL SELF-REGULATION

Self-regulation is the ability to adaptively regulate one’s attention, emotions, cognition, and behavior to respond effectively to internal as well as environmental demands.^{72–74} Self-regulation impairment is linked to poorer outcomes in school/academics^{75,76} and also to poorer physical^{77,78} and mental health.^{79–84} Self-regulation theory proposes several critical processes for people to initiate and maintain behavior change and to prevent self-regulatory failure,⁷³ including the capacity for standard setting, motivation to meet standards, self-monitoring, and willpower.^{73,74,85,86} We will build on the seminal neuroscience-based frameworks proposed by Hölzel and colleagues³⁸ and Vago and colleagues,⁴⁰ which outlined several relevant synergistic neurocognitive systems that underlie the effects of mindfulness on self-regulation, including *attentional control*, *cognitive control*, *emotion regulation*, and *self-related processes*. Importantly, we

will describe how these self-regulation mechanisms interact with basic neural mechanisms for *motivation and learning*, which are well studied in the addiction and behavioral medicine literature. We will describe how each of these systems may play a role in facilitating the way that MM affects a person's capacity for behavior change (Figure 1). The reviews aim to fill the gap in previous models of mindfulness and self-regulation by describing how, as MM practice develops, it may begin to enhance motivation for change by affecting processes of reward, associative and extinction learning, and habit formation, thereby shifting the balance from unhealthy to healthier behavioral repertoires.

Attention and Cognitive Control

Cognitive control is a fundamental capacity of human cognition that regulates access to specific goal-relevant information to facilitate the performance of specific behaviors.^{87–91} *Executive functions* are cognitive processes necessary for initiating and maintaining behavior change, and also for selecting and monitoring behavior toward attainment of goals; *mental set shifting, information updating and monitoring*, and *inhibition* are of particular importance.⁹² The subcomponents of cognitive control and executive function include *attentional control/regulation*,^{93,94} *cognitive inhibition, inhibitory control, working memory, decision making*, and *cognitive flexibility*.⁹⁵ Attentional control can be further subdivided into (1) *volitional orienting* of attention to task-relevant information (e.g., breath, body sensations), (2) *alerting* to and successfully sustaining attention and vigilance for unexpected stimuli, and (3) *conflict monitoring* among task-relevant internal thoughts, feelings, and responses.⁹⁶ These attentional processes contribute to recognizing both the consequences of ongoing unhealthy behavior and the affective precursors to avoidance or appetitive behaviors, contributing to adaptive behavior change.

MM may contribute to self-management of clinical symptoms, which are affected by unhealthy behaviors or maladaptive cognitive coping strategies, through explicit cognitive control processes. Two highly cited models^{38,40} suggest that mindfulness involves attention regulation (improvement in orienting, alerting/stability, bias, meta-awareness/monitoring)^{97–99} and inhibition/switching (decentering, inhibitory control, flexible engagement/disengagement).^{100,101} While individual studies support these models,¹⁰² meta-analyses of the effects of mindfulness on behavioral-cognitive assays demonstrate substantial heterogeneity,^{103,104} possibly related to diversity in mindfulness techniques, small sample sizes, nonspecific effects of common factors (e.g., group effects, empathic therapists), and multiple strategies for performing behavioral tasks. Moreover, populations with unique cognitive-dysfunction profiles (e.g., substance use, bipolar disorder, depression, age-related cognitive decline, or mild cognitive impairment)^{105–110} may experience improvement of disorder-specific dysfunction, whereas healthy participants may be sensitive to ceiling effects. In addition, the mindfulness effect may simply be making the brain more efficient, requiring less effort for a similar result.¹⁰⁶

Functional neuroimaging research demonstrates that these interrelated cognitive control processes generally engage overlapping higher-order brain networks, including a central executive network, salience network, dorsal and ventral attention networks, and default mode network (DMN).¹¹¹ These networks are anchored through a *frontoparietal network* that acts as a flexible hub interconnecting these networks, depending on the contextual and functional demands of cognition.^{112–116} This superordinate frontoparietal network, also referred to as the *cognitive control network*, includes the dorsolateral (dl) prefrontal cortex (PFC), posterior parietal cortex, anterior insula cortex, anterior cingulate cortex




Figure 1. Mindfulness influences on self-regulation and behavior change.

(ACC), and medial PFC.^{89–91,113,114} In the context of this flexible-hub framework, growing mechanistic evidence suggests that different types of MM engage associated frontoparietal network brain areas both structurally and functionally, differentially influencing psychological and cognitive outcomes.^{117,118} For example, studies show differences in brain activation and connectivity during focused-attention versus open-monitoring meditation,¹¹⁹ with focused attention engaging attentional control, whereas open-monitoring training engages labeling and emotional nonreactivity.¹²⁰ A key aspect of cognitive control is the capacity to monitor for conflicts in information processing and to transmit these signals to other executive functions.⁸⁷ This conflict-monitoring capacity has been associated with the ACC and seems to be enhanced with alterations in the ACC in the first few weeks of MM.^{121–123} Multiple fMRI studies suggest functional alterations between the ACC and insula in the context of monitoring for aversive stimuli during MM.^{123–126} Executive function is negatively affected by hippocampal dysfunction,¹²⁷ whereas MBSR has been associated with increases in hippocampal volume,¹²⁸ which can affect working memory performance.¹²⁹ Inhibitory control is commonly associated with the dorsal ACC (dACC) or inferior frontal gyrus, which may be affected by culture or by developing mental habits of behavioral consistency.¹³⁰ In meta-analyses, the dACC is often activated in studies of both focused-attention and open-monitoring meditation.¹¹⁷

Given that executive functions are critical for initiating and maintaining health behaviors, MBPs that enhance these processes may affect behavior-change capacity. By enhancing working memory and expanding attentional resources,¹³¹ MM may increase conscious awareness of behaviorally relevant external or internal stimuli (e.g., appetitive cues, stressor or craving-related body sensations), providing more time for higher-order cognitive functions, such as inhibitory control and other executive functions. These changes may help overcome prepotent motor responses, reduce the cognitive effort required for decision making about appetitive health behaviors, and diminish susceptibility to automatic behaviors in response to negative affect.^{106,132}

Emotion Regulation

Emotions influence our perception, thinking, and behavior, and are made up of subjective, physiological, and behavioral components.¹³³ *Emotion regulation* is the ability to modulate emotional experiences, enabling adaptive engagement with internal and external experience, which is essential for well-being and social adaptation.^{134,135} A variety of emotion-regulation strategies have been identified, including modifying the situation, altering attention toward it, modifying thoughts about it, and modulating the response to emotional experiences.¹³⁶ Effective strategies decrease the subjective experience of negative emotion and its physiological correlates, and are associated with long-term efficacy in daily life.¹³⁷ The best-studied strategy is cognitive *reappraisal*, the conscious reinterpretation

of a situation in a way that alters its meaning and lessens the emotional impact. Reappraisal may be one of the mechanisms through which MM improves emotion regulation and ameliorates anxiety and depression.^{138,139} According to one model with empirical support, mindfulness facilitates reappraisal by interrupting automatic reactions, allowing for conscious reflection.^{38,140} With ongoing practice, some theorize that MM practitioners transition from using reappraisal strategies to cultivating states of *nonappraisal* and equanimity, though longitudinal research is needed.^{33,38,141} The available evidence also provides some support for *exposure* and *extinction* as key mechanisms for mindfulness-related emotion regulation. Rather than engaging in experiential avoidance^{142–144} or thought suppression,¹⁴⁵ MBPs promote the ability to focus awareness on a difficult experience when it arises, creating a condition of exposure to negative emotional reactions.^{38,146} This change in focus is often aided by the capacity to shift experiential perspective—to step outside one’s immediate subjective experience to a more objective, non-identified awareness of one’s experience, which is called *decentering*,¹⁰⁰ *metacognitive monitoring*, or *meta-awareness*.^{147,148} With practice, repeated acceptance and awareness of emotional and physiological responses lead to reduction and even extinction of emotional reactivity.^{149–151} Therefore, whereas meditation often causes state changes with enhanced autonomic parasympathetic activity, exposure through MM practice may create greater trait *implicit* control over parasympathetic tone, possibly initially through ACC control,¹⁵² and then possibly, with ongoing practice, through ventromedial (vm) PFC modulation of the amygdala.^{148,153} Finally, as people learn to be mindful of emotions, the capacity for *emotion differentiation* may be enhanced, which is associated with an enhanced ability for emotion regulation.¹⁵⁴

Functional neuroimaging research implicates both distinct and overlapping networks involved in different emotion-regulation strategies. *Explicit* cognitive strategies like reappraisal are associated with activation of the frontoparietal executive network involved in selective attention, working memory, and response inhibition, and with deactivation of the amygdala.¹⁵⁵ Neural mechanisms of mindfulness-based emotion regulation appear to differ, depending on the subjects’ experience level. Naive subjects given brief training in mindfulness induction for a laboratory study appear to employ top-down control by PFC regions in fashion similar to that of subjects instructed to use cognitive reappraisal.¹⁵⁶ By contrast, for experienced meditators, downregulation of emotional reactivity when in a mindful state appears to involve deactivation of the DMN without deactivation of the amygdala.¹⁵⁷ MBSR completers show decreased amygdala reactivity accompanied by increased amygdala–vmPFC functional coupling, consistent with the finding that symptom improvement in anxiety patients correlates with increased amygdala–vmPFC coupling after MBSR.^{153,158} Long-term meditators, in contrast to eight-week MBSR completers, employ implicit emotion-regulation systems involving the insula and vmPFC to modulate emotional reactivity.¹⁵³ These findings

suggest that, over time, long-term mindfulness practice promotes greater awareness and acceptance of emotional experience rather than the suppression of emotions by top-down inhibitory control.^{148,153,156,157}

Healthy emotion regulation is important for behavior change to modulate emotions in alignment with one's goals.^{159,160} Without it, feelings of stress, anxiety, and depression may thwart one's intention to engage in health behaviors related, for example, to diet, exercise, or smoking—which promote positive health outcomes.^{161–164} Similarly, for patients with chronic illnesses facing a variety of physical and psychological challenges, emotion regulation is critical for maintaining optimal cognitive functioning and emotional balance.¹⁶⁵ Mindfulness may also improve the chances of successful behavior change by increasing awareness of how emotions influence decisions and behaviors.¹³⁷ A recent study of primary care patients with chronic illness reported that an eight-week MBI facilitated emotion regulation and catalyzed health behavior change.²⁰ Our recent meta-analysis found that subjects receiving MBPs, compared to controls, significantly improved on emotion-regulation measures,¹⁶⁶ suggesting that mindfulness may engage emotion-regulation processes that can improve behavioral outcomes.^{67,137,167,168}

Self-Related Processes

Cognitive science describes a continuum of *self-related processes* that aligns closely with large-scale neural networks discovered through functional connectivity studies—networks that are affected by MM.

At one end of the self-related process continuum, the *experiential self* is the self-as-subject, the embodied self that is supported by phenomenological body awareness unextended in time—that is, the sense of “I am in this moment.”^{169–171} Interoception is the sensory experience of homeostatic afferents related to the body's physiologic state, producing the feeling of present-moment self in the body.¹⁷² Broader definitions of *interoceptive awareness* include the tendency to listen to and trust body sensations as occurring within a dynamic relationship with appraisal and *interoceptive regulation* processes.^{173,174} Mindfulness training enhances interoceptive awareness and function.²⁰ Most MM starts with the body and seems to develop greater interoceptive accuracy.^{175,176} Interoceptive dysfunction may lead to difficulties predicting body states and to allostatic dyscontrol.¹⁷⁷ Interoceptive regulation may be enhanced through MM by increasing the capacity for perceptual-inference strategies (i.e., bringing the desired interoceptive state to what is sensed) instead of focusing only on active-inference strategies (i.e., trying to change the experience to the desired state).¹⁷⁸ Numerous recent studies suggest an impact of MM on improving interoceptive measures.^{20,179} Meta-analysis and mediation studies are needed to examine the full impact of mindfulness on interoceptive awareness and its effect on behavior change.

Mindfulness neuroscience describes a phenomenological experiential-self network,¹⁷¹ involving activation of the anterior

insula,⁴⁰ which is altered after MM training.¹⁸⁰ The insula provides an efficient means for processing large-scale interoceptive information in real time.¹⁸¹ Interoceptive activity and functional connectivity in the insula is disrupted in depression,¹⁸² which may be associated with a lack of emotional feeling in the body¹⁸³ and may be associated with alexithymia.^{184,185} Functional connectivity studies of large-scale networks propose a key role of the insula in a salience network, which is posited to shift attention between internal (e.g., internal body sensations) and external stimuli (e.g., external perception) with dorsal attentional and ventral emotional components.¹¹¹ Among MM practitioners, the experiential-self network may use moment-to-moment bodily experience in context to establish saliency for attention and regulation of cognitive resources.

At the other end of the self-related process continuum, a *narrative self*¹⁷⁰ emerges during development, starting in late latency,¹⁸⁶ then consolidating during adolescence¹⁸⁷ and emerging adulthood,¹⁸⁸ represented by autobiographical *self-monitoring* and social-cognitive and evaluative functions—in short, the self-as-object, “story of me as a person,” temporally extended into the past or future.¹⁷⁰ These cognitive science concepts align with cognitive clinical models of self-related processes, which have an emphasis on self-discrepancy¹⁸⁹ between the actual and idealized narrative self through positive (self-esteem and self-worth¹⁹⁰) and negative self-evaluation^{191,192} (self-devaluation, self-criticism, and self-critical rumination).^{74,193,194} Emerging therapeutic approaches also include self-schema^{195–197} or multiple self-like parts^{198,199} to explain the clinical complexity of narrative self-related processes. The narrative self overlaps with many functions of the DMN, the most prominent large-scale brain network underlying self-related processes.²⁰⁰ The DMN engages medial cortical regions involved in self-monitoring, self-judgment, self-referential mental activity, episodic-memory retrieval, autobiographical memory, self-related social-cognitive processes, and value-based decision making.^{201,202} The DMN connects during adolescent identity formation,²⁰³ increasingly potentiating self-criticism and negative self-evaluative rumination. Among adults, high levels of DMN connectivity are associated with depression.²⁰⁴ Importantly, DMN activity and connectivity are reduced among experienced mindfulness meditators.²⁰⁵ In a recent meta-analysis, the posterior cingulate cortex, which is a core DMN node involved in self-related processing,²⁰⁶ is reliably deactivated during both focused-attention and open-monitoring MM.¹¹⁸

Rumination is a response to distress that involves repetitively and passively focusing on symptoms of distress and the possible causes and consequences of these symptoms.^{207,208} Rumination predicts depression,²⁰⁹ bulimia, substance use, self-injurious behaviors, impaired problem solving,²⁰⁸ and aggressive behaviors.²¹⁰ One meta-analysis demonstrated that reductions in rumination partially mediated improved psychological functioning.¹⁵¹ Another recent meta-analysis found MBPs reduced negative self-related rumination, with a strong effect among six studies with inactive control groups, and potential

for benefit among two studies with active controls.¹⁶⁶ A meta-analysis of brain-imaging studies reports a strong association between DMN core regions, especially the dorsomedial (dm) PFC, and rumination.²¹¹ Reducing negative self-related rumination through MBPs may improve emotion regulation, reducing depressive symptoms and self-criticism. In this way, MM might prevent the detrimental repetition of negative global self-attributions that come when a person has a brief slip of unhealthy behavior after a period of sustained abstinence—which often leads persons to dive into a full-blown relapse, a process that Marlatt¹⁸ named the *abstinence violation effect*.

Other self-related processes on the continuum are the sense of agency²¹² related to the experience of one's actions and decision making as an individual agentic self, and *self-efficacy*, which is one's belief in one's ability to succeed in specific situations or accomplish a task.²¹³ A frontoparietal cognitive control network may provide flexible control in self-related processing²¹⁴—involving coupling between other frontal networks, depending on internally or externally directed task demands.^{91,112,114,215} The overarching cognitive control network is therefore recruited to support control of belief and goal-directed strategies, flexibility creating a sense of *self-agency* that contributes to resiliency.^{216,217} The dlPFC, which is a core node of the central executive network, is often activated during MM; MM enhances connectivity between the dlPFC and other executive-function regions.^{218,219} As MM practice continues over time, the central executive network and other frontal networks may regulate the DMN in control of cognition.²²⁰

Two small studies have demonstrated increases in task-specific *self-efficacy*, though the increases were not statistically significant compared with controls.^{221,222} An uncontrolled, prospective cohort study reported increased chronic illness self-efficacy after eight weeks of MBSR but less self-efficacy at one year.²²³ Two larger RCTs of an eight-week MBP with self-efficacy as a secondary outcome reported within-group improvements in self-efficacy ($d = 0.3 - 0.43$), though this result was not significantly different from a low-dose mindfulness comparator.^{20,224} The ability to practice mindfulness may lead to a greater sense of one's ability to make health behavior change; however, it is also possible that initial difficulty attaining expected levels of mindfulness practice may cause an opposite, deleterious effect on global self-efficacy and that the feedback from this failure may affect the capacity to succeed in self-regulation itself.^{225,226} Closer analysis of self-efficacy and self-agentic beliefs are needed to understand their role in mindful self-regulation models.

Finally, the warmth with which one relates to one's self is a self-related process that affects self-regulation and behavior. *Self-compassion* involves responding with a warm, kind, and understanding orientation toward oneself, as one would to a close friend, when we suffer, fail, or feel inadequate.²²⁷ Self-compassion may be a mechanism through which MM supports behavior change^{228,229} and engagement in health-promoting behaviors^{230,231} in the context of diabetes,²³² disordered eating,²³³ exercise-related goals in the face of

setbacks,²³⁴ and sexual HIV risk behaviors.²³⁵ Self-compassion has been proposed to have a direct effect on self-regulation²³⁶ in part by neurobiologically mimicking a supportive compassionate other, activating the soothing-affiliation system.²³⁷ A meta-analysis demonstrated a strong effect of MBPs for increasing self-compassion in four studies with inactive controls ($d = 0.73$), though the result was insignificant versus active comparators.¹⁶⁶ A rigorous RCT comparing an eight-week MBP versus a 60-minute mindfulness comparator reported significant effects of mindfulness dose on self-compassion ($d = 0.41$).²⁰ MM lowered levels of self-criticism, which was associated with reduced activation in dmPFC nodes of the DMN,²³⁸ initially through dlPFC-mediated self-compassionate reappraisal strategies.²³⁹ Kindness toward self may clarify that one is worthy of being cared about, thereby enhancing motivation for self-care. MBPs focused primarily on attentional aspects of mindfulness in initial manuals/manuscripts, but a beneficial implicit “hidden curriculum” emerged for cultivating warmth and self-kindness.²⁴⁰ More recently, Kabat-Zinn²⁴¹ clarified that “mindfulness” is the same as “heartfulness,” acknowledging the importance of love, kindness, and compassion. In this way, mindfulness practice may be described as “warmly being with present moment experience.” “Warm” mindfulness infused with self-compassion may be a more effective rapid emotion-regulation strategy than a response of “cool” mindful acceptance alone.²⁴² Interventions and programs that focus explicitly on cultivating inner compassion, which includes and extends beyond self-kindness, may help facilitate behavior change, particularly for individuals who are prone to excess self-criticism, shame, or unworthiness.²⁴³

Motivation and Learning

In self-regulation theories, motivation relies on an anticipatory proactive system (i.e., setting standards) and a reactive negative feedback system (i.e., evaluative judgment) for reducing the discrepancy between goals/standards and behavior.⁷²⁻⁷⁴ Early self-regulation theories, based on control theory⁷² or homeostasis (feedback error correction),²⁴⁴ emphasize the role of evaluative judgment to generate motivation for change. Importantly, control of motivation can be *stimulus driven* in response to an external stimulus or *control driven*, which is motivated by, and directed toward, a specific outcome (e.g., intentions, goal setting).²⁴⁵ The motivation for unhealthy behavior is often driven by external and interoceptive stimuli²⁴⁶ through *associative learning*—for example, operant conditioning with positive or negative reinforcement. While MBPs may initially engage negative feedback systems utilizing discrepancy through evaluative judgment or predictive error to generate motivation for practice (e.g., “I hate being depressed, so I am motivated to practice mindfulness”), evaluative judgment is deemphasized during MBPs in favor of developing a focus on acceptance, intention setting, and attention toward experiential monitoring rather than self-evaluation.²⁴⁷⁻²⁴⁹

Evaluative judgment and feedback occur at least partially through the phasic activity of midbrain dopamine neurons in the ventral tegmental area that encode a *prediction error* used to guide associative learning throughout the frontal cortex and the basal ganglia.²⁵⁰ Activity in ventral tegmental area dopaminergic neurons that project to the nucleus accumbens (NAc) may signal that a person's estimate of the value of current and future events is in error, and also indicate the error magnitude.²⁵¹ This dopaminergic signaling process for *reward-based learning* begins with the acquisition of a reward after a behavior and leads to operant conditioning (e.g., incentive to repeat a pleasant, rewarding state). This gives way to wanting through incentive salience for cue-elicited behaviors,²⁵² which causes automaticity and habit formation with the behavior eventually becoming epigenetically encoded in the dorsal striatum,^{253–255} making it harder to unlearn.²⁵⁶

Aversive learning happens through several mechanisms. *Avoidance action learning*, also referred to as negative reinforcement (e.g., action to avoid an unpleasant negative affect or aversive withdrawal state), depends on dopaminergic signaling in the NAc, though possibly through a different mechanism than approach learning.^{246,257} Pavlovian fear conditioning²⁵⁸ is due to signaling in the basolateral amygdala, which gets transmitted to the central amygdala, leading to the startle/freeze response to a stimulus.^{246,259–261} In cases where successful avoidance action takes place, however, a signal from the vmPFC blocks signal transmission to the central amygdala; instead of freezing, there is a predictive error signal in the NAc.²⁴⁶ Therefore, with aversive experiential learning, the freeze response and fear memory get conditioned first, but then successful avoidance action gets conditioned through engagement from the NAc and vmPFC, priming people to initiate avoidance action when a cue related to the previously aversive stimuli is experienced.²⁴⁶ Thus, the conditioned action urges that are at the heart of reward and experiential avoidance seem to be encoded in the prediction error in the NAc modulated by the vmPFC, which is the cortical region involved in subjective self-related valuation and self-regulatory goals.^{262–264}

Fear memories either get continually reconsolidated after cue exposure or move toward extinction if the feared conditioned effects do not arise through a NMDA receptor-mediated process in the basolateral amygdala.^{265,266} *Extinction learning* is the process of inhibiting conditioned fear responses and developing new learning that competes with prior conditioning.^{266,267} It also involves prediction error-related vmPFC activity,²⁶⁸ which may explain why approach and avoidance conditioning often interact when trying to reverse conditioning.^{269,270} Therefore, both negative and positive reinforcement approaches may rely on this dopaminergic error-prediction system and modulation from the vmPFC self-related valuation system, which together are likely candidates for the negative feedback evaluative judgment system in self-regulation theory.

New theories based on recent neuroscientific developments also propose that prediction-error processes may be

widely distributed throughout the brain's architecture (including the insula) for interoceptive regulation and allostasis, with a key role in emotion and behavioral regulation.^{271,272} These neural models of allostasis propose that efficient self-regulation requires *anticipating* needs and preparing to satisfy them *before* they arise through a process of predictive error prevention.²⁵⁶

As described above, external rewards and punishments can lead to associative learning through Pavlovian and operant conditioning, and lead to habits that persist with ongoing positive and negative reinforcement. Exposure and associative learning can also be employed both for removing unhealthy habitual behavior and for developing and reinforcing healthy habitual behavior.⁴¹ Mindfulness practice, in particular, is itself a unique healthy behavior that may have the capacity to accelerate the process of extinguishing and replacing unhealthy behavioral repertoires.^{273,274} Traditionally, MM has been effectively taught in retreat settings lacking most behavior-activating cues,⁴⁶ allowing interoceptive and cognitive exposure with reduced risk for actual harmful behaviors.²⁷⁵ In community MBPs, however, participants live in high-risk, cue-laden contexts,²⁷⁶ which can be more challenging. Despite this increased challenge, the social context of the group helps people spend time away from unhealthy behavior-activating cues and in a social environment that values collective learning of mindfulness.²⁷⁷ As community MBP practice continues, autonomic stability increases,¹⁵² allowing for a “window of tolerance,”^{278,279} within which exposure, response prevention, reconsolidation, associative learning, and extinction learning processes^{150,280} may begin to unwind the habit learning²⁷³ and fear conditioning²⁸¹ that were maintaining unhealthy habits.

At around four weeks of practice, nonclinical MBI participants generally begin to automatically pay attention to the present moment.²⁸² MM may be utilizing the same neural circuitry and associative-learning mechanisms to establish the habit of mindfulness, which can lead to a process of *therapeutic staged neuroplasticity*.²⁸³ Mindfulness practice associates readily available cues (e.g., the breath, present-moment sensations in the body, the physical experience of stress) with the internal mindful action of warmly being with present-moment experience with constant, accepting awareness of the experience's changing nature—a process traditionally called *Sampajanna* in the Pali language, meaning “clear comprehension of impermanence.”²⁶ This then sets in motion an automatic, internal behavioral repertoire that uses exposure and extinction processes to uproot deeply programmed, unhealthy conditioned behavioral patterns, while developing insight and ability to distinguish which behavioral patterns are healthier and do no harm.^{197,274,284} By combining this mindful acceptance with deepening interoceptive awareness, MM then begins to sidestep standard interoceptive feedback processes that drive automatic behaviors, thereby shifting interoceptive regulation toward use of the perceptual-inference instead of active-inference strategies. This change provides

an efficient pathway for discrepancy resolution that reduces predictive errors and refines allostatic regulation.^{173,285} The stimulus-driven approach to learning and motivation dissipates as motivation for mindfulness becomes fully internalized and intrinsic motivation for healthy behaviors awakens.

Self-determination theory suggests motivation is derived from competence, relatedness, and autonomy.²⁸⁶ The motivation for initiating and maintaining behavior change exists on a continuum from amotivation to extrinsic (external pressure), introjected (internal pressure/guilt/shame), identified (feels useful/important), internalized (aligns with deeply held values), and most optimally intrinsic (interesting/enjoyable). Initially, mindfulness and paying attention may come from introjected or identified motivation (e.g., feeling I should meditate or that meditation is important for me). As practice continues and one witnesses the moment-to-moment consequences of unhealthy behaviors and identifies what is deeply valued, motivation for healthy behaviors become increasingly *internalized*, and people become self-motivated.⁷⁴ As behavior aligns with internal standards, one develops more proactive systems (e.g., intention setting), reducing the burden from reliance on overreactive, judgment-driven negative feedback systems.²⁸⁷ Competence and autonomy are enhanced, and intrinsic motivations for wellness are progressively uncovered. *Curiosity*, which is a key aspect of the mindful state,¹⁰¹ may also modulate midbrain dopaminergic systems.^{288,289} Mindfully taking interest in health behaviors and its effects may generate greater levels of *intrinsic motivation*,^{287,290} enhancing overall motivational vigor and cognitive control through tonic dopamine release from the ventral tegmental area to the frontal central executive network regions.^{291–293} As mindfulness practice proceeds, mindful savoring of healthy, pleasant experiences may restructure reward processes and dopaminergic tone to reinstate value to natural rewards.²⁹⁴ Negative emotions have specific action urges, while positive emotions tend to broaden and build a person's repertoire of thought-action patterns.²⁹⁵ Therefore, emotions are often experienced as intrinsic motivation, and emotion differentiation²⁹⁶ and reappraisal through mindfulness may help reduce emotionally driven unhealthy behaviors and build internalized and intrinsic motivation for healthy behaviors. Finally, kindness and compassion increase *relatedness* with others and potentially make people more sensitive to affiliative motivational systems, through which oxytocin may drive dopaminergic learning and reward in the ventral tegmental area and NAc.^{297,298}

Mindful Self-Regulation: Synergy and Integration

Mindful self-regulation integrates attentional/cognitive control, emotion regulation, and self-related processes in synergy with mechanisms for motivation and learning as the practice of MM deepens. Mindful self-regulation starts with attentional control and curiosity about present-moment experience, leading to the development of interoceptive awareness and alternatives to self-critical rumination. Reappraisal of

mental content, decentering, and acceptance downregulates autonomic reactivity, allowing for exposure to aversive internal stimuli and ultimately developing equanimity. Yet, for people with high levels of limbic dysregulation (e.g., unresolved trauma, marginalized or disempowered status) or baseline attentional/cognitive impairments, this standard, “cool” MM pathway offers some challenges. In this context, a “warm” MM pathway that starts with establishing safety through a “window of tolerance,” while cultivating self-compassion and inner warmth, may be more effective.²⁹⁹ By reducing self-criticism, autonomic reactivity, and internal conflict, the focus shifts from unhealthy patterns toward self-care.³⁰⁰ By slowly reducing the limbic load on cognitive control resources, attentional training becomes more accessible and effective. The most effective MM teachers often engage both pathways in tandem, helping practitioners more quickly come into smooth regulation and potentially reducing adverse experiences;³⁰¹ yet, this key perspective has not been fully appreciated in previously proposed models of mindful self-regulation. As one begins to pay attention, consistent curiosity and kind awareness allow greater goal-driven control based on values, increased levels of internalized motivation, greater access to intrinsic motivation, and less reliance on stimulus-driven conditioning and evaluative negative feedback systems, eventually unwinding associative learning related to harmful behaviors. Behavior begins to emerge that puts positive cues in the environment, potentially activating healthy behaviors and supporting the behavior of mindful noticing itself, which leads to a positive feedback loop in favor of healthy behavior change. In this way, over time MM may cause an evolution toward greater efficiency in the human self-regulation system within a socio-environmental context, leading to greater feelings of interconnection and relatedness, and ultimately even supporting experiences of self-transcendence.⁴⁰

MBIs' EFFECTS ON PSYCHIATRICALLY RELEVANT BEHAVIOR CHANGE

Substance Use Disorders

Evidence supporting the impact of MBIs for alcohol (AUD) and substance use (SUD) disorders has been growing,^{302–310} supporting a few key mechanisms underlying the effect.^{41,273,311,312} A meta-analysis of 42 studies in 2017 reported a small effect of MBIs on substance misuse (standardized mean difference = -0.33) and a moderate effect on substance-related craving (standardized mean difference = -0.68).²³ A three-arm RCT enrolled abstinent intensive-outpatient and inpatient residential treatment program completers with SUD ($n = 286$), comparing mindfulness-based relapse prevention versus relapse prevention versus treatment as usual on substance use outcomes over 12 months. This study demonstrated mindfulness-based relapse prevention had fewer drug use and heavy-drinking days at 12 months compared with relapse prevention alone.³¹³ Mindfulness-based relapse prevention is the most commonly studied MBI for SUD, while other

MBIs and mindfulness-informed interventions are also being studied for various substances and stages of treatment (e.g., mindfulness-oriented recovery enhancement).^{274,314–318} While the strength of evidence for mindfulness effects on certain SUDs during specific stages of treatment is strong (e.g., AUD),³¹⁹ evidence for mindfulness and other SUDs is still emerging (e.g., methamphetamine, opioid use disorder).^{320–324}

MM studies for AUD/SUD support the general self-regulation mechanisms above while also focusing on disorder-specific deficits (e.g., reward processing, executive function, cue reactivity). For instance, mindfulness reduces impulsivity among people with opioid use disorder receiving methadone.^{197,325} MM may strengthen top-down cognitive control and repair executive-function deficits³²⁶ among people with SUD (e.g., working memory, inhibitory control)^{106,326} by increasing ACC/PFC activity and restoring frontostriatal connectivity.³¹¹ Mindfulness practice appears to partially decrease substance use through reductions in craving.^{327,328} Craving reduction may be from mindfulness strategies for decentering, acceptance, and attentional control that reduce intrusive cognitive elaboration (i.e., dwelling in positive recollections or attempts to suppress them) and that prevent cascades of emotional and physiologic reactivity.³²⁹ Mindfulness enhances emotion regulation, decoupling depression/negative affect from craving.^{327,328,330} MM encourages acceptance, reducing thought suppression,³²⁹ which may reduce craving since the suppression of thoughts/urges often amplifies craving.³³¹ Mindfulness dampens limbic reactivity and enhances vagal tone,³³² reducing the amplitude and elaboration of physiologic responses to stress and drug cues associated with craving.^{333,334} Mindfulness also seems to facilitate physiological recovery (e.g., high-frequency heart rate variability) with faster attentional disengagement after substance-related cues^{105,335} and stress.³³⁶ Evidence is also emerging that MM and mindful savoring practice may lead to restructuring reward processing,^{273,337} potentially remediating opioid use disorder–related hedonic dysregulation.³³⁸ By exposing practitioners to imaginal and interoceptive substance-related cues during practice and providing rewarding experiences (e.g., feelings of tranquility, curiosity about the novelty of present-moment experience), MM may allow extinction learning and a reversal of operant conditioning, unwinding years of habit formation.²⁷³ Despite the meta-analyses and efficacy and mechanistic studies supporting a role for mindfulness in AUD/SUD treatment, critical issues remain: establishing an evidence base to support real-world implementation in clinical settings; adapting interventions for various levels of care, patient readiness, and stages of treatment; determining the necessary dose of training at each level; and investigating issues of scalability for diverse populations.^{305–307}

Eating Behavior/Diet

The efficacy of MBIs on food-related behavior and weight loss has been documented across multiple reviews and meta-analyses.^{339–346} Among RCTs, MBIs have large effects on the reduction of binge eating, emotional eating, and eating when not feeling physically hungry,^{339,340,344} and moderate

effects on the improvement of eating attitudes.³⁴⁴ Small to moderate effects were noted for weight loss.^{339,344,347} MBIs that include both formal and informal practices (e.g., mindful eating)³³⁹ or that are combined with cognitive-behavioral therapy appear particularly beneficial.^{341,343}

MBIs may promote enhanced self-regulation and interoceptive regulation through increased ability to notice automatic thoughts or affective cues, accurate appraisal of internal stimuli, or increased awareness of craving, hunger, fullness, and other factors influencing eating patterns.^{38,348–350} For example, mindful eating promotes present-moment enjoyment of food while reducing both overall caloric intake and craving-related eating.^{351,352} Across studies of individuals who are obese or overweight, MBIs reduced levels of depression and anxiety with medium effect.³⁴⁴ This improvement in psychological health may disrupt cycles of reactive or emotional eating, foster self-compassion, and improve self-concept.^{348,353,354} Overall, MBIs promote healthier behaviors around food and enhance psychological well-being, resulting in weight loss or weight maintenance. More research is needed to examine the efficacy of MBIs on sustained weight loss.

In line with broader research on eating-related behaviors and obesity, MBIs, including mindfulness-based eating awareness training (MB-EAT),^{354,355} appear useful for the treatment of binge eating disorder via similar mechanisms of action.^{340,356,357} In a meta-analysis, studies of persons with binge eating disorder demonstrated large effects for the reduction of eating pathology, emotional eating, negative affect, and body dissatisfaction, along with a small effect for weight loss.³⁵⁷ Additional research is needed to compare the efficacy of MBIs to traditional approaches (e.g., cognitive-behavioral therapy) for this pathology.³⁵⁸ Little is known about MBIs in bulimia nervosa and anorexia nervosa.^{356,358–362} Initial meta-analytical reports are favorable, showing reduced eating pathology, emotional eating, and body dissatisfaction with medium effect, as well as weight gain in some underweight participants.³⁵⁷ Across eating disorders, including binge eating disorder, compassion-focused techniques and mindfulness-informed interventions (e.g., DBT) may be useful.^{358,361} Given the unique clinical characteristics of restrictive eating disorders, mindful eating may be distressing for some individuals, especially with acute symptoms. While MBIs show promise, more rigorous research is needed to understand their utility for eating disorder treatment.

Tobacco Smoking

The past decade has seen an evolution of treatments and studies on MBIs for tobacco smoking cessation, with mixed evidence. Two meta-analyses, both of which included four smoking-specific RCTs, found that MBIs were superior to other evidence-based treatments. One reported a moderate effect ($d = 0.42$),²² and the other a relative risk of abstinence of 1.88 (1.04–3.40).³⁶³ A 2017 meta-analysis with ten RCTs found that mindfulness-informed interventions (including

brief and mobile interventions) did not differ from comparators.³⁶⁴ In a now somewhat dated 2015 systematic review, only 13 of 198 articles on mindfulness and smoking were controlled empirical studies, with the majority being pilot or feasibility trials.³⁶⁵ The first RCT in 2011 compared mindfulness training to cognitive-behavioral therapy (American Lung Association's Freedom from Smoking), in which they found a five-times greater cessation rate with mindfulness,³⁶⁶ which a secondary analysis demonstrated was moderated by baseline levels of nonjudgment, with less self-judgment of inner experience supporting greater cessation with mindfulness training.³⁶⁷ Another analysis showed mindfulness training decoupled key links in the smoking reinforcement-learning pathway (craving and smoking).³⁶⁸ As treatment has moved digital, app-based paradigms have been tested, reporting similar decoupling mechanisms.^{369,370} Recent work demonstrated that app-based mindfulness training serves to target DMN brain networks involved in "getting caught up" in craving^{206,371} in a dose-dependent manner, with the degree of brain activity reduction in the posterior cingulate cortex predicting smoking reductions, especially among women.³⁷² More work is needed to replicate and extend findings.

Chronic Disease Self-Management

Self-management of chronic illness^{373,374} for common conditions³⁷⁵ is a priority for improving health care.^{376,377} As chronic physical illness is highly comorbid with mental illness,³⁷⁸ helping patients develop skills to self-manage chronic illness is essential in psychiatry. Chronic illness is often comorbid with anxiety, depression, trauma, and stress.^{379–381} Meta-analysis shows small effects for MBIs on enhancing quality of life and reducing anxiety, depression, and stress during chronic illness.²¹ Reducing mental health symptoms through MM may support greater chronic disease self-management. For instance, an eight-week MBI—integrated as health care treatment for primary care patients with comorbid mental and physical chronic illnesses—demonstrated increased rates of health-related action-plan initiation compared with participants randomized to a low-dose mindfulness comparator (OR = 2.91).²⁰ A pilot eight-week MBSR group adapted for hypertension demonstrated significant changes in modifiable determinants of blood pressure—physical activity, diet, and alcohol consumption—for those who were nonadherent to American Heart Association guidelines at baseline, and changes were maintained at one year.³⁸² Patients with chronic obstructive pulmonary disease who participated in an eight-week group program followed by ten monthly sessions reported increased health behavior changes and improvement in coping with illness.³⁸³ Qualitative findings from 41 individuals with long-term conditions participating in MBSR emphasized that "starting where I am" facilitated changes related to coping with and managing their illnesses.³⁸⁴ Mindfulness training helped lower blood glucose levels³⁸⁵ and hemoglobin A1c³⁸⁶ among those with type 2 diabetes but not among emerging adults with type 1 diabetes,³⁸⁷ suggesting further research is required.³⁸² Longer-term

continuation groups beyond eight weeks and the addition of kindness and compassion elements may be an important adaptation for helping those living with chronic illness.³⁸⁸

Violence, Suicide/Self-Injury, and Other High-Risk Behaviors

Impulsive violence and aggression are critical high-risk behaviors commonly monitored in psychiatry. A systematic review on the effect of mindfulness on aggression and violence included 22 adult studies (4 RCTs), concluding that MBIs and mindfulness-informed interventions (not including DBT) had a significant impact, with effect sizes from 0.21 to 0.87.³⁸⁹ Mindfulness skills may decrease impulsive thought and aggressive behavior through improved awareness of internal experience, reduction of experiential avoidance, and attentional shifting.^{390–392} An RCT with 58 patients with borderline personality disorder found decreased impulsive behavior after DBT versus treatment as usual.³⁹³ An RCT with 101 adults with recent partner aggression comparing ACT versus support group reported ACT was more efficacious in reducing physical and psychological aggression.³⁹⁴ An RCT with 56 healthy adults comparing MBSR versus waitlist reported MBSR decreased anger suppression and aggressive anger expression.³⁹⁵ In conclusion, empirical evidence suggests MBIs and mindfulness-informed interventions may reduce impulsive and aggressive behaviors. More methodologically rigorous research and meta-analyses are needed to confirm this hypothesis.

Dispositional mindfulness is negatively associated with suicidal ideation (SI)^{396,397} and nonsuicidal self-injury.^{398,399} DBT has the most evidence for decreasing suicidal and self-injurious behavior.^{400,401} In their systematic review, Chesin and colleagues⁴⁰² found six studies of MBIs and suicidal behavior.⁴⁰³ Four studies reported inferential statistics on the effect of MBIs on SI, with three reporting significant reductions. The study that did not find a significant reduction had relatively low baseline rates of SI, suggesting a possible basement effect.⁴⁰⁴ Since that review, other studies have also shown salubrious effects of MBIs on SI,^{405,406} with MBIs appearing to uncouple the association between depressive symptoms and SI.⁴⁰⁷ MBIs remediate some cognitive factors and self-related processes shown to be risk factors for SI—for example, self-critical rumination, hopelessness, and cognitive reactivity to suicide⁴⁰⁸—but they do not mediate the improvement in SI.⁴⁰⁸ Thus, while MBIs may help reduce SI, more needs to be studied regarding potential mechanisms. The strongest evidence for nonsuicidal self-injury has been for DBT, which is effective in reducing nonsuicidal self-injury urges and behaviors in RCTs and meta-analyses, with particularly consistent and strong evidence among adolescents.^{401,409,410} Studies of the effects of MBIs on nonsuicidal self-injury have been proposed, but results have not yet been published to our knowledge. One RCT, however, and several open-label trials for ACT show promise.^{411,412}

Dispositional mindfulness is positively correlated with inhibitory control and negatively correlated with impulsivity and with compulsive and high-risk sexual behavior.^{391,413–420} One RCT with 28 adults comparing 12 individual ACT sessions to a waitlist control condition to evaluate sexual impulsivity related to pornography showed a positive effect of the intervention on compulsive sexual behaviors.⁴²¹ Cross-sectional data suggest that mindfulness training could reduce hypersexual behavior.^{420,422} Spiritual self-schema therapy, a psychotherapy that includes MM as its primary component, showed increased motivation to reduce HIV risk behavior (i.e., sex without a condom) in an RCT with 72 drug users,²⁷⁴ and completers of the intervention had decreased impulsiveness on the Barratt Impulsiveness Scale.¹⁹⁷

DISCUSSION

Limitations of Current MBP Research

Many of the populations who may benefit from health-related behavior change—those with a disproportionate burden and comorbidity of chronic illness, substance use, and high-risk behaviors—are underrepresented in current research on mindfulness and compassion-based programs. These populations include trauma survivors,^{423,424} ethno-racial minorities^{425,426} (particularly those who identify as African American, Latinx, or indigenous), sexual or gender minorities,^{166,427,428} refugee/immigrant populations,⁴²⁹ and those at the intersection of these marginalized groups.⁴³⁰ While MBPs may be beneficial for addressing difficulties experienced by these populations (e.g., mindfulness-based relapse prevention may be more effective when delivered within groups composed primarily of ethnic/racial minorities),^{431,432} the current research base disproportionately represents well-educated, Caucasian, economically advantaged individuals.^{166,425} The paucity of ethno-racial diversity in both MBP researchers⁴³³ and MBP research participants^{166,425} mirrors the lack of diversity in psychological and biomedical research.⁴³⁴

While the level of MM practice engagement is often predictive of outcomes,⁴³⁵ trait mindfulness can be measured in the general non-meditating population, where it contributes to self-regulated behavior and positive affect.⁴³⁶ Moreover, even though eight weeks of MBP group training with a silent day of practice (23 hours in total) was more effective than a one-hour introduction to MM in catalyzing health behavior change, evidence supporting what dose intensity is required for behavior change is not yet definitive.^{20,437} Many doorways to discovering mindfulness likely exist, and those that are culturally inclusive of marginalized populations are more likely to be experienced as open doors, supporting greater access.⁴³⁸ While MBPs are well researched, with reliable dose-related effects within certain demographic groups,^{435,439} MBPs as currently designed are unlikely to be the exclusive optimal format for all. Innovations like rolling admissions in mindfulness-based relapse prevention represent an initial step toward accessibility.⁴⁴⁰ Overall, demands for the consistency

and integrity of MBPs need to be balanced with cultural accessibility in clinical dissemination research.

The future accessibility, relevance, and ethical dissemination of existing MBPs will depend on adequate training of diverse MBP teachers/facilitators and on research agendas that prioritize overlapping areas of cultural humility, cultural accessibility, and trauma-sensitive delivery of MBPs, particularly for those affected by historical and intergenerational traumas.⁴⁴¹ Given the potential relationship between interoception and social power,⁴⁴² internalized oppression,⁴⁴³ and chronic illness,^{173,444,445} future research is needed on interventions that cultivate embodiment and interoception,²⁸⁵ including movement,⁴⁴⁶ drumming,^{447,448} music,^{449–452} and dance⁴⁵³—which may be more culturally accessible and specifically helpful for people with intergenerational trauma,⁴⁵⁴ and serve to foster social connectedness for marginalized populations with chronic illnesses.⁴⁵³

Future Directions: Adverse Effects, Trauma-Informed Training, Dose Effects

Adverse effects have been largely undocumented in the MBP research field. A systematic review of mindfulness-based stress-reduction or cognitive-therapy RCTs reported that only 15.6% of 231 studies included any statement about adverse events.⁴⁵⁵ Yet, potential adverse effects may include panic, physical pain, reexperiencing of traumatic memories, and dissociative symptoms.³⁰¹ More needs to be understood about how program, participant, and teacher/leader factors contribute to adverse events.⁴⁵⁶ Britton⁴⁵⁷ and Grant and Schwartz⁴⁵⁸ propose a research agenda that takes into account that unwanted effects may emerge when aspects of mindfulness, such as decentering, focus on body sensations, and present-moment focus, are overemphasized. As the forms of adverse effects and reactions to symptoms vary based on individual and context, teachers and leaders should train in how to best monitor for, and respond to, adverse effects in MBP research.^{459,460}

Program design and training must be sensitive to potential adverse effects experienced by trauma survivors. A majority of individuals will experience a traumatic life event, and a subset of these will develop posttraumatic stress disorder (PTSD).⁴⁶¹ During MBPs, trauma survivors may be more likely to experience certain adverse effects such as traumatic flashbacks³⁰¹ and dissociation.⁴⁶² Trauma-informed training assumes that everyone might have a trauma history and focuses on offering choice, de-shaming adverse effects, supporting individual agency, and responding skillfully to trauma-related effects when they arise.^{424,460} MBPs have the potential to substantially reduce PTSD symptomology and to support neurobiological changes in networks implicated in PTSD psychopathology.^{111,463–469} Future research is needed on interventions that incorporate self-compassion, as it may mitigate and transform the shame-based nature of PTSD.⁴⁷⁰ When patients with PTSD are immersed within a safe, validating, healing community, such as well-designed trauma-informed

MBP groups featuring self-compassion, they are likely to successfully initiate meaningful behavior changes.²⁰

Finally, future clinical trials should include validated assays in clinical settings measuring the impact of dose effects on the self-regulation components described herein, and the extent to which each component mediates health behavior change.⁴⁷¹ More dismantling and dose-response studies could help clarify the most active components and their individual contributions.^{120,472} Care should be taken with this investigative approach, however, since it carries risks of oversimplifying MBPs to just a collection of techniques in a toolbox.⁴⁷³ When mindfulness programs are offered solely from this instrumental dimension of a skill to be acquired,²⁴¹ then the synergistic, holistic, and seemingly paradoxical approach to cultivating mindfulness, which many patients experience as engaging and deeply healing in MBPs, may be unwittingly sacrificed.

CONCLUSION

A growing evidence base supports the benefits of mindfulness for behavior change. A mindful self-regulation model based on an integration of neuroscientific findings describes the complex and synergistic effects of attention/cognitive control, emotion regulation, and self-related processes, as well as motivation and learning mechanisms that may provide a unique pathway toward sustainable behavior change. While evidence supports the impact of mindfulness on behavior change for key health behaviors related to psychiatric practice, more high-quality research is needed, especially with objective measures, larger samples, replication studies, active controls, and formal monitoring of adverse events.⁴⁷⁴ The field will also benefit from additional research on the impact of integrating compassion practices and from a focus on trauma-sensitive adaptations for diverse populations.

Declaration of interest: Dr. Brewer is the founder of MindSciences, the company that developed the mindfulness app for smoking cessation described in this article. He owns stock in, and serves as a noncompensated scientist for, the company and has previously served on the board of directors. This financial interest has been disclosed to and is being managed by Brown University, in accordance with its Conflict of Interest and Conflict of Commitment policies.

We would like to thank Willoughby Britton, Marcelo Demarzo, Gaelle Desbordes, Rebecca Ferrer, Elizabeth Hoge, Jean King, Jared Lindahl, Ethan Moitra, Neusa Rocha, and Jason Samlin for their contributions to this project.

REFERENCES

1. Tombor I, Michie S. Methods of health behavior change. In: Braddick O, ed. *Oxford research encyclopedias: psychology*. New York: Oxford University Press, 2017.
2. Nielsen L, Riddle M, King JW, et al. The NIH Science of Behavior Change Program: transforming the science through a focus on mechanisms of change. *Behav Res Ther* 2018;101:3–11.
3. Mokdad AH, Marks JS, Stroup DF, Gerberding JL. Actual causes of death in the United States, 2000. *JAMA* 2004;291:1238–45.
4. Danaei G, Ding EL, Mozaffarian D, et al. The preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS Med* 2009;6:e1000058.
5. Crimmins EM, Preston SH, Cohen B. International differences in mortality at older ages: dimensions and sources. Washington, DC: National Academies, 2011.
6. Schroeder SA. We can do better—improving the health of the American people. *N Engl J Med* 2007;357:1221–8.
7. Loef M, Walach H. The combined effects of healthy lifestyle behaviors on all cause mortality: a systematic review and meta-analysis. *Prev Med* 2012;55:163–70.
8. Noar SM, Zimmerman RS. Health behavior theory and cumulative knowledge regarding health behaviors: are we moving in the right direction? *Health Educ Res* 2005;20:275–90.
9. Michie H, West S, Campbell R, Brown R, Gainforth J, eds. *The behaviour change wheel: a guide to designing interventions*. London: Silverback, 2014.
10. Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: toward an integrative model of change. *J Consult Clin Psychol* 1983;51:390–5.
11. Rosenstock IM. Why people use health services. *Milbank Mem Fund Q* 1966;44:94–124.
12. Bandura A. Social cognitive theory: an agentic perspective. *Annu Rev Psychol* 2001;52:1–26.
13. Painter JE, Borba CPC, Hynes M, Mays D, Glanz K. The use of theory in health behavior research from 2000 to 2005: a systematic review. *Ann Behav Med* 2008;35:358–62.
14. Glanz K, Rimer B. *Theory at a glance: a guide for health promotion practice*. San Francisco, CA: U.S. Department of Health and Human Services, Public Health Service, National Institutes of Health, National Cancer Institute, 2005.
15. Ajzen I. Theories of cognitive self-regulation: the theory of planned behavior. *Organ Behav Hum Decis Process* 1991;50:179–211.
16. Perkins HW. Social norms and the prevention of alcohol misuse in collegiate contexts. *J Stud Alcohol Suppl* 2002;(14):164–72.
17. Miller WR, Rollnick S. *Motivational interviewing: helping people change*. 3rd ed. New York: Guilford, 2013.
18. Marlatt GA, Donovan DM. *Relapse prevention: maintenance strategies in the treatment of addictive behaviors*. 2nd ed. New York: Guilford, 2005.
19. Salmoirago-blotcher E, Hunsinger M, Morgan L, Fischer D, Carmody J. Mindfulness-based stress reduction and change in health-related behaviors. 2013;18:243–7.
20. Gawande R, To MN, Pine E, et al. Mindfulness training enhances self-regulation and facilitates health behavior change for primary care patients: a randomized controlled trial. *J Gen Intern Med* 2019;34:293–302.
21. Carlson LE. Mindfulness-based interventions for physical conditions: a narrative review evaluating levels of evidence. *ISRN Psychiatry* 2012;2012:651583.
22. Goldberg SB, Tucker RP, Greene PA, et al. Mindfulness-based interventions for psychiatric disorders: a systematic review and meta-analysis. *Clin Psychol Rev* 2018;59:52–60.
23. Li W, Howard MO, Garland EL, McGovern P, Lazar M. Mindfulness treatment for substance misuse: a systematic review and meta-analysis. *J Subst Abuse Treat* 2017;75:62–96.
24. Kabat-Zinn J. *Full catastrophe living: using the wisdom of your body and mind to face stress, pain and illness*. New York: Delacorte, 1990.
25. Sun J. Mindfulness in context: a historical discourse analysis. *Contemp Buddhism* 2014;15:394–415.

26. Bodhi B. What does mindfulness really mean? A canonical perspective. *Contemp Buddhism* 2011;12:19–39.
27. Nyanaponika T. *The heart of Buddhist meditation (Satipatthāna): a handbook of mental training based on the Buddha's way of mindfulness, with an anthology of relevant texts translated from the Pali and Sanskrit*. London: Rider, 1962.
28. Cahn BR, Polich J. Meditation states and traits: EEG, ERP, and neuroimaging studies. *Psychol Bull* 2006;132:180–211.
29. Bishop SR, Lau M, Shapiro S, et al. Mindfulness: a proposed operational definition. *Clin Psychol Sci Pract* 2004;11:230–41.
30. Baer RA. Mindfulness training as a clinical intervention: a conceptual and clinical review. *Clin Psychol Sci Pract* 2003;10:125–43.
31. Lutz A, Slagter HA, Dunne JD, Davidson RJ. Attention regulation and monitoring in meditation. *Trends Cogn Sci* 2008;12:163–9.
32. Hofmann SG, Grossman P, Hinton DE. Loving-kindness and compassion meditation: potential for psychological interventions. *Clin Psychol Rev* 2011;31:1126–32.
33. Desbordes G, Gard T, Hoge EA, et al. Moving beyond mindfulness: defining equanimity as an outcome measure in meditation and contemplative research. *Mindfulness (N Y)* 2014;(January):356–72.
34. Benson H, Beary JF, Carol MP. The relaxation response. *Psychiatry* 1974;37:37–46.
35. Edenfield TM, Saeed SA. An update on mindfulness meditation as a self-help treatment for anxiety and depression. *Psychol Res Behav Manag* 2012;5:131–41.
36. Shapiro SL, Carlson LE, Astin JA, Freedman B. Mechanisms of mindfulness. *J Clin Psychol* 2006;62:373–86.
37. Brown KW, Ryan RM, Creswell JD, et al. Mindfulness: theoretical foundations and evidence for its salutary effects. *Psychol Inq* 2007;18:211–37.
38. Hölzel BK, Lazar SW, Gard T, Schuman-Olivier Z, Vago DR, Ott U. How does mindfulness meditation work? Proposing mechanisms of action from a conceptual and neural perspective. *Perspect Psychol Sci* 2011;6:537–59.
39. Grabovac AD, Lau MA, Willett BR. Mechanisms of mindfulness: a Buddhist psychological model. *Mindfulness (N Y)* 2011;2:154–66.
40. Vago DR, Silbersweig DA. Self-awareness, self-regulation, and self-transcendence (S-ART): a framework for understanding the neurobiological mechanisms of mindfulness. *Front Hum Neurosci* 2012;6:296.
41. Brewer JA, Elwafi HM, Davis JH. Craving to quit: psychological models and neurobiological mechanisms of mindfulness training as treatment for addictions. *Psychol Addict Behav* 2013;27:366–79.
42. Garland EL, Froeliger B, Howard MO. Mindfulness training targets neurocognitive mechanisms of addiction at the attention-appraisal-emotion interface. *Front Psychiatry* 2014;4:1–16.
43. Alsubaie M, Abbott R, Dunn B, et al. Mechanisms of action in mindfulness-based cognitive therapy (MBCT) and mindfulness-based stress reduction (MBSR) in people with physical and/or psychological conditions: a systematic review. *Clin Psychol Rev* 2017;55:74–91.
44. Creswell JD. Mindfulness interventions. *Annu Rev Psychol* 2017;68:491–516.
45. Creswell JD, Lindsay EK, Villalba DK, Chin B. Mindfulness training and physical health: mechanisms and outcomes. *Psychosom Med* 2019;81:224–32.
46. Bowen S, Witkiewitz K, Dillworth TM, et al. Mindfulness meditation and substance use in an incarcerated population. *Psychol Addict Behav* 2006;20:343–7.
47. Kabat-Zinn J. Some reflections on the origins of MBSR, skillful means, and the trouble with maps. *Contemp Buddhism* 2011;12:281–306.
48. Husgafvel V. On the Buddhist roots of contemporary non-religious mindfulness practice: moving beyond sectarian and essentialist approaches. *Temenos* 2016;52:87–126.
49. Kabat-Zinn J, Lipworth L, Burney R. The clinical use of mindfulness meditation for the self-regulation of chronic pain. *J Behav Med* 1985;8:163–90.
50. Kabat-Zinn J, Massion AO, Kristeller J, et al. Effectiveness of a meditation-based stress reduction program in the treatment of anxiety disorders. *Am J Psychiatry* 1992;149:936–43.
51. Kabat-Zinn J, Wheeler E, Light T, et al. Influence of a mindfulness meditation-based stress reduction intervention on rates of skin clearing in patients with moderate to severe psoriasis undergoing phototherapy (UVB) and photochemotherapy (PUVA). *Psychosom Med* 1998;60:625–32.
52. Davidson RJ, Kabat-Zinn J, Schumacher J, et al. Alterations in brain and immune function produced by mindfulness meditation. *Psychosom Med* 2003;65:564–70.
53. Segal ZV, Williams JMG, Teasdale JD. *Mindfulness-based cognitive therapy for depression: a new approach to preventing relapse*. New York: Guilford, 2002.
54. Teasdale JD, Segal ZV, Williams JM, Ridgeway VA, Soulsby JM, Lau MA. Prevention of relapse/recurrence in major depression by mindfulness-based cognitive therapy. *J Consult Clin Psychol* 2000;68:615–23.
55. Williams JM, Teasdale JD, Segal ZV, Soulsby J. Mindfulness-based cognitive therapy reduces overgeneral autobiographical memory in formerly depressed patients. *J Abnorm Psychol* 2000;109:150–5.
56. Segal ZV, Williams JMG, Teasdale JD. *Mindfulness-based cognitive therapy for depression*. 2nd ed. New York: Guilford, 2013.
57. Crane RS, Brewer J, Feldman C, et al. What defines mindfulness-based programs? The warp and the weft. *Psychol Med* 2016;47:990–9.
58. Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clin Psychol Sci Pract* 2003;10:144–56.
59. Shapiro BG, Greenberg J, Pedrelli P, de Jong M, Desbordes G. Mindfulness-based interventions in psychiatry. *Focus (Madison)* 2018;16:32–9.
60. Hayes SC. Acceptance and commitment therapy and the new behavior therapies: mindfulness, acceptance, and relationship. In: Hayes SC, Follette VM, Linehan MM, eds. *Mindfulness and acceptance: expanding the cognitive-behavioral tradition*. New York: Guilford, 2004.
61. Linehan MM, Schmidt H 3rd, Dimeff LA, Craft JC, Kanter J, Comtois KA. Dialectical behavior therapy for patients with borderline personality disorder and drug-dependence. *Am J Addict* 1999;8:279–92.
62. Hayes SC, Strosahl K, Wilson KG. *Acceptance and commitment therapy: the process and practice of mindful change*. 2nd ed. New York: Guilford, 2012.
63. Cullen M. Mindfulness-based interventions: an emerging phenomenon. *Mindfulness (N Y)* 2011;2:186–93.
64. Keng S-L, Smoski MJ, Robins CJ. Effects of mindfulness on psychological health: a review of empirical studies. *Clin Psychol Rev* 2011;31:1041–56.
65. Barnes N, Hattan P, Black DS, Schuman-Olivier Z. An examination of mindfulness-based programs in US medical schools. *Mindfulness (N Y)* 2017;8:489–94.
66. Crane RS, Kuyken W. The mindfulness-based interventions: teaching assessment criteria (MBI:TAC): reflections on implementation and development. *Curr Opin Psychol* 2019;28:6–10.

67. Loucks EB, Schuman-Olivier Z, Britton WB, et al. Mindfulness and cardiovascular disease risk: state of the evidence, plausible mechanisms, and theoretical framework. *Curr Cardiol Rep* 2015;17:112.
68. Tang Y-Y, Holzel BK, Posner MI. The neuroscience of mindfulness meditation. *Nat Rev Neurosci* 2015;16:213–25.
69. Neff KD, Germer CK. A pilot study and randomized controlled trial of the mindful self-compassion program. *J Clin Psychol* 2013;69:28–44.
70. Tang Y-Y, Posner MI, Rothbart MK. Meditation improves self-regulation over the life span. *Ann N Y Acad Sci* 2014;1307:104–11.
71. Dimidjian S, Arch JJ, Schneider RL, Desormeau P, Felder JN, Segal ZV. Considering meta-analysis, meaning, and metaphor: a systematic review and critical examination of “third wave” cognitive and behavioral therapies. *Behav Ther* 2016;47:886–905.
72. Carver CS, Scheier MF. *On the self-regulation of behavior*. New York: Cambridge University Press, 1998.
73. Baumeister RF, Heatherton TF. Self-regulation failure: an overview. *Psychol Inq* 1996;7:1–15.
74. Bandura A. Social cognitive theory of self-regulation. *Organ Behav Hum Decis Process* 1991;50:248–87.
75. Blair C, Razza RP. Relating effortful control, executive function, and false belief understanding to emerging math and literacy ability in kindergarten. *Child Dev* 2007;78:647–63.
76. Duckworth AL, Tsukayama E, May H. Establishing causality using longitudinal hierarchical linear modeling: an illustration predicting achievement from self-control. *Soc Psychol Personal Sci* 2010;1:311–7.
77. Evans GW, Fuller-Rowell TEDS. Childhood cumulative risk and obesity: the mediating role of self-regulatory ability. *Pediatrics* 2012;129:e68–73.
78. Trindade A. The longitudinal effects of emotion regulation on physical and psychological health: a latent growth analysis exploring the role of cognitive fusion in inflammatory bowel disease. *Br J Health Psychol* 2017;23:171–85.
79. Forbes EE, Dahl RE. Neural systems of positive affect: relevance to understanding child and adolescent depression? *Dev Psychopathol* 2005;17:827–50.
80. Amstadter A. Emotion regulation and anxiety disorders. *J Anxiety Disord* 2008;22:211–21.
81. Shiels K, Hawk LH Jr. Self-regulation in ADHD: the role of error processing. *Clin Psychol Rev* 2010;30:951–61.
82. Kuntsche E, Knibbe R, Engels R, Gmel G. Drinking motives as mediators of the link between alcohol expectancies and alcohol use among adolescents. *J Stud Alcohol Drugs* 2007;68:76–8.
83. Deblois ME, Kubzansky LD. Childhood self-regulatory skills predict adolescent smoking behavior. *Psychol Health Med* 2016;21:138–51.
84. Zlotnick C, Wolfsdorf BA, Johnson B, et al. Impaired self-regulation and suicidal behavior among adolescent and young adult psychiatric inpatients. *Arch Suicide Res* 2003;7:149–57.
85. Wagner DD, Heatherton TF. Self-regulation and its failure: the seven deadly threats to self-regulation. In: Mikulincer M, Shaver PR, Borgida E, Bargh JA, eds. *APA handbook of personality and social psychology, vol. 1: Attitudes and social cognition*. Washington DC: American Psychological Association, 2014.
86. Baumeister RF, Tice DM, Vohs KD. The strength model of self-regulation: conclusions from the second decade of willpower research. *Perspect Psychol Sci* 2018;13:141–5.
87. Botvinick MM, Braver TS, Barch DM, Carter CS, Cohen JD. Conflict monitoring and cognitive control. *Psychol Rev* 2001;108:624–52.
88. Miller EK. The prefrontal cortex and cognitive control. *Nat Rev Neurosci* 2000;1:59–65.
89. Cole MW, Schneider W. The cognitive control network: integrated cortical regions with dissociable functions. *Neuroimage* 2007;37:343–60.
90. Botvinick M, Braver T. Motivation and cognitive control: from behavior to neural mechanism. *Annu Rev Psychol* 2015;66:83–113.
91. Breukelaar IA, Antees C, Grieve SM, et al. Cognitive control network anatomy correlates with neurocognitive behavior: a longitudinal study. *Hum Brain Mapp* 2017;38:631–43.
92. Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD. The unity and diversity of executive functions and their contributions to complex “frontal lobe” tasks: a latent variable analysis. *Cogn Psychol* 2000;41:49–100.
93. Eysenck MW, Derakshan N, Santos R, Calvo MG. Anxiety and cognitive performance: attentional control theory. *Emotion* 2007;7:336–53.
94. Randall JG, Oswald FL, Beier ME. Mind-wandering, cognition, and performance: a theory-driven meta-analysis of attention regulation. *Psychol Bull* 2014;140:1411–31.
95. Diamond A. Executive functions. *Annu Rev Psychol* 2013;64:135–68.
96. Posner MI, Petersen SE. The attention system of the human brain. *Annu Rev Neurosci* 1990;13:25–42.
97. Jha AP, Krompinger J, Baime MJ. Mindfulness training modifies subsystems of attention. *Cogn Affect Behav Neurosci* 2007;7:109–19.
98. Flavell JH. Metacognition and cognitive monitoring: a new area of cognitive-developmental inquiry. *Am Psychol* 1979;34:906–11.
99. Hussain D. Meta-cognition in mindfulness: a conceptual analysis. *Psychol Thought* 2015;8:132–41.
100. Safran J, Segal ZV. *Interpersonal process in cognitive therapy*. Oxford: Jason Aronson, 1996.
101. Lau MA, Bishop SR, Segal ZV, et al. The Toronto Mindfulness Scale: development and validation. *J Clin Psychol* 2006;62:1445–67.
102. Heeren A, Van Broeck N, Philippot P. The effects of mindfulness on executive processes and autobiographical memory specificity. *Behav Res Ther* 2009;47:403–9.
103. Chiesa A, Serretti A. Mindfulness based cognitive therapy for psychiatric disorders: a systematic review and meta-analysis. *Psychiatry Res* 2011;187:441–53.
104. Lao S-A, Kissane D, Meadows G. Cognitive effects of MBSR/MBCT: a systematic review of neuropsychological outcomes. *Conscious Cogn* 2016;45:109–23.
105. Garland EL, Boettiger CA, Gaylord S, Chanon VW, Howard MO. Mindfulness is inversely associated with alcohol attentional bias among recovering alcohol-dependent adults. *Cogn Ther Res* 2012;36:441–50.
106. Andreu CI, Cosmelli D, Slagter HA, Franken IHA. Effects of a brief mindfulness-meditation intervention on neural measures of response inhibition in cigarette smokers. *PLoS One* 2018;13:e0191661.
107. Stange JP, Eisner LR, Holzel BK, et al. Mindfulness-based cognitive therapy for bipolar disorder: effects on cognitive functioning. *J Psychiatr Pract* 2011;17:410–9.
108. Shapero BG, Greenberg J, Mischoulon D, Pedrelli P, Meade K, Lazar SW. Mindfulness-based cognitive therapy improves cognitive functioning and flexibility among individuals with elevated depressive symptoms. *Mindfulness (NY)* 2018;9:1457–69.
109. Gard T, Hölzel BK, Lazar SW. The potential effects of meditation on age-related cognitive decline: a systematic review. *Ann N Y Acad Sci* 2014;1307:89–103.
110. Wong WP, Coles J, Chambers R, Wu DB-C, Hassed C. The effects of mindfulness on older adults with mild cognitive impairment. *J Alzheimers Dis Rep* 2017;1:181–93.

111. Bressler SL, Menon V. Large-scale brain networks in cognition: emerging methods and principles. *Trends Cogn Sci* 2010;14:277–90.
112. Cole MW, Reynolds JR, Power JD, Repovs G, Anticevic A, Braver TS. Multi-task connectivity reveals flexible hubs for adaptive task control. *Nat Neurosci* 2013;16:1348–55.
113. Schultz DH, Ito T, Solomyak LI, et al. Global connectivity of the fronto-parietal cognitive control network is related to depression symptoms in the general population. *Netw Neurosci* 2018;3:107–23.
114. Zanto TP, Gazzaley A. Fronto-parietal network: flexible hub of cognitive control. *Trends Cogn Sci* 2013;17:602–3.
115. Amer T, Campbell KL, Hasher L. Cognitive control as a double-edged sword. *Trends Cogn Sci* 2016;20:905–15.
116. Dixon ML, De La Vega A, Mills C, et al. Heterogeneity within the frontoparietal control network and its relationship to the default and dorsal attention networks. *Proc Natl Acad Sci U S A* 2018;115:E1598–E1607.
117. Fox KCR, Nijeboer S, Dixon ML, et al. Is meditation associated with altered brain structure? A systematic review and meta-analysis of morphometric neuroimaging in meditation practitioners. *Neurosci Biobehav Rev* 2014;43:48–73.
118. Fox KCR, Dixon ML, Nijeboer S, et al. Functional neuroanatomy of meditation: a review and meta-analysis of 78 functional neuroimaging investigations. *Neurosci Biobehav Rev* 2016;65:208–28.
119. Marzetti L, Di Lanzo C, Zappasodi F, Chella F, Raffone A, Pizzella V. Magnetoencephalographic alpha band connectivity reveals differential default mode network interactions during focused attention and open monitoring meditation. *Front Hum Neurosci* 2014;8:832.
120. Britton WB, Davis JH, Loucks EB, et al. Dismantling mindfulness-based cognitive therapy: creation and validation of 8-week focused attention and open monitoring interventions within a 3-armed randomized controlled trial. *Behav Res Ther* 2018;101:92–107.
121. Botvinick MM, Cohen JD, Carter CS. Conflict monitoring and anterior cingulate cortex: an update. *Trends Cogn Sci* 2004;8:539–46.
122. Pozuelos JP, Mead BR, Rueda MR, Malinowski P. Short-term mindful breath awareness training improves inhibitory control and response monitoring. *Prog Brain Res* 2019;244:137–63.
123. Tang Y-Y, Lu Q, Feng H, Tang R, Posner MI. Short-term meditation increases blood flow in anterior cingulate cortex and insula. *Front Psychol* 2015;6:212.
124. Hölzel BK, Ott U, Hempel H, et al. Differential engagement of anterior cingulate and adjacent medial frontal cortex in adept meditators and non-meditators. *Neurosci Lett* 2007;421:16–21.
125. Nakata H, Sakamoto K, Kakigi R. Meditation reduces pain-related neural activity in the anterior cingulate cortex, insula, secondary somatosensory cortex, and thalamus. *Front Psychol* 2014;5:1489.
126. Haase L, Thom NJ, Shukla A, et al. Mindfulness-based training attenuates insula response to an aversive interoceptive challenge. *Soc Cogn Affect Neurosci* 2014;11:182–90.
127. Frodl T, Schaub A, Banac S, et al. Reduced hippocampal volume correlates with executive dysfunctioning in major depression. *J Psychiatry Neurosci* 2006;31:316–23.
128. Hölzel BK, Carmody J, Vangel M, et al. Mindfulness practice leads to increases in regional brain gray matter density. *Psychiatry Res* 2011;191:36–43.
129. Greenberg J, Romero VL, Elkin-Frankston S, Bezdek MA, Schumacher EH, Lazar SW. Reduced interference in working memory following mindfulness training is associated with increases in hippocampal volume. *Brain Imaging Behav* 2019;13:366–76.
130. Pornpattananangkul N, Hariri AR, Harada T, et al. Cultural influences on neural basis of inhibitory control. *Neuroimage* 2016;139:114–26.
131. Slagter HA, Lutz A, Greischar LL, et al. Mental training affects distribution of limited brain resources. *PLoS Biol* 2007;5:e138.
132. Garland EL, Bryan MA, Priddy SE, Riquino MR, Froeliger B, Howard MO. Effects of mindfulness-oriented recovery enhancement versus social support on negative affective interference during inhibitory control among opioid-treated chronic pain patients: a pilot mechanistic study. *Ann Behav Med* 2019;53:865–76.
133. Gross JJ, Thompson R. Emotion regulation: conceptual foundations. In: Gross JJ, ed. *Handbook of emotion regulation*. New York: Guilford, 2007.
134. Gross J, John O. Individual differences in two emotion regulation processes: implications for affect, relationships, and well-being. *J Pers Soc Psychol* 2003;85:348–62.
135. Ng W, Diener E. Personality differences in emotions: does emotion regulation play a role? *J Individ Differ* 2009;30:100–6.
136. Gross JJ. The emerging field of emotion regulation: an integrative review. *Rev Gen Psychol* 1998;2:271–99.
137. Werner K, Gross JJ. Emotion regulation and psychopathology: a conceptual framework. In: Kring AM, Sloan DM, eds. *Emotion regulation and psychopathology: a transdiagnostic approach to etiology and treatment*. New York: Guilford, 2010;13–37.
138. Hofmann SG, Sawyer AT, Witt AA, Oh D. The effect of mindfulness-based therapy on anxiety and depression: a meta-analytic review. *J Consult Clin Psychol* 2010;78:169–83.
139. Goyal M, Singh S, Sibinga EMS, et al. Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Intern Med* 2014;174:357–68.
140. Garland EL, Hanley AW, Goldin PR, Gross JJ. Testing the mindfulness-to-meaning theory: evidence for mindful positive emotion regulation from a reanalysis of longitudinal data. *PLoS One* 2017;12:e0187727.
141. Garland EL, Hanley A, Farb NA, Froeliger BE. State mindfulness during meditation predicts enhanced cognitive reappraisal. *Mindfulness (N Y)* 2015;6:234–42.
142. Hayes SC, Strosahl K, Wilson KG, et al. Measuring experiential avoidance: a preliminary test of a working model. *Psychol Rev* 2004;54:553–78.
143. Abramowitz JS, Blakey SM, eds. *Clinical handbook of fear and anxiety: maintenance processes and treatment mechanisms*. Washington, DC: American Psychological Association, 2020.
144. Chawla N, Ostafin B. Experiential avoidance as a functional dimensional approach to psychopathology: an empirical review. *J Clin Psychol* 2007;63:871–90.
145. Bowen S, Witkiewitz K, Dillworth TM, Marlatt GA. The role of thought suppression in the relationship between mindfulness meditation and alcohol use. *Addict Behav* 2007;32:2324–8.
146. Treanor M. The potential impact of mindfulness on exposure and extinction learning in anxiety disorders. *Clin Psychol Rev* 2011;31:617–25.
147. Bernstein A, Hadash Y, Lichtash Y, Tanay G, Shepherd K, Fresco DM. Decentering and related constructs: a critical review and metacognitive processes model. *Perspect Psychol Sci* 2015;10:599–617.
148. Guendelman S, Medeiros S, Ramples H. Mindfulness and emotion regulation: insights from neurobiological, psychological, and clinical studies. *Front Psychol* 2017;8:220.
149. Björkstrand J, Schiller D, Li J, et al. The effect of mindfulness training on extinction retention. *Sci Rep* 2019;9:19896.
150. Sevinc G, Hölzel BK, Greenberg J, et al. Strengthened hippocampal circuits underlie enhanced retrieval of extinguished fear memories following mindfulness training. *Biol Psychiatry* 2019;86:693–702.

151. Gu J, Strauss C, Bond R, Cavanagh K. How do mindfulness-based cognitive therapy and mindfulness-based stress reduction improve mental health and wellbeing? A systematic review and meta-analysis of mediation studies. *Clin Psychol Rev* 2015;37:1–12.
152. Tang Y-Y, Ma Y, Fan Y, et al. Central and autonomic nervous system interaction is altered by short-term meditation. *Proc Natl Acad Sci U S A* 2009;106:8865–70.
153. Kral TRA, Schuyler BS, Mumford JA, Rosenkranz MA, Lutz A, Davidson RJ. Impact of short- and long-term mindfulness meditation training on amygdala reactivity to emotional stimuli. *Neuroimage* 2018;181:301–13.
154. Barrett LF, Gross J, Christensen TC, Benvenuto M. Knowing what you're feeling and knowing what to do about it: mapping the relation between emotion differentiation and emotion regulation. *Cogn Emot* 2011;15:713–24.
155. Etkin A, Büchel C, Gross JJ. The neural bases of emotion regulation. *Nat Rev Neurosci* 2015;16:693–700.
156. Lutz J, Herwig U, Opialla S, et al. Mindfulness and emotion regulation—an fMRI study. *Soc Cogn Affect Neurosci* 2013;9:776–85.
157. Taylor VA, Grant J, Daneault V, et al. Impact of mindfulness on the neural responses to emotional pictures in experienced and beginner meditators. *Neuroimage* 2011;57:1524–33.
158. Hölzel BK, Hoge EA, Greve DN, et al. Neural mechanisms of symptom improvements in generalized anxiety disorder following mindfulness training. *Neuroimage Clin* 2013;2:448–58.
159. Gross JJ. Emotion regulation in adulthood: timing is everything. *Curr Dir Psychol Sci* 2001;10:214–9.
160. Gross JJ, Muñoz RF. Emotion regulation and mental health. *Clin Psychol Sci Pract* 1995;2:151–64.
161. Smyth JM, Arigo D. Recent evidence supports emotion-regulation interventions for improving health in at-risk and clinical populations. *Curr Opin Psychiatry* 2009;22:205–10.
162. Groesz LM, McCoy S, Carl J, et al. What is eating you? Stress and the drive to eat. *Appetite* 2012;58:717–21.
163. Chew B-H, Shariff-Ghazali S, Fernandez A. Psychological aspects of diabetes care: effecting behavioral change in patients. *World J Diabetes* 2014;5:796–808.
164. Appleton AA, Loucks EB, Buka SL, Kubzansky LD. Divergent associations of antecedent- and response-focused emotion regulation strategies with midlife cardiovascular disease risk. *Ann Behav Med* 2014;48:246–55.
165. de Ridder D, Geenen R, Kuijper R, van Middendorp H. Psychological adjustment to chronic disease. *Lancet* 2008;372:246–55.
166. Britton W, Loucks E, King J, et al. Meta-analytic review of self-regulation processes in mindfulness-based cognitive therapy and mindfulness-based stress reduction. PROSPERO 2016: CRD42016051765. https://www.crd.york.ac.uk/PROSPERO/display_record.php?ID=CRD42016051765
167. Brewer JA, Elwafi HM, Davis JH. Craving to quit: psychological models and neurobiological mechanisms of mindfulness training as treatment for addictions. *Psychol Addict Behav* 2013;27:366–79.
168. Fulwiler C, Brewer JA, Sinnott S, Loucks EB. Mindfulness-based interventions for weight loss and CVD risk management. *Curr Cardiovasc Risk Rep* 2015;9:46.
169. Zahavi D. The experiential self: objections and clarifications. In: Siderits M, Thompson E, Zahavi D, eds. *Self, no self? Perspectives from analytical, phenomenological, and Indian traditions*. Oxford: Oxford University Press, 2011.
170. Gallagher S. Philosophical conceptions of the self: implications for cognitive science. *Trends Cogn Sci* 2000;4:14–21.
171. Farb NA, Segal ZV, Mayberg H, et al. Attending to the present: mindfulness meditation reveals distinct neural modes of self-reference. *Soc Cogn Affect Neurosci* 2007;2:313–22.
172. Craig AD. Interoception: the sense of the physiological condition of the body. *Curr Opin Neurobiol* 2003;13:500–5.
173. Khoury NM, Lutz J, Schuman-Olivier Z. Interoception in psychiatric disorders: a review of randomized, controlled trials with interoception-based interventions. *Harv Rev Psychiatry* 2018;26:250–63.
174. Mehling WE, Acree M, Stewart A, Silas J, Jones A. The multi-dimensional assessment of interoceptive awareness, version 2 (MAIA-2). *PLoS One* 2018;13:e0208034.
175. Kerr C, Sacchet M, Lazar S, Moore C, Jones S. Mindfulness starts with the body: somatosensory attention and top-down modulation of cortical alpha rhythms in mindfulness meditation. *Front Hum Neurosci* 2013;7:12.
176. Bornemann B, Herbert BM, Mehling WE, Singer T. Differential changes in self-reported aspects of interoceptive awareness through 3 months of contemplative training. *Front Psychol* 2014;5:1504.
177. Harshaw C. Interoceptive dysfunction: toward an integrated framework for understanding somatic and affective disturbance in depression. *Psychol Bull* 2015;141:311–63.
178. Farb N, Daubenmier J, Price CJ, et al. Interoception, contemplative practice, and health. *Front Psychol* 2015;6:763.
179. Loucks EB, Nardi WR, Gutman R, et al. Mindfulness-based blood pressure reduction (MB-BP): stage 1 single-arm clinical trial. *PLoS One* 2019;14:e0223095.
180. Farb NA, Segal ZV, Anderson AK. Mindfulness meditation training alters cortical representations of interoceptive attention. *Soc Cogn Affect Neurosci* 2013;8:15–26.
181. Damasio A, Damasio H, Tranel D. Persistence of feelings and sentience after bilateral damage of the insula. *Cereb Cortex* 2012;23:833–46.
182. Avery JA, Drevets WC, Moseman SE, Bodurka J, Barcalow JC, Simmons WK. Major depressive disorder is associated with abnormal interoceptive activity and functional connectivity in the insula. *Biol Psychiatry* 2014;76:258–66.
183. Nummenmaa L, Glerean E, Hari R, Hietanen JK. Bodily maps of emotions. *Proc Natl Acad Sci U S A* 2014;111:646–51.
184. Murphy J, Brewer R, Hobson H, Catmur C, Bird G. Is alexithymia characterised by impaired interoception? Further evidence, the importance of control variables, and the problems with the heartbeat counting task. *Biol Psychol* 2018;136:189–97.
185. Herbert BM, Herbert C, Pollatos O. On the relationship between interoceptive awareness and alexithymia: is interoceptive awareness related to emotional awareness? *J Pers* 2011;79:1149–75.
186. Reese E, Yan C, Jack F, Hayne H. Emerging identities: narrative and self from early childhood to early adolescence. In: McLean KC, Pasupathi M, eds. *Narrative development in adolescence: creating the storied self*. New York: Springer, 2010.
187. Habermas T, Bluck S. Getting a life: the emergence of the life story in adolescence. *Psychol Bull* 2000;126:748–69.
188. McLean KC, Breen AV. Selves in a world of stories during emerging adulthood. In: Arnett JJ, ed. *The Oxford handbook of emerging adulthood*. New York: Oxford University Press, 2016.
189. Higgins ET. Self-discrepancy: a theory relating self and affect. *Psychol Rev* 1987;94:319–40.
190. Pelham BW, Swann WB. From self-conceptions to self-worth: on the sources and structure of global self-esteem. *J Pers Soc Psychol* 1989;57:672–80.
191. Goldin PR, Jazaieri H, Ziv M, Kraemer H, Heimberg R, Gross JJ. Changes in positive self-views mediate the effect of cognitive-behavioral therapy for social anxiety disorder. *Clin Psychol Sci* 2013;1:301–10.
192. Clore J, Gaynor S. Self-statement modification techniques for distressed college students with low self-esteem and depressive symptoms. *Int J Behav Consult Ther* 2006;2:314–31.

193. Blatt SJ, Quinlan DM, Chevron ES, McDonald C, Zuroff D. Dependency and self-criticism: psychological dimensions of depression. *J Consult Clin Psychol* 1982;50:113–24.
194. Watkins E, Teasdale JD. Rumination and overgeneral memory in depression: effects of self-focus and analytic thinking. *J Abnorm Psychol* 2001;110:353–7.
195. Markus H. Self-schemata and processing information about the self. *J Pers Soc Psychol* 1977;35:63–78.
196. Rafaeli E, Bernstein DP, Young J. Schema therapy: distinctive features. New York: Routledge, 2011.
197. Margolin A, Schuman-Olivier Z, Beitel M, Arnold RM, Fulwiler CE. A preliminary study of spiritual self-schema (3-S+) therapy for reducing impulsivity in HIV-positive drug users. *J Clin Psychol* 2007;63:979–99.
198. Shadick NA, Sowell NF, Frits ML, et al. A randomized controlled trial of an internal family systems-based psychotherapeutic intervention on outcomes in rheumatoid arthritis: a proof-of-concept study. *J Rheumatol* 2013;40:1831–41.
199. Schwartz RC. Moving from acceptance toward transformation with internal family systems therapy (IFS). *J Clin Psychol* 2013;69:805–16.
200. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Natl Acad Sci U S A* 2001;98:676–82.
201. Buckner RL, Andrews-Hanna JR, Schacter DL. The brain's default network: anatomy, function, and relevance to disease. *Ann N Y Acad Sci* 2008;1124:1–38.
202. Andrews-Hanna JR. The brain's default network and its adaptive role in internal mentation. *Neuroscientist* 2012;18:251–70.
203. Fair DA, Cohen AL, Dosenbach NU, et al. The maturing architecture of the brain's default network. *Proc Natl Acad Sci U S A* 2008;105:4028–32.
204. Lois G, Wessa M. Differential association of default mode network connectivity and rumination in healthy individuals and remitted MDD patients. *Soc Cogn Affect Neurosci* 2016;11:1792–801.
205. Brewer JA, Worhunsky PD, Gray JR, Tang YY, Weber J, Kober H. Meditation experience is associated with differences in default mode network activity and connectivity. *Proc Natl Acad Sci U S A* 2011;108:20254–9.
206. Brewer JA, Garrison KA, Whitfield-Gabrieli S. What about the “self” is processed in the posterior cingulate cortex? *Front Hum Neurosci* 2013;7:647.
207. Nolen-Hoeksema S. Responses to depression and their effects on the duration of depressive episodes. *J Abnorm Psychol* 1991;100:569–82.
208. Nolen-Hoeksema S, Wisco BE, Lyubomirsky S. Rethinking rumination. *Perspect Psychol Sci* 2008;3:400–24.
209. Wilkinson PO, Croudace TJ, Goodyer IM. Rumination, anxiety, depressive symptoms and subsequent depression in adolescents at risk for psychopathology: a longitudinal cohort study. *BMC Psychiatry* 2013;13:250.
210. Borders A, Giancola PR. Trait and state hostile rumination facilitate alcohol-related aggression. *J Stud Alcohol Drugs* 2011;72:545–54.
211. Zhou H-X, Chen X, Shen Y-Q, et al. Rumination and the default mode network: meta-analysis of brain imaging studies and implications for depression. *Neuroimage* 2020;206:116287.
212. Moore JW. What is the sense of agency and why does it matter? *Front Psychol* 2016;7:1272.
213. Bandura A. Self-efficacy: toward a unifying theory of behavioral change. *Psychol Rev* 1977;84:191–215.
214. Vago DR, Zeidan F. The brain on silent: mind wandering, mindful awareness, and states of mental tranquility. *Ann N Y Acad Sci* 2016;1373:96–113.
215. Spreng RN, Sepulcre J, Turner GR, Stevens WD, Schacter DL. Intrinsic architecture underlying the relations among the default, dorsal attention, and frontoparietal control networks of the human brain. *J Cogn Neurosci* 2013;25:74–86.
216. Miller GE, Chen E, Armstrong CC, et al. Functional connectivity in central executive network protects youth against cardiometabolic risks linked with neighborhood violence. *Proc Natl Acad Sci* 2018;115:12063–8.
217. Neiss MB, Stevenson J, Sedikides C, Kumashiro M, Finkel EJ, Rusbult CE. Executive self, self-esteem, and negative affectivity: relations at the phenotypic and genotypic level. *J Pers Soc Psychol* 2005;89:593–606.
218. Tomasino B, Fabbro F. Increases in the right dorsolateral prefrontal cortex and decreases the rostral prefrontal cortex activation after 8 weeks of focused attention based mindfulness meditation. *Brain Cogn* 2016;102:46–54.
219. Taren AA, Gianaros PJ, Greco CM, et al. Mindfulness meditation training and executive control network resting state functional connectivity: a randomized controlled trial. *Psychosom Med* 2017;79:674–83.
220. Bauer CCC, Whitfield-Gabrieli S, Díaz JL, Pasaye EH, Barrios FA. From state-to-trait meditation: reconfiguration of central executive and default mode networks. *eNeuro* 2019;6:ENEURO.0335–18.2019.
221. Wells RE, Burch R, Paulsen RH, Wayne PM, Houle TT, Loder E. Meditation for migraines: a pilot randomized controlled trial. *Headache* 2014;54:1484–95.
222. Gross CR, Kreitzer MJ, Reilly-Spong M, et al. Mindfulness-based stress reduction versus pharmacotherapy for chronic primary insomnia: a randomized controlled clinical trial. *Explore (NY)* 2011;7:76–87.
223. McCubbin T, Dimidjian S, Kempe K, Glassey MS, Ross C, Beck A. Mindfulness-based stress reduction in an integrated care delivery system: one-year impacts on patient-centered outcomes and health care utilization. *Perm J* 2014;18:4–9.
224. Gawande R, Pine E, Griswold T, et al. Insurance-reimbursable mindfulness for safety-net primary care patients: a pilot randomized controlled trial. *Mindfulness (N Y)* 2019;10:1744–59.
225. Burnette JL, O'Boyle EH, VanEpps EM, Pollack JM, Finkel EJ. Mind-sets matter: a meta-analytic review of implicit theories and self-regulation. *Psychol Bull* 2013;139:655–701.
226. Bandura A, Locke EA. Negative self-efficacy and goal effects revisited. *J Appl Psychol* 2003;88:87–99.
227. Germer CK, Neff KD. Self-compassion in clinical practice. *J Clin Psychol* 2013;69:856–67.
228. Kuyken W, Watkins E, Holden E, et al. How does mindfulness-based cognitive therapy work? *Behav Res Ther* 2010;48:1105–12.
229. Van Dam NT, Sheppard SC, Forsyth JP, Earleywine M. Self-compassion is a better predictor than mindfulness of symptom severity and quality of life in mixed anxiety and depression. *J Anxiety Disord* 2011;25:123–30.
230. Sirois FM, Kitner R, Hirsch JK. Self-compassion, affect, and health-promoting behaviors. *Health Psychol* 2015;34:661–9.
231. Dunne S, Sheffield D, Chilcot J. Brief report: self-compassion, physical health and the mediating role of health-promoting behaviours. *J Health Psychol* 2016;23:993–9.
232. Ferrari M, Dal Cin M, Steele M. Self-compassion is associated with optimum self-care behaviour, medical outcomes and psychological well-being in a cross-sectional sample of adults with diabetes. *Diabet Med* 2017;34:1546–53.
233. Kelly AC, Carter JC. Self-compassion training for binge eating disorder: a pilot randomized controlled trial. *Psychol Psychother* 2015;88:285–303.
234. Semenchuk BN, Strachan SM, Fortier M. Self-compassion and the self-regulation of exercise: reactions to recalled exercise setbacks. *J Sport Exerc Psychol* 2018;40(1):31–9.

235. Dawson Rose C, Weibel A, Sullivan KM, et al. Self-compassion and risk behavior among people living with HIV/AIDS. *Res Nurs Health* 2014;37:98–106.
236. Terry ML, Leary MR. Self-compassion, self-regulation, and health. *Self Identity* 2011;10:352–62.
237. Gilbert P. Compassion and cruelty. In: *Compassion and cruelty: a biopsychosocial approach*. London: Brunner-Routledge, 2005.
238. Lutz J, Bruhl AB, Doerig N, et al. Altered processing of self-related emotional stimuli in mindfulness meditators. *Neuroimage* 2016;124(Pt A):958–67.
239. Lutz J, Berry MP, Napadow V, et al. Neural activations during self-related processing in patients with chronic pain and effects of a brief self-compassion training—a pilot study. *Psychiatry Res Neuroimaging* 2020;304:111155.
240. Elana R. *The heart of mindfulness-based stress reduction: a MBSR guide for clinicians and clients*. Ashland, OR: Pesi, 2017.
241. Kabat-Zinn J. Too early to tell: the potential impact and challenges—ethical and otherwise—inherent in the mainstreaming of dharma in an increasingly dystopian world. *Mindfulness (N Y)* 2017;8:1125–35.
242. Diedrich A, Hofmann SG, Cuijpers P, Berking M. Self-compassion enhances the efficacy of explicit cognitive reappraisal as an emotion regulation strategy in individuals with major depressive disorder. *Behav Res Ther* 2016;82:1–10.
243. Gilbert P, Procter S. Compassionate mind training for people with high shame and self-criticism: overview and pilot study of a group therapy approach. *Clin Psychol Psychother* 2006;13:353–79.
244. Cannon WB. *The wisdom of the body*. 2nd ed. Oxford: Norton, 1939.
245. O'Doherty JP, Cockburn J, Pauli WM. Learning, reward, and decision making. *Annu Rev Psychol* 2017;68:73–100.
246. LeDoux JE, Moscarello J, Sears R, Campese V. The birth, death and resurrection of avoidance: a reconceptualization of a troubled paradigm. *Mol Psychiatry* 2017;22:24–36.
247. Shapiro SL, Schwartz GE. The role of intention in self-regulation: toward intentional systemic mindfulness. In: Boekaerts M, Pintrich PR, Zeidner M. *Handbook of self-regulation*. San Diego, CA: Academic, 2000.
248. Schwartz GE. Psychobiology of health: a new synthesis. In: Hammonds BL, Scheirer CJ, eds. *Master lecture series, vol. 3: Psychology and health*. Washington, DC: American Psychological Association, 1984.
249. Davidson RJ, Schwartz GE, Rothman LP. Attentional style and the self-regulation of mode-specific attention: an electroencephalographic study. *J Abnorm Psychol* 1976;85:611–21.
250. Morris G, Nevet A, Arkadir D, Vaadia E, Bergman H. Mid-brain dopamine neurons encode decisions for future action. *Nat Neurosci* 2006;9:1057–63.
251. Glimcher PW. Understanding dopamine and reinforcement learning: the dopamine reward prediction error hypothesis. *Proc Natl Acad Sci U S A* 2011;108(suppl 3):15647–54.
252. Berridge KC, Robinson TE. What is the role of dopamine in reward: hedonic impact, reward learning, or incentive salience? *Brain Res Brain Res Rev* 1998;28:309–69.
253. Brewer JA, Potenza MN. The neurobiology and genetics of impulse control disorders: relationships to drug addictions. *Biochem Pharmacol* 2008;75:63–75.
254. Malvaez M, Wassum KM. Regulation of habit formation in the dorsal striatum. *Curr Opin Behav Sci* 2018;20:67–74.
255. Malvaez M, Greenfield VY, Matheos DP, et al. Habits are negatively regulated by histone deacetylase 3 in the dorsal striatum. *Biol Psychiatry* 2018;84:383–92.
256. Sterling P. Allostasis: a model of predictive regulation. *Physiol Behav* 2012;106:5–15.
257. Castro DC, Bruchas MR. A motivational and neuropeptidergic hub: anatomical and functional diversity within the nucleus accumbens shell. *Neuron* 2019;102:529–52.
258. Hunt WA, Matarazzo JD, Weiss SM, Gentry WD. Associative learning, habit, and health behavior. *J Behav Med* 1979;2:111–24.
259. Maren S, Fanselow MS. The amygdala and fear conditioning: has the nut been cracked? *Neuron* 1996;16:237–40.
260. Sengupta A, Yau JOY, Jean-Richard-Dit-Bressel P, et al. Basolateral amygdala neurons maintain aversive emotional salience. *J Neurosci* 2018;38:3001–12.
261. Skelly MJ, Ariwodola OJ, Weiner JL. Fear conditioning selectively disrupts noradrenergic facilitation of GABAergic inhibition in the basolateral amygdala. *Neuropharmacology* 2017;113(Pt A):231–40.
262. D'Argembeau A. On the role of the ventromedial prefrontal cortex in self-processing: the valuation hypothesis. *Front Hum Neurosci* 2013;7:372.
263. Hiser J, Koenigs M. The multifaceted role of the ventromedial prefrontal cortex in emotion, decision making, social cognition, and psychopathology. *Biol Psychiatry* 2018;83:638–47.
264. Berkman ET, Livingston JL, Kahn LE. Finding the “self” in self-regulation: the identity-value model. *Psychol Inq* 2017;28:77–98.
265. Lee JLC, Milton AL, Everitt BJ. Reconsolidation and extinction of conditioned fear: inhibition and potentiation. *J Neurosci* 2006;26:10051–6.
266. Myers KM, Davis M. Mechanisms of fear extinction. *Mol Psychiatry* 2007;12:120–50.
267. Milad MR, Quirk GJ. Fear extinction as a model for translational neuroscience: ten years of progress. *Annu Rev Psychol* 2012;63:129–51.
268. Lange I, Goossens L, Michiels S, et al. Neural responses during extinction learning predict exposure therapy outcome in phobia: results from a randomized-controlled trial. *Neuropsychopharmacology* 2020;45:534–41.
269. Nasser HM, McNally GP. Neural correlates of appetitive-aversive interactions in Pavlovian fear conditioning. *Learn Mem* 2013;20:220–8.
270. Bulganin L, Bach D, Wittmann B. Prior fear conditioning and reward learning interact in fear and reward networks. *Front Behav Neurosci* 2014;8:67.
271. Barrett LF, Quigley KS, Hamilton P. An active inference theory of allostasis and interoception in depression. *Philos Trans R Soc Lond B Biol Sci* 2016;371:20160011.
272. Barrett LF. The theory of constructed emotion: an active inference account of interoception and categorization. *Soc Cogn Affect Neurosci* 2017;12:1–23.
273. Brewer J. Mindfulness training for addictions: has neuroscience revealed a brain hack by which awareness subverts the addictive process? *Curr Opin Psychol* 2019;28:198–203.
274. Margolin A, Beitel M, Schuman-Olivier Z, Avants SK. A controlled study of a spirituality-focused intervention for increasing motivation for HIV prevention among drug users. *AIDS Educ Prev* 2006;18:311–22.
275. Khoury B, Knäuper B, Schlosser M, Carrière K, Chiesa A. Effectiveness of traditional meditation retreats: a systematic review and meta-analysis. *J Psychosom Res* 2017;92:16–25.
276. Marlatt GA, Gordon JR. *Relapse prevention: maintenance strategies in the treatment of addictive behaviors*. New York: Guilford, 1985.
277. Cormack D, Jones FW, Maltby M. A “collective effort to make yourself feel better”: the group process in mindfulness-based interventions. *Qual Health Res* 2017;28:3–15.
278. Siegel DJ. *The developing mind: toward a neurobiology of interpersonal experience*. New York: Guilford, 1999.

279. Corrigan FM, Fisher JJ, Nutt DJ. Autonomic dysregulation and the Window of Tolerance model of the effects of complex emotional trauma. *J Psychopharmacol* 2011;25:17–25.
280. Kummar AS. Mindfulness and fear extinction: a brief review of its current neuropsychological literature and possible implications for posttraumatic stress disorder. *Psychol Rep* 2017;121:792–814.
281. Hölzel BK, Brunsch V, Gard T, et al. Mindfulness-based stress reduction, fear conditioning, and the uncinat fasciculus: a pilot study. *Front Behav Neurosci* 2016;10:124.
282. Demarzo MMP, Montero-Marin J, Cuijpers P, et al. The efficacy of mindfulness-based interventions in primary care: a meta-analytic review. *Ann Fam Med* 2015;13:573–82.
283. Kalivas PW, O'Brien C. Drug addiction as a pathology of staged neuroplasticity. *Neuropsychopharmacology* 2008;33:166–80.
284. Marcotte D, Avants SK, Margolin A. Spiritual self-schema therapy, drug abuse, and HIV. *J Psychoactive Drugs* 2003;35:389–91.
285. Farb N, Daubenmier J, Price CJ, et al. Interoception, contemplative practice, and health. *Front Psychol* 2015;6:763.
286. Deci EL, Ryan RM. The “what” and “why” of goal pursuits: human needs and the self-determination of behavior. *Psychol Inq* 2000;11:227–68.
287. Schultz P, Ryan R. The “why,” “what,” and “how” of healthy self-regulation: mindfulness and well-being from a self-determination theory perspective. In: Ostafin BD, Robinson MD, Meier BP, eds. *Handbook of mindfulness and self-regulation*. New York: Springer, 2015.
288. Gruber MJ, Ranganath C. How curiosity enhances hippocampus-dependent memory: the prediction, appraisal, curiosity, and exploration (PACE) framework. *Trends Cogn Sci* 2019;23:1014–25.
289. Gruber MJ, Gelman BD, Ranganath C. States of curiosity modulate hippocampus-dependent learning via the dopaminergic circuit. *Neuron* 2014;84:486–96.
290. Deci EL, Ryan RM, Schultz PP, Niemiec CP. Being aware and functioning fully: mindfulness and interest taking within self-determination theory. In: Brown KW, Creswell JD, Ryan RM, eds. *Handbook of mindfulness: theory, research, and practice*. New York: Guilford, 2015.
291. Manohar SG, Finzi RD, Drew D, Husain M. Distinct motivational effects of contingent and noncontingent rewards. *Psychol Sci* 2017;28:1016–26.
292. Beierholm U, Guitart-Masip M, Economides M, et al. Dopamine modulates reward-related vigor. *Neuropsychopharmacology* 2013;38:1495–503.
293. Yee DM, Braver TS. Interactions of motivation and cognitive control. *Curr Opin Behav Sci* 2018;19:83–90.
294. Garland EL, Froeliger B, Howard MO. Effects of mindfulness-oriented recovery enhancement on reward responsiveness and opioid cue-reactivity. *Psychopharmacol* 2014;231:3229–38.
295. Fredrickson BL. The broaden-and-build theory of positive emotions. *Philos Trans R Soc Lond B Biol Sci* 2004;359:1367–78.
296. Vandercammen L, Hofmans J, Theuns P. Relating specific emotions to intrinsic motivation: on the moderating role of positive and negative emotion differentiation. *PLoS One* 2014;9:e115396.
297. Bora E, Yucel M, Allen NB. Neurobiology of human affiliative behaviour: implications for psychiatric disorders. *Curr Opin Psychiatry* 2009;22:320–5.
298. Engen HG, Singer T. Compassion-based emotion regulation up-regulates experienced positive affect and associated neural networks. *Soc Cogn Affect Neurosci* 2015;10:1291–301.
299. da Silva J. Trauma-sensitive mindfulness: practices for safe and transformative healing. *Trans Anal J* 2018;48:288–90.
300. Renpenning K, Taylor SG, eds. *Self-care theory in nursing: selected papers of Dorothea Orem*. New York: Springer, 2003.
301. Lindahl JR, Fisher NE, Cooper DJ, Rosen RK, Britton WB. The varieties of contemplative experience: a mixed-methods study of meditation-related challenges in Western Buddhists. *PLoS One* 2017;12:e0176239.
302. Dakwar E, Levin FR. The emerging role of meditation in addressing psychiatric illness, with a focus on substance use disorders. *Harv Rev Psychiatry* 2009;17:254–67.
303. Hoppes K. The application of mindfulness-based cognitive interventions in the treatment of co-occurring addictive and mood disorders. *CNS Spectr* 2006;11:829–51.
304. Zgierska A, Rabago D, Chawla N, Kushner K, Koehler R, Marlatt A. Mindfulness meditation for substance use disorders: a systematic review. *Subst Abuse* 2009;30:266–94.
305. Sancho M, De Gracia M, Rodriguez RC, et al. Mindfulness-based interventions for the treatment of substance and behavioral addictions: a systematic review. *Front Psychiatry* 2018;9:95.
306. Priddy SE, Howard MO, Hanley AW, Riquino MR, Friberg-felsted K, Garland EL. Mindfulness meditation in the treatment of substance use disorders and preventing future relapse: neurocognitive mechanisms and clinical implications. *Subst Abuse Rehabil* 2018;16:103–14.
307. Spears CA. Mindfulness-based interventions for addictions among diverse and underserved populations. *Curr Opin Psychol* 2019;30:11–6.
308. Byrne SP, Haber P, Baillie A, Costa DSJ, Fogliati V, Morley K. Systematic reviews of mindfulness and acceptance and commitment therapy for alcohol use disorder: should we be using third wave therapies? *Alcohol Alcohol* 2019;54:159–66.
309. Brewer JA, Bowen S, Smith JT, Marlatt GA, Potenza MN. Mindfulness-based treatments for co-occurring depression and substance use disorders: what can we learn from the brain? *Addiction* 2010;105:1698–706.
310. Black DS. Mindfulness-based interventions: an antidote to suffering in the context of substance use, misuse, and addiction. *Subst Use Misuse* 2014;49:487–91.
311. Tang YY, Posner MI, Rothbart MK, Volkow ND. Circuitry of self-control and its role in reducing addiction. *Trends Cogn Sci* 2015;19:439–44.
312. Garland EL, Howard MO. Mindfulness-based treatment of addiction: current state of the field and envisioning the next wave of research. *Addict Sci Clin Pract* 2018;13:14.
313. Bowen S, Witkiewitz K, Clifasefi SL, et al. Relative efficacy of mindfulness-based relapse prevention, standard relapse prevention, and treatment as usual for substance use disorders: a randomized clinical trial. *JAMA Psychiatry* 2014;71:547–56.
314. Garland EL, Schwarz NM, Kelly A, Whitt A, Howard MO. Mindfulness-oriented recovery enhancement for alcohol dependence: therapeutic mechanisms and intervention acceptability. *J Soc Work Pract Addict* 2012;12:242–63.
315. Price CJ, Thompson EA, Crowell S, Pike K. Longitudinal effects of interoceptive awareness training through mindful awareness in body-oriented therapy (MABT) as an adjunct to women’s substance use disorder treatment: a randomized controlled trial. *Drug Alcohol Depend* 2019;198:140–9.
316. Vidrine JL, Spears CA, Heppner WL, et al. Efficacy of mindfulness-based addiction treatment (MBAT) for smoking cessation and lapse recovery: a randomized clinical trial. *J Consult Clin Psychol* 2016;84:824–38.
317. Tang YY, Tang R, Posner MI. Brief meditation training induces smoking reduction. *Proc Natl Acad Sci US A* 2013;110:13971–5.
318. Luoma JB, Kohlenberg BS, Hayes SC, Fletcher L. Slow and steady wins the race: a randomized clinical trial of acceptance

- and commitment therapy targeting shame in substance use disorders. *J Consult Clin Psychol* 2012;80:43–53.
319. Bowen S, Witkiewitz K, Clifasefi SL, et al. Relative efficacy of mindfulness-based relapse prevention, standard relapse prevention, and treatment as usual for substance use disorders: a randomized clinical trial. *JAMA Psychiatry* 2015;71:547–56.
 320. Glasner-Edwards S, Mooney LJ, Ang A, et al. Mindfulness based relapse prevention for stimulant dependent adults: a pilot randomized clinical trial. *Mindfulness (N Y)* 2017;8:126–35.
 321. Hamidi F, Kheiran S. Mindfulness-based relapse prevention to reduce high risk behaviors of people addicted to methamphetamine. *Int J High Risk Behav Addict* 2019;8:e92609.
 322. Chen XJ, Wang DM, Zhou LD, et al. Mindfulness-based relapse prevention combined with virtual reality cue exposure for methamphetamine use disorder: study protocol for a randomized controlled trial. *Contemp Clin Trials* 2018;70:99–105.
 323. Bloom-Foster J, Mehl-Madrona L. An ultra-brief mindfulness-based intervention for patients in treatment for opioid addiction with buprenorphine: a primary care feasibility pilot study. *J Altern Complement Med* 2020;26:34–43.
 324. Sokol R, LaVertu AE, Morrill D, Albanese C, Schuman-Olivier Z. Group-based treatment of opioid use disorder with buprenorphine: a systematic review. *J Subst Abuse Treat* 2018;84:78–87.
 325. Yaghubi M, Zargar F, Akbari H. Comparing effectiveness of mindfulness-based relapse prevention with treatment as usual on impulsivity and relapse for methadone-treated patients: a randomized clinical trial. *Addict Health* 2017;9:156–65.
 326. Valls-Serrano C, Caracul A, Verdejo-Garcia A. Goal management training and mindfulness meditation improve executive functions and transfer to ecological tasks of daily life in polysubstance users enrolled in therapeutic community treatment. *Drug Alcohol Depend* 2016;165:9–14.
 327. Witkiewitz K, Bowen S. Depression, craving, and substance use following a randomized trial of mindfulness-based relapse prevention. *J Consult Clin Psychol* 2010;78:362–74.
 328. Garland EL, Roberts-Lewis A, Kelley K, Tronnier C, Hanley A. Cognitive and affective mechanisms linking trait mindfulness to craving among individuals in addiction recovery. *Subst Use Misuse* 2014;49:525–35.
 329. Bowen S, Witkiewitz K, Dillworth TM, Marlatt GA. The role of thought suppression in the relationship between mindfulness meditation and alcohol use. *Addict Behav* 2007;32:2324–8.
 330. Levin ME, Luoma JB, Haeger JA. Decoupling as a mechanism of change in mindfulness and acceptance: a literature review. *Behav Modif* 2015;39:870–911.
 331. Moss AC, Erskine JAK, Albery IP, Allen JR, Georgiou GJ. To suppress, or not to suppress? That is repression: controlling intrusive thoughts in addictive behaviour. *Addict Behav* 2015;44:65–70.
 332. Price CJ, Merrill JO, McCarty RL, Pike KC, Tsui JI. A pilot study of mindful body awareness training as an adjunct to office-based medication treatment of opioid use disorder. *J Subst Abuse Treat* 2020;108:123–8.
 333. Brewer JA, Sinha R, Chen JA, et al. Mindfulness training and stress reactivity in substance abuse: results from a randomized, controlled stage I pilot study. *Subst Abuse* 2009;30:306–17.
 334. Westbrook C, Creswell JD, Tabibnia G, Julson E, Kober H, Tindle HA. Mindful attention reduces neural and self-reported cue-induced craving in smokers. *Soc Cogn Affect Neurosci* 2013;8:73–84.
 335. Garland EL. Trait mindfulness predicts attentional and autonomic regulation of alcohol cue-reactivity. *J Psychophysiol* 2011;25:180–9.
 336. Carroll H, Lustyk MKB. Mindfulness-based relapse prevention for substance use disorders: effects on cardiac vagal control and craving under stress. *Mindfulness (N Y)* 2018;9:488–99.
 337. Garland EL. Restructuring reward processing with mindfulness-oriented recovery enhancement: novel therapeutic mechanisms to remediate hedonic dysregulation in addiction, stress, and pain. *Ann N Y Acad Sci* 2016;1373:25–37.
 338. Garland EL, Atchley RM, Hanley AW, Zubieta J-K, Froeliger B. Mindfulness-oriented recovery enhancement mediates hedonic dysregulation in opioid users: neural and affective evidence of target engagement. *Sci Adv* 2019;5:eax1569.
 339. Carrière K, Khoury B, Günak MM, Knäuper B. Mindfulness-based interventions for weight loss: a systematic review and meta-analysis. *Obes Rev* 2017;19:164–77.
 340. Godfrey KM, Gallo LC, Afari N. Mindfulness-based interventions for binge eating: a systematic review and meta-analysis. *J Behav Med* 2014;38:348–62.
 341. Katterman SN, Kleinman BM, Hood MM, Nackers LM, Corsica JA. Mindfulness meditation as an intervention for binge eating, emotional eating, and weight loss: a systematic review. *Eat Behav* 2014;15:197–204.
 342. Mantzios M, Wilson JC. Mindfulness, eating behaviours, and obesity: a review and reflection on current findings. *Curr Obes Rep* 2015;4:141–6.
 343. O'Reilly GA, Cook L, Spruijt-Metz D, Black DS. Mindfulness-based interventions for obesity-related eating behaviours: a literature review. *Obes Rev* 2014;15:453–61.
 344. Rogers JM, Ferrari M, Mosely K, Lang CP, Brennan L. Mindfulness-based interventions for adults who are overweight or obese: a meta-analysis of physical and psychological health outcomes. *Obes Rev* 2016;18:51–67.
 345. Olson KL, Emery CF. Mindfulness and weight loss: a systematic review. *Psychosom Med* 2015;77:59–67.
 346. Warren JM, Smith N, Ashwell M. A structured literature review on the role of mindfulness, mindful eating and intuitive eating in changing eating behaviours: effectiveness and associated potential mechanisms. *Nutr Res Rev* 2017;30:272–83.
 347. Roche AL, Kroska EB, Denburg NL. Acceptance- and mindfulness-based interventions for health behavior change: systematic reviews and meta-analyses. *J Context Behav Sci* 2019;13(April):74–93.
 348. Brewer JA, Ruf A, Beccia AL, et al. Can mindfulness address maladaptive eating behaviors? Why traditional diet plans fail and how new mechanistic insights may lead to novel interventions. *Front Psychol* 2018;9:1418.
 349. Reed JR, Yates BC, Houfek J, Pullen CH, Briner W, Schmid KK. Eating self-regulation in overweight and obese adults: a concept analysis. *Nurs Forum* 2015;51:105–16.
 350. Baer RA, Fischer S, Huss DB. Mindfulness and acceptance in the treatment of disordered eating. *J Ration Emot Cogn Behav Ther* 2005;23:281–300.
 351. Arch JJ, Brown KW, Goodman RJ, Della Porta MD, Kiken LG, Tillman S. Enjoying food without caloric cost: the impact of brief mindfulness on laboratory eating outcomes. *Behav Res Ther* 2016;79:23–34.
 352. Mason AE, Jhaveri K, Cohn M, Brewer JA. Testing a mobile mindful eating intervention targeting craving-related eating: feasibility and proof of concept. *J Behav Med* 2018;41:160–73.
 353. Caldwell K, Baime M, Wolever R. Mindfulness based approaches to obesity and weight loss maintenance. *J Ment Health Couns* 2012;34:269–82.
 354. Kristeller JL, Wolever RQ. Mindfulness-based eating awareness training for treating binge eating disorder: the conceptual foundation. *Eat Disord* 2011;19:49–61.
 355. Kristeller JL, Epel E. Mindful eating and mindless eating: the science and the practice. In: Amanda I, Ngnoumen CT,

- Langer EJ, eds. *The Wiley Blackwell handbook of mindfulness*. Malden, MA: Wiley, 2014.
356. Godsey J. The role of mindfulness based interventions in the treatment of obesity and eating disorders: an integrative review. *Complement Ther Med* 2013;21:430–9.
 357. Turgon R, Ruffault A, Juneau C, Blatier C, Shankland R. Eating disorder treatment: a systematic review and meta-analysis of the efficacy of mindfulness-based programs. *Mindfulness (N Y)* 2019;10:2225–44.
 358. Linardon J, Fairburn CG, Fitzsimmons-Craft EE, Wilfley DE, Brennan L. The empirical status of the third-wave behaviour therapies for the treatment of eating disorders: a systematic review. *Clin Psychol Rev* 2017;58:125–40.
 359. Dunne J. Mindfulness in anorexia nervosa: an integrated review of the literature. *J Am Psychiatr Nurses Assoc* 2017;24:109–17.
 360. Masuda A, Hill ML. Mindfulness as therapy for disordered eating: a systematic review. *Neuropsychiatry (London)* 2013;3:433–47.
 361. Pisetsky EM, Schaefer LM, Wonderlich SA, Peterson CB. Emerging psychological treatments in eating disorders. *Psychiatr Clin North Am* 2019;42:219–29.
 362. Wanden-Berghe RG, Sanz-Valero J, Wanden-Berghe C. The application of mindfulness to eating disorders treatment: a systematic review. *Eat Disord* 2010;19:34–48.
 363. Oikonomou MT, Arvanitis M, Sokolove RL. Mindfulness training for smoking cessation: a meta-analysis of randomized-controlled trials. *J Health Psychol* 2017;22:1841–50.
 364. Maglione MA, Maher AR, Ewing B, et al. Efficacy of mindfulness meditation for smoking cessation: a systematic review and meta-analysis. *Addict Behav* 2017;69:27–34.
 365. de Souza ICW, de Barros VV, Gomide HP, et al. Mindfulness-based interventions for the treatment of smoking: a systematic literature review. *J Altern Complement Med* 2015;21:129–40.
 366. Brewer JA, Mallik S, Babuscio TA, et al. Mindfulness training for smoking cessation: results from a randomized controlled trial. *Drug Alcohol Depend* 2011;119:72–80.
 367. Schuman-Olivier Z, Hoepfner BB, Evins AE, Brewer JA. Finding the right match: mindfulness training may potentiate the therapeutic effect of nonjudgment of inner experience on smoking cessation. *Subst Use Misuse* 2014;49:586–94.
 368. Elwafi HM, Witkiewitz K, Mallik S, Thornhill TA, Brewer JA. Mindfulness training for smoking cessation: moderation of the relationship between craving and cigarette use. *Drug Alcohol Depend* 2013;130:222–9.
 369. Garrison KA, Pal P, Rojiani R, Dallery J, O'Malley SS, Brewer JA. A randomized controlled trial of smartphone-based mindfulness training for smoking cessation: a study protocol. *BMC Psychiatry* 2015;15:83.
 370. Garrison KA, Pal P, O'Malley SS, Pittman BP, Gueorguieva R, Brewer JA. Craving to quit: a randomized controlled trial of smartphone app-based mindfulness training for smoking cessation. *Nicotine Tob Res* 2018;22:324–31.
 371. Garrison KA, Santoyo JF, Davis JH, Thornhill TA, Kerr CE, Brewer JA. Effortless awareness: using real time neurofeedback to investigate correlates of posterior cingulate cortex activity in meditators' self-report. *Front Hum Neurosci* 2013;7:440.
 372. Janes AC, Datko M, Roy A, et al. Quitting starts in the brain: a randomized controlled trial of app-based mindfulness shows decreases in neural responses to smoking cues that predict reductions in smoking. *Neuropsychopharmacology* 2019;44:1631–8.
 373. Swendeman D, Ingram BL, Rotheram-Borus MJ. Common elements in self-management of HIV and other chronic illnesses: an integrative framework. *AIDS Care* 2009;21:1321–34.
 374. Grady PA, Gough LL. Self-management: a comprehensive approach to management of chronic conditions. *Am J Public Health* 2014;104:e25–31.
 375. Ornstein SM, Nietert PJ, Jenkins RG, Litvin CB. The prevalence of chronic diseases and multimorbidity in primary care practice: a PPRNet report. *J Am Board Fam Med* 2013;26:518–24.
 376. Bodenheimer T, Lorig K, Holman H, Grumbach K. Patient self-management of chronic disease in primary care. *JAMA* 2002;288:2469–75.
 377. Kanaan SB. *Promoting effective self-management approaches to improve chronic disease care: lessons learned*. Oakland, CA: California HealthCare Foundation, 2008.
 378. Scott KM, Lim C, Al-Hamzawi A, et al. Association of mental disorders with subsequent chronic physical conditions: world mental health surveys from 17 countries. *JAMA Psychiatry* 2016;73:150–8.
 379. Cohen S, Janicki-Deverts D, Miller GE. Psychological stress and disease. *JAMA* 2007;298:1685–7.
 380. Chapman DP, Perry GS, Strine TW. The vital link between chronic disease and depressive disorders. *Prev Chronic Dis* 2005;2:A14.
 381. Kessler RC, DuPont RL, Berglund P, Wittchen HU. Impairment in pure and comorbid generalized anxiety disorder and major depression at 12 months in two national surveys. *Am J Psychiatry* 1999;156:1915–23.
 382. Loucks EB, Gilman SE, Britton WB, Gutman R, Eaton CB, Buka SL. Associations of mindfulness with glucose regulation and diabetes. *Am J Health Behav* 2016;40:258–67.
 383. Benzo RP. Mindfulness and motivational interviewing: two candidate methods for promoting self-management. *Chron Respir Dis* 2013;10:175–82.
 384. Ludwig K, Haindl A, Laufs R, et al. Self-regulation in preschool children's everyday life: exploring day-to-day variability and the within- and between-person structure. *J Self Regul Regul* 2016;2:99–117.
 385. Youngwanichsetha S, Phumdoung S, Ingkathawornwong T. The effects of mindfulness eating and yoga exercise on blood sugar levels of pregnant women with gestational diabetes mellitus. *Appl Nurs Res* 2014;27:227–30.
 386. Armani Kian A, Vahdani B, Noorbala AA, et al. The impact of mindfulness-based stress reduction on emotional wellbeing and glycemic control of patients with type 2 diabetes mellitus. *J Diabetes Res* 2018;2018:1986820.
 387. Ellis DA, Carcone AI, Slatcher R, et al. Efficacy of mindfulness-based stress reduction in emerging adults with poorly controlled, type 1 diabetes: a pilot randomized controlled trial. *Pediatr Diabetes* 2019;20:226–34.
 388. Homan KJ, Sirois FM. Self-compassion and physical health: exploring the roles of perceived stress and health-promoting behaviors. *Health Psychol Open* 2017;4:2055102917729542.
 389. Gillions A, Cheang R, Duarte R, Kingdom U, Ireland N. The effect of mindfulness practice on aggression and violence levels in adults: a systematic review. *Aggress Violent Behav* 2019;48:104–15.
 390. Stratton KJ. Mindfulness-based approaches to impulsive behaviors. *New School Psychol Bull* 2006;4:49–71.
 391. Peters JR, Erisman SM, Upton BT, Baer RA, Roemer L. A preliminary investigation of the relationships between dispositional mindfulness and impulsivity. *Mindfulness* 2011;2:228–35.
 392. Singh NN, Wahler RG, Adkins AD, Myers RE. Soles of the feet: a mindfulness-based self-control intervention for aggression by an individual with mild mental retardation and mental illness. *Res Dev Disabil* 2003;24:158–69.
 393. Verheul R, Van den Bosch L, Koeter M, de Ridder M. Dialectical behaviour therapy for women with borderline personality disorder: 12-month, randomised clinical trial in the Netherlands. *Br J Psychiatry* 2003;182:135–40.

394. Zarling A, Lawrence E, Marchman J. A randomized controlled trial of acceptance and commitment therapy for aggressive behavior. *J Consult Clin Psychol* 2015;83:199–212.
395. Robins CJ, Keng S, Ekblad AG, Brantley JG. Effects of mindfulness-based stress reduction on emotional experience and expression: a randomized controlled trial. *J Clin Psychol* 2012;68:117–31.
396. Cheng S, Banks K, Bartlett BA, Miguel GS, Vujanovic AA. Posttraumatic stress and mindfulness facets in relation to suicidal ideation severity among psychiatric inpatients. *Mindfulness (N Y)* 2017;9:761–72.
397. Zeng W, Ma Z, Li H. Mindfulness and suicidal ideation in Chinese older adults: perceived stress as mediator. *Soc Behav Pers* 2017;45:733–40.
398. Garisch JA, Wilson MS. Prevalence, correlates, and prospective predictors of non-suicidal self-injury among New Zealand adolescents: cross-sectional and longitudinal survey data. *Child Adolesc Psychiatry Ment Health* 2015;9:28.
399. Heath NL, Carsley D, De Riggi ME, Mills D, Mettler J. The relationship between mindfulness, depressive symptoms, and non-suicidal self-injury amongst adolescents. *Arch Suicide Res* 2016;20:635–49.
400. Méndez-Bustos P, Calati R, Rubio-Ramírez F, Olié E, Courtet P, Lopez-Castroman J. Effectiveness of psychotherapy on suicidal risk: a systematic review of observational studies. *Front Psychol* 2019;10:277.
401. DeCou CR, Comtois KA, Landes SJ. Dialectical behavior therapy is effective for the treatment of suicidal behavior: a meta-analysis. *Behav Ther* 2019;50:60–72.
402. Chesin M, Interian A, Kline A, Benjamin-Phillips C, Latorre M, Stanley B. Reviewing mindfulness-based interventions for suicidal behavior. *Arch Suicide Res* 2016;20:507–27.
403. Le TN, Gobert JM. Translating and implementing a mindfulness-based youth suicide prevention intervention in a native American community. *J Child Fam Stud* 2013;24:12–23.
404. Barnhofer T, Crane C, Hargus E, Amarasinghe M, Winder R, Williams JMG. Mindfulness-based cognitive therapy as a treatment for chronic depression: a preliminary study. *Behav Res Ther* 2009;47:366–73.
405. Lu R, Zhou Y, Wu Q, et al. The effects of mindfulness training on suicide ideation among left-behind children in China: a randomized controlled trial. *Child Care Health Dev* 2019;45:371–9.
406. Forkmann T, Brakemeier E-L, Teismann T, Schramm E, Michalak J. The effects of mindfulness-based cognitive therapy and cognitive behavioral analysis system of psychotherapy added to treatment as usual on suicidal ideation in chronic depression: results of a randomized-clinical trial. *J Affect Disord* 2016;200:51–7.
407. Barnhofer T, Crane C, Brennan K, et al. Mindfulness-based cognitive therapy (MBCT) reduces the association between depressive symptoms and suicidal cognitions in patients with a history of suicidal depression. *J Consult Clin Psychol* 2015;83:1013–20.
408. Chesin MS, Benjamin-Phillips CA, Keilp J, Fertuck EA, Brodsky BS, Stanley B. Improvements in executive attention, rumination, cognitive reactivity, and mindfulness among high-suicide risk patients participating in adjunct mindfulness-based cognitive therapy: preliminary findings. *J Altern Complement Med* 2016;22:642–9.
409. McCauley E, Berk MS, Asarnow JR, et al. Efficacy of dialectical behavior therapy for adolescents at high risk for suicide a randomized clinical trial. *JAMA Psychiatry* 2018;75:777–85.
410. Mehlum L, Ramberg M, Tørmoen AJ, et al. Dialectical behavior therapy compared with enhanced usual care for adolescents with repeated suicidal and self-harming behavior: outcomes over a one-year follow-up. *J Am Acad Child Adolesc Psychiatry* 2016;55:295–300.
411. Rees CS, Hasking P, Breen LJ, Lipp OV, Mamotte C. Group mindfulness based cognitive therapy vs group support for self-injury among young people: study protocol for a randomised controlled trial. *BMC Psychiatry* 2015;15:154.
412. Tighe J, Nicholas J, Shand F, Christensen H. Efficacy of acceptance and commitment therapy in reducing suicidal ideation and deliberate self-harm: systematic review. *JMIR Ment Health* 2018;5:e10732.
413. Oberle E, Schonert-reichl KA, Lawlor MS, Kimberly C. Mindfulness and inhibitory control in early adolescence. *J Early Adolesc* 2012;32:565–88.
414. Eisenlohr-Moul TA, Peters JR, Pond RS, DeWall CN. Both trait and state mindfulness predict lower aggressiveness via anger rumination: a multilevel mediation analysis. *Mindfulness (N Y)* 2016;7:713–26.
415. Kelley TM, Lambert EG. Mindfulness as a potential means of attenuating anger and aggression for prospective criminal justice professionals. *Mindfulness (N Y)* 2012;3:261–74.
416. Feldman G, Greeson J, Renna M, Robbins-Monteith K. Mindfulness predicts less texting while driving among young adults: examining attention- and emotion-regulation motives as potential mediators. *Pers Individ Dif* 2011;51:856–61.
417. Murphy C, Mackillop J. Living in the here and now: interrelationships between impulsivity, mindfulness, and alcohol misuse. *Psychopharmacology (Berl)* 2012;219:527–36.
418. Black DS, Sussman S, Johnson CA, Milam J. Testing the indirect effect of trait mindfulness on adolescent cigarette smoking through negative affect. *J Subst Abuse* 2012;17:417–29.
419. Lakey CE, Campbell WK, Warren K, Goodie AS. Dispositional mindfulness as a predictor of the severity of gambling outcomes. *Pers Individ Dif* 2007;43:1698–710.
420. Shorey RC, Elmquist J, Gawrysiak MJ, Anderson S, Stuart GL. The relationship between mindfulness and compulsive sexual behavior in a sample of men in treatment for substance use disorders. *Mindfulness (N Y)* 2016;7:866–73.
421. Crosby JM, Twohig MP, Crosby JM. Acceptance and commitment therapy for problematic internet pornography use: a randomized trial. *Behav Ther* 2016;47:355–66.
422. Reid RC, Bramen JE, Anderson A, Cohen MS. Mindfulness, emotional dysregulation, impulsivity, and stress proneness among hypersexual patients. *J Clin Psychol* 2014;70:313–21.
423. Brattström O, Eriksson M, Larsson E, Oldner A. Socio-economic status and co-morbidity as risk factors for trauma. *Eur J Epidemiol* 2014;30:151–7.
424. Goodman A, Sommers-Flanagan J, Treleaven DA. Trauma-sensitive mindfulness: practices for safe and transformative healing. *J Contemp Psychother* 2018;49:127–8.
425. Waldron EM, Hong S, Moskowitz JT, Burnett-Zeigler I. A systematic review of the demographic characteristics of participants in US-based randomized controlled trials of mindfulness-based interventions. *Mindfulness (N Y)* 2018;9:1671–92.
426. Cook BL, Trinh N-H, Li Z, Hou SS-Y, Progovac AM. Trends in racial-ethnic disparities in access to mental health care, 2004–2012. *Psychiatr Serv* 2017;68:9–16.
427. Budge SL, Adelson JL, Howard KAS. Anxiety and depression in transgender individuals: the roles of transition status, loss, social support, and coping. *J Consult Clin Psychol* 2013;81:545–57.
428. Gilman SE, Cochran SD, Mays VM, Hughes M, Ostrow D, Kessler RC. Risk of psychiatric disorders among individuals reporting same-sex sexual partners in the National Comorbidity Survey. *Am J Public Health* 2001;91:933–9.
429. Khullar D, Chokshi DA. Challenges for immigrant health in the USA—the road to crisis. *Lancet* 2019;393:2168–74.
430. Crenshaw KW. Twenty years of critical race theory: looking back to move forward. *Conn Law Rev* 2011;43:1253–354.

431. Witkiewitz K, Greenfield BL, Bowen S. Mindfulness-based relapse prevention with racial and ethnic minority women. *Addict Behav* 2013;38:2821–4.
432. Greenfield BL, Roos C, Hagler KJ, Stein E, Bowen S, Witkiewitz KA. Race/ethnicity and racial group composition moderate the effectiveness of mindfulness-based relapse prevention for substance use disorder. *Addict Behav* 2018;81:96–103.
433. Proulx J, Croff R, Oken B, et al. Considerations for research and development of culturally relevant mindfulness interventions in American minority communities. *Mindfulness (N Y)* 2018;9:361–70.
434. George S, Duran N, Norris K. A systematic review of barriers and facilitators to minority research participation among African Americans, Latinos, Asian Americans, and Pacific Islanders. *Am J Public Health* 2014;104:e16–31.
435. Carmody J, Baer RA. Relationships between mindfulness practice and levels of mindfulness, medical and psychological symptoms and well-being in a mindfulness-based stress reduction program. *J Behav Med* 2008;31:23–33.
436. Brown KW, Ryan RM. The benefits of being present: mindfulness and its role in psychological well-being. *J Pers Soc Psychol* 2003;84:822–48.
437. Carmody J, Baer RA. How long does a mindfulness-based stress reduction program need to be? A review of class contact hours and effect sizes for psychological distress. *J Clin Psychol* 2009;65:627–38.
438. Davidson RJ, Kaszniak AW. Conceptual and methodological issues in research on mindfulness and meditation. *Am Psychol* 2015;70:581–92.
439. Norris CJ, Creem D, Hendlar R, Kober H. Brief mindfulness meditation improves attention in novices: evidence from ERPs and moderation by neuroticism. *Front Hum Neurosci* 2018;12:315.
440. Roos C, Kirouac M, Stein E, Wilson A, Bowen S, Witkiewitz K. An open trial of rolling admission mindfulness-based relapse prevention (Rolling MBRP): feasibility, acceptability, dose-response relations, and mechanisms. *Mindfulness (N Y)* 2019;10:1062–73.
441. Mohatt NV, Thompson AB, Thai ND, Tebes JK. Historical trauma as public narrative: a conceptual review of how history impacts present-day health. *Soc Sci Med* 2014;106:128–36.
442. Moeini-Jazani M, Knoeferle K, de Molière L, Gatti E, Warlop L. Social power increases interoceptive accuracy. *Front Psychol* 2017;8:1322.
443. Krieger N. Living and dying at the crossroads: racism, embodiment, and why theory is essential for a public health of consequence. *Am J Public Health* 2016;106:832–3.
444. Quadt L, Critchley HD, Garfinkel SN. The neurobiology of interoception in health and disease. *Ann N Y Acad Sci* 2018;1428:112–28.
445. Khalsa SS, Adolphs R, Cameron OG, et al. Interoception and mental health: a roadmap. *Biol Psychiatry Cogn Neurosci Neuroimaging* 2018;3:501–13.
446. van der Kolk BA, Stone L, West J, et al. Yoga as an adjunctive treatment for posttraumatic stress disorder. *J Clin Psychiatry* 2014;75:e559–65.
447. Geller S. Cultivation of therapeutic presence: therapeutic drumming and mindfulness practices. *Tijdschr Clientgerichte Psychother* 2009;47:273–87.
448. Fancourt D, Perkins R, Ascenso S, Carvalho LA, Steptoe A, Williamon A. Effects of group drumming interventions on anxiety, depression, social resilience and inflammatory immune response among mental health service users. *PLoS One* 2016;11:e0151136.
449. Travis R, Gann E, Crooke AHD, Jenkins SM. Hip hop, empowerment, and therapeutic beat-making: potential solutions for summer learning loss, depression, and anxiety in youth. *J Hum Behav Soc Environ* 2019;29:744–65.
450. Levy I, Emdin C, Adjapong ES. Hip-hop cypher in group work. *Soc Work Groups* 2018;41:103–10.
451. Mindful. Hip-hop state of mind. 2020. <https://www.mindful.org/hip-hop-state-mind/>
452. Mindful Life Project. MLP resources. ND. <http://mindfullifeproject.org/resources/>
453. Grasser LR, Al-Saghir H, Wanna C, Spinei J, Javanbakht A. Moving through the trauma: dance/movement therapy as a somatic-based intervention for addressing trauma and stress among Syrian refugee children. *J Am Acad Child Adolesc Psychiatry* 2019;58:1124–6.
454. Stanek D. Bridging past and present: embodied intergenerational trauma and the implications for dance/movement therapy. *Body Mov Dance Psychother* 2014;10:94–105.
455. Wong SYS, Chan JYC, Zhang D, Lee EKP, Tsoi KKF. The safety of mindfulness-based interventions: a systematic review of randomized controlled trials. *Mindfulness (N Y)* 2018;9:1344–57.
456. Baer R, Crane C, Miller E, Kuyken W. Doing no harm in mindfulness-based programs: conceptual issues and empirical findings. *Clin Psychol Rev* 2019;71:101–14.
457. Britton WB. Can mindfulness be too much of a good thing? The value of a middle way. *Curr Opin Psychol* 2019;28:159–65.
458. Grant AM, Schwartz B. Too much of a good thing: the challenge and opportunity of the inverted U. *Perspect Psychol Sci* 2011;6:61–76.
459. Bent S, Padula A, Avins AL. Brief communication: better ways to question patients about adverse medical events. *Ann Intern Med* 2006;144:257–61.
460. Treleaven D. Trauma-sensitive mindfulness: practices for safe and transformative healing. *Trans Anal J* 2018;48:288–90.
461. Afifi T, Asmundson G, Sareen J. Epidemiology of traumatic events and posttraumatic stress disorder. In: Nutt DJ, Stein MB, Zohar J, eds. *Post traumatic stress disorder: diagnosis, management and treatment*. Boca Raton, FL: Taylor & Francis, 2009.
462. Compson J. Meditation, trauma and suffering in silence: raising questions about how meditation is taught and practiced in Western contexts in the light of a contemporary trauma resiliency model. *Contemp Buddhism* 2014;15:274–97.
463. Boyd JE, Lanius RA, McKinnon MC. Mindfulness-based treatments for posttraumatic stress disorder: a review of the treatment literature and neurobiological evidence. *J Psychiatry Neurosci* 2018;43:7–25.
464. Polusny MA, Erbes CR, Thuras P, et al. Mindfulness-based stress reduction for posttraumatic stress disorder among veterans: a randomized clinical trial. *JAMA* 2015;314:456–65.
465. Kearney DJ, McDermott K, Malte C, Martinez M, Simpson TL. Effects of participation in a mindfulness program for veterans with posttraumatic stress disorder: a randomized controlled pilot study. *J Clin Psychol* 2012;69:14–27.
466. Banks K, Newman E, Saleem J. An overview of the research on mindfulness-based interventions for treating symptoms of posttraumatic stress disorder: a systematic review. *J Clin Psychol* 2015;71:935–63.
467. King AP, Erickson TM, Giardino ND, et al. A pilot study of group mindfulness-based cognitive therapy (MBCT) for combat veterans with posttraumatic stress disorder (PTSD). *Depress Anxiety* 2013;30:638–45.
468. Kearney DJ, Malte CA, McManus C, Martinez ME, Felleman B, Simpson TL. loving-kindness meditation for posttraumatic stress disorder: a pilot study. *J Trauma Stress* 2013;26:426–34.

469. Bormann JE, Oman D, Walter KH, Johnson BD. Mindful attention increases and mediates psychological outcomes following mantram repetition practice in veterans with posttraumatic stress disorder. *Med Care* 2014;52:S13–8.
470. Herman JL. Posttraumatic stress disorder as a shame disorder. In: *Shame in the therapy hour*. Dearing RL, Tangney JP, eds. Washington, DC: American Psychological Association, 2011.
471. Dimidjian S, Segal ZV. Prospects for a clinical science of mindfulness-based intervention. *Am Psychol* 2015;70:593–620.
472. Stein E, Witkiewitz K. Dismantling mindfulness-based programs: a systematic review to identify active components of treatment. *Mindfulness* (N Y) 2020;25 Jul [Epub ahead of print].
473. Powell SK. Mindfulness: another tool in the case managers' toolbox. *Prof Case Manag* 2014;19:159–60.
474. Van Dam NT, van Vugt MK, Vago DR, et al. Mind the hype: a critical evaluation and prescriptive agenda for research on mindfulness and meditation. *Perspect Psychol Sci* 2018;13:36–61.