

Research



Prevalence and correlates of psychiatric morbidity, comorbid anxiety and depression among medical students in public and private tertiary institutions in a Nigerian state: a cross-sectional analytical study

Joshua Falade, Adedayo Hakeem Oyebanji, Adefunke Olarinre Babatola, Olusola Olawumi Falade, Temitope Ojo Olumuyiwa

Corresponding author: Adefunke Olarinre Babatola, Department of Pediatrics, Ekiti State University, Ado Ekiti, Ekiti State, Nigeria. mamafunky@yahoo.co.uk

Received: 14 Jul 2020 - **Accepted:** 01 Aug 2020 - **Published:** 14 Sep 2020

Keywords: Comorbid anxiety and depression, psychiatric morbidity, medical students, private and public institutions, Nigeria

Copyright: Joshua Falade et al. Pan African Medical Journal (ISSN: 1937-8688). This is an Open Access article distributed under the terms of the Creative Commons Attribution International 4.0 License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Cite this article: Joshua Falade et al. Prevalence and correlates of psychiatric morbidity, comorbid anxiety and depression among medical students in public and private tertiary institutions in a Nigerian state: a cross-sectional analytical study. Pan African Medical Journal. 2020;37(53). 10.11604/pamj.2020.37.53.24994

Available online at: <https://www.panafrican-med-journal.com//content/article/37/53/full>

Prevalence and correlates of psychiatric morbidity, comorbid anxiety and depression among medical students in public and private tertiary institutions in a Nigerian state: a cross-sectional analytical study

Joshua Falade¹, Adedayo Hakeem Oyebanji², Adefunke Olarinre Babatola^{3,&}, Olusola Olawumi Falade⁴, Temitope Ojo Olumuyiwa⁵

¹Department of Mental Health, Afe Babalola University, Ado Ekiti, Ekiti State, Nigeria,

²Department of Pediatrics, Afe Babalola University, Ado Ekiti, Ekiti State, Nigeria, ³Department of

Pediatrics, Ekiti State University, Ado Ekiti, Ekiti State, Nigeria, ⁴School of Nursing Osun state Hospitals' Management Board Osogbo, Osogbo, Osun State Nigeria, ⁵Department of Community Health, Obafemi Awolowo University, Ile-Ife, Nigeria

&Corresponding author

Adefunke Olarinre Babatola, Department of Pediatrics, Ekiti State University, Ado Ekiti, Ekiti State, Nigeria

Abstract

Introduction: the study assessed the prevalence and factors associated with psychiatric morbidity (an array of psychological disorders), and comorbid anxiety and depression among medical students in Ekiti State, Nigeria. **Methods:** a cross-sectional study of medical students in two universities (one public and one private) in Ekiti state was conducted. A semi-structured questionnaire with adapted questions from the General Health Questionnaire and Hospital Anxiety and Depression Scale was used to obtain information on socio-demographic characteristics, psychiatric morbidity and comorbid anxiety and depression. Data was collected from April 2019 to August 2019. Data was analyzed using Statistical Package for Social Sciences (SPSS) version 21. **Results:** a total of 944 medical students participated in the survey. The overall prevalence of psychiatric morbidity and comorbid anxiety and depression among the respondents was 25.0% (CI = 22.1-27.8) and 14.3% (CI = 12.3-16.5) respectively. The factors independently associated with psychiatric morbidity included being a student of a private institution [adjusted odds ratio [AOR] =6.533, [95% confidence interval [C.I] =3.298-12.940], average academic performance [AOR =1.711, [95% C.I =1.173-2.496], below average academic performance [AOR =2.425, [95% C.I =1.313-4.478], and having a father or a mother with highest level of formal education below first degree [AOR =3.147, [95% C.I =1.579-6.272] and [AOR =2.053, [95% C.I =1.074-3.927] respectively.

The factors independently associated with comorbid anxiety and depression were being a student receiving less than one dollar equivalent per day as allowance [AOR = 1.953, [95% C.I = 1.135-3.360] and being a student from the Igbo ethnic group [AOR = 0.533, [95% C.I = 0.333-0.853]. **Conclusion:** the prevalence of psychiatry morbidity and comorbid anxiety and depression was high among medical students in Ekiti State, Nigeria. Periodic medical [mental health] screening for medical students may be appropriate to screen, detect and manage psychiatric comorbidities. This will help to ensure optimal mental health for this group of university students.

Introduction

Medical education encompasses training to become a medical practitioner; or a specialist. It requires mastering large volume of information, a longer duration of training and more intensive method of assessment [1]. Many medical students struggle with their own capacity to meet the demands of the curriculum [2-4]. It is also imperative for them to be balanced psychologically to ensure success in the training and future practice of medicine [5]. Medical schools are known to be stressful environments for students and hence medical students have been believed to experience greater incidences of psychological disorders [6]. Depression and anxiety are among the common mental health problems among medical students and are associated with poor academic performance, disability and poor quality of life. A better understanding of the magnitude and correlates of depression and anxiety is essential for planning appropriate intervention for those population groups. Globally, studies have reported high prevalence of psychiatric morbidity anxiety and depression among medical students [7-9]. The global prevalence of depression amongst medical students was 28.0% [6]. In the developed countries Becker found that 37% of young adults between the ages of 15 to 24 years in the United States of America had a diagnosable mental

disorder, and many of these individuals were college students [10].

In Nigeria a study in Ibadan indicated that 12% of pre-clinical students [10] and 21% of clinical students had psychiatric morbidity [10], while Makanjuola *et al.* reported that 14.7% of the medical students in Ilorin had psychiatric morbidity [11], Oshodi *et al.* reported 42.2% of medical students in Lagos to have psychiatric morbidity [12] while Oku *et al.* reported that 39.2% of medical student in Calabar had poor mental health [13]. Previous research has identified long hours of study, and the impositions of emotional burden [14], high workload [15] and considerable financial pressure as the principal stressors. Omigbodun *et al.* found that the prevalence of psychological distress was be higher among male students. Other socio-demographic factors associated with psychological morbidity include being in a transition year of study, reporting financial distress and not being a 'Pentecostal Christian', excessive school work, congested classrooms, strikes by faculty, lack of laboratory equipment, family problems, insecurity, and financial and health problems. They also noticed that medical and dental students had significantly higher GHQ scores than the physiotherapy and nursing students [16]. There is paucity of data on psychiatric morbidity and comorbid anxiety and depression among students in medical schools in Nigeria especially private medical schools. The study therefore assessed the prevalence and factors associated with psychiatric morbidity, and comorbid anxiety and depression among medical students in Ekiti State, Nigeria.

Methods

Study settings, participants and design: this was a cross sectional study which involved public and private medical schools in Ekiti State southwestern, Nigeria. There are two medical schools in the state; Ekiti State University a public non-residential institution owned by the state government and Afe Babalola University, a private residential university owned by Chief Afe Babalola.

Both universities are located in the State Capital (Ado Ekiti) of Ekiti state. The 2 medical schools in Ekiti State were purposively selected and all 965 students in second year through sixth year in both schools were invited to participate in the study. All second to sixth year-medical students in both universities who gave informed consent were enrolled. Having retrieved the students' class lists, each participant was approached before the lecture hours or the hospital's Ground Round for the students in clinical postings to explain the purpose of the study, give assurance of confidentiality and to explain the benefits of the study. Thereafter, the informed consent form and the self-administered questionnaire were given to each participant. Each questionnaire was checked by the researchers and trained research assistants for adequate completion. A few students were asked to fill identified blank spaces in the questionnaire before final submission. The study spanned from April 2019 to August 2019.

Study instruments

The socio-demographic schedule/proforma: this section contained information on the socio-demographic profiles of respondents. It contained the following items: age, gender, religion, marital status, a self-rated academic performance grouped as follows: -below average [1-4], average (5 and 6) and above average (7-10). The schedule also includes sources of finance, marital status of the parent, family type and parents' level of education and occupation.

The general health questionnaire (GHQ): the David Goldberg designed GHQ: a self-administered screening instrument for detecting non-psychotic psychiatric disorders, focusing on the inability to carry out normal function. The 12-item General Health Questionnaire (GHQ-12) has been extensively used as a short screening instrument, producing results that are comparable to longer versions of the GHQ. In a study, the validity of the GHQ-12 was determined against the Composite International Diagnostic Interview (CIDI). It has a

sensitivity of 68% and specificity of 70% [17, 18]. The 12-item version of the instrument was used in this study. The GHQ-12 is scored using the binary method (0-0-1-1). Each item has four possible responses, typically being 'not at all', 'no more than usual', 'rather more than usual' and 'much more than usual'. Therefore a score of 3 and above will be used as an indication of psychiatric morbidity [19]. In a study by Gureje, the alpha coefficient of the GHQ-12 was 0.82 [18].

Hospital anxiety and depression scale (HADS): the HADS is a self-report instrument efficiently used to assess depression and anxiety [17]. In a Nigerian study, the sensitivity for the anxiety sub-scale ranged from 85.0% to 92.9%, while sensitivity for the depression sub-scale ranged from 89.5% to 92.1% [18]. The HADS is considered to be unaffected by coexisting general medical conditions [19], unlike GHQ items where symptoms may refer to physical cause like insomnia and weight loss. For GHQ 12, using 3 as cut-off point, ≥ 3 is "possible psychiatric morbidity [i.e. 'GHQ 12 cases'] and < 3 as no morbidity [i.e. 'GHQ 12 non-cases']". The Hospital Anxiety and Depression Scale will be scored accordingly 0-7 = normal, 8-10 = borderline abnormal, 11-21 = abnormal. Respondents with borderline abnormal and abnormal cases were considered as having anxiety or depression.

Ethical consideration: ethical approval was obtained from the Research Ethics Committee of the Ekiti State University Teaching Hospital Ethics and Research Committee. [Protocol Number-EKSUTH/A67/2019/01/005] and Afe Babalola University Participation was voluntary and informed consent was obtained from the study participants.

Data Analysis: the Statistical Package for Social Sciences (SPSS version 21) was used for Data analysis. The socio-demographic details of respondents were reported using descriptive statistics. Categorical variables were summarized using frequencies and percentages while continuous variables were summarized using

means and standard deviation. Pearson's chi-square test was used to determine the association between comorbid anxiety and depression, psychiatric morbidity and socio-demographic variables. The independent variables entered into the binary logistic regression model were those that were significant at 5% ($p < 0.05$) on bivariate analysis and other factors that have been found to be significant predictors from literature (e.g. age and sex) [20-22]. Confidence interval was set at 95% and all tests were two-tailed. Statistical significance was considered at a p-value less than 0.05.

Results

There were 965 eligible medical students in both institutions, out of which 944 (97.8%) of them participated in the study. Five hundred and twelve (54.2%) of the respondents were aged 10-20 years while 432 (45.8%) were aged 21 years and above. The age range of the medical students was between 16 and 32 years, overall mean was 21 ± 3.0 years, with mean age of 19.8 ± 1.8 years and 25.25 ± 2.7 years in the Private and Public institutions respectively. Most of the respondents (78.4%) attend the private tertiary institution. Female, Christian, and single were 68.0%, 81% and 95.6% respectively. The majority, (94.0%) are financed by their parents, over half (51.2%) of the respondents were from Yoruba ethnicity, 449 (47.6%) were from the Igbo ethnicity and 12 (1.2%) were Hausas. Five hundred and forty-six (57.8%) were in pre-clinical stage (second and third year medical students) of the medical training while 398 (42.2%) were in the clinical years. Five hundred and twenty-five (55.6%) described themselves as academically average, 347 (36.8%) above average and 72 (7.6%) below average. About one-tenth (100) spend equivalent of one dollar or less per day while the rest 884 (89.4%) spend more than one dollar daily. Six hundred and forty-five (68.3%) were satisfied with monthly allowance. A significant proportion (91.8%) were from monogamous background, With regard to the respondents' parents highest

level of formal education; 853 (90.4%) of their mothers had highest level of education being tertiary education while the rest 91(9.6%) as no tertiary education (none, primary and secondary education). Eight hundred and seventy four (92.6%) of their fathers had highest level of education being tertiary education while the rest 70 (7.4%) had no tertiary education.

Prevalence of psychiatric morbidity and comorbid anxiety and depression of the respondents: the overall prevalence of psychiatric morbidity and comorbid anxiety and depression among the respondents was 25.0% (21.1-27.8) and 14.3% (12.1-16.5) respectively. The prevalence of psychiatric morbidity among respondents from the public institution was 11.8% (7.5-16.6) while it was 28.6% (25.3-32.2) among respondents from the private institution. The prevalence of comorbid anxiety and depression among respondents from public and private institution was 27% (20.8-33.2) and 10.8% (8.6-13.2) respectively.

Association of comorbid anxiety and depression with socio-demographic characteristics of the respondents: Table 1 shows comparison between 135 subjects with anxiety and depression and 809 without. More participants aged 21 years and above (59.3%) had comorbid anxiety and depression compared to those below 21 years, this difference was statistically significant [$p = 0.001$]. A higher proportion of medical students in public university had comorbid anxiety and depression compared to those in the private tertiary institution, this was statistically significant [$p < 0.001$]. A significant number who were Muslims had comorbid anxiety and depression compared to others of other religion, this was also statistically significant [$p = 0.004$]. Respondents who were of the Igbo ethnic group were less frequently affected by anxiety and depression compared to other ethnicities, this difference was statistically significant [$p < 0.001$]. More respondents who described themselves as being below average academically were more likely to have had anxiety and depression compared to others, this difference was statistically significant [$P < 0.001$].

Respondent that spent one dollar or less per day had significant comorbid anxiety and depression than those with more than one dollar daily [< 0.001].

Association of psychiatric morbidity with socio demographic characteristics of the respondents: Table 2 shows comparison between 212 subjects with psychiatric morbidity and 528 without. A significant proportion who attends private tertiary institutions (28.6%) had psychiatric morbidity compared to respondents in a public institution, difference was statistically significant [$p < 0.001$]. One hundred and seventy-five (27.3%) female respondents had psychiatric morbidity compared with 61 (20.2%) male respondents, this was statistically significant [$p = 0.020$]. Traditional worshippers (50.0%) had statistically significant psychiatric morbidity compared to other religions [$p < 0.001$]. Hausa respondents were more likely to be affected by psychiatric morbidity, this difference was statistically significant [$p = 0.013$]. Additionally, more respondents who described themselves as being below average in academic performance had significant psychiatric morbidity than those who described themselves as being average or above average academically. This was statistically significant [$p < 0.001$]. One hundred and four (34.8%) respondents, not satisfied with their monthly allowance had psychiatric morbidity compared to one hundred and thirty-two (20.5%) who were satisfied, this was statistically significant [$p \text{ value} < 0.001$]. Respondents with parents (father or mother) with highest level of formal education as tertiary education were less frequently affected by psychiatric morbidity compared to those with parents with no tertiary education, the difference was statistically significant, [$p \text{ value} < 0.001$ and $p = 0.001$ respectively]. Respondents who were not satisfied with their monthly allowance were more likely to be affected by psychiatry morbidity, the difference was also statistically significant [$p < 0.001$].

Socio-demographic and clinical variables independently associated with psychiatric morbidity by logistic regression analysis: the

factors independently associated with psychiatric morbidity included academic performance below average and average which was about 2.4 folds and 1.7folds respectively increased odd of psychiatric morbidity [Adjusted odds ratio [AOR] =2.425, [95% confidence interval [CI] =1.313-4.478] and [AOR] =1.711, [95% C.I = 1.173-2.496] respectively. Besides, being a student at a private institution has about 6.5 odd increase of psychiatric morbidity compared with their counterparts in the public institution [AOR = 6.533, [95% C.I = 3.298-12.940]. Students whose fathers didn't have tertiary education [AOR = 3.147, [95% C.I = 1.579-6.272], students whose mothers didn't have tertiary education [AOR = 2.053, [95% C.I = 1.074-3.927] and students who were not satisfied with their monthly allowance [AOR =2.2135, [95% C.I = 1.495-3.049] were more likely to have psychiatric morbidity. However, Muslims were less likely to have psychiatric morbidity [AOR = 0.348, [95% C.I = 0.194-0.626] (Table 3).

Association between comorbid anxiety and depression and other variables by logistic regression analysis: medical students who spent equivalent of less than one dollar per day were approximately 2 times more likely to report symptoms of comorbid anxiety and depression [AOR] =1.953, [95% C.I = 1.135-3.360] while respondents of Igbo ethnic groups were less likely to report symptoms of anxiety and depression [AOR] = 0.533, [95% C.I = 0.333-0.853] (Table 4).

Discussion

The mean age of the medical students in Ekiti State was 20.9 years, those in the public institution were 5.45 years older than those in the private institution. Most of the respondents were in pre-clinical years and less than 20 years. The mean age of the medical students in Ekiti State is lower than 26.9 years and 23.91 years reported by previous studies in the public institutions [23,24]. The mean age of medical students in the public institution in this study is however similar to previous reports. There is a dearth of study among students in private institutions. More females

were found in this study compared to earlier reports for public schools [23,25,26]. This may be a new trend in gender distribution or due to the recent national campaign and advocacy for girl child education. Most of the respondents were single probably due to the age range, high parental literacy level and tradition i.e the Yorubas prefers females to complete first degrees before marriage to honour the parents. The socio-economic instability of the nation and parental viewpoint may be responsible for dissatisfaction with monthly allowances or dissatisfaction may be from comparison with fellow rich medical students. Poor remuneration from parents may be responsible for spending less than one dollar per day. Most of the respondents were average students. Since 50% score is required for tests and examinations, most students will therefore describe themselves as average. This observation is similar to a previous report in southwestern Nigeria [27].

The prevalence of psychiatric morbidity among the medical students in this study was 25%, a potential for undesirable consequences now and in future for the students [28]. A prevalence of 11.8% in this study among medical students in the public institution is comparable with 12.0% reported among preclinical students but lower than 21.0% and 45% among clinical students in two public medical institutions, in Ibadan and Lagos, Nigeria [14-16]. These two towns are cosmopolitan cities when compared with this study location. Other public medical institutions in Nigeria reported high prevalences [29, 30]. The prevalence of psychiatric morbidity among medical students in the private institution was 28.6% and this is higher than the 11.8% reported in the public institution. It is lower than 46.2% reported in a private medical institution in Malaysia [31]. There is paucity of studies among medical students in private institutions in Nigeria, this is probably because many private institutions may think that these reports can tarnish the image of their university.

Factors such as academic performance, religion, satisfaction with monthly allowance and institutional type were individual predictors of psychiatric morbidity identified in this study. Medical education requires mastering of tremendous amount of information, spending more years in training, scoring higher grades with ability to recall information from previous courses [32]. Inability to handle necessary information may result in struggles with the medical curriculum and psychiatric morbidity [16]. Respondents who described themselves as average and below-average were more likely to report symptoms of psychiatric morbidity [33]. Religion is an important concept in mental health commonly used as coping strategy [27]. In this study Muslims were less likely to report symptoms of psychiatric morbidity, probably due to an underlining moral necessity for protection and care of the vulnerable individuals inherent within it [34]. Parental education influences children's manner, actions and motivation. In this study, respondents whose parents highest level of formal education were below first degree were more likely to be affected by psychiatric morbidity like in earlier studies [35]. The level of education may affect the rate understanding, relationship and financial capacity. Parents who do not have tertiary education are less likely to understand what it takes to be a doctor even emotionally.

In this study respondents not satisfied with their monthly allowances were more likely to report symptoms of psychiatric morbidity. This non-satisfaction may be relative to the student's lifestyle, ostentatious background, future expectations and other psychosocial factors. Financial satisfaction may be improved with holiday jobs, scholarships, management skills training and subsidized tuition. Respondents in the private institution were more likely to report symptoms of psychiatric morbidity in this study. Medical students are relatively younger in private institution than in public university. College life for teenagers may impose major challenges to studying and socializing and failure to manage these challenges may lead to poor mental health

with repercussions [36]. Private universities have stricter rules, restriction of movement and freedom and stiffer punishments for disobedience. Private universities in Nigeria are very expensive with a myriad of fees. The whole process of raising funds for school may be stressful for both parents and students. This may be a source of psychiatric morbidity.

In this study the prevalence of comorbid anxiety and depression among the respondents was 14.3%, it was 27.0% in public institution against 10.8% in private institution. Medical students have been found to experience higher levels of depression and anxiety compared to other students or the general population [4]. In Pakistan, a study conducted among private medical students found prevalence of anxiety and depression using the Aga Khan University Anxiety and Depression Scale [AKUADS] of 4th year, 3rd year, 2nd year and 1st year to be 49%, 47%, 73%, and 66% respectively [37]. A higher prevalence of anxiety and depression [43.89%] was found amongst students of Nishtar Medical College, Multan with prevalence of anxiety and depression specifically found to be 45.86%, 52.58%, 47.14%, 28.75% and 45.10% among students of first, second, third, fourth and final years respectively [38]. The consensus from all the previous studies, responsible for high prevalence of anxiety and depression among private medical students were competitiveness and demand for high grades thus students are pushed to work harder. This may be to justify their high tuition and to attract future potential students. The majority of medical students have financial indebtedness to school and financiers. There are also poor leisure activities and exercise, especially for fresher and final year students. These may be due to pressure from parents as a result of higher tuition [39]. Additionally, high parental expectations also cause undue stress [40].

In this study, more respondents from public institution reported symptoms of comorbid anxiety and depression. Tuition fees are lower in public institutions than in private institutions,

however incessant industrial actions relating to staff salaries and infrastructure issues prolong students' stay in school. The vast majority of students in public universities often come from low socio-economic backgrounds, may be exposed to chronic poverty with large families. Respondents who spend less than one dollar equivalent per day in this study were more likely to be affected by comorbid anxiety and depression. Medical education is financially demanding, non-affordability of basic needs from living below the WHO poverty line is stressful and a template for comorbid anxiety and depression. Respondents from the Igbo tribe in this study were less likely to have comorbid anxiety and depression than the Yoruba tribe. The Igbos are predominantly businessmen and women who may have the financial strength to make their wards comfortable in the school.

Conclusion

Comorbid depression and anxiety, and psychiatric morbidity are common among Nigeria medical students, both in private and public institutions. In this study, a variety of predictors as well as protective factors were identified. An effective model for prevention and early detection of depression, anxiety and psychiatric morbidity among medical students need to be developed. Policymakers, school administrators, government, parents, academic governing body and students should work together to reduce these high prevalence rates to improve the mental health of our future doctors.

What is known about this topic

- *Prevalence of psychiatric morbidity anxiety and depression is high among medical students globally;*
- *Long hours of study, high workload, the impositions of emotional burden and considerable financial pressure are principal stressors.*

What this study adds

- *The prevalence of comorbid anxiety and depression is higher among students in public medical school than private medical school;*
- *The factors independently associated with psychiatric morbidity included being a student with self-reported description of being below and average academically, and being a student in a private institution;*
- *The factor independently associated with comorbid anxiety and depression was being a student receiving less than one dollar equivalent per day.*

Competing interests

The authors declare no competing interests.

Authors' contributions

Falade Joshua conceived the research, analysed the data and wrote the first draft of the manuscript. Adedayo Hakeem Oyebanji supervised the project, methods and proofread the manuscript. Adefunke Olarinre Babatola supervised the project, methods and proofread the manuscript. Olusola Olawumi Falade supervised the project, methods and proofread the manuscript. Temitope Olumuyiwa Ojo analysed the data and proofread the manuscript. All the authors read and approved the final version of the manuscript.

Acknowledgments

We acknowledge the participants, research assistance and the school authorities for the opportunity to carry out the study.

Tables

Table 1: association of comorbid anxiety and depression with socio-demographic characteristics of the respondents

Table 2: association of psychiatric morbidity with socio-demographic characteristics of the respondents

Table 3: the socio-demographic and clinical variables independently associated with psychiatric morbidity by logistic regression analysis

Table 4: association between comorbid anxiety and depression and other variables using logistic regression

References

1. Malu A. Universities and medical education in Nigeria. *Nigerian Medical Journal*. 2010;51(2): 84. **Google Scholar**
2. Caverzagie KJ, Nousiainen MT, Ferguson PC, Ten Cate O, Ross S, Harris KA *et al*. Overarching challenges to the implementation of competency-based medical education. *Med Teach*. 2017;39(6): 588-93. **PubMed** | **Google Scholar**
3. Singla M, Sharma A, Aggarwal N. To compare psychological morbidity among students appearing for final professional MBBS examination and civil services examination. *Indian Journal of Positive Psychology*. 2018;9(4): 436-42. **Google Scholar**
4. Tabalipa FdO, Souza MFd, Pfützenreuter G, Lima VC, Traebert E, Traebert J. Prevalence of anxiety and depression among medical students. *Revista Brasileira de Educação Médica*. 2015;39 (3): 388-94. **Google Scholar**
5. Bhugra D, Sauerteig S-O, Bland D, Lloyd-Kendall A, Wijesuriya J, Singh G *et al*. A descriptive study of mental health and wellbeing of doctors and medical students in the UK. *Int Rev Psychiatry*. Nov-Dec 2019;31(7-8): 563-568. **PubMed** | **Google Scholar**
6. Puthran R, Zhang MW, Tam WW, Ho RC. Prevalence of depression amongst medical students: A meta-analysis. *Med Educ*. 2016;50 (4): 456-68. **PubMed** | **Google Scholar**
7. Al-Busaidi Z, Bhargava K, Al-Ismaïly A, Al-Lawati H, Al-Kindi R, Al-Shafae M *et al*. Prevalence of depressive symptoms among university students in Oman. *Oman Med J*. 2011 Jul;26(4): 235-9. **PubMed** | **Google Scholar**
8. Chatterjee C, Mandal PK, Mallik S, Manna N, Sardar JC, Dasgupta S. A study on mental distress among MBBS students in a Medical College, Kolkata, India. *Annals of Tropical Medicine and Public Health*. 2012;5 (5): 453. **Google Scholar**
9. Zaki N, Ibrahim J. Psychiatric morbidity among third year medical students at the Ain Shams University, Cairo, Egypt. *Middle East Current Psychiatry*. 2011;18 (1): 51-6. **Google Scholar**
10. Becker M, Martin L, Wajeer E, Ward J, Shern D. Students mental illnesses in university setting: faculty and student attitudes, beliefs, knowledge, and experiences. *Psychiatr Rehabil J*. Spring 2002;25(4): 359-68. **PubMed** | **Google Scholar**
11. Omokhodion F. Psychosocial problems of pre-clinical students in the University of Ibadan Medical School. *Afr J Med Med Sci*. 2003 Jun;32(2): 135-8. **PubMed** | **Google Scholar**
12. Makanjuola A, Abiodun O, Sajo S. Alcohol and psychoactive substance use among medical students of the University of Ilorin, Nigeria. *European Scientific Journal*. 2014;10(8).
13. Oshodi O, Erinfolami A, Akinbode A. Problematic Internet Usage Among Students In a Medical School In Lagos, Nigeria. *Nigerian Journal of Psychiatry*. 2012;10(1): 48-55. **Google Scholar**
14. Oku A, Oku O, Owoaje E, Monjok E. An assessment of mental health status of undergraduate medical trainees in the University of Calabar, Nigeria: a cross-sectional study. *Open Access Maced J Med Sci*. 2015 Jun 15;3(2): 356-62. **PubMed** | **Google Scholar**
15. Rosenthal JM, Okie S. White coat, mood indigo-depression in medical school. *N Engl J Med*. 2005 Sep 15;353(11): 1085-8. **PubMed** | **Google Scholar**
16. Adams J. Straining to describe and tackle stress in medical students. *Med Educ*. 2004;38(5): 463-4. **PubMed** | **Google Scholar**

17. Omigbodun OO, Odukogbe A-TA, Omigbodun AO, Yusuf OB, Bella TT, Olayemi O. Stressors and psychological symptoms in students of medicine and allied health professions in Nigeria. *Soc Psychiatry Psychiatr Epidemiol.* 2006;41(5): 415-21. **PubMed | Google Scholar**
18. Gureje O, Obikoya B. The GHQ-12 as a screening tool in a primary care setting. *Soc Psychiatry Psychiatr Epidemiol.* 1990 Sep;25(5): 276-80. **PubMed | Google Scholar**
19. Yusuf AD, Issa BA, Ajiboye PO, Buhari OIN. The correlates of stress coping styles and psychiatry morbidity in the first year of Medical education at a Nigerian University. *Afr J Psychiatry (Johannesbg).* 2013 May;16(3): 206-15. **PubMed | Google Scholar**
20. Giusti EM, Jonkman A, Manzoni GM, Castelnuovo G, Terwee CB, Roorda LD *et al.* Proposal for Improvement of the Hospital Anxiety and Depression Scale for the Assessment of Emotional Distress in Patients With Chronic Musculoskeletal Pain: A Bifactor and Item Response Theory Analysis. *J Pain.* Mar-Apr 2020;21(3-4): 375-389. **PubMed | Google Scholar**
21. Abiodun O. A validity study of the Hospital Anxiety and Depression Scale in general hospital units and a community sample in Nigeria. *Br J Psychiatry.* 1994 Nov;165(5): 669-72. **PubMed | Google Scholar**
22. Snaith R. The concepts of mild depression. *Br J Psychiatry.* 1987 Mar;150: 387-93. **PubMed | Google Scholar**
23. James BO, Thomas IF, Omoaregba JO, Okogbenin EO, Okonoda KM, Ibrahim AW *et al.* Psychosocial correlates of perceived stress among undergraduate medical students in Nigeria. *Int J Med Educ.* 2017 Oct 26;8: 382-388. **PubMed | Google Scholar**
24. Okoye O, Maduka-Okafor F, Eze B. What does the medical student know about eye donation/corneal transplant? The University of Nigeria scenario. *West Indian Med J.* 2010 Jan;59(1): 41-4. **PubMed | Google Scholar**
25. Ngasa SN, Sama C-B, Dzekem BS, Nforchu KN, Tindong M, Aroke D *et al.* Prevalence and factors associated with depression among medical students in Cameroon: a cross-sectional study. *BMC psychiatry.* 2017;17(1): 216. **PubMed | Google Scholar**
26. Opakunle T, Aloba O, Opakunle O, Oyewole A, Osokoya O. Prevalence and correlates of obsessive-compulsive symptoms in a sample of undergraduate clinical medical students in Osogbo, Southwestern Nigeria. *Nigerian Journal of Health Sciences.* 2017;17(2): 66. **Google Scholar**
27. Falade J, Eegunranti B, Tobih J, Ogundiran A, Ibigbami O, Akinsulore A *et al.* Prevalence, Pattern and Correlates of Study Difficulty among Students of College Of Health Sciences Ladoke Akintola University of Technology Osogbo. *Advances in Social Sciences Research Journal.* 2019;6(9): 322-38. **Google Scholar**
28. Zeng W, Chen R, Wang X, Zhang Q, Deng W. Prevalence of mental health problems among medical students in China: A meta-analysis. *Medicine (Baltimore).* 2019 May;98(18): e15337. **PubMed | Google Scholar**
29. Imo UO. Burnout and psychiatric morbidity among doctors in the UK: a systematic literature review of prevalence and associated factors. *BJPsych Bull.* 2017;41(4): 197-204. **PubMed | Google Scholar**
30. Yussuf A, Issa B, Ajiboye P, Buhari O. The correlates of stress, coping styles and psychiatric morbidity in the first year of medical education at a Nigerian University. *Afr J Psychiatry (Johannesbg).* 2013 May;16(3): 206-15. **PubMed | Google Scholar**
31. Zaid Z, Chan S, Ho J. Emotional disorders among medical students in a Malaysian private medical school. *Singapore Med J.* 2007 Oct;48(10): 895-9. **PubMed | Google Scholar**
32. Rotenstein LS, Ramos MA, Torre M, Segal JB, Peluso MJ, Guille C *et al.* Prevalence of depression, depressive symptoms, and suicidal ideation among medical students: a systematic review and meta-analysis. *JAMA.* 2016 Dec 6;316(21): 2214-2236 **PubMed | Google Scholar**

33. Khan AA, Abid Sultan FAJ. Effects of perceived stressors and psychiatric morbidity on academic performance of university students. *Computer Science*. 2018;389: 125. **Google Scholar**
34. Tzeferakos GA, Douzenis AI. Islam, mental health and law: a general overview. *Ann Gen Psychiatry*. 2017 Jul 6;16: 28. **PubMed | Google Scholar**
35. Benkaakaa A. The Influence of the Socio-Cultural Background of Parents on Child's Education. 2019.
36. bin Othman MNA, Rashid MAA. Stress and Mental Health of Undergraduate Students at a Private Higher Learning Institution in Malaysia. *People: International Journal of Social Sciences*. 2018;4(2). **Google Scholar**
37. Inam S, Saqib A, Alam E. Prevalence of anxiety and depression among medical students of private university. *J Pak Med Assoc*. 2003 Feb;53(2): 44-7. **PubMed | Google Scholar**
38. Jadoon NA, Yaqoob R, Raza A, Shehzad MA, Zeshan SC. Anxiety and depression among medical students: a cross-sectional study. *J Pak Med Assoc*. 2010;60(8): 699-702. **PubMed | Google Scholar**
39. Yusoff MSB, Rahim AFA, Yaacob MJ. Prevalence and sources of stress among Universiti Sains Malaysia medical students. *Malays J Med Sci*. 2010 Jan;17(1): 30-7. **PubMed | Google Scholar**
40. Sreeramareddy CT, Shankar PR, Binu V, Mukhopadhyay C, Ray B, Menezes RG. Psychological morbidity, sources of stress and coping strategies among undergraduate medical students of Nepal. *BMC Med Educ*. 2007 Aug 2;7: 26. **PubMed | Google Scholar**

Table 1: association of comorbid anxiety and depression with socio-demographic characteristics of the respondents			
Variable	Anxiety and depressionabsentn [%]	Anxiety and depression presentn [%]	P-value
Age group (years)			
10-20	457 [56.5]	55 [40.7]	0.001
21 and above	352 [43.5]	80 [59.3]	
Institution type			
Private	660 [81.6]	80 [78.4]	<0.001
Public	149 [18.4]	55 [21.6]	
Gender			
Male	256 [31.6]	46 [34.1]	0.575
Female	553 [68.4]	89 [65.9]	
Marital status			
Single	775 [95.8]	127 [94.1]	0.369
Married	34 [4.2]	8 [5.9]	
Ethnicity			
Yoruba	388 [48.0]	102 [71.8]	<0.001
Igbo	413 [51.0]	36 [25.4]	
Hausa	8 [1.0]	4 [2.8]	
Religion			
Christianity	670 [82.8]	98 [72.6]	0.004
Islam	131 [16.2]	37 [27.4]	
Traditional	8 [1.0]	0 [0.0]	
Stage			
Pre-clinical	476 [58.8]	70 [51.9]	0.128
Clinical	333 [41.2]	65 [48.1]	
Academic performance			
Below average	58 [7.2]	30 [19.9]	<0.001
Average	461 [57.0]	64 [42.4]	
Above average	290 [35.8]	57 [37.8]	
Source of finance			
Parents	762 [94.2]	125 [92.6]	0.602
Other family members	3 [0.4]	1 [0.7]	
Self	18 [2.2]	3 [2.2]	
Spouse	26 [3.2]	6 [4.4]	
Average amount spent			
≤ one dollar per day	69 [8.5]	31 [23.0]	<0.001
> one dollar per day	740 [91.5]	104 [77.0]	
Satisfied with monthly allowance			
Yes	554 [68.5]	91 [67.4]	0.804
No	255 [31.5]	44 [32.6]	
Parents marital status			
Single parent	32 [4.0]	15 [6.5]	0.165
Married	713 [88.1]	206 [88.8]	
Divorce/separated	26 [3.2]	4 [1.7]	
Widow/widower	38 [4.7]	7 [3.0]	
Family type			
Monogamy	747 [92.3]	120 [88.9]	0.176
Polygamy	62 [7.7]	15 [11.1]	
Fathers educational level			
No tertiary education	60 [7.4]	8 [5.9]	0.531
Tertiary education	747 [92.6]	127 [94.1]	
Mothers educational level			
No tertiary education	81 [89.0]	10 [11.0]	0.342
Tertiary education	728 [85.3]	125 [14.7]	

Table 2: association of psychiatric morbidity with socio-demographic characteristics of the respondents

Variable	No Psychiatric morbidityn [%]	Psychiatric morbidity presentn [%]	P value
Institution type			
Private	528 [74.6]	212 [89.8]	<0.001
Public	180 [25.4]	24 [10.2]	
Gender			
Male	241[34.0]	61 [25.9]	0.020
Female	467[66.0]	175[74.2]	
Marital status			
Single	674 [95.2]	228 [96.6]	0.362
Married	34 [4.8]	8 [3.4]	
Ethnicity			
Yoruba	387 [54.2]	102 [43.2]	0.013
Igbo	319 [44.7]	130 [55.1]	
Hausa	8 [1.1]	4 [1.7]	
Religion			
Christianity	554 [78.2]	214 [90.7]	<0.001
Islam	150 [21.2]	18 [7.6]	
Traditional	4 [0.6]	4 [1.7]	
Stage			
Preclinical	419 [59.3]	127 [53.8]	0.136
Clinical	287 [40.7]	109 [46.2]	
Academic performance			
Below average	42 [5.9]	30 [12.7]	<0.001
Average	373 [52.7]	152 [64.4]	
Above average	293 [41.4]	54 [22.9]	
Average monthly allowance			
≤ One dollar per day	82 [11.6]	18 [7.6]	0.087
>One dollar per day	626 [88.4]	218 [92.4]	
Satisfaction with allowance			
Yes	513 [72.5]	132 [55.9]	<0.001
No	195 [27.5]	104 [44.1]	
Parents marital status			
Single parent	37 [5.2]	10 [4.2]	0.794
Married	616 [87.0]	206 [87.3]	
Divorce/separated/widowed	55 [4.2]	20 [0.0]	
Fathers educational level			
No tertiary education	37 [5.2]	33 [14.0]	<0.001
Tertiary education	671 [94.7]	203 [86.0]	
mothers educational level			
No tertiary education	55 [7.8]	36 [15.3]	0.001
Tertiary education	653 [95.2]	200 [84.7]	

Table 3: the socio-demographic and clinical variables independently associated with psychiatric morbidity by logistic regression analysis

Variable	P-value	Odd ratio (95% Confidence Interval)
Tribe		
Yoruba [ref]		
Hausa	0.480	1.646 [0.413-6.561]
Igbo	0.081	0.725 [0.505-1.040]
Religion		
Christianity [ref]		
Islam	<0.001	0.348 [0.194-0.626]
Traditional	0.315	2.163 [0.480-9.744]
Marital status		
Married[ref]		
Single	0.565	1.324 [0.509-3.442]
Institution		
Public[ref]		
Private	<0.001	6.533 [3.298-12.940]
Academic performance		
Above average [ref]		
Below average	0.005	2.425 [1.313-4.478]
Average	0.005	1.711 [1.173-2.496]
Average Satisfaction		
Yes [ref]		
No	<0.001	2.135 [1.495-3.049]
Fathers level of education		
Tertiary education [ref]		
No tertiary education	0.001	3.147 [1.579-6.272]
Mothers level of education		
Tertiary education [ref]		
No tertiary education	0.030	2.053 [1.074-3.927]
Gender		
Female [ref]		
Male	0.241	0.802 [0.555-1.159]
Age (years)		
10-20 [ref]		
21 and above	0.104	0.733 [0.505-1.066]

Table 4: association between comorbid anxiety and depression and other variables using logistic regression

Variable	P-value	Odds ratio (95% Confidence Interval)
Age group (in years)		
10-20 [ref]		
21 and above	0.594	1.138 [0.708-1.830]
Gender		
Male [ref]		
Female	0.745	1.069 [0.714-1.601]
Tribe		
Yoruba[ref]		
Hausa	0.109	2.835 [0.792-10.145]
Igbo	0.009	0.533 [0.333-0.853]
Religion		
Christianity [ref]		
Islam	0.395	1.226 [0.766-1.963]
Traditional	0.999	0.000 [0.000-.]
Marital status		
Married [ref]		
Single	0.805	1.115 [0.468-2.661]
Institution		
Private [ref]		
Public	0.121	1.588 [0.885-2.849]
Monthly allowance		
More than a dollar per day [ref]		
Less than one dollar per day	0.016	1.953 [1.135-3.360]