

Effect of 7,12-Dimethylbenz(α)anthracene on the Expression of miR-330, miR-29a, miR-9-1, miR-9-3 and the mTORC1 Gene in CBA/Ca Mice

ANDRAS TOMESZ^{1,2}, LASZLO SZABO^{1,2}, RICHARD MOLNAR^{1,2}, ARPAD DEUTSCH¹,
RICHARD DARAGO¹, DOMOKOS MATHE³, FERENC BUDAN^{2,4},
NOWRASTEH GHODRATOLLAH², TIMEA VARJAS², BALAZS NEMETH² and ISTVAN KISS²

¹Doctoral School of Health Sciences, Faculty of Health Sciences, University of Pécs, Pécs, Hungary;

²Department of Public Health Medicine, Medical School, University of Pécs, Pécs, Hungary;

³Department of Biophysics and Radiation Biology, Semmelweis University, Budapest, Hungary;

⁴Institute of Environmental Engineering, University of Pannonia, Veszprém, Hungary

Abstract. *Background/Aim: Development of malignant tumors is preceded by molecular biological events. Our aim was to establish an assay panel by using miRNAs and other genes for the rapid screening of potential carcinogens or chemopreventive agents. Materials and Methods: Six male and 6 female CBA/Ca mice received 20 mg/bwkg 7,12-dimethylbenz(α)anthracene (DMBA) intraperitoneally, and 24 h later RNA was isolated from parenchymal organs. Expression of miR-330, miR-29a, miR-9-1, miR-9-3 and mTORC1 was analysed by real time polymerase chain reaction and compared to non-treated controls. Results: DMBA caused significant alterations in the expression of the studied genes. The most profound changes were the strongly elevated miR-9-3 and mTORC1 expressions in female mice in all organs studied. Conclusion: miR-9-3 and mTORC1 expression in female mice were found to be the most suitable biomarkers for rapid identification of possible carcinogenic effects.*

According to estimates, in 2018 there were 18.1 million new malignant cancer cases and 9.6 million cancer deaths (1). According to WHO data, 30-50 % of all malignant tumors could be prevented (2), and early diagnosis would usually increase the chance of successful treatment.

This article is freely accessible online.

Correspondence to: András Tomesz, Department of Public Health Medicine, Medical School, University Of Pécs, H-7624 Pécs, Szigeti Street 12, Hungary. Tel: +36 207772812, e-mail: tomesza@gmail.com

Key Words: Gene expression, microRNA, carcinogenicity, 7,12-dimethylbenz(α)anthracene, DMBA, miR-330, miR-29a, miR-9-1, miR-9-3, mTORC1.

The development of most malignant tumors is preceded by molecular biological events such as changes in gene expression patterns. During this period clinical symptoms do not appear, so, it is difficult to identify any signs of the carcinogenic process. Identification of tumor development based on mutations, chromosome aberrations and/or the phenotypic appearance of the disease is costly, time-consuming and often delayed. Therefore, development of comprehensive biomarker assays to detect the harmful effects of carcinogenic substances as early as possible is of great interest.

Conventional carcinogenicity examination methods - like the Ames test, that is applicable to detect mutations - are based on genotoxicity and show mutagenicity at chromosome or gene levels (3). However, changes in gene expression can signal damages caused by a much wider range of carcinogenic compounds (as well as by ionizing radiation) both in cell cultures and *in vivo* animal models. Within this, mapping of the effects of microRNAs (miRNAs) provides the possibility to recognize details of relevant regulatory pathways and the functions of mutually regulated miRNAs, as well as the modifications and the physiological processes reflected by their patterns of expression, even as biomarkers. Biomarkers providing early (24-48 h) identification of possibly tumorigenic agents may play an important role in the deeper understanding of carcinogenic processes. Finally, using them as tools for primary, secondary and tertiary prevention they can open up new possibilities in further reduction of cancer incidence or mortality rates.

The miRNAs are small, 19-22 nucleotide length single stranded RNA molecules. *Via* their binding to the corresponding 3' UTR of mRNAs, they influence the translation process and protein synthesis. miRNAs regulate the expression of approximately 30-50% of the human genes (4, 5), thereby influencing important biological processes

like cell cycle (6), differentiation or apoptosis (7). It is noteworthy, that many miRNAs are present in fragile genomic regions and in malignant tissues they mostly display altered expression patterns (8, 9). Certain miRNAs are direct targets of oncoproteins or tumor suppressor proteins, for example the *p53* tumor suppressor protein regulates the expression of *miR-34* through a *p53*-responsive element upstream of the *miR-34* locus (10, 11). Others directly regulate the expression of oncogenes and/or tumor suppressor genes, such as *miR-199a* that regulates the *MET* proto-oncogene (12). Furthermore, several microRNAs, e.g. *miR-134*, are involved in the proliferation of tumor cells, in apoptosis, invasion and metastasis (13). Evidence based on experimental studies and clinical histological sample analysis suggests that miRNAs in the serum are relatively stable and could serve as potential biomarkers for the detection of different tumors and other diseases (14).

A wide variety of potentially carcinogenic and anti-carcinogenic agents are known. 7,12-dimethylbenz(α) anthracene (DMBA) is a commonly occurring environmental polycyclic aromatic hydrocarbon that, acting as a mutagen, is capable of exerting a carcinogenic effect, that can be tested both in cell culture and *in vivo* animal models (15, 16). DMBA is activated by *cytochrome P450* metabolizing enzymes (17, 18), and then it may participate in the early stages of carcinogenesis, *i.e.* in the initiation and the promotion (19). Previous studies have demonstrated that DMBA affects the expression of *H-RAS*, *C-MYC* and *TP53* genes even 24-48 h after treatment, which can thus be used as early biomarkers (20).

Our aim was to establish an assay panel by using miRNAs and other genes, that can be used for the rapid screening of potential carcinogens, or anticarcinogenic substances. Therefore, in the present study experimental animals were exposed to DMBA and the expression levels of *miR-330*, *miR-29a*, *miR-9-1*, *miR-9-3* were examined. These miRNAs have already been reported to exhibit altered expression in several malignant tumors and through different signaling pathways they play a role in the regulation of oncogenes or tumor suppressor genes, cell growth, proliferation, invasion, migration, metastasis or apoptosis (21-26). Besides the mentioned microRNAs, we also examined the expression of the *mTORC1* gene. Activating mutations in *mTORC1* have been identified in a wide range of human cancers and as one of the central regulators of cell growth, it has a key role in the development of malignant tumors, cardiovascular diseases, obesity or diabetes (27, 28). Our hypothesis was that the altered expression patterns of the studied miRNAs and *mTORC1*, as biomarkers, may indicate early damages caused by DMBA. In order to explore this, parenchymal organs of CBA/CA mice, *i.e.* the spleen, liver and kidneys were examined after exposure to DMBA. Thus, we can further understand the supposed roles of these miRNAs and *mTORC1* in tumorigenesis and we can evaluate their potential

use as early biomarkers by developing an assay panel for identification of early signs of carcinogenic processes.

Materials and Methods

During the test we used two groups of 6-8 weeks old CBA/Ca mice. Both the control and the DMBA treated group consisted of 12 mice (6 males and 6 females). The DMBA treated group received 20 mg/kg/bw DMBA intraperitoneally dissolved in 0.1 ml corn oil (Sigma), while the control group received 0.1 ml corn oil only. Following a 24 h exposure to DMBA, the mice were sacrificed by cervical dislocation, the liver, kidneys and spleen were removed and RNA was isolated from them as described below. Mice received humane care in accordance with the appropriate guidelines concerning laboratory animals, and the experiment was conducted according to the current ethical permission (Ethical permission no.: BA02/2000-79/2017).

RNA isolation. Total cellular RNA was isolated using TRIZOL reagent (Thermo Fisher Scientific, Waltham, MA, USA) according to the manufacturer's instructions. The quality of RNA was examined by NanoDrop absorption photometry and for RT-PCR process only RNA fractions with $A_{260/280} > 2.0$ were used.

Reverse transcription polymerase chain reaction (RT-PCR). One-step PCR, including reverse transcription and target amplification, was performed using Kapa SYBR FAST One-step qPCR kit (Kapa Biosystems, Wilmington, MA, USA) in a 96-well plate, on a LightCycler 480 qPCR (Roche Diagnostics, Indianapolis, IN, USA) platform.

The temperature program was set as follows: 5 minutes incubation at 42°C followed by a 3 min incubation at 95°C, then 45 cycles were performed (95°C – 5 s, 56°C – 15 s, 72°C – 5 s) and a fluorescent reading was made at the end of each cycle. In each run melting curve analysis (95°C – 5 s, 65°C – 60 s, 97°C ∞) was performed to strengthen the specificity of the amplification. The reaction mixture was included: 10 μ l KAPA SYBR FASTqPCR Master Mix, 0.4 μ l KAPA RT Mix, 0.4 dUTP, 0.4 μ l primers, 5 μ l miRNA template supplemented with sterile double-distilled water to a total volume of 20 μ l.

Primer sequences of the *mTORC1* gene and of the examined miRNAs (*miR-330*, *miR-29a*, *miR-9-1*, *miR-9-3*) and the internal control (mouse *U6* gene) are summarized in Table I. Primers were synthesized by Integrated DNA Technologies (Leuven, Belgium), sequences are from previous publications (29, 30).



Calculations and statistical analysis. Relative miRNA expression levels were calculated and compared using the $2^{-\Delta\Delta CT}$ method. During the statistical analysis distributions and variances were tested using the Kolmogorov-Smirnov Levene's test, and T-probe to compare averages. IBM SPSS 21 (Armonk, NY, USA) statistical software was used for calculations and analysis. We determined the level of statistical significance at p -value < 0.05.

Results

Changes in miRNA and mTORC1 gene expression in the liver after DMBA treatment. Twenty-four hours after treatment, DMBA-induced changes could be observed in the expression of the examined miRNAs and the *mTORC1* gene.

Table I. Primer sequences (5'-3') of the *mTORC1* gene, the examined miRs (*miR-330*, *miR-29a*, *miR-9-1*, *miR-9-3*) and the internal control gene (mouse *U6*).

	Forward	Reverse
miR-330	GACCCTTTGGCGATCTCTG	CTGTGCTTTGCTCGTTGGAT
miR-29a	CCCCTTAGAGGATGACTGATTC	AACCGATTTCAGATGGTGCT
miR-9-1	CGGGGTTGGTTGTTATCTTT	TGGGGTTATTTTACTTTCGGTTA
miR-9-3	GCCCGTTTCTCTCTTTGGTT	TCTAGCTTTATGACGGCTCTGTGG
<i>mTORC1</i>	AAGGCCTGATGGGATTTGG	TGTCAAGTACACGGGGCAAG
mouse <i>U6</i>	CGCTTCGGCAGCACATATAC	TTCACGAATTTGCGTGTCAAT

Figure 1. Gene expression results after 24 h in the liver of females compared to non-treated mice. In case of *miR-29a* and *mTORC1*, $p < 0.05$, as well as in *miR-9-1* and *miR-9-3*, $p < 0.001$. * $p < 0.05$, *** $p < 0.001$.Figure 2. Gene expression results after 24 h in the liver of males compared to non-treated mice. In case of *miR-330*, *miR-9-1* and *miR-9-3*, $p < 0.05$. * $p < 0.05$.

In comparison with the control group, we found statistically significant ($p < 0.05$) changes in the expression of *miR-29a*, *miR-9-1*, *miR-9-3* and *mTORC1* in the liver of DMBA-treated female mice. DMBA treatment resulted in an increase in the expression of *miR-29a*, *miR-9-3* and *mTORC1*, and in a decrease in the expression of *miR-9-1*, as it is shown in Figure 1. Compared to controls the biggest difference was observed for *miR-9-3* (329%).

In males, the changes in *miR-330*, *miR-9-1* and *miR-9-3* were statistically significant ($p < 0.05$). While *miR-330* showed an increased expression, *miR-9-1* and *miR-9-3* showed decreased expression (Figure 2). Compared to controls the biggest difference was observed for *miR-330* (184%).

Changes in miRNA and mTORC1 gene expression in the spleen following DMBA treatment. In the spleen of female mice, DMBA induced a statistically significant change in the expression ($p < 0.05$) of *miR-9-3* and *mTORC1* genes compared with the control group (Figure 3).

As it is shown in Figure 3, both the examined miRNAs and *mTORC1* showed an increased expression as the result

of DMBA treatment, and the biggest difference was found in *miR-9-3* (130%).

The results on the expression of *miR-330* and *miR-29a* were statistically significant in the spleen of males ($p < 0.05$). While *miR-330* showed increased expression, *miR-29a* showed decreased expression upon DMBA treatment (Figure 4). The strongest effect was observed for *miR-330* (456%).

Changes in miRNA and mTORC1 gene expression in the kidney following DMBA treatment. Figure 5 illustrates that DMBA treatment lead to statistically significant changes in the expression ($p < 0.05$) of *mTORC1* and *miR-330*, *miR-9-1* and *miR-29a* in the kidney of females. We found an elevated expression of both miRNAs and *mTORC1* as a result of DMBA treatment, compared with the control group. The difference was particularly strong in the case of *miR-9-1* (332%).

With the exception of *miR-29a*, decreased expressions were observed in the kidney of males, particularly *miR-9-1* where the difference was the highest (-88%). However,




Figure 3. Gene expression results after 24 h in the spleen of females compared to non-treated mice. In case of miR-9-3 and mTORC1, $p < 0.001$. *** $p < 0.001$.




Figure 4. Gene expression results after 24 h in the spleen of males compared to non-treated mice. In case of miR-330, $p < 0.05$, while in case of miR-29a, $p < 0.001$. * $p < 0.05$, ** $p < 0.001$.

statistically significant results were found only in the cases of miR-330 and miR-9-1 (Figure 6).

Discussion

miR-330: As a result of DMBA treatment a smaller increase in the expression of miR-330 tumor suppressor was observed in the spleen and liver of female mice than in male mice. This may be due to the high expression of the proapoptotic *Phosphatase and tensin homolog (PTEN)* tumor suppressor gene in the absence of testosterone (31), the gene product of which reduces the oncogenic effect of MYC proteins highly expressed due to DMBA treatment (19). The decrease in the oncogenic effect of the MYC protein family is known regarding the *Inhibitor of growth protein 4 (ING4)* (32), however, miR-330 inhibits the expression of ING4 gene and – from this point of view – it may even have a tumor promoting effect (33).

The increase in the expression of miR-330 tumor suppressor, measured in the kidney of female mice, is consistent with the observation, that it counteracts the estrogen-induced risk increase for renal cell carcinoma (RCC) (34). However, it is known that in DMBA-induced hypercholesterolemia (35) the levels of low-density lipoprotein (LDL) cholesterol increase, which may consequently lead to decreased expression of the *LDL receptor gene (LDLR)*, whereas LDLR and miR-330 cross-regulate each other's expression (36).

Based on our results and literature data, it can be concluded that the expression levels of miR-330 could be used as a biomarker mainly in male animals. However, according to literature data due to the multifaceted effects of sex hormones and because of the regulation by a negative feedback mechanism, the use of miR-330 expression as a biomarker is questionable.

miR-29a: DMBA increases the levels of *transforming growth factor beta 1 (TGF-β1)* which in turn, decreases the expression of miR-29a (37, 38). However, in addition to the oncogenic effect of TGF-β1, the opposite effect (observed e.g. in hepatocytes) as tumor suppressor has also been described (39, 40). For example, Zhang *et al.* have found in a mouse model that spontaneous tumor formation occurs in the absence of TGF-β1 (41). The various manifestations of the effects of TGF-β1 are explained by the ability of miRNAs to regulate TGF-β1 levels (42), which in the case of miR-29a tumor suppressor presumably occurs in the form of mutual cross-regulation. This shows that regulatory factors other than miR-29a may have a stronger effect on the TGF-β1-mediated effects of DMBA.

According to data from the literature, miR-29a is involved in a wide range of cellular biological processes which are still not fully explored. Considering that a significant decrease in miR-29a expression was observed only in the spleen of males, and that the mechanism of regulation is complex, apparently contradictory and is still unexplored, the expression of miR-29a, as a biomarker, is not relevant in the animal model we describe in this article.

miR-9-1 and miR-9-3: According to literature data, miR-9 inhibits the progression of hepatocellular carcinoma (HCC) as tumor suppressor, and furthermore, the expression of miR-9-1 has been shown to be significantly reduced in HCC tissues and negatively correlated with the overall survival of HCC (43, 44). It is known, that silencing of miR-9-1 enhances the expression of oncogenes induced by the RUNX1-RUNXIT1 transcription factor complex regulated by miR-9-1 (23). Furthermore, the miR-9 decrease has been shown to be an early biomarker for the development of various malignancies, such as breast cancer (45). Acting as transcription factors, MYC and MYCN oncoproteins directly bind to a MYC binding site in the miR-9-




Figure 5. Gene expression results after 24 h in the kidney of females compared to non-treated mice. In case of *miR-9-1*, $p < 0.05$, as well as in *miR-330*, *miR-9-3* and *mTORC1*, $p < 0.001$. * $p < 0.05$, *** $p < 0.001$.




Figure 6. Gene expression results after 24 h in the kidney of males compared to non-treated mice. In case of *miR-330*, $p < 0.05$, while in *miR-9-1*, $p < 0.001$. * $p < 0.05$, *** $p < 0.001$.

3 locus (with an increased expression due to DMBA) and may increase *miR-9* expression in tumor cells, that (in this case acting as promoter of oncogenes) induces a further increase of *c-MYC* through the amplification of *E-cadherin*, thus promoting the development of hepatocellular carcinoma (46, 24).

This finding is supported by the fact that we found significant difference between *miR-9-1* and *miR-9-3* expression in the liver tissue of female subjects.

Twenty-four hours after DMBA treatment expression of *miR-9-3* was increased in each tested organs of female mice, despite the protective effect of estrogen, while that of *miR-9-1* was decreased in the liver. As it could be expected, *miR-9-3* can be well used as a biomarker in the case of female mice according to our own study, as well. Thus, by studying *miR-9-3*, and extending the experimental design by complementing DMBA treatment with presumably chemopreventive agents, new molecular epidemiological relationships are likely to be discovered.

mTORC1: Mutations in *mTOR* can be found in a wide range of human malignant tumorous diseases, in agreement with the proven tumor promoting effect of *mTOR* (27, 47). The activation of *mTOR* signaling induces several oncogenic processes, including promotion of the growth, survival and proliferation of cancerous cells (48). *mTORC1* mediates cell proliferation stimulated by growth factors such as *insulin-like growth factor (IGF)* (28). In DMBA-induced breast tumors high levels of *transforming growth factor α (TGFA)*, *IGF-1* and probably *IGF-2* at the mRNA levels can be related to the effect of estrogen (49). The potential oncogenic role of *mTORC1* has also been demonstrated in our model. We also found that due to the effect of DMBA, *mTORC1* expression increased in female mice, in contrast to expected decrease due to the *PTEN* gene expression (31), as *PTEN* is an inhibitor of *mTORC1*. This inhibition lacks a direct

negative feedback (28), so a more distant negative feedback regulatory mechanism could exist.

Despite our expectations, *mTORC1* gene - whose cancer promoting mechanisms are supported by literature data - has not shown significant changes in male mice. However, the expression of *mTORC1* gene showed significant increase in the liver, spleen and kidneys of females as a result of DMBA treatment. Therefore, expression of the *mTORC1* gene in females in the model used in the present study appears to be applicable for testing carcinogens.

In summary, consistent and significant changes toward the same direction in all the three studied organs were observed in the case of *miR-9-3* and *mTORC1* genes in female mice. The expression patterns were somewhat different in males which may be due to sex-specific hormonal differences (such as the effects of estrogen and testosterone) that require further investigation. Thus, the interpretation of *miR-330*, *miR-29a* and *miR-9-1* gene expression changes seems to be rather complex, while *miR-9-3* and *mTORC1* genes in females were found to be suitable for the rapid screening of potential carcinogens. The same model can be used later to study chemopreventive effects as well by the simultaneous use of DMBA and supposed chemopreventive agents, measuring to what extent these agents are able to reduce/prevent the DMBA-induced gene expression changes.

While gene expressions are sensitive biomarkers, and in contrast to classic genotoxicity tests they are able to detect minor/early signs of carcinogenic exposures, sometimes they might be influenced by potential confounders. Our results also emphasize that early carcinogenicity testing requires a careful selection of the test system, since hormonal differences can substantially influence gene expression patterns. In order to create a robust assay panel designed for our purposes, additional genes need probably to be tested.

Conflicts of Interest

The Authors have no conflicts of interest to declare.

Authors' Contributions

Study design and conception: AT, LS, RM, IK; Experiments: AT, LS, RM, NG, TV, AD, RD; Data analysis and interpretation: AT, LS, IK; Manuscript preparation: AT, IK; Critical reviewing and revising the manuscript: DM, FB, BN, IK.

References

- Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A: Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 68(6): 394-424, 2018. PMID: 30207593. DOI: 10.3322/caac.21492
- Danaei G, Vander Hoorn S, Lopez AD, Murray CJ and Ezzati M and Comparative Risk Assessment collaborating group (Cancers): Causes of cancer in the world: comparative risk assessment of nine behavioural and environmental risk factors. *Lancet* 366(9499): 1784-1793, 2005. PMID: 16298215. DOI: 10.1016/S0140-6736(05)67725-2
- Mortelmans K and Zeiger E: The Ames Salmonella/microsome mutagenicity assay. *Mutat Res* 455(1-2): 29-60, 2000. PMID: 11113466. DOI: 10.1016/S0027-5107(00)00064-6
- Bartel DP: MicroRNAs: target recognition and regulatory functions. *Cell* 136(2): 215-233, 2009. PMID: 19167326. DOI: 10.1016/j.cell.2009.01.002
- Rajewsky N: microRNA target predictions in animals. *Nat Genet* 38: S8-13, 2006. PMID: 16736023. DOI: 10.1038/ng1798
- Bueno MJ and Malumbres M: MicroRNAs and the cell cycle. *Biochim Biophys Acta* 1812(5): 592-601, 2011. PMID: 21315819. DOI: 10.1016/j.bbdis.2011.02.002
- Bueno MJ, Pérez de Castro I and Malumbres M: Control of cell proliferation pathways by microRNAs. *Cell Cycle* 7(20): 3143-3148, 2008. PMID: 18843198. DOI: 10.4161/cc.7.20.6833
- Calin GA, Sevignani C, Dumitru CD, Hyslop T, Noch E, Yendamuri S, Shimizu M, Rattan S, Bullrich F, Negrini M and Croce CM: Human microRNA genes are frequently located at fragile sites and genomic regions involved in cancers. *Proc Natl Acad Sci USA* 101(9): 2999-3004, 2004. PMID: 14973191. DOI: 10.1073/pnas.0307323101
- Visone R and Croce CM: MiRNAs and cancer. *Am J Pathol* 174(4): 1131-1138, 2009. PMID: 19264914. DOI: 10.2353/ajpath.2009.080794
- Corney DC, Flesken-Nikitin A, Godwin AK, Wang W and Nikitin AY: MicroRNA-34b and MicroRNA-34c are targets of p53 and cooperate in control of cell proliferation and adhesion-independent. *Growth Cancer Res* 67(18): 8433-8438, 2007. PMID: 17823410. DOI: 10.1158/0008-5472.CAN-07-1585
- He C, Xiong J, Xu X, Lu W, Liu L, Xiao D and Wang D: Functional elucidation of MiR-34 in osteosarcoma cells and primary tumor samples. *Biochem Biophys Res Commun* 388(1): 35-40, 2009. PMID: 19632201. DOI: 10.1016/j.bbrc.2009.07.101
- Kim S, Lee UJ, Kim MN, Lee E-J, Kim JY, Lee MY, Choung S, Kim YJ and Choi Y-C: MicroRNA miR-199a* regulates the MET Proto-oncogene and the downstream extracellular signal-regulated kinase 2 (ERK2). *J Biol Chem* 283(26): 18158-18166, 2008. PMID: 18456660. DOI: 10.1074/jbc.M800186200
- Pan JY, Zhang F, Sun CC, Li SJ, Li G, Gong FY, Bo T, He J, Hua RX, Hu WD, Yuan ZP, Wang X, He QQ and Li DJ: miR-134: A human cancer suppressor? *Mol Ther Nucleic Acids* 6: 140-149, 2017. PMID: 28325280. DOI: 10.1016/j.omtn.2016.11.003
- Chen X, Ba Y, Ma L, Cai X, Yin Y, Wang K, Guo J, Zhang Y, Chen J, Guo X, Li Q, Li X, Wang W, Zhang Y, Wang J, Jiang X, Xiang Y, Xu C, Zheng P, Zhang J, Li R, Zhang H, Shang X, Gong T, Ning G, Wang J, Zen K, Zhang J and Zhang CY: Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and other diseases. *Cell Res* 18(10): 997-1006, 2008. PMID: 18766170. DOI: 10.1038/cr.2008.282
- Bosland MC and Prinsen MK: Induction of dorsolateral prostate adenocarcinomas and other accessory sex gland lesions in male Wistar rats by a single administration of N-methyl-N-nitrosourea, 7,12-dimethylbenz(a)anthracene, and 3,2'-dimethyl-4-aminobiphenyl after sequential treatment with cyproterone acetate and testosterone propionate. *Cancer Res* 50(3): 691-699, 1990. PMID: 2137026.
- El-Sohemy A and Archer MC: Inhibition of N-methyl-N-nitrosourea- and 7,12-dimethylbenz[a]anthracene-induced rat mammary tumorigenesis by dietary cholesterol is independent of Ha-Ras mutations. *Carcinogenesis* 21(4): 827-831, 2000. PMID: 10753223. DOI: 10.1093/carcin/21.4.827
- Gao J, Lauer FT, Dunaway S and Burchiel SW: Cytochrome P450 1B1 is required for 7,12-dimethylbenz(a)-anthracene (DMBA) induced spleen cell immunotoxicity. *Toxicol Sci* 86(1): 68-74, 2005. PMID: 15843505. DOI: 10.1093/toxsci/kfi176
- Lindhe O, Granberg L and Brandt I: Target cells for cytochrome p450-catalysed irreversible binding of 7,12-dimethylbenz[a]anthracene (DMBA) in rodent adrenal glands. *Arch Toxicol* 76(8): 460-466, 2002. PMID: 12185414. DOI: 10.1007/s00204-002-0367-1
- Budán F, Varjas T, Nowrasteh G, Prantner I, Varga Z, Ember E, Cseh J, Gombos K, Pázsit E, Gobel G, Bauer M, Gracza T, Arany I, Perjési P, Ember I and Kiss I: Early modification of c-myc, Ha-ras and p53 expressions by chemical carcinogens (DMBA, MNU). *In Vivo* 23(4): 591-598, 2009. PMID: 19567395.
- Ember I, Kiss I and Pusztai Z: Effect of 7,12-dimethylbenz(a)anthracene on onco/suppressor gene action *in vivo*: a short-term experiment. *Anticancer Res* 18(1A): 445-447, 1998. PMID: 9568117.
- Cheng CW, Yu JC, Hsieh YH, Liao WL, Shieh JC, Yao CC, Lee HJ, Chen PM, Wu PE and Shen CY: Increased cellular levels of microRNA-9 and microRNA-221 correlate with cancer stemness and predict poor outcome in human breast cancer. *Cell Physiol Biochem* 48(5): 2205-2218, 2018. PMID: 30110679. DOI: 10.1159/000492561
- Wang LQ, Kwong YL, Kho CS, Wong KF, Wong KY, Ferracin M, Calin GA and Chim CS: Epigenetic inactivation of miR-9 family microRNAs in chronic lymphocytic leukemia—implications on constitutive activation of NFκB pathway. *Mol Cancer* 12: 173, 2013. PMID: 24373626. DOI: 10.1186/1476-4598-12-173
- Fu L, Shi J, Liu A, Zhou L, Jiang M, Fu H, Xu K, Li D, Deng A, Zhang Q, Pang Y, Guo Y, Hu K, Zhou J, Wang Y, Huang W, Jing Y, Dou L, Wang L, Xu K, Ke X, Nervi C, Li Y and Yu L: A minicircuitry of microRNA-9-1 and RUNX1-RUNX1T1 contributes to leukemogenesis in t(8;21) acute myeloid

- leukemia. *Int J Cancer* 140(3): 653-661, 2017. PMID: 27770540. DOI: 10.1002/ijc.30481
- 24 Ma L, Young J, Prabhala H, Pan E, Mestdagh P, Muth D, Teruya-Feldstein J, Reinhardt F, Onder TT, Valastyan S, Westermann F, Speleman F, Vandesompele J and Weinberg RA: miR-9, a MYC/MYCN-activated microRNA, regulates E-cadherin and cancer metastasis. *Nat Cell Biol* 12(3): 247-256, 2010. PMID: 20173740. DOI: 10.1038/ncb2024
- 25 Wang JY, Zhang Q, Wang DD, Yan W, Sha HH, Zhao JH, Yang SJ, Zhang HD, Hou JC, Xu HZ, He YJ, Hu JH, Zhong SL and Tang JH: MiR-29a: a potential therapeutic target and promising biomarker in tumors. *Biosci Rep* 38(1), 2018. PMID: 29217524. DOI: 10.1042/BSR20171265
- 26 Tréhoux S, Lahdaoui F, Delpu Y, Renaud F, Leteurtre E, Torrisani J, Jonckheere N and Van Seuning I: Micro-RNAs miR-29a and miR-330-5p function as tumor suppressors by targeting the MUC1 mucin in pancreatic cancer cells. *Biochim Biophys Acta* 1853(10 Pt A): 2392-2403, 2015. PMID: 26036346. DOI: 10.1016/j.bbamcr.2015.05.033
- 27 Sato T, Nakashima A, Guo L, Coffman K and Tamanoi F: Single amino-acid changes that confer constitutive activation of mTOR are discovered in human cancer. *Oncogene* 29(18): 2746-2752, 2010. PMID: 20190810. DOI: 10.1038/onc.2010.28
- 28 Hall MN: mTOR-what does it do? *Transplant Proc* 40(10 Suppl): S5-8, 2008. PMID: 19100909. DOI: 10.1016/j.transproceed.2008.10.009
- 29 Shor B, Cavender D and Harris C: A kinase-dead knock-in mutation in mTOR leads to early embryonic lethality and is dispensable for the immune system in heterozygous mice. *BMC Immunol* 10: 28, 2009. PMID: 19457267. DOI: 10.1186/1471-2172-10-28
- 30 Uchida S, Hara K, Kobayashi A, Funato H, Hobara T, Otsuki K, Yamagata H, McEwen BS and Watanabe Y: Early life stress enhances behavioral vulnerability to stress through the activation of REST4-mediated gene transcription in the medial prefrontal cortex of rodents. *J Neurosci* 30(45): 15007-15018, 2010. PMID: 21068306. DOI: 10.1523/JNEUROSCI.1436-10.2010
- 31 55 Ma J, Fan Y, Zhang J, Feng S, Hu Z, Qiu W, Long K, Jin L, Tang Q, Wang X, Zhou Q, Gu Y, Xiao W, Liu L, Li X and Li M: Testosterone-dependent miR-26a-5p and let-7g-5p act as signaling mediators to regulate sperm apoptosis *via* targeting PTEN and PMAIP1. *Int J Mol Sci* 19(4), 2018. PMID: 29670053. DOI: 10.3390/ijms19041233
- 32 Berger PL, Frank SB, Schulz VV, Nollet EA, Edick MJ, Holly B, Chang TT, Hostetter G, Kim S and Miranti CK: Transient induction of ING4 by Myc drives prostate epithelial cell differentiation and its disruption drives prostate tumorigenesis. *Cancer Res* 74(12): 3357-3368, 2014. PMID: 24762396. DOI: 10.1158/0008-5472.CAN-13-3076
- 33 Hu X, Feng Y, Sun L, Qu L and Sun C: Roles of microRNA-330 and its target gene *ing4* in the development of aggressive phenotype in hepatocellular carcinoma cells. *Dig Dis Sci* 62(3): 715-722, 2017. PMID: 28050784. DOI: 10.1007/s10620-016-4429-2
- 34 Moyad MA: Obesity, interrelated mechanisms, and exposures and kidney cancer. *Semin Urol Oncol* 19(4): 270-279, 2001. PMID: 11769879.
- 35 Iqbal J, Minhajuddin M and Beg ZH: Suppression of 7,12-dimethylbenz[α]anthracene-induced carcinogenesis and hypercholesterolaemia in rats by tocotrienol-rich fraction isolated from rice bran oil. *Eur J Cancer Prev* 12(6): 447-453, 2003. PMID: 14639121. DOI: 10.1097/01.cej.0000102802.33147.4b
- 36 Zhang GM, Wang MY, Liu YN, Zhu Y, Wan FN, Wei QY and Ye DW: Functional variants in the low-density lipoprotein receptor gene are associated with clear cell renal cell carcinoma susceptibility. *Carcinogenesis* 38(12): 1241-1248, 2017. PMID: 29029037. DOI: 10.1093/carcin/bgx098
- 37 Johnson AT, Kaufmann YC, Luo S, Todorova V and Klimberg VS: Effect of glutamine on glutathione, IGF-I, and TGF-beta 1. *J Surg Res* 111(2): 222-228, 2003. PMID: 12850466. DOI: 10.1016/s0022-4804(03)00083-0
- 38 Kriegel AJ, Liu Y, Fang Y, Ding X and Liang M: The miR-29 family: genomics, cell biology, and relevance to renal and cardiovascular injury. *Physiol Genomics* 44(4): 237-244, 2012. PMID: 22214600. DOI: 10.1152/physiolgenomics.00141.2011
- 39 Akhurst RJ: TGF-beta antagonists: why suppress a tumor suppressor? *J Clin Invest* 109(12): 1533-1536, 2002. PMID: 12070299. DOI: 10.1172/JCI15970
- 40 Carr BI, Hayashi I, Branum EL and Moses HL: Inhibition of DNA synthesis in rat hepatocytes by platelet-derived type beta transforming growth factor. *Cancer Res* 46(5): 2330-2334, 1986. PMID: 3008986.
- 41 Zhang Y, Wen G, Shao G, Wang C, Lin C, Fang H, Balajee AS, Bhagat G, Hei TK and Zhao Y: TGFBI deficiency predisposes mice to spontaneous tumor development. *Cancer Res* 69(1): 37-44, 2009. PMID: 19117985. DOI: 10.1158/0008-5472.CAN-08-1648
- 42 Suzuki HI: MicroRNA control of TGF- β signaling. *Int J Mol Sci* 19(7), 2018. PMID: 29958433. DOI: 10.3390/ijms19071901
- 43 Zhang J, Cheng J, Zeng Z, Wang Y, Li X, Xie Q, Jia J, Yan Y, Guo Z, Gao J, Yao M, Chen X and Lu F: Comprehensive profiling of novel microRNA-9 targets and a tumor suppressor role of microRNA-9 *via* targeting IGF2BP1 in hepatocellular carcinoma. *Oncotarget* 6(39): 42040-42052, 2015. PMID: 26547929. DOI: 10.18632/oncotarget.5969
- 44 Zhang J, Chong CC, Chen GG and Lai PB: A seven-microRNA expression signature predicts survival in hepatocellular carcinoma. *PLoS One* 10(6): e0128628, 2015. PMID: 26046780. DOI: 10.1371/journal.pone.0128628
- 45 Orangi E and Motovali-Bashi M: Evaluation of miRNA-9 and miRNA-34a as potential biomarkers for diagnosis of breast cancer in Iranian women. *Gene* 687: 272-279, 2019. PMID: 30468908. DOI: 10.1016/j.gene.2018.11.036
- 46 Morin PJ: Beta-catenin signaling and cancer. *Bioessays* 21(12): 1021-1030, 1999. PMID: 10580987. DOI: 10.1002/(SICI)1521-1878(199912)22:1<1021::AID-BIES6>3.0.CO;2-P
- 47 Grabiner BC, Nardi V, Birsoy K, Possemato R, Shen K, Sinha S, Jordan A, Beck AH and Sabatini DM: A diverse array of cancer-associated MTOR mutations are hyperactivating and can predict rapamycin sensitivity. *Cancer Discov* 4(5): 554-563, 2014. PMID: 24631838. DOI: 10.1158/2159-8290.CD-13-0929
- 48 Laplante M and Sabatini DM: mTOR signaling in growth control and disease. *Cell* 149(2): 274-293, 2012. PMID: 22500797. DOI: 10.1016/j.cell.2012.03.017
- 49 Ruggeri BA, Klurfeld DM, Kritchevsky D and Frick KK: Steady-state mRNA expression for growth factors in DMBA-induced rat mammary tumors. *Cancer Lett* 55(2): 89-93, 1990. PMID: 2124948. DOI: 10.1016/0304-3835(90)90016-q

Received June 12, 2020

Revised July 2, 2020

Accepted July 6, 2020