



Letter to the Editor

COVID-19 and acute mesenteric ischemia: A review of literature



Dear Editor,

Coronavirus disease-2019 (COVID-19) caused by SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2) has caused global health crisis. Initially considered a respiratory tract pathogen, it can cause multiple organ dysfunction. It has also been described to predispose to venous and arterial thromboembolism; however, limited published data is available regarding mesenteric thrombosis COVID-19. We conducted a rapid review of current scientific literature available in PubMed to identify cases of AMI in COVID-19 patients- total of 13 cases were found. We delineated clinical characteristics and outcome in these patients. Clinicians should be aware of the life-threatening situation in COVID-19 patients.

A novel coronavirus termed as SARS-CoV-2 (severe acute respiratory syndrome coronavirus 2) has been the causative agent of a pandemic that originated in Wuhan China in December 2019. Coronavirus disease-2019 (COVID-19) can present with a wide variety of complications during infection. For optimal management of these patients, understanding of various systemic manifestations and complications of SARS-CoV2 is vital. Although in COVID-19 respiratory symptoms predominate, both arterial and venous thrombosis can occur with COVID-19. Arterial thrombosis reported so far include stroke, acute limb ischemia, acute mesenteric ischemia and acute coronary syndrome.^{1–4} Limited literature is available regarding acute mesenteric ischemia (AMI). We did an extensive literature review on COVID-19 associated mesenteric thrombosis.

We searched PubMed for this literature review using search terms 'COVID-19 and mesenteric thrombosis', 'COVID-19 and mesenteric ischemia', and 'COVID-19 and bowel ischemia'. All the case reports who had COVID-19 associated mesenteric thrombosis so far is reviewed, and relevant data abstracted from these studies in Table 1. COVID-19 diagnosis was made by PCR assay except in one patient it was negative (suspected COVID-19).

Clinical characteristics of the COVID-19 patients with AMI are summarized in Table 1.^{5–15} The median age of the patient was 56 years (range 9–79 years). We found total of 13 patients-

9 were male, 3 female and for 1 patient sex was not reported. AMI can occur as a presenting feature or a late complication of COVID-19 during hospitalization (median 7 days). 6 patients had pre-existing comorbidities while 7 patients had none. The pre-existing conditions reported were hypertension, diabetes, obesity, obstructive sleep apnea, anxiety, idiopathic medullar aplasia, chronic bronchitis, essential thrombocytoysis, and cardiac transplantation. Presenting symptoms were nausea, vomiting, abdominal pain, diarrhea, fever, cough, shortness of breath, eructation, pain in throat and stroke. The diagnosis of AMI was made by contrast enhanced computed tomography. 4 patients had concurrent thrombosis at other sites – case 3 had stroke, case 7 had portal and mesenteric vein thrombosis, case 8 had splenic and renal infarcts and case 11 had superior mesenteric and portal vein thrombosis. 10 patients had surgery, 2 patients had conservative management and 1 was started on therapeutic anticoagulation with heparin. Out of 13 patients, 4 patients died.

Acute mesenteric ischemia is a rare abdominal emergency and is associated with high rates of morbidity and mortality. Prompt diagnosis requires a high index of suspicion and early contrast computed tomography imaging. The exact pathological mechanism leading to the complication of AMI in COVID-19 is not well understood at present, possibilities include - direct invasion of bowel tissue by the virus given expression of angiotensin converting enzyme 2 on enterocytes, the target receptor for SAR-CoV-2 or viral infection of the endothelial cell leading to diffuse endothelial inflammation or increased procoagulant factors like factor VIII, von Willebrand factor, fibrinogen or virus induced cytokine storm leading to coagulation and fibrinolysis activation.^{16–18} Additional explanations for the hypercoagulability may be the presence of high numbers of prothrombotic circulating microvesicles which are cytoplasmic microparticles stemming from platelets or monocytes and Neutrophil extracellular traps (NETs) released from activated neutrophils, constitute a mixture of nucleic DNA, histones and nucleosomes.¹⁸ Treatment of this life-threatening condition includes surgical resection of the necrotic bowel, restoration of blood flow to the ischemic intestine and supportive measure - gastrointestinal decom-

Table 1 – Summarizing Clinical characteristics of the COVID-19 patients with AMI.

	Age	Medical history	Presenting signs and symptoms	Timing of AMI diagnoses	Imaging findings	Other site of thrombosis	Treatment	Rx of COVID-19	Outcome
Case 1 ⁵	55 M	HTN	Nausea, generalized abdominal pain, diarrhea	Day 7	CT scan of the abdomen and pelvis with IV contrast showed thrombus 1.6 cm in length in SMA	None	Laparotomy and SMA thromboembolectomy	HCQ, azithromycin and ceftriaxone	NR
Case 2 ⁶	47 M	Anxiety, obesity, OSA	Fever, dry cough and vomiting	Day 8	CT of the abdomen revealed diffuse small-bowel distension with widespread pneumatisis, circumferential mural thickening, free fluid, mesenteric free air and portal venous gas	None	Therapeutic heparin	NR	Discharged
Case 3 ⁷	56 NR	None	Stroke, next day developed abdominal pain and vomiting	Day 2	CT scan showed a free-floating thrombus in the aortic arch associated with an occlusion of the superior mesenteric artery	Stroke	Endovascular thrombectomy and laparotomy with the resection of two meters of the small bowel	NR	NR
Case 4 ⁸	69 M	None	Epigastric pain, constipation and eructation	Day 1	CT angiogram demonstrated a thrombus in the proximal segment of the superior mesenteric artery with complete occlusion in the right ileocolic branches	None	Small bowel resection and superior mesenteric artery thromboembolectomy	NR	Discharged
Case 5 ⁹	52 M	None	Cough and fever	Day 13	CT scan showed arterial thrombosis of vessels efferent of the superior mesenteric artery with bowel distension	None	An intestinal resection with stapled side-to side anastomosis was performed	NR	Discharged

Table 1 (Continued)

	Age	Medical history	Presenting signs and symptoms	Timing of AMI diagnoses	Imaging findings	Other site of thrombosis	Treatment	Rx of COVID-19	Outcome
Case 6 ¹⁰	75 M	None	Abdominal pain, vomiting, cough and SOB	Day 1	CT angiography showed thrombus in the descending thoracic aorta with embolic occlusion of the superior mesenteric artery	None	Catheter-directed thrombolysis was commenced but the patient developed worsening abdominal symptoms and underwent laparotomy, requiring resection of 150 cm of ischemic small bowel	NR	NR
Case 7 ¹¹	79 F	None	Fever, epigastric abdominal pain	Day 1	CT scan of the chest, abdomen, and pelvis at the arterial and portal phases, showed right-portal vein thrombosis, thrombosis of the distal part of the upper mesenteric vein, proximal thrombosis of the superior mesenteric artery and jejunal artery.	Portal vein and mesenteric vein thrombosis	Superior mesenteric artery thrombectomy and laparotomy with resection of a meter of necrotic ileum and right colon.	NR	Died
Case 8 ¹⁹	58 M	None	SOB and abdominal pain	Day 1	CT scan showed dilated small bowel loops, signs of bowel wall ischemia, splenic and renal infarcts without macrovascular arterial occlusion	Concurrent splenic and renal infarcts and 3 weeks later digital necrosis of bilateral feet	Laparotomy was performed, and a partial small bowel resection was done.	NR	SIH
Case 9 ¹³	9 F	Idiopathic medullary aplasia	Fever, abdominal pain, vomiting, diarrhea	NR	NR	NR	Resection of the ischemic bowel loop with double ileostomy was performed.	NR	Died

Table 1 (Continued)

	Age	Medical history	Presenting signs and symptoms	Timing of AMI diagnoses	Imaging findings	Other site of thrombosis	Treatment	Rx of COVID-19	Outcome
Case 10 ¹⁵	70 M	None	Abdominal pain, nausea, fever, pain in throat and cough	Day 1	Contrast-enhanced CT of the abdomen showed acute small bowel hypoperfusion	None	Conservative management	NR	Died
Case 11 ¹⁴	28 F	ET	Abdominal pain and vomiting	Day 5	Abdominal CT scan showed segmental small bowel ischemia	Superior mesenteric and portal vein thrombosis	Laparotomy -Bowel resection	NR	Discharged
Case 12 ¹⁴	56 M	HTN, DM, obesity	ARDS	Day 9	CT scan showed bowel ischemia and mesenteric venous gas in proximal jejunum	None	Laparotomy -Bowel resection	NR	SIH
Case 13 ¹⁴	67 M	Chronic bronchitis, diabetes, and heart transplantation	ARDS	Day 6	CT scan showed inflammatory segmental ileitis with a localized thickening of one small bowel loop and edema	None	Conservative management	NR	Died

M: male; F: female; NR: not reported; HTN: hypertension; OSA: obstructive sleep apnea; ET: essential thrombocytosis; DM: diabetes; SOB: shortness of breath; ARDS: acute respiratory distress syndrome; CT: computed tomography; SMA: superior mesenteric artery; HCQ: hydroxychloroquine; SIH: Still in hospital (at the time of writing of respective manuscript).

pression, fluid resuscitation, hemodynamic support. Health care providers should have high index of suspicion regarding this life-threatening complication of COVID-19 so that timely intervention can be done.

REFERENCES

1. Kaur P, Qaqa F, Ramahi A, Shamoon Y, Singhal M, Shamoon F, et al. Acute upper limb ischemia in a patient with COVID-19. *Hematology/Oncology and Stem Cell Therapy*. 2020.
2. Kaur S, Bansal R, Kollimuttathuillam S, Gowda AM, Singh B, Mehta D, et al. The looming storm: Blood and cytokines in COVID-19. *Blood Reviews*. 2020;100743.
3. Kaur P, Posimreddy S, Singh B, Qaqa F, Habib HA, Maroules M, et al. COVID-19 Presenting as Acute Limb Ischaemia. *Eur J Case Rep Intern Med*. 2020;7(6), 001724-.
4. Singh B, Kaur P, Ajdir N, Gupta S, Maroules M. Covid-19 Presenting as Acute Limb Ischemia. *Cureus*. 2020;12(7):e9344.
5. Cheung S, Quiwa JC, Pillai A, Onwu C, Tharayil ZJ, Gupta R. Superior Mesenteric Artery Thrombosis and Acute Intestinal Ischemia as a Consequence of COVID-19 Infection. *Am J Case Rep*. 2020;21:e925753.
6. Kiely J, Duggan WP, O'Dwyer M. Extensive pneumatisis intestinalis and portal venous gas mimicking mesenteric ischaemia in a patient with SARS-CoV-2. *Ann R Coll Surg Engl*. 2020;102(6):e145-7.
7. Azouz E, Yang S, Monnier-Cholley L, Arrivé L. Systemic arterial thrombosis and acute mesenteric ischemia in a patient with COVID-19. *Intensive Care Med*. 2020;46(7):1464-5.
8. Mitchell JM, Rakheja D, Gopal P. SARS-CoV-2-related Hypercoagulable State Leading to Ischemic Enteritis Secondary to Superior Mesenteric Artery Thrombosis. *Clin Gastroenterol Hepatol*. 2020.
9. AB L, Pacioni C, Ponton S, Francavilla S, Cuzzoli A. Arterial Mesenteric Thrombosis as a Complication of SARS-CoV-2 Infection. *Eur J Case Rep Intern Med*. 2020;7(5):001690.
10. Vulliamy P, Jacob S, Davenport RA. Acute aorto-iliac and mesenteric arterial thromboses as presenting features of COVID-19. *Br J Haematol*. 2020;189(6):1053-4.
11. de Barry O, Mekki A, Diffre C, Seror M, El Hajjam M, Carlier RY. Arterial and venous abdominal thrombosis in a 79-year-old woman with COVID-19 pneumonia. *Radiol Case Rep*. 2020;15(7):1054-7.
12. Levolger S, Bokkers RPH, Wille J, Kropman RHJ, de Vries J-PPM. Arterial thrombotic complications in COVID-19 patients. *J Vasc Surg Cases Innov Tech*. 2020;6(3):454-9.
13. Khesrani LS, Chana K, Sadar FZ, Dahdouh A, Ladjadj Y, Bouguermouh D. Intestinal ischemia secondary to Covid-19. *Pediatr Surg Case Rep*. 2020;101604.
14. Ignat M, Philouze G, Aussenauc-Belle L, Faucher V, Collange O, Mutter D, et al. Small bowel ischemia and SARS-CoV-2 infection: an underdiagnosed distinct clinical entity. *Surgery*. 2020;168(1):14-6.
15. Farina D, Rondi P, Botturi E, Renzulli M, Borghesi A, Guelfi D, et al. Gastrointestinal: Bowel ischemia in a suspected coronavirus disease (COVID-19) patient. *J Gastroenterol Hepatol*. 2020.
16. Parry AH, Wani AH, Yaseen M. Acute Mesenteric Ischemia in Severe Coronavirus-19 (COVID-19): Possible Mechanisms and Diagnostic Pathway. *Acad Radiol*. 2020;27(8):1190.
17. Varga Z, Flammer AJ, Steiger P, Haberecker M, Andermatt R, Zinkernagel AS, et al. Endothelial cell infection and endotheliitis in COVID-19. *Lancet*. 2020;395(10234):1417-8.
18. Panigada M, Bottino N, Tagliabue P, Grasselli G, Novembrino C, Chantarangkul V, et al. Hypercoagulability of COVID-19 patients in intensive care unit: A report of thromboelastography findings and other parameters of hemostasis. *J Thromb Haemost*. 2020;18(7):1738-42.
19. Levolger S, Bokkers RPH, Wille J, Kropman RHJ, de Vries J. Arterial thrombotic complications in COVID-19 patients. *J Vasc Surg Cases Innov Tech*. 2020;6(3):454-9.

Balraj Singh  *

Department of Hematology & Oncology Saint Joseph's University Medical Center Paterson, New Jersey, United States

Parminder Kaur 

Department of Cardiology, Saint Joseph's University Medical Center Paterson, New Jersey, United States

* Corresponding author at: Saint Joseph's University Medical Center, 703 Main Street, Paterson, New Jersey 07503, United States.

E-mail address: bsriar9@gmail.com (B. Singh).

Received 18 September 2020

Available online 12 November 2020

2531-1379/

© 2020 Associação Brasileira de Hematologia, Hemoterapia e Terapia Celular. Published by Elsevier Editora Ltda. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).
<https://doi.org/10.1016/j.htct.2020.10.959>