

LETTER

COVID-19 mortality increases with northerly latitude after adjustment for age suggesting a link with ultraviolet and vitamin D

Dear Editors,

We read with interest the review by Dr Kohlmeier in which he reported a correlation between COVID-19 mortality among African-Americans across the USA and northern latitude.¹ We previously reported a north–south gradient in global COVID-19 mortality but were conscious that lack of ultraviolet exposure and consequent vitamin D insufficiency was not the only possible explanation.² We have now investigated the relationships between latitude, age of population, population density and pollution with COVID-19 mortality.

COVID-19 mortality per million by country was downloaded from <https://www.worldometers.info/coronavirus/> on 18 May 2020.³ We included all 117 countries with population >1 million and ≥150 COVID-19 cases. Data by country for population %≥65 years, population density and air pollution (particles of matter <2.5 µm diameter µg/m³) were obtained from public sources.^{4–6} Latitude was entered for each country's capital city. The hypothesis was that there was no

relationship between mortality and latitude below a threshold and that thereafter mortality increased with latitude. Mortality data were log transformed, and piecewise linear modelling was used to explore the relationship with latitude. This was adjusted for %≥65, and pollution and population density were investigated to see if they further explained variability in mortality.

The analysis supported the hypothesis with a threshold of 28° north and a model of zero slope below the threshold, and a linear model above the threshold was fitted. The age adjustment was highly significant ($p<0.0005$), with an estimated mortality increase of 13.7% (95% CI 7.4% to 20.3%) for each 1% increase in %≥65. Latitude was also significant ($p=0.031$) with an estimated 4.4% (95% CI 0.4% to 8.5%) increase in mortality for each 1° further north (table 1, figure 1). Countries with higher pollution included many with younger populations, and pollution was negatively associated with mortality but added no significant explanatory power to a model containing latitude and age. Population density expressed per country was not significantly associated with mortality.

The proportion of older people in each country impacts greatly on COVID-19 mortality, but after adjustment for this, a strong association remains across the Northern hemisphere between latitude and higher COVID-19 mortality. This association exists above 28° north not far from the latitude, usually stated as 35°

north, beyond which populations commonly get insufficient ultraviolet B to maintain normal vitamin D blood levels throughout winter. There are exceptions, but COVID-19 mortality correlates with reported vitamin D levels across Europe,⁷ and in sunnier Brazil, where mortality is rising, 28% prevalence of vitamin D deficiency is reported.⁸ An association between vitamin D insufficiency and COVID-19 severity is supported by substantial evidence of its impact on cytokine response to pathogens.⁷ A direct effect of ultraviolet light on the environmental survival of severe acute respiratory syndrome coronavirus 2 is also possible but would not explain the association between mortality and ethnicity,⁹ whereas people with dark skin need more ultraviolet exposure for equivalent vitamin D synthesis.

This analysis supports the link between latitude and COVID-19 mortality reported within the USA by Dr Kohlmeier.¹ Evidence linking vitamin D deficiency with COVID-19 severity is circumstantial but growing. Obtaining more direct evidence may be difficult as people could be reluctant to trial a placebo in place of a vitamin supplement. If the association between vitamin D deficiency and COVID-19 severity is causative, the disease should prove seasonal, since more severely affected individuals are infectious for longer. We agree that very high vitamin D doses >4000 IU/day should only be taken in the context of clinical trials¹⁰ but urge that vitamin D supplementation at more moderate dose should

Table 1 Associations between COVID-19 mortality by country, latitude and % of population ≥65 years

Variable	Regression coefficient	SE	P value	% of variation explained	Effect size (95% CI)*
Univariate models					
Latitude	0.1074	0.0142	<0.0005	33.1	11.3% (8.3% to 14.5%)
%≥65	0.1766	0.0199	<0.0005	40.4	19.3% (14.8% to 24.1%)
Multivariate model					
Latitude	0.0428	0.0196	0.031	43.0	4.4% (0.4% to 8.5%)
%≥65	0.1281	0.0291	<0.0005		13.7% (7.4% to 20.3%)

*The effect size is, for latitude, the percentage increase in mortality from one location, situated at least 28° north, to another location 1° further north and, for %≥65, the percentage increase in mortality for each one % increase in %≥65.




Figure 1 A. COVID-19 mortality per 1 million population by country compared with latitude of capital cities. Fitted values are derived from a piecewise linear model of the logarithm of mortality on latitude. This was based on a threshold of 28° north that explained the greatest amount of variation. B. Logarithm of COVID-19 mortality per 1 million compared with latitude with and without adjustment for age (%≥65 years).

be taken by all those at risk of deficiency, including people with darker skin or living in institutions.

Jonathan Rhodes ,¹ Frank Dunstan,² Eamon Laird,³ Sreedhar Subramanian,¹ Rose A Kenny⁴

¹Department of Cellular and Molecular Physiology, University of Liverpool Institute of Translational Medicine, Liverpool, UK

²School of Medicine, Cardiff University, Cardiff, UK

³The Irish Longitudinal Study on Ageing, University of Dublin Trinity College School of Medicine, Dublin, Ireland

⁴Department of Medical Gerontology, St James Hospital, Dublin, Ireland

Correspondence to Professor Jonathan Rhodes, Cellular and molecular physiology, University of Liverpool Institute of Translational Medicine, Liverpool L69 3BX, UK; rhodesjm@liverpool.ac.uk

Correction notice The paper has been corrected since it was published online. The data in figure 1 and

table 1 have been updated.

Contributors All authors contributed to writing and revision and approved the final version.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests JR with the University of Liverpool and Provox UK holds a patent for use of a soluble fibre preparation as maintenance therapy for Crohn's disease plus a patent for its use in antibiotic-associated diarrhoea. Patent also held with the University of Liverpool and others in relation to use of modified heparins in cancer therapy. SS has received speaker fees from MSD, Actavis, Abbvie, Dr Falk Pharmaceuticals and Shire and received educational grants from MSD, Abbvie and Actavis and is an advisory board member for Abbvie, Dr Falk Pharmaceuticals and Vifor pharmaceuticals. FD, EL and RAK have no conflicts to declare.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; internally peer reviewed.



OPEN ACCESS

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.



To cite Rhodes J, Dunstan F, Laird E, *et al*. COVID-19 mortality increases with northerly latitude after adjustment for age suggesting a link with ultraviolet and vitamin D. *BMJ Nutrition, Prevention & Health* 2020;3:e000110

Received 29 May 2020

Accepted 3 June 2020

bmjnph 2020;3:118–120.

doi:10.1136/bmjnph-2020-000110

ORCID iD

Jonathan Rhodes <http://orcid.org/0000-0002-1302-260X>

REFERENCES

- 1 Kohlmeier M. Avoidance of vitamin D deficiency to slow the COVID-19 pandemic. *BMJNPH* 2020;3:e000096.
- 2 Rhodes JM, Subramanian S, Laird E, *et al*. Editorial: low population mortality from COVID-19 in countries South of latitude 35 degrees North supports vitamin D as

- a factor determining severity. *Aliment Pharmacol Ther* 2020;51:1434–7.
- 3 Coronavirus cases, 2020. Available: <https://www.worldometers.info/coronavirus/> [Accessed 18 May 2020].
 - 4 Population ages 65 and above (% of total population), 2020. Available: <https://data.worldbank.org/indicator/SP.POP.65UP.TO.ZS> [Accessed 18 May 2020].
 - 5 List of countries and dependencies by population density, 2020. Available: https://en.wikipedia.org/wiki/List_of_countries_and_dependencies_by_population_density [Accessed 18 May 2020].
 - 6 Outdoor air pollution, 2020. Available: <https://ourworldindata.org/outdoor-air-pollution> [Accessed 18 May 2020].
 - 7 Laird E, Rhodes J, Kenny RA, *et al*. Vitamin D and inflammation: potential implications for severity of COVID-19. *Irish Med J* 2020;113:P81.
 - 8 Pereira-Santos M, Santos JYGD, Carvalho GQ, *et al*. Epidemiology of vitamin D insufficiency and deficiency in a population in a sunny country: Geospatial meta-analysis in Brazil. *Crit Rev Food Sci Nutr* 2019;59:2102–9.
 - 9 Coronavirus (COVID-19) related deaths by ethnic group, England and Wales: 2 March 2020 to 10 April 2020, 2020. Available: <https://www.ons.gov.uk/peoplepopulationandcommunity/birthsdeathsandmarriages/deaths/articles/coronavirusrelateddeathsbyethnicgroupenglandandwales/2march2020to10april2020> [Accessed 29 May 2020].
 - 10 Lanham-New SA, Webb AR, Cashman KD, *et al*. Vitamin D and SARS-CoV-2 virus/COVID-19 disease. *BMJNPH* 2020;3:e000089.