

# Lifestyle modification for the management of atrial fibrillation

Yehia Fanous MD, Paul Dorian MD

■ Cite as: *CMAJ* 2020 November 16;192:E1469. doi: 10.1503/cmaj.201084

## 1 Weight loss, achieved and sustained through diet and exercise, is associated with fewer atrial fibrillation recurrences<sup>1</sup>

Trials showing the benefit of weight loss (in combination with other strategies for managing risk factors) in atrial fibrillation have primarily used high-protein, low-glycemic-index diets, along with low-intensity exercise, to achieve reductions in body mass index (BMI).<sup>1,2</sup> Patients with intermittent atrial fibrillation and BMI  $\geq 27$  kg/m<sup>2</sup> achieved the greatest increase in arrhythmia-free survival with a sustained weight loss of  $\geq 10\%$  of body weight (number need to treat [NNT] = 4.8).<sup>1-3</sup>

## 2 Recreational exercise confers cardioprotective benefits in atrial fibrillation independently of weight loss<sup>1-3</sup>

Observational studies show that exercise, at any recreational intensity, is associated with reduced risk of developing atrial fibrillation, and improved symptom control in patients with persistent atrial fibrillation.<sup>1,3,4</sup> A useful goal is to progressively increase to 200 min/wk at moderate intensity (1000 metabolic equivalents [METs]), which is associated with an approximately 10% decrease in risk of incident atrial fibrillation.<sup>1,3</sup>

## 3 Reducing alcohol intake, in individuals who consume $\geq 14$ drinks per week, is associated with a decreased burden of atrial fibrillation and decreased related hospital admissions<sup>1-3,5</sup>

There is conflicting evidence as to whether a linear dose-response relationship or a threshold effect exists between alcohol consumption and risk of atrial fibrillation.<sup>1,3</sup> In a recent randomized controlled trial, abstinence in moderate drinkers ( $17 \pm 8$  drinks/wk) with intermittent atrial fibrillation was associated with reduced recurrence (NNT = 5).<sup>1</sup> For patients who consume moderate to heavy amounts of alcohol, especially with binge drinking, it is appropriate to counsel them to reduce their intake.

## 4 Smoking cessation is recommended as part of strategies for preventing and managing atrial fibrillation<sup>1</sup>

A dose-dependent relationship has been suggested between smoking and atrial fibrillation.<sup>1</sup> Moreover, smoking is strongly associated with the development of chronic obstructive pulmonary disease, an important risk factor for atrial fibrillation.<sup>1</sup>

## 5 Limiting caffeine intake is unlikely to be useful in reducing atrial fibrillation incidence or burden<sup>1,3</sup>

No evidence exists to suggest that limiting habitual caffeine intake confers any benefit in reducing the risk of atrial fibrillation.<sup>1</sup>

## References

1. Chung MK, Eckhardt LL, Chen LY, et al.; American Heart Association Electrocardiography and Arrhythmias Committee and Exercise, Cardiac Rehabilitation, and Secondary Prevention Committee of the Council on Clinical Cardiology; Council on Arteriosclerosis, Thrombosis and Vascular Biology; Council on Cardiovascular and Stroke Nursing; Council on Lifestyle and Cardio-metabolic Health. Lifestyle and risk factor modification for reduction of atrial fibrillation: a scientific statement from the American Heart Association. *Circulation* 2020;141:e750-72.
2. Pathak RK, Middeldorp ME, Meredith M, et al. Long-term effect of goal-directed weight management in an atrial fibrillation cohort: a long-term follow-up study (LEGACY). *J Am Coll Cardiol* 2015;65:2159-69.
3. Wingerter R, Steiger N, Burrows A, et al. Impact of lifestyle modification on atrial fibrillation. *Am J Cardiol* 2020;125:289-97.
4. Svedberg N, Sundström J, James S, et al. Long-term incidence of atrial fibrillation and stroke among cross-country skiers. *Circulation* 2019;140:910-20.
5. Gémes K, Malmo V, Laugsand LE, et al. Does moderate drinking increase the risk of atrial fibrillation? The Norwegian HUNT (Nord-Trøndelag Health) study. *J Am Heart Assoc* 2017;6:e007094.

**Competing interests:** None declared.

This article has been peer reviewed.

**Affiliations:** Department of Medicine (Fanous), University of Western Ontario, London, Ont.; Division of Cardiology (Dorian), Department of Medicine, St. Michael's Hospital; Faculty of Medicine (Dorian), University of Toronto, Toronto, Ont.

**Correspondence to:** Paul Dorian, pauldorian@unityhealth.to

*CMAJ* invites submissions to "Five things to know about ..." Submit manuscripts online at <http://mc.manuscriptcentral.com/cmaj>