



Published in final edited form as:

*Chromosome Res.* 2015 December ; 23(4): 767–779. doi:10.1007/s10577-015-9495-3.

## Identification of the centromeric repeat in the threespine stickleback fish (*Gasterosteus aculeatus*)

Jennifer N. Cech<sup>1,2</sup>, Catherine L. Peichel<sup>1,\*</sup>

<sup>1</sup>Divisions of Human Biology and Basic Sciences, Fred Hutchinson Cancer Research Center, Seattle, WA 98109, USA

<sup>2</sup>Graduate Program in Molecular and Cellular Biology, University of Washington, Seattle, WA 98195, USA

### Abstract

Centromere sequences exist as gaps in many genome assemblies due to their repetitive nature. Here we take an unbiased approach utilizing CENP-A chromatin immunoprecipitation followed by high throughput sequencing to identify the centromeric repeat sequence in the threespine stickleback fish (*Gasterosteus aculeatus*). A 186 bp, AT-rich repeat was validated as centromeric using both FISH (fluorescence in situ hybridization) and IF-FISH (immunofluorescence combined with FISH) on interphase nuclei and metaphase spreads. This repeat hybridizes strongly to the centromere on all chromosomes, with the exception of weak hybridization to the Y chromosome. Together, our work provides the first validated sequence information for the threespine stickleback centromere.

### Keywords

Centromere; CENP-A; ChIP-seq; *Gasterosteus aculeatus*; Threespine stickleback

### Introduction

Proper segregation of chromosomes is essential for the faithful and equal separation of each chromosome during both mitosis and meiosis. Mis-segregation of chromosomes can lead to aneuploidy, which is often associated with cancer, miscarriage, and birth defects such as viable trisomies (Kops et al. 2005; Morales et al. 2007; Hunt and Hassold 2010; Lister et al. 2010; Revenkova et al. 2010; Gordon et al. 2012; Ricke and van Deursen 2013). Physical separation of chromosomes occurs by the attachment of microtubule spindle fibers via the kinetochore to a primary constriction on each chromosome, called the centromere. Because of the vital role centromeres play in cell division, each chromosome must contain only a single functional centromere. Functional centromeres are defined by the presence of a centromeric histone variant known as either CenH3 or CENP-A (Palmer et al. 1991; Sullivan et al. 1994). CENP-A replaces histone H3 at functional centromeres and localizes to the

\* Author for correspondence: Catherine L. Peichel, Divisions of Human Biology and Basic Sciences, Fred Hutchinson Cancer Research Center, 1100 Fairview Ave North, Mailstop C2-023, Seattle, WA 98109, USA, phone: 206-667-1628, fax: 206-667-6523, cpeichel@fhcrc.org.

primary constriction on metaphase chromosome spreads (Earnshaw and Rothfield 1985; Palmer et al. 1989; Sullivan and Schwartz 1995; Warburton et al. 1997; Amor et al. 2004).

While the presence of CENP-A at functional centromeres is almost entirely conserved among eukaryotes (Henikoff et al. 2001; Malik and Henikoff 2009; Drinnenberg et al. 2014), the genetic sequence at centromeres varies dramatically between species (Henikoff et al. 2001; Alkan et al. 2011; Melters et al. 2013) and even among chromosomes of the same species or closely related species (Shang et al. 2010; Tek et al. 2010; Piras et al. 2010; Gong et al. 2012). Despite variation at the sequence level, most centromere sequences have conserved characteristics, such as being highly repetitive and AT-rich (Melters et al. 2013). In addition, many mammalian centromere repeats contain a conserved CENP-B box, a 17 bp DNA motif that recruits the CENP-B protein, thought to aid in de novo centromere formation (Masumoto et al. 1989; Ohzeki et al. 2002; Alkan et al. 2011). Previous methods used to identify centromere sequences include isolation of satellite DNA, restriction digest approaches, and BAC (bacterial artificial chromosome) cloning (Maio 1971; Manuelidis 1978; Haaf et al. 1993; Garrido-Ramos et al. 1994; Crollius et al. 2000; Edwards and Murray 2005; Tek et al. 2010; Shang et al. 2010). However, these approaches can be limited by the restriction enzymes used, by biases against cloning repetitive DNA, and assumptions that the most repetitive sequences in the genome will be the centromere. Chromatin immunoprecipitation (ChIP) with a CENP-A antibody followed by cloning of the ChIP DNA has recently been used to both confirm putative centromere sequences (Zhong 2002; Nagaki et al. 2003; Nagaki et al. 2004; Nagaki and Murata 2005; Houben et al. 2007), and to identify novel repeats (Lee et al. 2005; Edwards and Murray 2005; Nagaki et al. 2008; Tek et al. 2010; Shang et al. 2010; Alkan et al. 2011). However, CENP-A ChIP cloning is limited both by the number of clones that can be sequenced and biases against cloning repetitive DNA. Chromatin immunoprecipitation with a CENP-A antibody followed by high throughput sequencing (ChIP-seq) is the most unbiased approach to identify functional centromeric associated DNA (Henikoff et al. 2015), but it has not yet been extensively used (Gong et al. 2012; Henikoff et al. 2015).

Here we sought to identify the centromere sequence in threespine stickleback fish (*Gasterosteus aculeatus*), an emerging model organism used to study the genetic, genomic, and molecular basis for evolution (Kingsley and Peichel 2007). The threespine stickleback has an assembled genome (Jones et al. 2012), yet the centromere sequence is still unknown. Gaps in the genome assembly correspond with the cytological constriction on most chromosomes (Urton et al. 2011), suggesting that the stickleback centromere is comprised of repetitive sequences (Henikoff 2002; Rudd and Willard 2004). To identify the threespine stickleback centromere sequence, we first identified the complete threespine stickleback CENP-A coding sequence, and then performed ChIP-seq using a threespine stickleback specific CENP-A antibody. We confirmed that this 186 bp, AT-rich sequence is centromeric using both fluorescence in situ hybridization (FISH) and CENP-A immunofluorescence coupled with FISH (IF-FISH).

## Materials and Methods

### Fish use and care

Three populations of *G. aculeatus* were used in this study: wild caught fish from Lake Union (Seattle, Washington, USA), laboratory-reared Pacific Ocean fish, and laboratory-reared Japan Sea fish. Both of these laboratory populations were derived from wild-caught fish collected in Akkeshi on Hokkaido Island, Japan (Kitano and Mori 2007; Kitano et al. 2009). Fish were housed in 29-gallon aquarium tanks in summer lighting conditions (16 h light, 8 h dark) at approximately 16°C in 0.35% saltwater (3.5 g/L Instant Ocean salt (Spectrum Brands, USA); 0.4 ml/l sodium bicarbonate). Fish were fed with live brine shrimp nauplii and frozen *Mysis* shrimp twice daily. All institutional and national guidelines for the care and use of laboratory animals were followed, and all procedures were approved by the Fred Hutchinson Cancer Research Center Institutional Animal Care and Use Committee (protocol 1575). Fish were caught from Lake Union with permission of the Washington Department of Fish and Wildlife (Scientific Collection permits 12-057, 13-039, 14-065, and 15-033).

### Determining the *Cenpa* coding sequence

To find the putative *G. aculeatus* *Cenpa* gene, the *Danio rerio* CENP-A protein sequence was used to search for a homologous sequence in the threespine stickleback genome assembly (Ensembl BROAD S1; Feb 2006) using blastp. The putative threespine stickleback CENP-A protein (ENSGACP00000024534) is a product of the gene (ENSGACG00000018561.1). However, this predicted gene was missing a stop codon. To determine the full *Cenpa* coding sequence, RNA was extracted using Trizol (Life Technologies, USA) from a Japan Sea male stickleback kidney. An Invitrogen 3' RACE (rapid amplification of cDNA ends) kit (Life Technologies, USA) was used to make cDNA. Internal primer JC06 (5' GAGAGGTGTGCCAGAGCTTCTC 3') was used in conjunction with the 3' RACE kit primer AUAP to amplify and sequence the entire 3' end of the *Cenpa* gene. This coding sequence was used to design primers (5' ATGCGTCACAATTCATCTACC 3' and 5' TTACAGGTTGTCCACCCC 3') to amplify the entire cDNA. To determine whether the Japan Sea and Pacific Ocean *Cenpa* cDNA sequences are the same, Pacific Ocean male RNA was extracted from liver tissue using Trizol, cDNA was synthesized using the SuperScript III First-Strand Synthesis System kit (Life Technologies, USA), and the *Cenpa* cDNA was Sanger sequenced following polymerase chain reaction (PCR) amplification with the primers above.

### Antibody design

A *G. aculeatus* CENP-A specific antibody was designed to amino acids 1-22 (MRHNSSTSRKKGKTPQHRPPLA) of the N-terminal end of the threespine stickleback CENP-A protein (Fig. S1b). The rabbit IgG affinity purified polyclonal antibody was produced by Covance Research Products (USA).

### Western blot

50 mg of Pacific Ocean female kidney and liver tissue was dissected in 400 µl RIPA (150 mM sodium chloride, 1% NP-40, 0.5 % sodium deoxycholate, 0.1% sodium dodecyl sulfate

(SDS), 50  $\mu$ M Tris pH 8.0, 0.3 mM phenylmethanesulfonylfluoride (PMSF)). Using a pellet pestle, tissue was homogenized in 400  $\mu$ l RIPA, 0.3 mM PMSF on ice. The pestle was rinsed with an additional 600  $\mu$ l RIPA with PMSF, then the sample was rotated 2 h at 4°C. Cells were spun for 20 min at 12,000 rpm and 4°C. The pellet was resuspended in 30  $\mu$ l of 2% SDS, and 6  $\mu$ l of 6X Laemmli buffer was added. The samples were boiled at 95°C for 5 min and then run on a 15% sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) gel in running buffer (25 mM Tris, 0.192 M glycine, 0.2% SDS). Proteins were transferred from the gel to a polyvinylidene difluoride membrane for 1 h at 14 volts in transfer buffer (25 mM Tris, 192 mM glycine, 10% methanol). The membrane was blocked in I-BLOCK (Applied Biosystems, USA) for 1 h. The membrane was incubated with the primary antibody (rabbit, anti-stickleback CENP-A) at 1:1000 in I-BLOCK plus 5% bovine serum albumin (BSA) on a shaking nutator overnight. The membrane was washed 3 $\times$ 5 min with phosphate-buffered saline Tween-20 (PBST), then incubated with alkaline phosphatase conjugated goat, anti-rabbit secondary antibody (Applied Biosystems, USA) at 1:10,000 in I-BLOCK for 1 h. The membrane was washed 2 $\times$ 5 min with PBST, then 2 $\times$ 5 min in assay buffer (20 mM Tris pH 9.8, 1 mM magnesium chloride). A few drops of CDP-star Chemiluminescent substrate (Sigma-Aldrich, USA) was added to membrane. Film was exposed for 5 sec, then developed.

### **Immunofluorescence on metaphase spreads and interphase nuclei**

Immunofluorescence was conducted using a protocol adapted from (Blower et al. 2002). For metaphase spreads, Lake Union male spleen tissue was homogenized in 1 ml of 0.56% potassium chloride (KCl) with a glass dounce. For interphase nuclei, 15 Pacific Ocean embryos at 48 days post fertilization (dpf) were homogenized in 10 ml of 0.56% KCl with a glass dounce. Cells were incubated for 45 min on ice to swell, and 350  $\mu$ l (metaphase spreads) or 250  $\mu$ l (interphase nuclei) of the cell suspension was spun for 10 min at 1950 rpm onto Fisherfinest Premium Superfrost microscope slides through a Fisher single cytology funnel (Fisher Scientific, USA) using a Cytospin 3 (Shandon, USA). Slides were immediately placed in ice-cold methanol for 20 min followed by a 10 min room temperature incubation in 4% formaldehyde in phosphate-buffered saline (PBS). Slides were washed 10 min in PBST (PBS, 0.05% Tween-20) and then incubated with block solution (PBST, 1% BSA, 0.02% sodium azide) for 20 min at room temperature. Primary antibody (rabbit, anti-stickleback CENP-A) was added at 1:300 in block solution for 4 h at room temperature. Slides were washed 3 $\times$ 5 min in PBST at room temperature. A secondary Alexa Fluor 488 chicken, anti-rabbit IgG antibody (Life Technologies, USA) was added at 1:500 in block solution and incubated overnight at 4°C. Slides were washed 3 $\times$ 5 min in PBST, then counterstained with vectashield 4',6'-Diamidino-2-phenylindole (DAPI) stain (Vector Laboratories, USA) for 10 min before imaging as described in the Microscopy section.

### **CENP-A chromatin immunoprecipitation (ChIP)**

**Chromatin preparation from tissue:** The ChIP protocol was performed using the SimpleChIP Plus Enzymatic Chromatin IP kit (Magnetic Beads) (Cell Signaling Technology, USA). Approximately 0.1 g of kidney and liver tissue was dissected from two Pacific Ocean females for two independent samples. The tissue was placed into 2 ml of 1X protease inhibitor cocktail (PIC) buffer. The tissue was dounced 3 times with a glass dounce,

and the dounce was rinsed with an additional 2 ml of 1X PIC buffer. 4 ml of the tissue suspension was incubated with 180  $\mu$ l of 37% formaldehyde for 20 min at room temperature on a nutator. From this point, the Simple ChIP protocol was followed with the following modifications: cells were incubated for 30 min at 37°C with 1.6  $\mu$ l of micrococcal nuclease (MNase) to create mono and dinucleosomes. MNase treatment was stopped with addition of 40  $\mu$ l 0.5 M ethylenediaminetetraacetic acid (EDTA). Cells were spun at 13,000 rpm for 1 min at 4°C and resuspended in 400  $\mu$ l 1X ChIP buffer. Nuclear membranes were disrupted with 3 $\times$ 30 sec sonication using a probe sonicator on high, with a 30 sec incubation on wet ice between pulses. Lysates were clarified by spinning at 10,000 rpm for 10 min at 4°C. 30  $\mu$ l of the supernatant was taken for chromatin analysis and quantification, and the rest was stored at -80°C until immunoprecipitation (IP).

**Immunoprecipitation:** 2  $\times$  10  $\mu$ g of chromatin was diluted in 500  $\mu$ l 1X ChIP buffer. For each sample, 10  $\mu$ l was taken and stored at -20°C for use as the input samples. For each 10  $\mu$ g IP, chromatin was incubated with 4.64  $\mu$ g of rabbit, anti-stickleback CENP-A antibody in 1X ChIP buffer overnight at 4°C with rotation. 30  $\mu$ l of magnetic beads were added to each 10  $\mu$ g IP. Following a 2 h incubation at 4°C with rotation, experimental methods were carried out following the Simple ChIP kit instructions. Input and IP DNA was purified using a MinElute PCR purification kit (Qiagen, USA). Two IP and two input samples were each eluted in 11  $\mu$ l EB buffer. Paired end 150 bp sequencing was performed on the Illumina HiSeq 2500 (Illumina, USA) at the Fred Hutchinson Cancer Research Center Genomics Shared Resource. Fastq files containing the raw sequencing data have been deposited to the NCBI Sequence Read Archive (study accession SRP063504).

**ChIP analysis:** CENP-A ChIP data was analyzed using the pipeline described by (Henikoff et al. 2015). Paired reads were merged using SeqPrep (<https://github.com/jstjohn/SeqPrep>). We then discarded any CENP-A ChIP reads that aligned to the BROADS1 assembly of the *G. aculeatus* genome (Jones et al. 2012), using bwa v0.7.12 (Li and Durbin 2009). The remaining merged pairs (9,182,498 from Pacific Ocean female 1 (POF1) and 10,208,973 from Pacific Ocean female 2 (POF2)) were run through CD-HIT-EST (Li and Godzik 2006, Fu et al. 2012) to create a set of the most abundant IP clusters, each cluster sharing at least 90% sequence identity. The longest read for each cluster was used a reference sequence for that cluster. The clusters were ranked by the number of reads, and the top 500 cluster reference sequences were identified. For each cluster reference sequence, we first counted the number of reads in both the total IP and input samples (all mapped and unmapped reads), and then normalized these counts by calculating the reads per million (RPM) for both the IP and input samples. We calculated the fold enrichment of each cluster reference sequence in the IP relative to input by dividing the IP RPM by the input RPM. The top 500 most abundant cluster references were then ranked by order of fold enrichment. Sequences with a fold enrichment of less than one were discarded. The total average fold enrichment for all of the top enriched clusters combined was calculated by dividing the total IP RPM by the total input RPM.

The cluster reference sequences determined from above were aligned to the most abundant cluster sequence (Fig. S2). The Geneious (Biomatters, New Zealand) sequence global

alignment program (parameter: cost matrix 65% similarity, gap open penalty = 12, gap extension penalty = 3) was used for alignments. From this alignment, a consensus sequence repeating unit was determined using the >50% identity “strict” parameter in Geneious. A consensus repeating unit was determined for each independent IP sample. The two consensus repeats were then aligned to create a single consensus centromeric repeat sequence (CEN) (Fig. S3).

To find the percentage of reads from each ChIP experiment that contained sequences with homology to the CEN, we used BWA-MEM (Li 2013) to align all reads from either the input or the IP to the consensus repeat sequence. Only reads with a score >30 were counted where the score is dependent on: B INT ([4]), O INT [,INT] ([6,6]), E INT [,INT] (‘{O}+{E}\*k’ [1,1], and L INT [,INT] ([5,5]).

To find assembled scaffolds that contain sequences homologous to the centromeric repeat sequence, we conducted a BLAST search for the CEN sequence in the threespine stickleback genome assembly (Ensembl BROAD S1; Feb 2006). We identified sequences with homology to the CEN sequence in scaffolds from regions of the genome that were not assigned to chromosome assemblies and in scaffolds at the edges of gaps in chromosome assemblies that correspond to the putative position of centromeres as determined using the p:q chromosome arm length ratios (Urton et al. 2011). Up to 5 kb of sequence data from each region was extracted, and the consensus centromeric repeat was aligned to these sequences in Geneious (Biomatters, New Zealand) to identify tandem repeats and to determine percent identity between the consensus centromere repeat and the repeats present in the genome assembly.

### Fluorescence in situ hybridization (FISH) on metaphase spreads and interphase nuclei

**CEN FISH probe:** PCR primers JC103 (5’GGTGCTAGATTTAGGAAAACA 3’) and JC106 (5’GTGCATTCATGACTTTTAAGG 3’) were used to amplify the threespine stickleback centromeric repeat from genomic DNA. The PCR product appeared as a ladder. The entire ladder was extracted using the QIAquick gel extraction kit (Qiagen, USA). The purified PCR product was cloned into the PCR2.1 vector (Life Technologies, USA). The clone used as a template to make the FISH probe contained 1.5 copies of the centromeric repeat. This clone was amplified using primers JC103 and JC106 with PCR fluorescein labeling mix (Roche Applied Science, Switzerland) to make a fluorescein-12-dUTP labeled probe of 288 bp. The probe was lyophilized at 55°C for 3 hours, then resuspended in 10 µl hybridization buffer (50% formamide, 2X saline-sodium citrate (SSC), 10% dextran sulfate, 0.2 mM Ethylenediaminetetraacetic acid (EDTA), 2 mM Tris pH 8.0). The probe was denatured at 72°C for 5 min and then stored at 37°C until use.

**Sex chromosome FISH probe:** The *G. aculeatus* BAC clone CHORI-213 101E08 was used as a FISH probe to distinguish the X and Y chromosomes. This BAC clone hybridizes to the end of the p arm of the Y chromosome and to the middle of the p arm of the X chromosome (Ross and Peichel 2008). The BAC FISH probe was labeled with Alexa-568 following the protocol in (Urton et al. 2011).

**Metaphase spreads and interphase nuclei:** Wild-caught fish from Lake Union were injected with 10  $\mu$ l of 1% colchicine in PBS for 16 h to arrest mitotic cells. Spleen and liver tissue was dissected and homogenized in 2 ml of 0.56% KCl using a glass dounce. Metaphase spreads and interphase nuclei were prepared following the protocol described in (Ross and Peichel 2008).

**Hybridization:** Slides were washed in 2X SSC for 5 min, then in denature solution (2X SSC, 70% formamide) for 2 min in a 72°C water bath. Slides were dehydrated in 70% cold ethanol for 2 min, followed by a room temperature dehydration series for 2 min each in 85%, 90%, and 100% ethanol. After the slides were air dried for 5 min, 10  $\mu$ l of either the CEN probe, the sex chromosome probe, or both was added. Then, the slides were covered with a coverslip, sealed with rubber cement, and hybridized at 37°C overnight in a humid chamber. Slides were washed in a 42°C water bath 3 $\times$ 5 min in wash solution 1 (2X SSC, 50% formamide), followed by 3 $\times$ 5 min in 2X SSC. Slides were air dried 5 min, then counterstained and mounted with vectashield DAPI stain. Images were taken as described in the Microscopy section.

### Immunofluorescence-FISH

For IF-FISH, 15 Lake Union embryos at 48 dpf were dounced in 10 ml 0.56% KCl and 0.1% Tween-20 using a glass dounce. Metaphase and interphase nuclei slides were prepared following the protocol in the Immunofluorescence section, with the following modifications: primary rabbit, anti-stickleback CENP-A antibody was diluted 1:100 in block, and 100  $\mu$ l was added to the slide, which was covered with a glass coverslip and incubated in a humid chamber overnight at 4°C. Slides were washed 3 $\times$ 5 min with PBST. The secondary goat, anti-rabbit 568 antibody (Invitrogen, USA) was diluted 1:600 in block, and 100  $\mu$ l was added to the slide and incubated 4 h at room temperature in the dark. Slides were washed 3 $\times$ 5 min in PBST, then post fixed with 4% formaldehyde in PBS for 10 min at room temperature. 10  $\mu$ l of PCR labeled CEN probe was lyophilized at 55°C for 4 h and resuspended in 10  $\mu$ l of hybridization buffer (50% formamide, 2X SSC, 10% dextran sulfate, 20% TE pH 8). The probe was denatured at 72°C for 5 min in a water bath and then put at 37°C until use. Hybridization with the CEN probe was performed following the protocol in the Hybridization section.

### Microscopy

Slides were imaged using a Nikon Eclipse 80i microscope (Nikon, Japan) with an automated filter turret (Chroma filters 31000v2 (DAPI), 41001 (FITC), and 41004 (Texas Red); Chroma, USA). Images were taken using the 100X objective, using the Photometrics CoolSNAP ES2 camera (Photometrics, USA). NIS Elements imaging software (BR 3.00, SP7, Hotfix8, Build 548, Nikon, Japan) was used to pseudo-color the images. For CENP-A immunofluorescence images, the Alexa Fluor-488 labeled antibody and DAPI were pseudo-colored purple and grey, respectively. For FISH images, the Alexa Fluor-568 labeled BAC probe and DAPI were pseudo-colored purple and grey, respectively; the fluorescein labeled CEN probe is green. For IF-FISH images, the Alexa Fluor-568 labeled CENP-A antibody, and DAPI were pseudo-colored purple and grey, respectively.

## Results

### Identification of the threespine stickleback CENP-A coding sequence

Using 3' RACE, PCR, and cDNA sequencing, we identified the entire coding sequence of the threespine stickleback CENP-A gene, which is 447 bp and encodes a protein of 148 amino acids (Fig. S1a). Interestingly, the amino acid sequences between Pacific Ocean and Japan Sea populations differ by a single amino acid at position 36 (GenBank accession numbers [KT321854](#) (Pacific Ocean), [KT321855](#) (Japan Sea)). The Pacific Ocean sequence is used as the reference here. Like other known CenH3-like proteins, the threespine stickleback CENP-A protein has an N-terminal tail that is highly diverged from the threespine stickleback H3 protein (Fig. S1b). The N-terminal tail is also highly diverged among CENP-A proteins from different fish species, while the C-terminal histone fold domain is more conserved (Fountain and Kral 2011).

### Threespine stickleback CENP-A is centromere specific

To determine whether the threespine stickleback CENP-A protein is a centromeric histone variant, we designed an antibody that would recognize the N-terminal end of the CENP-A, but not the H3, protein (Fig. S1b). A Western blot on protein lysate from threespine stickleback tissue shows that the CENP-A antibody is specific to a prominent band of around 17 kD, consistent with its predicted atomic weight of 16.7 kD (Fig. S1c). The antibody hybridizes to punctate spots on interphase nuclei (Fig. 1a) and to the primary constriction of all chromosomes on metaphase spreads (Fig. 1b,c). Taken together, these data suggest that this is a bona fide centromeric histone.

### CENP-A associated sequences in threespine stickleback

To identify the centromere sequence, we performed CENP-A ChIP-seq in two independent female fish from the Pacific Ocean population. Using the cluster analysis pipeline described in (Henikoff et al. 2015), we identified the 500 most abundant sequences in each IP sample and then determined that the majority of these sequences (320 in POF1 and 299 in POF2) were enriched in the IP sequences relative to input sequences (Table 1). After aligning these enriched sequences (Fig. S2), we found a single consensus repeating unit for each individual IP sample. These two consensus repeats are 99.5% identical and were aligned to create a single consensus repeat (Fig. S3). The consensus repeat is 186 bp with an AT-content of 62.9% (Fig. 2a), and was 17.04 or 17.68-fold enriched in the IP relative to input in two independent ChIP-seq experiments (Table 1). This repeat made up 4.43 and 4.29% of total reads from each input sample, highlighting its overall abundance in the genome. Like many other centromeric repeats, this consensus repeat is AT-rich (Melters et al. 2013), contains a putative centromere protein B (CENP-B) box (Fig. 2b) (Masumoto et al. 2004; Edwards and Murray 2005; Henikoff et al. 2015), and has a sequence length consistent with wrapping around a single nucleosome (Willard 1991; Shelby et al. 1997; Henikoff et al. 2001).

Nearly all of the sequences (316/320 from POF1; 289/299 from POF2) that were enriched in the CENP-A IP relative to the input aligned to the consensus repeat (Table 1). The few enriched sequences that did not align to the consensus repeat (4 from POF1; 10 from POF2) did not align to each other and were low in abundance, with less than 150 total reads in the



IP and 84 total reads in the input. It is possible that these low abundance sequences are bound by CENP-A and are interspersed among long stretches of the consensus tandem repeat. However, our data suggests that the majority of sequences bound to CENP-A in the threespine stickleback are variants of the consensus repeat that we identified (Table 1).

### **The CENP-A associated sequence is centromere specific**

To determine whether the sequence is the bona fide centromere, we generated a fluorescently labeled probe of the consensus repeat sequence identified by ChIP-seq and found that it hybridizes to distinct loci in interphase nuclei (Fig. 3a), as well as to the constriction on 41 of 42 metaphase chromosomes in male threespine sticklebacks (Fig. 3b,c). IF-FISH also shows that the centromere repeat co-localizes with CENP-A in both interphase nuclei (Fig. 4a), as well as on metaphase spreads (Fig. 4b,c). All of these results suggest that the 186 bp repeat, now referred to as GacCEN (GenBank accession number [KT321856](#)), is the centromeric repeat in threespine stickleback fish.

### **The GacCEN probe shows weak hybridization to the Y chromosome**

In mammals, the Y chromosome centromere can differ from the centromere on the other chromosomes. For example, the house mouse Y centromere is comprised of a novel satellite repeat (Pertile et al. 2009), and the human Y centromere is a divergent alpha satellite repeat (Wolfe et al. 1985; Miga et al. 2014). Thus, we hypothesized that the single chromosome that did not hybridize to the GacCEN probe in males was the Y (Fig. 3b). Indeed, when we performed FISH with the GacCEN probe and a BAC probe that distinguishes the X and Y chromosomes (Ross and Peichel 2008), we found that the Y chromosome centromere shows very weak hybridization to this centromeric repeat, while the X chromosome shows strong hybridization (Fig. 5).

### **GacCEN repeat shows a decrease in percent identity outside of the core centromere**

We searched for sequences with homology to the consensus centromeric repeat in the current stickleback genome assembly (Jones et al. 2012; Ensembl BROADS1) to see if we could uncover any large scaffolds containing the GacCEN repeat. We found multiple copies of a similar repeat on four scaffolds that did not map to chromosome assemblies and at the edges of ten different scaffolds from nine assembled chromosomes, corresponding to the locations of centromeric constrictions in metaphase spreads (Urton et al. 2011). As predicted (Smith 1976; Schueler et al. 2001; Henikoff 2002; Shepelev et al. 2009; Henikoff et al. 2015), the repeats found at the edge of the assembled contigs show a decrease in percent identity to the GacCEN as they move away from the core centromere (Fig. S4). None of the GacCEN containing contigs appear to have any higher order repeat (HOR) structure. However, until longer centromere-containing reads are obtained, it will be difficult to determine whether stickleback centromeres in fact have a HOR structure.

## **Discussion**

By identifying the complete coding sequence of the threespine stickleback CENP-A protein and designing a species-specific antibody, we were able to perform an unbiased CENP-A ChIP-seq experiment to identify the threespine stickleback centromere repeat. We validated

this repeat as centromeric by performing FISH, as well as IF-FISH with the CENP-A antibody on metaphase spreads. This repeat shares similar characteristics to other centromere repeats in its size, AT content, and putative CENP-B box (Alkan et al. 2011, Melters et al. 2013). It should be noted that the putative *G. aculeatus* CENP-B box shows weak conservation to the evolutionarily conserved human CENP-B domains (Alkan et al. 2011), yet shows the same number of conserved bases as the putative dog, platypus, and *Xenopus* CENP-B boxes (Fig. 2b). The lack of a fully conserved CENP-B box was consistent across all individual cluster repeats, suggesting there are no small subsets of active CENP-B box containing repeats. Interestingly, we also find no evidence for a CENP-B gene in the *G. aculeatus* genome assembly. Although it is not clear whether the CENP-B box in the GacCEN is functional, it is interesting that the repeat shares similar sequence characteristics with other known centromeric repeats.

A previous study (Melters et al. 2013) used a bioinformatic pipeline to identify candidate centromere sequences in 282 species, including two putative centromere sequences of 313 bp and 186 bp in threespine stickleback (Table S4 in Melters et al. 2013). While they could not identify the source of the discordance between the two potential repeats, they assumed that the 313 bp sequence was the centromere because of its higher abundance in the genome. However, we tested this repeat using FISH and found that it does not hybridize to the centromere on metaphase chromosomes (data not shown). In fact, the 186 bp repeat that they identified is 93.0% similar to the GacCEN repeat identified in our study (D. Melters personal communication, July 7, 2015). These results underscore the importance of FISH and/or CENP-A ChIP validation to discern between multiple candidate sequences identified by other methods.

We also previously attempted to identify the *G. aculeatus* centromere sequence using methods that do not rely on a species-specific CENP-A antibody. We used the RepeatNet program (Alkan et al. 2011), which identifies the most abundant sequences in a sequenced genome. In addition, we used a restriction digest approach to identify abundant repeats in the genome. While both computational and restriction digest methods are easy to use and cheaper than ChIP-seq, they generated a large list of potential repeats. To narrow down our list of putative sequences, we assumed, as is common, that the centromeric repeat would not map to the assembled regions of the genome, and that it would be one of the most abundant repeats in the unassembled genome sequences (Alkan et al. 2011; Melters et al. 2013). Based on these criteria, we made a list of candidates to test by FISH, but none hybridized to the centromere. However, after performing the ChIP-seq experiment, we realized that the GacCEN repeat was actually identified using both the computational and the restriction digest methods. Because the edges of many genome assembly scaffold edges contained a few, more diverged, copies of this repeat, our analysis showed that the putative sequences mapped to distinct chromosomes. Furthermore, the GacCEN repeat was not as abundant as other repeats we identified. For these two reasons, the GacCEN repeat sequence was not among the first to be tested. Our experience highlights that other approaches can be used to identify centromeric repeats in organisms without a known CENP-A antibody or an available epitope-tagged CENP-A protein. However, CENP-A ChIP-seq is still the most unbiased way to identify or validate a centromeric repeat.

Our work also demonstrates that it is possible in some cases to identify remnants of centromere repeats on the edges of genome assembly gaps. We found some repeats on the edges of assembled scaffolds that were slightly more divergent than the core centromere sequences identified by ChIP-seq (Fig. S4). This is consistent with what has been seen in human CENP-A ChIP-seq experiments in which the sequence homology among  $\alpha$ -satellite repeats decreases with distance from the core functional centromere (Henikoff et al. 2015). Unequal crossing over at nearly identical repeats is thought to lead to the homogenization of the core centromere, with mutation leading to divergence and unique differences in repeats outside of the core centromere (Smith 1976; Schueler et al. 2001; Henikoff 2002; Shepelev et al. 2009; Henikoff et al. 2015). These differences allow for the assembly of the centromeric repeats. However, Henikoff et al. 2015 showed that more diverged repeats may lose their centromere function, implying that the core, unassembled centromere repeats make up functional centromeres. Our initial filtering step to remove all CENP-A ChIP reads that mapped to the assembled genome actually discarded the repeats we later found on the edges of genome assembly gaps. While these similar repeat may still have centromeric function, by discarding any repeats that mapped to the genome, we ensured that our methods enriched for the core, functional centromere sequences.

Finally, while we have determined that 41 out of the 42 threespine stickleback chromosomes show hybridization with the GacCEN probe in males, the Y centromere appears to show weak hybridization. This could be due to either a different or divergent Y specific centromere repeat (Pertile et al. 2009; Miga et al. 2014), or to a decrease in the number of repeats on the Y centromere array. Future work will use a similar unbiased ChIP-seq approach in both males and females to identify the Y-chromosome centromere repeat.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements

We thank Sue Biggins and Steve Henikoff for reading this manuscript. We thank Ryan Basom of the Fred Hutchinson Cancer Research Center Genomics Shared Resource for help with ChIP-seq data analysis, Jaki Braggin for help with the Western blots, and the Henikoff, Malik and Peichel labs for helpful discussions. This research was supported by a National Science Foundation Graduate Research Fellowship (DGE-1256082), the National Institutes of Health Chromosome Metabolism and Cancer Training Grant (T32 CA009657), and the Fred Hutchinson Cancer Research Center.

## Abbreviations

<b>BAC</b>	Bacterial artificial chromosome
<b>BSA</b>	Bovine serum albumin
<b>CEN</b>	Centromeric repeat sequence
<b>CenH3</b>	Centromeric histone H3
<b>CENP-A</b>	Centromere protein A

<b>CENP-B</b>	Centromere protein B
<b>ChIP</b>	Chromatin immunoprecipitation
<b>ChIP-seq</b>	Chromatin immunoprecipitation sequencing
<b>DAPI</b>	4',6'-Diamidino-2-phenylindole
<b>DPF</b>	Days post fertilization
<b>EDTA</b>	Ethylenediaminetetraacetic acid
<b>FISH</b>	Fluorescence in situ hybridization
<b>GacCEN</b>	Threespine stickleback ( <i>Gasterosteus aculeatus</i> ) centromeric repeat sequence
<b>H3</b>	Histone H3
<b>HOR</b>	Higher order repeat
<b>IF-FISH</b>	Immunofluorescence combined with FISH
<b>IP</b>	Immunoprecipitation
<b>KCl</b>	Potassium chloride
<b>MNase</b>	Micrococcal nuclease
<b>PBS</b>	Phosphate-buffered saline
<b>PBST</b>	Phosphate-buffered saline Tween-20
<b>PCR</b>	Polymerase chain reaction
<b>PMSF</b>	Phenylmethanesulfonylfluoride
<b>POF1</b>	Pacific Ocean female 1
<b>POF2</b>	Pacific Ocean female 2
<b>RPM</b>	Reads per million
<b>SDS</b>	Sodium dodecyl sulfate
<b>SDS-PAGE</b>	Sodium dodecyl sulfate polyacrylamide gel electrophoresis
<b>SSC</b>	Saline-sodium citrate
<b>3' RACE</b>	Rapid amplification of cDNA ends


## References

- Alkan C, Cardone MF, Catacchio CR, et al. (2011) Genome-wide characterization of centromeric satellites from multiple mammalian genomes. *Genome Res* 21:137–145 [PubMed: 21081712]

- Amor DJ, Bentley K, Ryan J, et al. (2004) Human centromere repositioning “in progress”. *Proc Natl Acad Sci USA* 101:6542–6547 [PubMed: 15084747]
- Blower MD, Sullivan BA, Karpen GH (2002) Conserved organization of centromeric chromatin in flies and humans. *Dev Cell* 2:319–330 [PubMed: 11879637]
- Crollius HR, Jaillon O, Dasilva C, et al. (2000) Characterization and repeat analysis of the compact genome of the freshwater pufferfish *Tetraodon nigroviridis*. *Genome Res* 10:939–949 [PubMed: 10899143]
- Drinnenberg IA, deYoung D, Henikoff S, Malik HS (2014) Recurrent loss of CenH3 is associated with independent transitions to holocentricity in insects. *eLife* 3: e03676
- Earnshaw WC, Rothfield N (1985) Identification of a family of human centromere proteins using autoimmune sera from patients with scleroderma. *Chromosoma* 91:313–321 [PubMed: 2579778]
- Edwards NS, Murray AW (2005) Identification of *Xenopus* CENP-A and an associated centromeric DNA repeat. *Mol Biol Cell* 16:1800–1810 [PubMed: 15673610]
- Fountain DM, Kral LG (2011) Isolation and characterization of the *Etheostoma tallapoosae* (Teleostei: Percidae) CENP-A gene. *Genes* 2:829–840 [PubMed: 24710294]
- Fu L, Niu B, Zhu Z, et al. (2012) CD-HIT: accelerated for clustering the next-generation sequencing data. *Bioinformatics* 28:3150–3152 [PubMed: 23060610]
- Garrido-Ramos MA, Jamilena M, Lozano R, et al. (1994) Cloning and characterization of a fish centromeric satellite DNA. *Cytogenet Cell Genet* 65:233–237 [PubMed: 8258296]
- Gong Z, Wu Y, Koblizkova A, et al. (2012) Repeatless and repeat-based centromeres in potato: implications for centromere evolution. *Plant Cell* 24:3559–3574 [PubMed: 22968715]
- Gordon DJ, Resio B, Pellman D (2012) Causes and consequences of aneuploidy in cancer. *Nat Rev Genet* 13:189–203 [PubMed: 22269907]
- Haaf T, Schmid M, Steinlein C, et al. (1993) Organization and molecular cytogenetics of a satellite DNA family from *Hoplias malabaricus* (Pisces, Erythrinidae). *Chromosome Res* 1:77–86 [PubMed: 8143092]
- Henikoff JG, Thakur J, Kasinathan S, Henikoff S (2015) A unique chromatin complex occupies young  $\alpha$ -satellite arrays of human centromeres. *Science Advances* 1:e1400234 [PubMed: 25927077]
- Henikoff S (2002) Near the edge of a chromosome’s “black hole”. *Trends Genet* 18:165–167 [PubMed: 11932007]
- Henikoff S, Ahmad K, Malik HS (2001) The centromere paradox: stable inheritance with rapidly evolving DNA. *Science* 293:1098–1102 [PubMed: 11498581]
- Houben A, Schroeder-Reiter E, Nagaki K, et al. (2007) CENH3 interacts with the centromeric retrotransposon *cereba* and GC-rich satellites and locates to centromeric substructures in barley. *Chromosoma* 116:275–283 [PubMed: 17483978]
- Hunt P, Hassold T (2010) Female meiosis: coming unglued with age. *Curr Biol* 20:R699–R702 [PubMed: 20833308]
- Jones FC, Grabherr MG, Chan YF, et al. (2012) The genomic basis of adaptive evolution in threespine sticklebacks. *Nature* 484:55–61 [PubMed: 22481358]
- Kingsley DM, Peichel CL (2007) The molecular genetics of evolutionary change in sticklebacks In: Östlund-Nilsson S, Mayer I, Huntingford FA, eds. *Biology of the Three-Spined Stickleback*. Boca Raton: CRC press, pp 41–81
- Kitano J, Mori S, Peichel CL (2007) Phenotypic divergence and reproductive isolation between sympatric forms of Japanese threespine sticklebacks. *Biol J Linn Soc* 91:671–685
- Kitano J, Ross JA, Mori S, et al. (2009) A role for a neo-sex chromosome in stickleback speciation. *Nature* 461:1079–1083 [PubMed: 19783981]
- Kops GJPL, Weaver BAA, Cleveland DW (2005) On the road to cancer: aneuploidy and the mitotic checkpoint. *Nat Rev Cancer* 5:773–785 [PubMed: 16195750]
- Lee H-R, Zhang W, Langdon T, et al. (2005) Chromatin immunoprecipitation cloning reveals rapid evolutionary patterns of centromeric DNA in *Oryza* species. *Proc Natl Acad Sci USA* 102:11793–11798 [PubMed: 16040802]
- Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM. arXiv:1303.3997

- Li H, Durbin R (2009) Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25:1754–1760 [PubMed: 19451168]
- Li W, Godzik A (2006) Cd-hit: a fast program for clustering and comparing large sets of protein or nucleotide sequences. *Bioinformatics* 22:1658–1659 [PubMed: 16731699]
- Lister LM, Kouznetsova A, Hyslop LA, et al. (2010) Age-related meiotic segregation errors in mammalian oocytes are preceded by depletion of cohesin and Sgo2. *Curr Biol* 20:1511–1521 [PubMed: 20817533]
- Maio JJ (1971) DNA strand reassociation and polyribonucleotide binding in the African green monkey, *Cercopithecus aethiops*. *J Mol Biol* 56:579–595 [PubMed: 4995480]
- Malik HS, Henikoff S (2009) Major evolutionary transitions in centromere complexity. *Cell* 138:1067–1082 [PubMed: 19766562]
- Manuelidis L (1978) Chromosomal localization of complex and simple repeated human DNAs. *Chromosoma* 66:23–32 [PubMed: 639625]
- Masumoto H, Masukata H, Muro Y, et al. (1989) A human centromere antigen (CENP-B) interacts with a short specific sequence in alphoid DNA, a human centromeric satellite. *J Cell Biol* 109:1963–1973 [PubMed: 2808515]
- Masumoto H, Nakano M, Ohzeki J-I (2004) The role of CENP-B and alpha-satellite DNA: de novo assembly and epigenetic maintenance of human centromeres. *Chromosome Res* 12:543–556 [PubMed: 15289662]
- Melters DP, Bradnam KR, Young HA, et al. (2013) Comparative analysis of tandem repeats from hundreds of species reveals unique insights into centromere evolution. *Genome Biol* 14:R10 [PubMed: 23363705]
- Miga KH, Newton Y, Jain M, et al. (2014) Centromere reference models for human chromosomes X and Y satellite arrays. *Genome Res* 24:697–707 [PubMed: 24501022]
- Morales C, Sánchez A, Bruguera J, et al. (2007) Cytogenetic study of spontaneous abortions using semi-direct analysis of chorionic villi samples detects the broadest spectrum of chromosome abnormalities. *Am J Med Genet* 146A:66–70
- Nagaki K, Cheng Z, Ouyang S, et al. (2004) Sequencing of a rice centromere uncovers active genes. *Nat Genet* 36:138–145 [PubMed: 14716315]
- Nagaki K, Kashihara K, Murata M (2008) A centromeric DNA sequence colocalized with a centromere-specific histone H3 in tobacco. *Chromosoma* 118:249–257 [PubMed: 19048266]
- Nagaki K, Murata M (2005) Characterization of CENH3 and centromere-associated DNA sequences in sugarcane. *Chromosome Res* 13:195–203 [PubMed: 15861308]
- Nagaki K, Talbert PB, Zhong CX, et al. (2003) Chromatin immunoprecipitation reveals that the 180-bp satellite repeat is the key functional DNA element of *Arabidopsis thaliana* centromeres. *Genetics* 163:1221–1225 [PubMed: 12663558]
- Ohzeki J-I, Nakano M, Okada T, Masumoto H (2002) CENP-B box is required for de novo centromere chromatin assembly on human alphoid DNA. *J Cell Biol* 159:765–775 [PubMed: 12460987]
- Palmer DKD, O'Day KK, Margolis RLR (1989) Biochemical analysis of CENP-A, a centromeric protein with histone-like properties. *Prog Clin Biol Res* 318:61–72 [PubMed: 2626439]
- Palmer DK, O'Day K, Trong HL, et al. (1991) Purification of the centromere-specific protein CENP-A and demonstration that it is a distinctive histone. *Proc Natl Acad Sci USA* 88:3734–3738 [PubMed: 2023923]
- Pertile MD, Graham AN, Choo KHA, Kalitsis P (2009) Rapid evolution of mouse Y centromere repeat DNA belies recent sequence stability. *Genome Res* 19:2202–2213 [PubMed: 19737860]
- Piras FM, Nergadze SG, Magnani E, et al. (2010) Uncoupling of satellite DNA and centromeric function in the genus *Equus*. *PLoS Genet* 6:e1000845 [PubMed: 20169180]
- Revenkova E, Herrmann K, Adelfalk C, Jessberger R (2010) Oocyte cohesin expression restricted to predictate stages provides full fertility and prevents aneuploidy. *Curr Biol* 20:1529–1533 [PubMed: 20817531]
- Ricke RM, van Deursen JM (2013) Aneuploidy in health, disease, and aging. *J Cell Biol* 201:11–21 [PubMed: 23547028]

- Ross JA, Peichel CL (2008) Molecular cytogenetic evidence of rearrangements on the Y chromosome of the threespine stickleback fish. *Genetics* 179:2173–2182 [PubMed: 18689886]
- Rudd MK, Willard HF (2004) Analysis of the centromeric regions of the human genome assembly. *Trends Genet* 20:529–533 [PubMed: 15475110]
- Schueler MG, Higgins AW, Rudd MK, et al. (2001) Genomic and genetic definition of a functional human centromere. *Science* 294:109–115 [PubMed: 11588252]
- Shang WH, Hori T, Toyoda A, et al. (2010) Chickens possess centromeres with both extended tandem repeats and short non-tandem-repetitive sequences. *Genome Res* 20:1219–1228 [PubMed: 20534883]
- Shelby RD, Vafa O, Sullivan KF (1997) Assembly of CENP-A into centromeric chromatin requires a cooperative array of nucleosomal DNA contact sites. *J Cell Biol* 136:501–513 [PubMed: 9024683]
- Shepelev VA, Alexandrov A, Yurov YB, et al. (2009) The evolutionary origin of man can be traced in the layers of defunct ancestral alpha satellites flanking the active centromeres of human chromosomes. *PLoS Genet* 5: e1000641 [PubMed: 19749981]
- Smith GP (1976) Evolution of repeated DNA sequences by unequal crossover. *Science* 191:528–535 [PubMed: 1251186]
- Sullivan BA, Schwartz S (1995) Identification of centromeric antigens in dicentric Robertsonian translocations: CENP-C and CENP-E are necessary components of functional centromeres. *Hum Mol Genet* 4:2189–2197 [PubMed: 8634687]
- Sullivan KF, Hechenberger M, Masri K (1994) Human CENP-A contains a histone H3 related histone fold domain that is required for targeting to the centromere. *J Cell Biol* 127:581–592 [PubMed: 7962047]
- Tek AL, Kashihara K, Murata M, Nagaki K (2010) Functional centromeres in soybean include two distinct tandem repeats and a retrotransposon. *Chromosome Res* 18:337–347 [PubMed: 20204495]
- Urton JR, McCann SR, Peichel CL (2011) Karyotype differentiation between two stickleback species (*Gasterosteidae*). *Cytogenet Genome Res* 135:150–159 [PubMed: 21921583]
- Warburton PE, Cooke CA, Bourassa S, et al. (1997) Immunolocalization of CENP-A suggests a distinct nucleosome structure at the inner kinetochore plate of active centromeres. *Curr Biol* 7:901–904 [PubMed: 9382805]
- Willard HF (1991) Evolution of alpha satellite. *Curr Opin Genet Dev* 1:509–514 [PubMed: 1822282]
- Wolfe J, Darling SM, Erickson RP, et al. (1985) Isolation and characterization of an alphoid centromeric repeat family from the human Y chromosome. *J Mol Biol* 182:477–485 [PubMed: 4040175]
- Zhong CX (2002) Centromeric retroelements and satellites interact with maize kinetochore protein CENH3. *Plant Cell* 14:2825–2836 [PubMed: 12417704]




**Fig. 1.** The threespine stickleback CENP-A antibody localizes to the centromere. Immunofluorescence reveals that the CENP-A antibody (*purple*) hybridizes to: **(a)** distinct puncta in interphase nuclei from Pacific Ocean embryos; **(b)** a single region on each chromosome in a metaphase spread from a Lake Union male; and **(c)** the primary constriction (*arrowheads*) on each chromosome in a higher magnification view of the boxed region in panel **(b)**. *Scale bar*, 5  $\mu$ m




A	
GAGGTGCTAGTTTTAGGAAAACACTGTTAACCAATGCATTCTTGTGTTCTGTGCGTTTTTCAG CTTCTCTCGGCCTGCAAACGCCTTAATAGTCAAGAATGCACCAAACTTAGTTTGAACAAA <b>AAAGGTTGGAAAAC TATT</b> CACAAACCATGATACCATCATAAAACAGATAAATGTACTTTCCT	
B	
	# conserved bases
<i>H. sapiens</i>	* * * * * * * * * *
<i>C. familiaris</i>	C T T C G T T G G A A A C G G G A
<i>O. anatinus</i>	C T T C A T A T G G A A A T T C G 8/17
<i>X. laevis</i>	C T T T G C T C C C G G C C T G A 8/17
<i>G. aculeatus</i>	T A G T G T T G G A A G C T A A T 8/17
	A A A G G T T G G A A A A C T A T 8/17


**Fig. 2.** Threespine stickleback centromere repeat sequence. **a** The consensus GacCEN repeat is 186 bp and has an AT-content of 62.9%. The putative CENP-B box is highlighted in bold. **b** The putative CENP-B box in the GacCEN shows sequence similarity to the CENP-B box in human (*Homo sapiens*), dog (*Canis familiaris*), platypus (*Ornithorhynchus anatinus*), and the African clawed frog (*Xenopus laevis*). Identical nucleotides to the human CENP-B box are in red and asterisks denote the evolutionary conserved domains in humans (Ohzeki et al. 2002, Alkan et al. 2011). This sequence has been deposited in GenBank (accession number [KT321856](#)).



**Fig. 3.** The threespine stickleback CEN repeat hybridizes to the centromere. FISH reveals that the CEN probe (*green*) hybridizes to: (a) distinct puncta in interphase nuclei from Lake Union embryos; (b) a single region on each chromosome except one (*arrow*) in a metaphase spread from a Lake Union male; and (c) the primary constriction (*arrowheads*) on each chromosome in a higher magnification view of the boxed region in panel (b). *Scale bar*, 5  $\mu\text{m}$



**Fig. 4.** The threespine stickleback centromere repeat colocalizes with CENP-A. IF-FISH with CENP-A antibody (*purple*) shows co-localization with the CEN probe (*green*) in: (a) interphase nuclei and (b, c) metaphase chromosomes from Lake Union embryos. Panel (c) shows a higher magnification view of the boxed region in panel (b). *Scale bar*, 5  $\mu$ m



**Fig. 5.** The threespine stickleback centromere repeat shows weak hybridization to the Y chromosome centromere. **a** FISH reveals that the CEN probe (*green*) shows weak hybridization to the Y chromosome but strong hybridization to the X chromosome on a metaphase spread from a Pacific Ocean male. The X and Y- chromosomes can be identified and distinguished by hybridization to a FISH probe from BAC 101E08 (*purple*) (Ross and Peichel 2008). Higher magnification views of hybridization to the X (**b**) and Y (**c**) chromosomes are shown. *Scale bar*, 5  $\mu$ m

**Table 1.**

Summary of CENP-A ChIP seq data from two independent Pacific Ocean females.

	<b>Pacific Ocean female 1</b>	<b>Pacific Ocean female 2</b>
Number enriched clusters in 500 most abundant	320	299
Number enriched clusters with GacCEN repeat	316	289
Average fold enrichment of IP/input	17.04	17.68
Range of fold enrichment across clusters	1.25 - 87.60	1.01 - 134.00
Total normalized IP RPM for enriched GacCEN containing clusters	307,600.96	308,096.65
Total normalized input RPM for enriched GacCEN containing clusters	18,046.51	17,426.30
Total IP reads	28,309,375	34,297,458
Total IP reads that map to GacCEN (% total)	20,245,271 (71.51%)	23,695,808 (69.09%)
Total input reads	26,102,993	30,137,027
Total input reads that map to GacGEN (% total)	1,155,758 (4.43%)	1,291,918 (4.29%)

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript