

# Sphingomyelin Acyl Chains Influence the Formation of Sphingomyelin- and Cholesterol-Enriched Domains

Oskar Engberg, Kai-Lan Lin, Victor Hautala, J. Peter Slotte, and Thomas K. M. Nyholm\*

(Biophysical Journal 119, 913–923; September 1, 2020)

In the originally published version of this article, Figure 2 was duplicated in Figure 3. Figure 3 has now been corrected online, and the correct Figure 3 appears below.




FIGURE 3 The effect of temperature on the solubility limits of different SMs in the  $L_d$  phase. The solubility of 16:0-SM (A) and 24:0-SM (B) in the  $L_d$  phase was measured in POPC bilayers containing 0 or 20 mol % cholesterol. Above these concentrations, the SMs formed ordered domains. The values are averages of  $\geq 3$  experiments  $\pm$  SD.

\*Correspondence: [tnyholm@abo.fi](mailto:tnyholm@abo.fi)

<https://doi.org/10.1016/j.bpj.2020.11.001>

© 2020 Biophysical Society.






FIGURE 3 The effect of temperature on the solubility limits of different SMs in the  $L_d$  phase. The solubility of 16:0-SM (A) and 24:0-SM (B) in the  $L_d$  phase was measured in POPC bilayers containing 0 or 20 mol % cholesterol. Above these concentrations, the SMs formed ordered domains. The values are averages of  $\geq 3$  experiments  $\pm$  SD.