

Exploring the Chemical Space of Benzothiazole-Based DNA Gyrase B Inhibitors

Žiga Skok,¹ Michaela Barančoková,¹ Ondřej Benek, Cristina Durante Cruz, Päivi Tammela, Tihomir Tomašič, Nace Zidar, Lucija Peterlin Mašič, Anamarija Zega, Clare E. M. Stevenson, Julia E. A. Mundy, David M. Lawson, Anthony Maxwell, Danijel Kikelj, and Janez Ilaš*



Cite This: *ACS Med. Chem. Lett.* 2020, 11, 2433–2440



Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: We designed and synthesized a series of inhibitors of the bacterial enzymes DNA gyrase and DNA topoisomerase IV, based on our recently published benzothiazole-based inhibitor bearing an oxalyl moiety. To improve the antibacterial activity and retain potent enzymatic activity, we systematically explored the chemical space. Several strategies of modification were followed: varying substituents on the pyrrole carboxamide moiety, alteration of the central scaffold, including variation of substitution position and, most importantly, modification of the oxalyl moiety. Compounds with acidic, basic, and neutral properties were synthesized. To understand the mechanism of action and binding mode, we have obtained a crystal structure of compound **16a**, bearing a primary amino group, in complex with the N-terminal domain of *E. coli* gyrase B (24 kDa) (PDB: 6YD9). Compound **15a**, with a low molecular weight of 383 Da, potent inhibitory activity on *E. coli* gyrase (IC₅₀ = 9.5 nM), potent antibacterial activity on *E. faecalis* (MIC = 3.13 μM), and efflux impaired *E. coli* strain (MIC = 0.78 μM), is an important contribution for the development of novel gyrase and topoisomerase IV inhibitors in Gram-negative bacteria.

KEYWORDS: DNA gyrase, topoisomerase IV, GyrB, ParE, antibacterial, benzothiazole

The treatment of infections has advanced beyond recognition since the first discoveries of antimicrobial compounds. Widespread distribution, low price, and high safety profile have established antimicrobial drugs as a cornerstone of modern medicine. However, due to rising resistance issues, the efficacy of existing infection treatments is rapidly decreasing. The dearth of new anti-infectives introduced to the market over recent years, and no discovery of novel classes, has deepened the problem.^{1–4} Antimicrobial resistance is now a serious global health concern; infections that are currently readily treatable could become untreatable in the near future.⁵

DNA topoisomerases are enzymes that alter DNA topology, being involved in important biological processes in the cell (replication, transcription, chromosome condensation, etc.).^{6–11} Bacterial DNA gyrase and DNA topoisomerase IV (topo IV) are attractive targets for antibacterial drug discovery due to their well-described structure and mechanism, as well as their absence from eukaryotes. Both enzymes are type II topoisomerases, sharing high functional and structural similarity. They change the topology of DNA by cleaving


both strands of the double-helix. Gyrase has the unique function of introducing negative supercoiling ahead of the replication fork, while topo IV is involved in chromosome decatenation. The enzymes consist of two pairs of subunits with different functions, forming heterotetrameric structures: A₂B₂ (gyrase) or C₂E₂ (topo IV). The main role of the GyrA and ParC subunits is to release torsional stress by breaking and rejoining strands of the DNA molecule. GyrB and ParE contain the ATP-binding site and provide the energy required for the enzyme function by ATP hydrolysis.^{12,13}

DNA gyrase inhibitors can influence the enzyme action at different levels.¹⁴ Inhibition of the GyrA subunit can cause stabilization of the DNA–enzyme complex, and thus broken

Received: July 29, 2020

Accepted: October 13, 2020

Published: October 15, 2020



strands of DNA cannot be rejoined. GyrA inhibitors, so-called gyrase poisons, are represented by fluoroquinolones, which are widely in clinical use, although struggling with side effects and growing resistance concerns.⁵ These problems have encouraged further research on inhibitors with different mechanisms of action. GyrB inhibitors are not currently in clinical use; novobiocin was withdrawn from the market in the 1960s due to significant side effects. Since then, no other GyrB inhibitor has passed clinical trials.⁷ Low cross-resistance between potential GyrB inhibitors and fluoroquinolones is a promising and motivating factor for further research. Novel gyrase inhibitors based on various chemical structures were discovered in recent years.⁸

Our research group has recently published several articles focusing on ATP-competitive inhibitors of the DNA gyrase B protein (GyrB).^{15–17} The compounds share a common pyrrole-2-carboxamide moiety, originating from marine alkaloids such as oroidin.¹⁸ Compounds are potent inhibitors of gyrase; however, they lack potent antimicrobial activity. The mode of action for this series of compounds was confirmed by solving the crystal structure of a complex of *E. coli* GyrB with a benzothiazole-type inhibitor **1** (PDB 5L3J).¹⁵

Our goal was to explore the chemical space of pyrrole-benzothiazole GyrB inhibitors by selecting those that retain potent enzyme inhibition while optimizing their physicochemical properties. Several strategies of modification illustrated in Figure 1 were followed: alteration of the pyrrole carboxamide




Figure 1. Structure of previously discovered compound **1** (PDB 5L3J) with highlighted positions selected for structure modifications.

moiety, alteration of the central scaffold, including variation of the substitution position, and most importantly modification of the oxalyl moiety.

The 4,5-dibromo-1*H*-pyrrole-2-carboxamide interacting with the protein through hydrogen bonding interactions of the pyrrole nitrogen and adjacent carbonyl with Asp73, either directly or via a network of hydrogen bonds involving conserved water molecule, respectively does not permit substantial changes. Modification to 3,4-dichloro-5-methyl-1*H*-pyrrole-2-carboxamide has several potential beneficial properties. The key role of the pyrrole nitrogen and nonsubstituted carboxamide moiety remains unchanged. Smaller chlorine atoms (compared to bromine) are likely to fit better into the *E. coli* topo IV hydrophobic pocket and the *Staphylococcus aureus* gyrase hydrophobic pocket, which has a smaller volume than the *E. coli* gyrase hydrophobic pocket.¹⁹ The 3,4-dichloro-5-methyl-1*H*-pyrrole-carboxamide moiety is present in the natural antibiotics kibelomycin^{20–22} and amycolamicin^{23,24} and was used in the Astra Zeneca pyrrolamide series of gyrase B inhibitors.^{25–27}


To explore the influence of the central scaffold we interchanged the substituents on positions 2 and 6 of benzothiazole, resulting in interchanging the position of the benzene ring and thiazole ring of the benzothiazole moiety²⁸ in the binding site and its π -cation interactions with Arg76, as well as influencing the acidity of the pyrrole amide proton. An obvious bioisosteric approach was the replacement of the ring

sulfur (benzothiazole) with nitrogen to give a benzimidazole central scaffold. Indeed, benzimidazole-ureas were presented as potent dual inhibitors of bacterial topoisomerases in a recent study by Vertex Pharmaceuticals.²⁹ In this manner, crucial cation- π interactions between the core benzene ring and Arg76 and the Glu50-Arg76 salt bridge interactions were preserved. Additional interactions might have been introduced by enlarging the central scaffold, but we did not plan to increase the scaffold MW.

The focus of the optimization strategy was the replacement of the oxalyl moiety, which is likely too acidic and might impair permeation into bacteria and thus diminish antimicrobial activity. As confirmed by the crystal structure of **1**, the oxalyl moiety extending out of the binding pocket and making only a single direct hydrogen bond to the protein, via the carbonyl group to Arg136, can be replaced without loss of inhibitory activity. We systematically changed the acidic oxalyl moiety to (i) a neutral small aliphatic acyl moiety (acetamide group), (ii) a neutral aromatic moiety (isonicotinamide moiety), (iii) a neutral polar group with H-bond donor/acceptor potential (urea derivative), and (iv) an aliphatic primary amino group (derivatives of glycine and beta alanine).

Synthesis of benzimidazole compounds, illustrated in Scheme 1, starts with coupling of 6-nitro-1*H*-benzo[*d*]-

Scheme 1^a




^aReagents and conditions: (a) 2,2,2-trichloro-1-(4,5-dibromo-1*H*-pyrrol-2-yl)ethan-1-one, Na_2CO_3 , DMF, 80 °C, 16 h; (b) $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$, EtOH, reflux, 12 h; (c) ethyl oxalyl chloride (for **4a**) or methyl malonyl chloride (for **4b**), Et_3N , 1,4-dioxane, rt, 12 h; (d) 1 M NaOH, 1,4-dioxane, rt, 16 h.

imidazol-2-amine with 2,2,2-trichloro-1-(4,5-dibromo-1*H*-pyrrol-2-yl)ethan-1-one and a weak base in DMF. The nitro group of intermediate **2** is reduced using tin(II) chloride dihydrate to give amino-derivative **3**, further acylated with respective acyl chloride to obtain esters **4a–b**. Corresponding carboxylic acids **5a–b** were obtained by alkaline hydrolysis.

Synthesis of benzothiazole-based compounds is illustrated in Scheme 2. Acylation of 6-nitro-benzo[*d*]thiazol-2-amine (**6**) with ethyl oxalyl chloride/methyl malonyl chloride/acetonyl chloride gives intermediates **7a**, **7b**, and **7c** respectively.

The nitro group in position 6 of the benzothiazole was then reduced using hydrogen and palladium catalyst to obtain compounds **8a–c**. Coupling with 3,4-dichloro-5-methyl-1*H*-pyrrole-2-carboxylic acid (activated in situ with oxalyl chloride) in DCM in the presence of pyridine as base gave compounds **9a–c**. The ethyl oxalyl derivative **9a** was further hydrolyzed to carboxylic acid **10**. Synthesis of the reversed isomers with the pyrrole moiety at position 2 is more

Scheme 2^a

^aReagents and conditions: (a) corresponding acyl chloride, Et₃N, 1,4-dioxane, rt, 4 h; (b) H₂, 10% Pd/C, EtOH (for **8a** and **8c**) or MeOH (for **8b**), rt, 24 h; (c) 3,4-dichloro-5-methyl-1H-pyrrole-2-carbonyl chloride, pyridine, DCM; (d) 1 M NaOH, MeOH, rt, 24 h; (e) 3,4-dichloro-5-methyl-1H-pyrrole-2-carbonyl chloride, toluene, reflux, 16 h; (f) SnCl₂·2H₂O, EtOH, reflux, 12 h; (g) ethyl oxalyl chloride, Et₃N, 1,4-dioxane, rt, 8 h (for the synthesis of **13**); (h) Ac₂O, Et₃N, DCM, rt, 2 h (for the synthesis of **15a**); or CDI, DMF, rt, 3 h; NH₃, rt, 16 h (for the synthesis of **15b**); or nicotinic acid, EDC, NMM, HOBT, DMF, rt, 12 h (for the synthesis of **15c**); or corresponding Boc-amino acid, EDC, NMM, HOBT, DMF, rt, 12 h (for the synthesis of **15d–e**); (h) 4 M HCl, 1,4-dioxane, rt, 5 h.

demanding. The coupling of benzothiazole **6** with the pyrrole-2-carbonyl chloride moiety was done in refluxing toluene without any base, to obtain intermediate **11**. The nitro group is

reduced using tin(II) chloride dihydrate to give key amino derivative **12**, which was further derivatized. Acylation with oxalyl chloride moiety gave ester **13**. The corresponding carboxylic acid **14** is isolated after hydrolysis with 1 M sodium hydroxide. Reaction of amino derivative **12** with acetic anhydride in the presence of triethylamine in DCM gave target compound **15a**. To obtain the urea derivative **15b**, aminobenzothiazole **12** was first stirred at RT with CDI in DMF for 3 h and heated to 50 °C overnight after addition of ammonia gas. Compounds **15c–e** were prepared by coupling of **12** with nicotinic acid (**15c**), Boc-β-alanine (**15d**), or Boc-glycine (**15e**) in the presence of coupling agents EDC, HOBT, or NMM, respectively, in DMF. The Boc protective group of **15d–e** was finally removed with HCl in dioxane, to obtain **16a–b**.

Compounds were evaluated in *E. coli* and *S. aureus* gyrase supercoiling assays as well as in *E. coli* and *S. aureus* topo IV relaxation assays. The results are presented in Tables 1–3 as IC₅₀ values or residual activity (RA) of the enzyme in a concentration of 10 μM of the inhibitor.

The benzimidazole-based series was shown to be only weakly active (Table 1), with inhibitory activity against *E. coli* gyrase in the micromolar range. Carboxylic acids **5a–b** were approximately 10-fold more active than the corresponding esters **4a–b**, due to possible ionic interactions additional to hydrogen bonds with Arg136. The most potent compound **5a** (IC₅₀ = 0.60 μM) showed weak activity against *S. aureus* gyrase and *E. coli/S. aureus* topo IV). Direct comparison of benzothiazole **1** with benzimidazole **5a** shows that replacement of sulfur with nitrogen resulted in 10-fold lower enzymatic inhibition against *E. coli* gyrase. Poor activity results and poor solubility of benzimidazoles pointed us back to the benzothiazole central scaffold, and the benzimidazole series was not further extended. Replacement of 4,5-dibromo-1H-pyrrole with 3,4-dichloro-5-methyl-1H-pyrrole indeed resulted in good inhibitory activity against *E. coli* gyrase, but even more importantly it introduced good inhibitory activity against *E. coli* topo IV (**14**: IC₅₀ topo IV = 75 nM) and potent inhibitory activity against *S. aureus* gyrase and topo IV, which was completely absent in the case of the dibromo analogue **1**. When comparing compounds with the pyrrole attached to position 2 (compound **14**) to a regioisomer with the pyrrole attached to position 6 (compound **10**), the inhibitory activity on *E. coli* gyrase is favorable for compound **10** and even more favorable regarding *E. coli* topo IV inhibitory activity as well as *S. aureus* gyrase and topo IV. Overall, compound **14** has


Table 1. Inhibitory Activity of Series I of Compounds with Benzimidazole Central Scaffold

4a: n = 0, R = Et
4b: n = 1, R = Me
5a: n = 0, R = H
5b: n = 1, R = H

Cmpd	n	R	IC ₅₀ (μM) or RA (%) ^a			
			<i>E. coli</i> gyrase	<i>E. coli</i> topo IV	<i>S. aureus</i> gyrase	<i>S. aureus</i> topo IV
4a	0	Et	4.0 ± 1.6 μM	100%	100%	100%
4b	1	Me	7.0 ± 3.4 μM	100%	100%	100%
5a	0	H	0.60 ± 0.32 μM	12 ± 2 μM	80 ± 23 μM	31 ± 1 μM
5b	1	H	1.5 ± 0.2 μM	100%	100%	100%
1			0.058 μM	13 μM	>100 μM	10 μM
novobiocin			0.17 μM	11 μM	0.041 μM	27 μM

^aResidual activity of the enzyme at 10 μM concentration of the compound.


Table 2. Inhibitory Activity of Benzothiazole Compounds with Acidic Terminal Functional Groups



Cmpd	n	R	IC ₅₀ (nM) or RA (%) ^a			
			<i>E. coli</i> gyrase	<i>E. coli</i> topo IV	<i>S. aureus</i> gyrase	<i>S. aureus</i> topo IV
9a	0	Et	290 ± 170 nM	100%	69%	100%
9b	1	Me	200 ± 180 nM	100%	59%	48%
10	0	H	29 ± 16 nM	6 400 ± 3 000 nM	250 ± 130 nM	910 ± 340 nM
13	0	Et	48 ± 12 nM	100%	100%	100%
14	0	H	4.8 ± 2.1 nM	75 ± 28 nM	38 ± 16 nM	290 ± 180 nM
1			58 nM	13 000 nM	>100 μM	10 000 nM
novobiocin			168 nM	11 000 nM	41 nM	27 000 nM

^aResidual activity of the enzyme at 10 μM concentration of the compound.

Table 3. Inhibitory Activity of Benzothiazole Compounds with Neutral or Basic Terminal Functional Groups



Cmpd	R	IC ₅₀ (nM) or RA (%) ^a			
		<i>E. coli</i> gyrase	<i>E. coli</i> topo IV	<i>S. aureus</i> gyrase	<i>S. aureus</i> topo IV
9c		66 ± 8 nM	100%	35 400 nM	100%
12		16 000 ± 4 000 nM	100%	60%	100%
15a	-CH ₃	9.5 ± 2.5 nM	4 600 ± 100 nM	400 ± 120 nM	1 600 ± 300 nM
15b	-NH ₂	26 ± 8 nM	5 200 ± 2 700 nM	1 300 ± 600 nM	5 800 ± 3 100 nM
15c	-pyridine-4-yl	2 500 ± 1 500 nM	100%	100%	100%
15d	-CH ₂ CH ₂ NHBoc	110 ± 20 nM	100%	780 nM	100%
15e	-CH ₂ NHBoc	29 ± 15 nM	60%	260 ± 120 nM	100%
16a	-CH ₂ CH ₂ NH ₃ ⁺ Cl ⁻	110 ± 50 nM	10 000 ± 2 000 nM	1 500 ± 600 nM	290 ± 180 nM
16b	-CH ₂ NH ₃ ⁺ Cl ⁻	280 ± 10 nM	100%	380 nM	100%

^aResidual activity of the enzyme at 10 μM concentration of the compound.

superior enzymatic activity against all four tested enzymes compared to novobiocin. Benzothiazole with 3,4-dichloro-5-methyl-1*H*-pyrrole on position 2 was thus selected to explore the possible replacements of the unfavorable oxalyl moiety. The results of the *E. coli* gyrase inhibitory activity assays reveal that the anionic center is not required for potent inhibitory activity. Compounds with the acetyl moiety, the urea derivative, and the glycine derivative with a free primary amino group all possess *E. coli* gyrase inhibitory activity in the low nanomolar range (10–25 nM). Having an aromatic moiety (15c) pointed to the water environment (and possibly having π–cationic interactions with Arg136) is clearly not optimal for this series of compounds, although such an approach was successful in tricyclic inhibitors of GyrB (PDB: 4KFG).³⁰ Investigation of the Boc-protected amino acid derivatives 15d and 16a reveals that the bulky lipophilic moiety can have favorable binding to *E. coli* GyrB. Although this might seem contradictory, it is known from thermodynamic evaluations that the binding of compound with “unfavorable” lipophilic moieties extending into a water environment can be beneficial as more polar/ionized groups can pay a high desolvation penalty, which contributes to net unfavorable binding.³¹ The

amino compound 12, lacking the carbonyl group, is a very weak binder, which indicates that a carbonyl moiety is a prerequisite for potent enzyme binding. Acetyl derivative 15a with lowest molecular weight in the series and single digit nanomolar binding with IC₅₀ = 9.5 nM seemed very interesting; therefore, regioisomer 9c with a pyrrole attached to position 6 and acetyl to position 2 of benzothiazole was prepared. The trend observed already from previous compounds was the same in this series: the isomer 15a with pyrrole attached to position 2 of benzothiazole was 7-fold more potent than regioisomer with pyrrole attached to position 6.

To confirm binding to the ATP-pocket of GyrB and to gain insight into molecular interactions, crystallization of selected ligands with the 24 kDa fragment (N-terminal subdomain) of GyrB was attempted. Among crystallization attempts with 15a, 16a, and 9c, cocrystals of 16a with *E. coli* GyrB24 were obtained and the structure was solved to a resolution of 1.60 Å (Figure 2). The 3,4-dichloro-5-methyl-1*H*-pyrrole moiety of 16a is bound to the adenine-binding pocket, making a hydrogen bond between the pyrrole NH group and Asp73 side chain, while interaction of the Asp73 side chain and




Figure 2. Crystal structure of the complex formed between **16a** and *E. coli* GyrB 24 kDa fragment (PDB: 6YD9). The protein is depicted in cartoon representation covered by a semitransparent molecular surface. Omit $mF_{\text{obs}} - DF_{\text{calc}}$ positive difference electron density for the ligand at 1.6 Å resolution is depicted at two contour levels: 3σ (magenta mesh) and 12σ (black mesh), with the latter highlighting the locations of the electron-dense chlorine and sulfur atoms. Shown below are the interactions between amino acid residues with ligand (red line: hydrogen bond, blue line: π -cation interactions).

pyrrolamide carbonyl oxygen is bridged by a coordinated water molecule (NHCO–H₂O H-bond distance is 2.61 Å). Comparison with the GyrB crystal structure of kibelomycin (PDB: 4URM)³² and Astra Zeneca's pyrrolamide (PDB: 3TTZ)³³ reveals identical interactions of the 3,4-dichloro-5-methyl-1*H*-pyrrole moiety. There is a strong π -cation interaction (3.28 Å) between the aromatic ring and Arg76. The carbonyl moiety of β -alanine interacts with Arg136

through a H-bond (2.87 Å), while the primary amino group of β -alanine is pointed toward the water environment, making no interactions with the protein. While interactions of the oxalyl moiety of **1** with Arg136 were not apparent (PDB code 5L3J, Resolution: 2.83 Å), improved resolution now offers clear insight into the binding mode. Hydrogen bonding of Arg136 with the carbonyl oxygen of β -alanine is evident, thus demonstrating that a carboxylic acid moiety is not prerequisite for binding, and explains strong enzymatic inhibitory activity of compounds having a neutral side chain.

Compounds were assayed on 4 wild type bacterial strains to determine their antibacterial activity (*E. coli*, *S. aureus*, *Enterococcus faecalis*, *Pseudomonas aeruginosa*) (Table 4). Additionally, we included two other *E. coli* strains, an *E. coli* strain with impaired outer membrane (*lpxC* deletion mutant) and an *E. coli* strain with a defective efflux pump (*tolC* deletion mutant), to evaluate if weak antibacterial activity on wild type could be due to weak permeability and/or strong efflux. Both benzimidazole and benzothiazole inhibitors with the oxalyl moiety (**5a**, **10**, and **14**) were inactive in all bacterial strains (MIC > 50 μM). Ester derivative **9b** which had good enzymatic inhibitory activity (IC₅₀ = 200 nM) showed also a good MIC (1.56 μM) in the *E. coli* strain with a defective efflux pump, indicating that the ester moiety is beneficial for entry into bacteria; however, such compounds are often the substrates for efflux pumps. Ester derivative **13** which had very good enzymatic inhibitory activity (IC₅₀ = 48 nM) showed only weak MIC (50 μM) in the *E. coli* strain with a defective efflux pump.

A significant improvement in MIC value was expected with the compound lacking the acidic oxalyl moiety. Compound **16b**, having a glycine moiety with a free primary amino group, was the only compound showing some antibacterial activity on *S. aureus* wild type (MIC = 25 μM), confirming that good enzymatic inhibitory activity (*S. aureus* gyrase IC₅₀ = 380 nM) with suitable physicochemical properties is needed for antibacterial activity. When looking at Gram-positive *E. faecalis*, the results were more encouraging, as four compounds showed good antibacterial activity. Both compounds with a free amino group (**16a** MIC = 12.5 μM and **16b** MIC = 6.25 μM) had good antibacterial activity. Boc-protected analogue **15e** and compound **15a** with an acetyl moiety had even better antibacterial activity with MIC = 3.13 μM , which also correlated with enzymatic inhibitory potency. When looking at more challenging Gram-negative *E. coli* and *P. aeruginosa*,

Table 4. Antibacterial Activity of Selected Compounds

Cmpd	MIC (μM) ^a					
	<i>E. coli</i> (ATCC 25922)	<i>S. aureus</i> (ATCC 29213)	<i>E. faecalis</i> (ATCC 29212)	<i>P. aeruginosa</i> (ATCC 27853)	<i>E. coli</i> (JD17464) ^b	<i>E. coli</i> (JWS503) ^c
4a , 4b , 5a , 5b , 9a , 9c , 10 , 14 , 15c , 15d	>50 μM	>50 μM	>50 μM	>50 μM	>50 μM	>50 μM
9b	>50 μM	>50 μM	>50 μM	>50 μM	>50 μM	1.56 μM
13	>50 μM	>50 μM	>50 μM	>50 μM	>50 μM	50 μM
15a	>50 μM	>50 μM	3.13 μM	>50 μM	>50 μM	0.78 μM
15b	>50 μM	>50 μM	>50 μM	>50 μM	>50 μM	1.56 μM
15e	>50 μM	>50 μM	3.13 μM	>50 μM	>50 μM	0.78 μM
16a	>50 μM	>50 μM	12.5 μM	>50 μM	>50 μM	12.5 μM
16b	>50 μM	25 μM	6.25 μM	>50 μM	>50 μM	3.13 μM
ciprofloxacin	0.05 μM	1.51 μM	3.02 μM	3.02 μM	0.121 μM	0.015 μM

^aMIC (minimum inhibitory concentration that inhibits the growth of bacteria by $\geq 90\%$) values against *E. coli* and *S. aureus*. ^b*E. coli* strain with impaired outer membrane, *lpxC* deletion mutant. ^c*E. coli* strain with defective efflux pump, *tolC* deletion mutant.

the results were less encouraging, as none of the compounds showed antibacterial activity.

Testing on the *E. coli* strain with impaired outer membrane (*lpxC* deletion mutant), revealed that permeability is not a main issue as tested compounds did not show any antibacterial activity. On the contrary, the compounds were very active when tested on the *E. coli* strain with a defective efflux pump. The most potent compounds were **15e** and **15a** with submicromolar MIC values (MIC = 0.78 μ M). Both are neutral molecules with lipophilic moieties (acetyl and Boc). Compounds with free amino group were also potent antibacterial compounds (**16a** and **16b**) but were not superior to the neutral compounds (**15a** and **15e**). This is somehow contrary to the recent proposal that the introduction of a primary amino group improves entry and causes accumulation in Gram-negative bacteria.³⁴ To confirm the mechanism of uptake for the reported compounds (porin pathway or direct diffusion),³⁵ additional studies should be performed in order to guide optimization.^{36,37} Urea derivative **15b** was also a potent antibacterial compound on the efflux impaired *E. coli* strain (MIC = 1.56 μ M), while it was inactive on all other strains.

In summary, three series of GyrB/ParE inhibitors were designed, synthesized, and evaluated in enzymatic and antibacterial assays. Chemical space was thoroughly explored; compounds with acidic, basic, and neutral properties were synthesized, possessing very potent inhibitory activity on *E. coli* and/or *S. aureus* gyrase and/or topo IV. The best compounds were active on Gram-positive bacterium *E. faecalis* with the best compound having MIC = 3.13 μ M. The compounds were inactive on Gram-negative bacteria because they are good substrates for bacterial efflux pumps, but **15a** and **15e** showed potent antibacterial activity on the efflux impaired *E. coli* strain (MIC = 0.78 μ M). Compound **15a** with low molecular weight 383 Da displayed potent inhibitory activity on *E. coli* gyrase (IC₅₀ = 9.5 nM) and potent antibacterial activity on *E. faecalis* (MIC = 3.13 μ M) and on the efflux impaired *E. coli* strain (MIC = 0.78 μ M) and thus makes an important contribution for the development of novel gyrase and topoisomerase inhibitors in Gram-negative bacteria.

■ ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsmchemlett.0c00416>.

- Experimental conditions for biochemical and microbiological assays,
- experimental conditions for cocrystallization, collection, and refinement statistics for compound **16a**,
- experimental procedures for the synthesis of all the compounds described in this manuscript, and
- representative analytical data (PDF)

■ AUTHOR INFORMATION

Corresponding Author

Janez Ilaš – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia; orcid.org/0000-0002-0124-0474; Phone: +386-1-4769576; Email: janez.ilas@ffa.uni-lj.si

Authors

Žiga Skok – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia

Michaela Barančoková – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia

Ondřej Benek – University of Hradec Králové, Faculty of Science, Department of Chemistry, 500 03 Hradec Králové, Czech Republic; orcid.org/0000-0001-6840-1933

Cristina Durante Cruz – Centre for Drug Research, Division of Pharmaceutical Biosciences, Faculty of Pharmacy, University of Helsinki, Helsinki FI-00014, Finland

Päivi Tammela – Centre for Drug Research, Division of Pharmaceutical Biosciences, Faculty of Pharmacy, University of Helsinki, Helsinki FI-00014, Finland; orcid.org/0000-0003-4697-8066

Tihomir Tomašič – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia; orcid.org/0000-0001-5534-209X

Nace Zidar – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia; orcid.org/0000-0003-1905-0158

Lucija Peterlin Mašič – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia

Anamarija Zega – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia; orcid.org/0000-0003-4065-0019

Clare E. M. Stevenson – Department of Biological Chemistry, John Innes Centre, Norwich NR4 7UH, U.K.

Julia E. A. Mundy – Department of Biological Chemistry, John Innes Centre, Norwich NR4 7UH, U.K.

David M. Lawson – Department of Biological Chemistry, John Innes Centre, Norwich NR4 7UH, U.K.

Anthony Maxwell – Department of Biological Chemistry, John Innes Centre, Norwich NR4 7UH, U.K.; orcid.org/0000-0002-5756-6430

Danijel Kikelj – University of Ljubljana, Faculty of Pharmacy, 1000 Ljubljana, Slovenia

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acsmchemlett.0c00416>

Author Contributions

[†]Ž.S. and M.B. contributed equally. The manuscript was written with contributions from all of the authors.

Funding

The work was funded by the EU H2020 ITN-ETN Project INTEGRATE (Project Reference: 642620), EU FP7 Integrated Project MAREX (Project No. FP7-KBBE-2009-3-245137), Academy of Finland (Grant No. 277001), and Slovenian Research Agency (Grant No. P1-0208). Work in A.M.'s lab was supported by the Biotechnology and Biosciences Research Council (UK) Institute Strategic Programme Grant BB/P012523/1, and the Wellcome Trust (Investigator Award 110072/Z/15/Z).

Notes

The authors declare no competing financial interest.

■ ACKNOWLEDGMENTS

We acknowledge Diamond Light Source for access to beamline I04 under proposal MX18565 and Sara Henderson for providing the protein sample.

■ ABBREVIATIONS

MIC, minimal inhibitory concentration; RA, residual activity

REFERENCES

- (1) Aminov, R. I. A brief history of the antibiotic era: lessons learned and challenges for the future. *Front. Microbiol.* **2010**, *1*, 134.
- (2) Nathan, C.; Cars, O. Antibiotic resistance—problems, progress, and prospects. *N. Engl. J. Med.* **2014**, *371*, 1761–1763.
- (3) Payne, D. J.; Gwynn, M. N.; Holmes, D. J.; Pompliano, D. L. Drugs for bad bugs: confronting the challenges of antibacterial discovery. *Nat. Rev. Drug Discovery* **2007**, *6*, 29–40.
- (4) Silver, L. L. Challenges of antibacterial discovery. *Clin Microbiol Rev.* **2011**, *24*, 71–109.
- (5) Brown, E. D.; Wright, G. D. Antibacterial drug discovery in the resistance era. *Nature* **2016**, *529*, 336–343.
- (6) Emmerson, A. M.; Jones, A. M. The quinolones: decades of development and use. *J. Antimicrob. Chemother.* **2003**, *51* (Suppl1), 13–20.
- (7) Bisacchi, G. S.; Manchester, J. I. A New-Class Antibacterial—Almost. Lessons in Drug Discovery and Development: A Critical Analysis of More than 50 Years of Effort toward ATPase Inhibitors of DNA Gyrase and Topoisomerase IV. *ACS Infect. Dis.* **2015**, *1*, 4–41.
- (8) Barančokova, M.; Kikelj, D.; Ilaš, J. Recent progress in the discovery and development of DNA gyrase B inhibitors. *Future Med. Chem.* **2018**, *10*, 1207–1227.
- (9) Bates, A.D.; Maxwell, A. *DNA Topology*; Oxford University Press, Oxford, 2005.
- (10) Chen, S. H.; Chan, N. L.; Hsieh, T. S. New mechanistic and functional insights into DNA topoisomerases. *Annu. Rev. Biochem.* **2013**, *82*, 139–170.
- (11) Vos, S. M.; Tretter, E. M.; Schmidt, B. H.; Berger, J. M. All tangled up: how cells direct, manage and exploit topoisomerase function. *Nat. Rev. Mol. Cell Biol.* **2011**, *12*, 827–841.
- (12) Bush, N.G.; Evans-Roberts, K.; Maxwell, A. DNA Topoisomerases. *EcoSal Plus* **2015**, *6*, DOI: 10.1128/ecosalplus.ESP-0010-2014.
- (13) Schoeffler, A. J.; Berger, J. M. DNA topoisomerases: harnessing and constraining energy to govern chromosome topology. *Q. Rev. Biophys.* **2008**, *41*, 41–101.
- (14) Collin, F.; Karkare, S.; Maxwell, A. Exploiting bacterial DNA gyrase as a drug target: current state and perspectives. *Appl. Microbiol. Biotechnol.* **2011**, *92*, 479–497.
- (15) Gjorgjieva, M.; Tomašič, T.; Barančokova, M.; Katsamakos, S.; Ilaš, J.; Tammela, P.; Mašič, L. P.; Kikelj, D. Discovery of Benzothiazole Scaffold-Based DNA Gyrase B Inhibitors. *J. Med. Chem.* **2016**, *59*, 8941–8954.
- (16) Zidar, N.; Macut, H.; Tomašič, T.; Brvar, M.; Montalvão, S.; Tammela, P.; Šolmajer, T.; Mašič, L. P.; Ilaš, J.; Kikelj, D. N-Phenyl-4,5-dibromopyrrolamides and N-Phenylindolamides as ATP Competitive DNA Gyrase B Inhibitors: Design, Synthesis, and Evaluation. *J. Med. Chem.* **2015**, *58*, 6179–6194.
- (17) Zidar, N.; Tomašič, T.; Macut, H.; Sirc, A.; Brvar, M.; Montalvão, S.; Tammela, P.; Ilaš, J.; Kikelj, D. New N-phenyl-4,5-dibromopyrrolamides and N-Phenylindolamides as ATPase inhibitors of DNA gyrase. *Eur. J. Med. Chem.* **2016**, *117*, 197–211.
- (18) Zidar, N.; Montalvão, S.; Hodnik, Ž.; Nawrot, D. A.; Žula, A.; Ilaš, J.; Kikelj, D.; Tammela, P.; Peterlin Mašič, L. Antimicrobial Activity of the Marine Alkaloids, Clathrocin and Oroidin, and Their Synthetic Analogues. *Mar. Drugs* **2014**, *12*, 940–963.
- (19) Lamut, A.; Skok, Ž.; Barančoková, M.; Gutierrez, L. J.; Cruz, C. D.; Tammela, P.; Draskovits, G.; Szili, P. É.; Nyerges, A.; Pál, C.; Molek, P.; Bratkovič, T.; Ilaš, J.; Zidar, N.; Zega, A.; Enriz, R. D.; Kikelj, D.; Tomašič, T. Second-generation 4,5,6,7-tetrahydrobenzo-[d]thiazoles as novel DNA gyrase inhibitors. *Future Med. Chem.* **2020**, *12* (4), 277–297.
- (20) Phillips, J. W.; Goetz, M. A.; Smith, S. K.; Zink, D. L.; Polishook, J.; Onishi, R.; Salowe, S.; Wiltsie, J.; Allocco, J.; Sigmund, J.; Dorso, K.; Lee, S.; Skwish, S.; de la Cruz, M.; Martin, J.; Vicente, F.; Genilloud, O.; Lu, J.; Painter, R. E.; Young, K.; Overbye, K.; Donald, R. G.; Singh, S. B. Discovery of kibelomycin, a potent new class of bacterial type II topoisomerase inhibitor by chemical-genetic profiling in *Staphylococcus aureus*. *Chem. Biol.* **2011**, *18*, 955–965.
- (21) Singh, S. B.; Goetz, M. A.; Smith, S. K.; Zink, D. L.; Polishook, J.; Onishi, R.; Salowe, S.; Wiltsie, J.; Allocco, J.; Sigmund, J.; Dorso, K.; de la Cruz, M.; Martin, J.; Vicente, F.; Genilloud, O.; Donald, R. G.; Phillips, J. W. Kibelomycin A, a congener of kibelomycin, derivatives and their antibacterial activities. *Bioorg. Med. Chem. Lett.* **2012**, *22*, 7127–7130.
- (22) Lu, J.; Patel, S.; Sharma, N.; Soisson, S. M.; Kishii, R.; Takei, M.; Fukuda, Y.; Lumb, K. J.; Singh, S. B. Structures of kibelomycin bound to *Staphylococcus aureus* GyrB and ParE showed a novel U-shaped binding mode. *ACS Chem. Biol.* **2014**, *9*, 2023–2031.
- (23) Tohyama, S.; Takahashi, Y.; Akamatsu, Y. Biosynthesis of amycolamicin: the biosynthetic origin of a branched alpha-aminoethyl moiety in the unusual sugar amycolose. *J. Antibiot.* **2010**, *63*, 147–149.
- (24) Sawa, R.; Takahashi, Y.; Hashizume, H.; Sasaki, K.; Ishizaki, Y.; Umekita, M.; Hatano, M.; Abe, H.; Watanabe, T.; Kinoshita, N.; Homma, Y.; Hayashi, C.; Inoue, K.; Ohba, S.; Masuda, T.; Arakawa, M.; Kobayashi, Y.; Hamada, M.; Igarashi, M.; Adachi, H.; Nishimura, Y.; Akamatsu, Y. Amycolamicin: a novel broad-spectrum antibiotic inhibiting bacterial topoisomerase. *Chem. - Eur. J.* **2012**, *18*, 15772–15781.
- (25) Sherer, B. A.; Hull, K.; Green, O.; Basarab, G.; Hauck, S.; Hill, P.; Loch, J. T.; Mullen, G.; Bist, S.; Bryant, J.; Boriack-Sjodin, A.; Read, J.; DeGrace, N.; Uria-Nickelsen, M.; Illingworth, R. N.; Eakin, A. E. Pyrrolamide DNA gyrase inhibitors: Optimization of antibacterial activity and efficacy. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 7416–7420.
- (26) Eakin, A. E.; Green, O.; Hales, N.; Walkup, G. K.; Bist, S.; Singh, A.; Mullen, G.; Bryant, J.; Embrey, K.; Gao, N.; Breeze, A.; Timms, D.; Andrews, B.; Uria-Nickelsen, M.; Demeritt, J.; Loch, J. T., 3rd; Hull, K.; Blodgett, A.; Illingworth, R. N.; Prince, B.; Boriack-Sjodin, P. A.; Hauck, S.; MacPherson, L. J.; Ni, H.; Sherer, B. Pyrrolamide DNA gyrase inhibitors: fragment-based nuclear magnetic resonance screening to identify antibacterial agents. *Antimicrob. Agents Chemother.* **2012**, *56*, 1240–1246.
- (27) Uria-Nickelsen, M.; Blodgett, A.; Kamp, H.; Eakin, A.; Sherer, B.; Green, O. Novel DNA gyrase inhibitors. *Int. J. Antimicrob. Agents* **2013**, *41*, 28–35.
- (28) Cascioferro, S.; Parrino, B.; Carbone, D.; Schillaci, D.; Giovannetti, E.; Cirrincione, G.; Diana, P. Thiazoles, Their Benzofused Systems, and Thiazolidinone Derivatives: Versatile and Promising Tools to Combat Antibiotic Resistance. *J. Med. Chem.* **2020**, *63*, 7923–7956.
- (29) Charifson, P. S.; Grillot, A.-L.; Grossman, T. H.; Parsons, J. D.; Badia, M.; Bellon, S.; Deininger, D. D.; Drumm, J. E.; Gross, C. H.; LeTiran, A.; Liao, Y.; Mani, N.; Nicolau, D. P.; Perola, E.; Ronkin, S.; Shannon, D.; Swenson, L. L.; Tang, Q.; Tessier, P. R.; Tian, S.-K.; Trudeau, M.; Wang, T.; Wei, Y.; Zhang, H.; Stamos, D. Novel Dual-Targeting Benzimidazole Urea Inhibitors of DNA Gyrase and Topoisomerase IV Possessing Potent Antibacterial Activity: Intelligent Design and Evolution through the Judicious Use of Structure-Guided Design and Structure–Activity Relationships. *J. Med. Chem.* **2008**, *51*, 5243–5263.
- (30) Tari, L. W.; Li, X.; Trzoss, M.; Bensen, D. C.; Chen, Z.; Lam, T.; Zhang, J.; Lee, S. J.; Hough, G.; Phillipson, D.; Akers-Rodriguez, S.; Cunningham, M. L.; Kwan, B. P.; Nelson, K. J.; Castellano, A.; Locke, J. B.; Brown-Driver, V.; Murphy, T. M.; Ong, V. S.; Pillar, C. M.; Finn, J. Tricyclic GyrB/ParE (TriBE) Inhibitors: A New Class of Broad-Spectrum Dual-Targeting Antibacterial Agents. *PLoS One* **2013**, *8* (12), e84409.
- (31) Cramer, J.; Krimmer, S. G.; Heine, A.; Klebe, G. Paying the Price of Desolvation in Solvent-Exposed Protein Pockets: Impact of Distal Solubilizing Groups on Affinity and Binding Thermodynamics in a Series of Thermolysin Inhibitors. *J. Med. Chem.* **2017**, *60*, 5791–5799.
- (32) Lu, J.; Patel, S.; Sharma, N.; Soisson, S. M.; Kishii, R.; Takei, M.; Fukuda, Y.; Lumb, K. J.; Singh, S. B. Structures of Kibelomycin Bound to *Staphylococcus aureus* GyrB and ParE Showed a Novel U-Shaped Binding Mode. *ACS Chem. Biol.* **2014**, *9*, 2023.

(33) Sherer, B. A.; Hull, K.; Green, O.; Basarab, G.; Hauck, S.; Hill, P.; Loch, J. T.; Mullen, G.; Bist, S.; Bryant, J.; Boriack-Sjodin, A.; Read, J.; Degrace, N.; Uria-Nickelsen, M.; Illingworth, R. N.; Eakin, A. E. Pyrrolamide DNA gyrase inhibitors: Optimization of antibacterial activity and efficacy. *Bioorg. Med. Chem. Lett.* **2011**, *21*, 7416–7420.

(34) Richter, M. F.; Drown, B. S.; Riley, A. P.; Garcia, A.; Shirai, T.; Svec, R. L.; Hergenrother, P. J. Predictive rules for compound accumulation yield a broad-spectrum antibiotic. *Nature* **2017**, *545* (7654), 299–304.

(35) Vergalli, J.; Bodrenko, I. V.; Masi, M.; Moynié, L.; Acosta-Gutiérrez, S.; Naismith, J. H.; Davin-Regli, A.; Ceccarelli, M.; van den Berg, B.; Winterhalter, M.; Pagès, J. M. Porins and small-molecule translocation across the outer membrane of Gram-negative bacteria. *Nat. Rev. Microbiol.* **2020**, *18* (3), 164–176.

(36) Winterhalter, M.; Ceccarelli, M. Physical methods to quantify small antibiotic molecules uptake into Gram-negative bacteria. *Eur. J. Pharm. Biopharm.* **2015**, *95*, 63–7.

(37) Vergalli, J.; Dumont, E.; Pajović, J.; Cinquin, B.; Maigre, L.; Masi, M.; Réfrégiers, M.; Pagès, J. M. Spectrofluorimetric quantification of antibiotic drug concentration in bacterial cells for the characterization of translocation across bacterial membranes. *Nat. Protoc.* **2018**, *13* (6), 1348–1361.