

Unusual allergen in a butcher with respiratory symptoms

Ingrid Sander¹, Claus Keller², Christina Czibor¹, Ursula Meurer¹, Rolf Merget¹ and Monika Raulf¹

©2020 Dustri-Verlag Dr. K. Feistle
ISSN 2512-8957

DOI 10.5414/ALX02126E
e-pub: December 2, 2020

¹Institute for Prevention and Occupational Medicine of the German Social Accident Insurance, Institute of the Ruhr University Bochum (IPA), Bochum, and ²Medical Practice for Pulmonary and Bronchial Medicine, Allergology and Environmental Medicine, Frankfurt am Main, Germany

Key words

allergy – respiratory symptoms – butcher – transglutaminase – *Streptomyces mobaraensis*

German version

published in
Allergologie,
Vol. 43, No. 1/2020,
pp. 15-19

Abstract. A 37-year-old butcher developed respiratory symptoms during sausage and chicken production in a large company. In addition to various spices, the enzyme transglutaminase was a possible cause. The lung function test showed mild partial reversible airway obstruction and severe bronchial hyperresponsiveness. The IgE test showed sensitizations to various spice mixtures, coriander (0.74 kU/L), and to the ImmunoCAP-bound transglutaminase preparation from the workplace (7.12 kU/L). The skin prick tests with this transglutaminase were also positive. In the immunoblot of this preparation, a 40-kD protein reacted with the patient's IgE and was identified as transglutaminase from *Streptomyces mobaraensis* by inhibition experiments. This is the first case of a butcher with an allergy to transglutaminase. After moving to a small enterprise without enzyme use, his symptoms improved. Sensitization and the course of the symptoms indicate a dominant role of transglutaminase in the patient's allergic asthma.

Introduction

While the risk of developing occupational allergic asthma is considered very high for bakers, this risk is considerably lower for butchers [2, 9]. In the scientific literature, there are only case reports of butchers with occupational asthma [3, 4, 5, 7, 14] or contact dermatitis [1, 16] caused by meat proteins, various vegetable spices, the red dye carmine, or the enzyme papain, which is used as a meat tenderizer.

With this knowledge, a butcher who had developed respiratory problems while working in a large meat processing company was medically examined in Frankfurt in 2017. He

had irritable cough and sneezing attacks in the morning during sausage production and the addition of spice mixtures with paprika and pepper. These symptoms became worse in the afternoon during chicken production, and he also suffered from shortness of breath and wheals on his skin. A possible trigger was the addition of very dusty binding agents which also contained the enzyme transglutaminase. The enzyme transglutaminase from *Streptomyces mobaraensis* catalyzes the cross-linking of proteins and thus leads to a more stable texture or joins pieces of meat. It has been increasingly used in the production of meat, fish, and dairy products since mid-2011.

Case report

The 37-year-old butcher suffered from pollen allergy and bronchial asthma since his youth. He had been a smoker for 20 years with 10 cigarettes a day. For 8 years he has additionally suffered from season-independent symptoms during his work in the meat factory. These could not be prevented even when wearing a breathing mask, possibly due to insufficient tightness. However, he was only given the breathing mask after the symptoms began, and he did not wear it permanently. He relieved his severe respiratory symptoms in the afternoon during the production of chicken nuggets by double dosing his asthma spray containing beclomethasone dipropionate and formeterol fumarate (200 µg/6 µg per stroke).

In the preliminary findings, there were indications of slightly elevated IgE antibodies

Received
August 8, 2018;
accepted in revised form
November 25, 2020

Correspondence to
Dr. Ingrid Sander
Institute for Prevention
and Occupational
Medicine of the German
Social Accident
Insurance, Institute of
the Ruhr University
Bochum, Center of
Allergology/Immunology,
Bürkle-de-la-Camp-Platz
1, 44789 Bochum,
Germany
sander@ipa-dguv.de




Figure 1. Silver-stained SDS-PAGE of transglutaminase preparations. M = molecular weight marker proteins, lane 1: transglutaminase from the workplace (May 2017), lane 2: transglutaminase from the workplace (June 2017), lane 3: Activa TI transglutaminase.

to a spice mixture with laurel, paprika, green pepper, and mustard (test system unknown). The skin prick tests revealed sensitization to pollen from birch, alder, hazel, mugwort, and house dust mites. However, the total IgE value was within the normal range. In the documented lung function tests, a moderate obstruction (FEV_1 69% of the target value) was found, which was either not reversible, or could only be reduced by a maximum of 15% with β -mimetics.

Laboratory Analysis

Two samples of the transglutaminase preparation from the workplace (manufacturer and composition unknown), and a

transglutaminase from *Streptomyces mobaraensis* (Activa TI Ajinomoto, Modernist Pantry, USA) were analyzed. According to Modernist Pantry, Activa TI transglutaminase is an enzyme preparation without added helper protein for stabilization.

All samples were first analyzed by SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and silver staining (Figure 1), with 0.5 μ g protein per lane applied [10]. Both workplace samples had identical protein patterns with bands at 67, 40, 33, 16, 15, 13, 12, and 10 kDa (estimated average 22 kDa). The Activa transglutaminase showed silver-stained bands at 67, 40, and 12 kDa (estimated average 30 kDa). With a 5-fold molar excess of NHS-biotin, one transglutaminase from the workplace and the Activa transglutaminase were biotinylated and bound to streptavidin ImmunoCAPs [12]. Specific IgE binding was quantified using CAP-FEIA (Thermo Fisher Scientific, Phadia AB, Sweden).

In addition, both transglutaminases (15 μ g protein each) were transferred to PVDF membranes after SDS-PAGE, which were then cut into strips [10]. One strip each was stained with India Ink (1 h) [6]. The remaining strips were used for IgE immunoblots [10] with the patient serum and a control serum and IgE inhibition experiments. The control serum had a similar total IgE value and a comparable s_{x1} value (basic test inhalation allergens) compared to the patient's serum. Each strip was incubated with 400 μ L patient serum (or control serum), 80 μ L 10% BSA (bovine serum albumin), and 320 μ L phosphate salt buffer overnight at 4 °C; the inhibitor strips additionally contained 0.7% or 2.8% (w/v) of the transglutaminase preparations.

Results

The patient's lung function test in the body plethysmograph revealed a slight obstruction (FEV_1 at 74% of target) and a highly hyperreactive bronchial system.

The skin prick test with transglutaminase from the workplace was positive (wheal as large as in the positive control with histamine). In vitro sensitization was shown in the ImmunoCAP IgE test for coriander (f317, 0.74 kU/l) as well as for the workplace




Figure 2. IgE immunoblot of transglutaminase from the workplace including inhibitions. M = molecular weight marker proteins, I = India-Ink staining of transglutaminase from the workplace (May 2017) on the blot membrane, lane 1: IgE binding of patient's serum, lane 2: reaction with control serum, lane 3: reaction of patient's serum inhibited with 2.8% Activa transglutaminase, lane 4: reaction of patient's serum inhibited with 0.7% Activa transglutaminase, lane 5: reaction of patient's serum inhibited with 2.8% transglutaminase from the workplace, track 6: Reaction patient serum inhibited with 0.7% transglutaminase from the workplace, lane 7: control with anti-IgE detection antibody.

transglutaminase bound to ImmunoCAPs (7.12 kU/l) and Activa transglutaminase from *Streptomyces mobaraensis* (7.48 kU/l). The spices mace (f266), nutmeg (f282), cardamom (f267), and bell pepper (f218) were negative in the ImmunoCAP IgE test. The total IgE value was 55.9 kU/l.

In the immunoblot, a 40-kD protein was identified as IgE-binding, which was present in the preparation from the workplace as well as in the Activa enzyme preparation. Figure 2 shows the immunoblot with the workplace transglutaminase; the immunoblot with the Activa transglutaminase looks almost identical (not shown). The IgE binding of the

patient serum to the 40-kDa protein could be completely inhibited with the workplace transglutaminase as well as with the Activa transglutaminase already at a concentration of 0.7%. The band at 40 kDa, which is still weakly recognizable after inhibition, has an intensity similar to that of the strips with the control serum or with the anti-IgE detection antibody.

The patient reacted positively in both the nasal and bronchial provocation tests after workplace simulation with paprika powder. Provocation with transglutaminase was not performed for safety reasons, due to the reactivity of this enzyme.

Discussion

With pre-existing pollen allergy, the butcher presented in this case report developed respiratory problems at work throughout the year. His symptoms were particularly severe in the afternoon while making chicken nuggets. In this process, the enzyme transglutaminase acts as a binder, linking pieces of meat by cross-linking proteins. Both the skin prick test and the specific IgE test showed a clear sensitization to the enzyme product used at the workplace. However, the composition of the enzyme product was unknown. In principle, sensitization to components added for stabilization, such as wheat or milk proteins, would also be possible. Therefore, an enzyme preparation with a known composition was additionally tested. This contained transglutaminase from the bacterium *Streptomyces mobaraensis* and no helper proteins. The sIgE test with this product was also clearly positive (slightly higher than with transglutaminase from the workplace). In the analysis by SDS-PAGE, the two transglutaminases differed only by some additional bands in the product used at the workplace. The IgE immunoblot then identified a 40-kDa protein in both products as the allergen. This fits well with the theoretical molecular weight of 44 kDa of the transglutaminase from *Streptomyces mobaraensis* [15]. The identity of the 40-kDa proteins of both products is supported by the complete inhibition of IgE binding of the patient's serum to this protein band.

The enzyme transglutaminase from *Streptomyces mobaraensis* is used in the production of meat, fish, and dairy products. Two previous cases have been described in the literature: a case of respiratory allergy to microbial transglutaminase in a person working in company where ingredients for the food industry were mixed together [11] and a case of the chef of a restaurant offering "molecular cuisine" who had developed urticaria, rhinoconjunctivitis, and asthma when using transglutaminase powder [8]. This is the first case of a butcher with an allergy to transglutaminase. After moving to a small enterprise where no enzymes are used, his symptoms improved. Sensitization and the course of the symptoms indicate a dominant role of transglutaminase in the patient's aller-

gic asthma. Transglutaminase from *Streptomyces mobaraensis* is therefore added to the list of industrially produced enzymes that are triggers of occupational respiratory allergies [13].

Funding

The study was financed by the German Social Accident Insurance (project IPA-14).

Conflict of interest

The authors state that there is no conflict of interest.

References

- [1] Anliker MD, Borelli S, Wüthrich B. Occupational protein contact dermatitis from spices in a butcher: a new presentation of the mugwort-spice syndrome. *Contact Dermat.* 2002; 46: 72-74. [CrossRef PubMed](#)
- [2] Coppeters Y, Piette D. Targeting pupils at risk of occupational asthma. *Patient Educ Couns.* 2004; 55: 136-141. [CrossRef PubMed](#)
- [3] Ferrer A, Carnés J, Marco FM, Andréu C, Fernández-Caldas E. Occupational allergic rhinoconjunctivitis and asthma to goat and cross-reactivity with cow epithelium. *Ann Allergy Asthma Immunol.* 2006; 96: 579-585. [CrossRef PubMed](#)
- [4] Ferrer A, Marco FM, Andreu C, Sempere JM. Occupational asthma to carmine in a butcher. *Int Arch Allergy Immunol.* 2005; 138: 243-250. [CrossRef PubMed](#)
- [5] Fraj J, Lezaun A, Colás C, Duce F, Domínguez MA, Alonso MD. Occupational asthma induced by aniseed. *Allergy.* 1996; 51: 337-339. [CrossRef PubMed](#)
- [6] Hancock K, Tsang VC. India ink staining of proteins on nitrocellulose paper. *Anal Biochem.* 1983; 133: 157-162. [CrossRef PubMed](#)
- [7] Jiang N, Yin J, Wen L. Papain Induced Occupational Asthma with Kiwi and Fig Allergy. *Allergy Asthma Immunol Res.* 2016; 8: 170-173. [CrossRef PubMed](#)
- [8] Johnsen CR, Callesen KT, Jensen BM, Poulsen LK. Occupational allergy to transglutaminase in a chef – a case story. <http://webcast.eaaci.cyim.com/mediatheque/media.aspx?mediaId=47931&channel=8518>.
- [9] Karjalainen A, Kurppa K, Virtanen S, Keskinen H, Nordman H. Incidence of occupational asthma by occupation and industry in Finland. *Am J Ind Med.* 2000; 37: 451-458. [CrossRef PubMed](#)
- [10] Kespohl S, Maryska S, Zahradnik E, Sander I, Brüning T, Raulf-Heimsoth M. Biochemical and immunological analysis of mould skin prick test solution: current status of standardization. *Clin*

- Exp Allergy. 2013; 43: 1286-1296. [CrossRef](#)
[PubMed](#)
- [11] De Palma G, Apostoli P, Mistrello G, Zanotta S, Bertorelli G. Microbial transglutaminase: a new and emerging occupational allergen. Ann Allergy Asthma Immunol. 2014; 112: 553-554. [CrossRef](#)
[PubMed](#)
- [12] Sander I, Kespohl S, Merget R, Goldscheid N, Degens PO, Brüning T, Raulf-Heimsoth M. A new method to bind allergens for the measurement of specific IgE antibodies. Int Arch Allergy Immunol. 2005; 136: 39-44. [CrossRef](#) [PubMed](#)
- [13] Sander I, Raulf M. Industriell hergestellte Enzyme als Verursacher von beruflichen Atemweg-sallergien – eine Übersicht. Allergologie. 2018; 41: 511-528. [CrossRef](#)
- [14] Sastre J, Olmo M, Novalvos A, Ibañez D, Lahoz C. Occupational asthma due to different spices. Allergy. 1996; 51: 117-120. [CrossRef](#) [PubMed](#)
- [15] UniProt Consortium. UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res. 2019; 47 (D1): D506-D515. [CrossRef](#) [PubMed](#)
- [16] Vanstreels L, Merk HF. [Protein contact dermatitis in a butcher]. Hautarzt. 2012; 63: 926-928. [Cross-Ref](#) [PubMed](#)