



An alternative miRISC targets a cancer-associated coding sequence mutation in FOXL2

Eunkyong Shin^{1,†} , Hanyong Jin^{2,†} , Dae-Shik Suh^{3,†}, Yongyang Luo¹ , Hye-Jeong Ha¹,
Tae Heon Kim⁴, Yoonsoo Hahn² , Seogang Hyun^{2,*} , Kangseok Lee^{2,**} & Jeehyeon Bae^{1,***}

Abstract

Recent evidence suggests that animal microRNAs (miRNAs) can target coding sequences (CDSs); however, the pathophysiological importance of such targeting remains unknown. Here, we show that a somatic heterozygous missense mutation (c.402C>G; p.C134W) in *FOXL2*, a feature shared by virtually all adult-type granulosa cell tumors (AGCTs), introduces a target site for miR-1236, which causes haploinsufficiency of the tumor-suppressor *FOXL2*. This miR-1236-mediated selective degradation of the variant *FOXL2* mRNA is preferentially conducted by a distinct miRNA-loaded RNA-induced silencing complex (miRISC) directed by the Argonaute3 (AGO3) and DHX9 proteins. In both patients and a mouse model of AGCT, abundance of the inversely regulated variant *FOXL2* with miR-1236 levels is highly correlated with malignant features of AGCT. Our study provides a molecular basis for understanding the conserved *FOXL2* CDS mutation-mediated etiology of AGCT, revealing the existence of a previously unidentified mechanism of miRNA-targeting disease-associated mutations in the CDS by forming a non-canonical miRISC.

Keywords allelic imbalance; Argonaute3; DHX9; metastasis; miR-1236

Subject Category RNA Biology

DOI 10.15252/emboj.2020104719 | Received 16 February 2020 | Revised 28

August 2020 | Accepted 6 September 2020 | Published online 20 November 2020

The EMBO Journal (2020) 39: e104719

Introduction

MicroRNAs (miRNAs) are endogenous, noncoding RNAs of ~22 nucleotides (nt) in length that suppress the stability or translational efficiency of mRNAs. Conventionally, miRNAs are known to target sequences in the 3'-untranslated regions (UTRs) of mRNAs. However, recent multiple high-throughput sequencing and proteomic analyses suggest that miRNAs can also bind sites within

mRNA coding sequences (CDSs; Chi *et al.*, 2009; Hafner *et al.*, 2010; Schnall-Levin *et al.*, 2010; Hausser *et al.*, 2013; Xue *et al.*, 2013; Broughton *et al.*, 2016), and the intracellular effects of miRNAs targeting CDS sites have been proposed (Forman *et al.*, 2008; Tay *et al.*, 2008; Elcheva *et al.*, 2009; Schnall-Levin *et al.*, 2011; Zhang *et al.*, 2018). However, the physiological relevance and the pathological consequences of miRNA binding to CDSs remain unclear.

Argonaute (AGO) clade proteins are essential components of miRNA-loaded RNA-induced silencing complexes (miRISCs) that select target mRNAs by directly interacting with mature miRNAs (Hock & Meister, 2008; Czech & Hannon, 2011). In mammals, four AGO paralogs (AGO1–4) are involved in miRNA pathways (Hock & Meister, 2008; Czech & Hannon, 2011), and they share ~80% amino acid-sequence identity (Sasaki *et al.*, 2003). AGO2 has been described as a specialized AGO that possesses slicer activity, enabling cleavage of target mRNAs by miRNAs and small-interfering RNAs (siRNAs; Liu *et al.*, 2004; Meister *et al.*, 2004). However, previous data suggested that all mammalian AGOs may serve overlapping and distinct functions in miRNA-mediated regulation (Su *et al.*, 2009). High-throughput pyrosequencing data showed that the profiles of miRNAs associated with AGO2 and AGO3 largely overlap, but preferential associations with AGO2 or AGO3 also occur for a small set of miRNAs (Azuma-Mukai *et al.*, 2008). However, the functional significances of mammalian AGO1, AGO3, and AGO4 in miRNA activity are poorly understood.

Conclusive evidence demonstrating clear pathophysiological consequences elicited by miRNA binding to the CDSs of disease-associated gene loci is lacking. Here, we investigated whether miRNA binding to the CDS of *FOXL2* contributes to adult-type granulosa cell tumor (AGCT) development. GCTs are malignant ovarian cancers comprising AGCTs and juvenile GCTs (JGCTs) (Schumer & Cannistra, 2003). *FOXL2* is evolutionarily conserved and encodes a forkhead-domain transcription factor essential for the ovary development and function (Crisponi *et al.*, 2001; Schmidt *et al.*, 2004; Uhlenhaut *et al.*, 2009). A highly prevalent heterozygous somatic missense mutation (c.402C>G; p.C134W) in *FOXL2* is exclusively

1 School of Pharmacy, Chung-Ang University, Seoul, Korea

2 Department of Life Science, Chung-Ang University, Seoul, Korea

3 Department of Obstetrics and Gynecology, Asan Medical Center, University of Ulsan College of Medicine, Seoul, Korea

4 Department of Pathology, Bundang CHA Hospital, CHA University, Seongnam, Korea

*Corresponding author. Tel: +82 2 8205805; E-mail: sghyun@cau.ac.kr

**Corresponding author. Tel: +82 2 8205241; E-mail: kangseok@cau.ac.kr

***Corresponding author. Tel: +82 2 8205604; E-mail: jeehyeon@cau.ac.kr

†These authors contributed equally to this work

found in > 97% of patients with AGCT and is considered the main cause of AGCT (Shah *et al*, 2009). However, the etiological nature of the 402C>G mutation remains largely unknown. Previously, we showed that the FOXL2 protein acts as a tumor suppressor and directs DNA double-strand repair pathways in granulosa cells, whereas C134W FOXL2 was defective due to accelerated MDM2-mediated ubiquitination and proteasomal degradation (Kim *et al*, 2011; Kim *et al*, 2014; Jin *et al*, 2020). However, a relatively moderate change in FOXL2 protein stability by the C134W mutation does not appear to wholly account for haploinsufficiency of the mutant FOXL2 (Kim *et al*, 2011; Kim *et al*, 2014).

Here, we identified allelic imbalance in FOXL2 mRNAs in patients with AGCT arising from recognition of the 402C>G locus as a target site of miR-1236 that drives degradation of this variant FOXL2 mRNA, which explains the etiology of this conserved mutation in AGCTs.

Results

Allelic imbalance of FOXL2 transcripts in AGCT samples

To study allelic imbalance of heterozygous FOXL2 mRNAs, we analyzed the relative levels of wild-type (WT) and variant FOXL2 (402C>G) mRNAs from complementary DNA (cDNA) samples from the individual AGCT tissues by high-throughput ultra-deep RNA sequencing. Ultra-deep RNA sequencing analysis of AGCT tissues showed that decreased proportion of variant FOXL2 mRNA level compared with WT FOXL2 mRNA in 20 AGCT patients with an average ratio of 62:38 for WT to 402C>G ($P = 0.004$), with a variable relative abundance, was observed among samples (Appendix Fig S1A). A previous study, which identified this conserved mutation, reported relative abundance of WT versus variant FOXL2 mRNA levels in four AGCT patients, where no uniformed trend was observed (Shah *et al*, 2009). For these reasons, we recruited additional AGCT patients from two independent hospitals and performed high-throughput pyrosequencing analysis ($n = 46$) that enables amplification and detection of both alleles from the same pyrosequencing reaction using common primers designed to bind FOXL2 cDNA. As shown in Fig 1A and Appendix Fig S1B, the relative abundance of FOXL2 mRNA analyzed by pyrosequencing was 72:28 for WT:402C>G in 46 AGCTs including 20 corresponding AGCTs analyzed for RNA sequencing presented in Appendix Fig S1A. In addition, allele-specific real-time and semi-quantitative RT-PCR analyses of 46 AGCTs were performed using primers presented in Appendix Fig S1C, and we observed consistent lower steady-state levels of FOXL2 variant mRNA compared with WT FOXL2 mRNA (Appendix Fig S1D and E). For these analyses, we used paired genomic DNA (gDNA) levels of both alleles for the normalization of data, where the gDNA levels of both alleles were similar in all AGCTs (Appendix Fig S1F). These results indicate that contamination of non-cancerous stromal cells in preparation of total RNA from AGCT tissues for these analyses was minimal.

We obtained analogous results using AGCT-derived KGN cells by pyrosequencing, allele-specific semi-quantitative RT-PCR, and allele-specific real-time RT-PCR, which are heterozygous for the 402C>G mutation (Fig 1B). The relative abundance of variant

FOXL2 mRNA was ~22% of WT FOXL2 mRNA levels, while the gDNA levels of both alleles were similar in KGN cells (Fig 1B). When COV434, a cell line derived from JGCTs lacking the 402C>G mutation (Jamieson *et al*, 2010), was tested as a control using the mutant allele-specific primer, no amplicons containing the 402C>G mutation were detected (Fig 1B). We also observed a comparable allelic imbalance by allele-specific real-time RT-PCR on full-length FOXL2 mRNA generated by the cap analysis of gene expression (CAGE) method from KGN cells (Appendix Fig S1G).

Next, we tested whether the decreased steady-state levels of the variant FOXL2 mRNA resulted from alterations in mRNA stability. WT and variant FOXL2 mRNA abundances were measured in KGN cells at several time points using allele-specific real-time RT-PCR after adding actinomycin D (ActD), which blocks transcription. The level of variant FOXL2 mRNA decreased more rapidly than that of WT mRNA ($t_{1/2} = 3.68$ h for variant FOXL2 versus $t_{1/2} = 15.43$ h for WT), indicating that the lower steady-state level of the variant FOXL2 mRNA resulted from decreased mRNA stability (Fig 1C). Variant FOXL2 mRNA instability was unlikely due to its altered secondary structure, as the 402C>G mutation was not predicted to affect the mRNA structure, based on M-fold analysis (<http://www.bioinfo.rpi.edu/applications/mfold>; Zuker, 2003; Appendix Fig S2A and B).

Selective degradation of the variant FOXL2 mRNA by miR-1236-3p

Next, we determined whether selective degradation of the variant FOXL2 mRNA was due to miRNAs targeting the mutation site in the CDS. By analyzing the variant FOXL2 mRNA sequence with an miRNA-prediction tool (<https://genie.weizmann.ac.il/pubs/mir07/index.html>; Kertesz *et al*, 2007), we identified miRNAs predicted to bind the sequence surrounding the mutation (Appendix Table S1). We selected five miRNAs predicted to preferentially bind the mutant allele over the WT allele for further analysis. Using DNA oligonucleotides complementary to these miRNAs (anti-miRNAs), which effectively inhibited the respective miRNAs based on upregulation of their known target mRNAs (Appendix Fig S3A), we found that anti-miR-1236 specifically increased the variant FOXL2 mRNA-expression level without affecting that of WT FOXL2 in KGN cells (Fig 2A). The remaining four anti-miRNAs did not affect the mRNA levels of WT FOXL2 or the variant (Fig 2A). Consistent with this effect at the mRNA level, anti-miR-1236 also increased FOXL2 protein expression (Fig 2B). Conversely, transfection of an miR-1236 RNA mimic decreased FOXL2 protein expression (Fig 2C). In contrast to our observations with KGN cells, transfecting miR-1236 into COV434 cells (which lack the 402C>G mutation) did not affect FOXL2 protein expression (Fig 2C). To further validate the specificity of miR-1236 on the FOXL2 variant, we cotransfected a WT- or variant FOXL2-expression plasmid together with miR-1236 RNA into 293T cells expressing minimal WT FOXL2 (but not the variant), and changes in FOXL2 expression were monitored by Western blotting. The miR-1236 mimic specifically decreased the variant FOXL2 level without affecting that of WT FOXL2 (Appendix Fig S3B). The effective depletion or overexpression of miR-1236 in cells transfected with anti-miRNAs or mimic oligonucleotides, respectively, was confirmed with a TaqMan® microRNA assay for miR-1236 (Appendix Fig S3C).




Figure 1.

Figure 1. Allelic imbalance of heterozygous FOXL2 transcripts in AGCT cells.

- A Bar graph and box-and-whisker plots are presented, which show the allelic proportions of WT *FOXL2* mRNA and 402C>G *FOXL2* mRNA in AGCT tissues from 46 patients analyzed by pyrosequencing. The box plot represents the minimum value, first quartile, median, third quartile, and maximum value of a data set. X-axis indicates mRNAs of WT *FOXL2* and 402C>G *FOXL2*. The whiskers extend to the most extreme data points not considered outliers, and the outliers are represented as dots. Comparisons between groups were performed using Student's *t*-test, and *P* values are presented.
- B The relative abundances of WT and variant *FOXL2* mRNA were analyzed in KGN and COV434 cells by pyrosequencing (left graph), allele-specific RT-PCR (middle graph), and real-time RT-PCR (right graph). gDNA was detected as a positive control. The relative abundances of the variant *FOXL2* mRNA were normalized to that of WT mRNA (set to 1). *FOXL2* mRNA levels detected by real-time RT-PCR were normalized to matching gDNA levels. The pyrosequencing data are presented from two independent experiments. The allele-specific semi-quantitative and real-time RT-PCR data are presented as the mean \pm SEM from three independent experiments. The *P* values were analyzed by unpaired, two-tailed Student's *t*-test ($***P < 0.001$). n.d. not detected.
- C RNA-decay rates of WT and 402C>G *FOXL2* mRNAs in KGN cells were determined after treatment with 5 μ g/ml ActD for the indicated times. The estimated half-lives of each transcript are presented. The data are presented as the mean \pm SEM from three independent experiments.

Source data are available online for this figure.

Moreover, we generated miR-1236-knockout (KO) KGN and COV434 cell lines using a clustered regularly interspaced short palindromic repeat (CRISPR)/CRISPR-associated protein 9 (Cas9)-nickase-based system (Appendix Fig S4A–H) that is known to exert minimal off-target effects (Ran *et al.*, 2013). Using extracted total RNA from the KO cells, we performed TaqMan® microRNA analyses and confirmed miR-1236 depletion (Appendix Fig S4B and F). Northern blotting showed that miR-1236 was detectable in control KGN cells, but not in miR-1236^{-/-} cells (Appendix Fig S4I). Then, we evaluated WT and variant *FOXL2* mRNA levels in the KO cells. Variant *FOXL2* mRNA expression reverted to the level of WT mRNA when both alleles (miR-1236^{-/-}) were excised and was partially restored when a single allele (miR-1236^{-/+}) of miR-1236 was disrupted, without affecting WT *FOXL2* mRNA expression (Fig 2D). Re-introducing miR-1236 mimic via transfection in miR-1236-KO cells downregulated variant *FOXL2* mRNA expression without altering WT *FOXL2* expression (Fig 2D), demonstrating that restoration of variant *FOXL2* mRNA level resulted from miR-1236 KO. The observation that known miR-1236 target genes, *AFP* and *ZEB1* (Wang *et al.*, 2014; Gao *et al.*, 2015), were upregulated in these KO cells also suggested that efficient and effective excision of miR-1236 occurred (Appendix Fig S4J and K). Consistently, elevated *FOXL2* protein expression was confirmed in miR-1236-KO KGN cells, whereas no change in *FOXL2* expression occurred in miR-1236-KO COV434 cells (Fig 2E). These *in vivo* genome-editing results further corroborated miR-1236 as an endogenous functional miRNA that selectively acted on variant *FOXL2* mRNA.

Allele-specific downregulation of the *FOXL2* mRNA variant with miR-1236 via targeting the 402C>G mutation site

Reporter constructs were developed to confirm that miR-1236 targets the 402C>G locus in the variant *FOXL2* mRNA CDS. A 231-bp DNA fragment containing the 402C>G locus of the *FOXL2* variant or the corresponding fragment of WT *FOXL2* was inserted, in-frame, into the CDS of the luciferase gene (Fig 3A). miR-1236 transfection decreased the activity of the luciferase reporter harboring the 402C>G *FOXL2* variant sequence by 70%, without affecting the activity of the luciferase reporter containing the WT *FOXL2* sequence (Fig 3B). Similarly, we observed allele-specific repression when the 231-bp *FOXL2* fragments were inserted in the 3'-UTR of the luciferase reporter gene (Fig 3C and D). The specificity of the interaction between miR-1236 and the variant locus in *FOXL2* mRNA was further tested using two miR-1236 mutants. The miR-

1236-G mutant shifted the 7-mer seed match from the 402C>G *FOXL2* mRNA to the WT *FOXL2* mRNA (Fig 3E), whereas the seed sequence of the miR-1236-U mutant did not match either 402C>G or WT *FOXL2* mRNA (Fig 3F). Notably, the selectivity of the miR-1236-G mutant was reversed, as it repressed the WT reporter without affecting the 402C>G mutant reporter (Fig 3E). In contrast, the miR-1236-U mutant did not suppress the 402C>G mutant or WT *FOXL2* reporter (Fig 3F).

In addition, the preferential binding of miR-1236 to 402C>G over the WT *FOXL2* transcript was confirmed by performing *in vitro*-binding assays. Synthetic 230-nt transcripts of WT or 402C>G *FOXL2* mRNA were incubated with radiolabeled miR-1236, and RNA-duplex formation was monitored. miR-1236 preferentially duplexed with the 402C>G *FOXL2* transcript, with predicted dissociation constants (K_d) of 1,589 and 194 nM for the WT and 402C>G *FOXL2* transcripts, respectively (Fig 3G). Thus, these data indicate that the 402C>G locus was critical for distinguishing the effects of miR-1236 on *FOXL2* expression.

AGO3 as the major miRISC component for miR-1236-mediated *FOXL2* variant mRNA degradation

Because AGO2 is known to act primarily on the 3'-UTR of target mRNAs (Hafner *et al.*, 2010) and a recent study demonstrated that AGO3 is also associated with slicer activity (Park *et al.*, 2017), we investigated the possibility that AGO3 can regulate RISC activity against the variant *FOXL2* mRNA by recognizing the mutated site in its CDS. Each AGO was knocked down using siRNAs, and changes in the levels of WT and variant *FOXL2* mRNAs were examined in KGN cells. Of interest, we found that AGO3 knockdown preferentially increased mRNA expression of the *FOXL2* variant without affecting that of WT *FOXL2* (Fig 4A). In contrast, AGO2 knockdown increased both the WT and variant mRNAs (Fig 4A), indicating that WT *FOXL2* mRNA is degraded by AGO2-mediated miRNAs targeting the 3'-UTR, as previously described (Dai *et al.*, 2013; Luo *et al.*, 2015; Wang *et al.*, 2015). Depletion of either AGO1 or AGO4 did not affect the levels of the WT and variant *FOXL2* mRNAs (Fig 4A). Consistent with the effects on the mRNA levels, increased *FOXL2* protein levels were observed in KGN cells following the depletion of either AGO2 or AGO3 (Fig 4B). In addition, we determined whether the slicer activity of AGO3 promoted the decay of the variant *FOXL2* using a slicer-incompetent E638A mutant of AGO3 (Park *et al.*, 2017). AGO3 knockdown consistently elevated the level of variant *FOXL2* mRNA and transfecting WT-AGO3 in AGO3-depleted cells effectively




Figure 2. Selective downregulation of the variant FOXL2 mRNA allele by miR-1236.

A, B Changes in WT and variant FOXL2 mRNA expression were assessed by RT-PCR (top) and real-time RT-PCR (bottom) (A) or by Western blot analysis (B) after transfecting KGN cells with anti-miRNAs for 48 h.

C FOXL2 protein levels after transfection of control or miR-1236 were assessed in KGN and COV434 cells.

D The mRNA levels of WT and variant FOXL2 in control, miR-1236^{+/+}, and miR-1236^{-/-} KGN cells, with or without miR-1236 transfection, were determined by RT-PCR (top) or real-time RT-PCR (bottom).

E FOXL2 protein expression in control, miR-1236^{+/+}, and miR-1236^{-/-} KGN cells and two independent miR-1236^{-/-} (#1 and #2) COV434 cell lines were determined by Western blotting.

Data information: Representative gel images are also shown. All quantified results (mean ± SEM) are from at least three independent experiments. Different letters ($P < 0.0001$; Student–Newman–Keuls test) or asterisks ($*P < 0.05$; Student's *t*-test) denote significant differences.

Source data are available online for this figure.

restored the level of variant FOXL2 mRNA to that in cells without any treatment (Appendix Fig S5A). Transfected E638A-AGO3 also decreased the expression of variant FOXL2 mRNA to a similar extent as transfected WT-AGO3 (Appendix Fig S5A). Consistently, the

transfected WT-AGO3 or E638A mutant-AGO3 similarly decreased the level of FOXL2 protein (Appendix Fig S5B). Thus, these results indicate that AGO3-mediated destabilization of the variant FOXL2 mRNA does not require its slicer activity. Based on this result, we




Figure 3.

Figure 3. Allele-specific downregulation of *FOXL2* mRNA by *miR-1236*, which targets 402C>G.

- A Schematic representation of the luciferase reporter constructs used to assay miR-1236 activity against a CDS target site in *FOXL2* mRNA. The 231-bp human *FOXL2* segments harboring either the C402 (WT) or the G402 (mutant) nucleotide were inserted in-frame into the CDS of the *luciferase* gene in the pGL3 control vector.
- B Luciferase activity of the reporter constructs shown in (A) was measured in KGN cells after transfection with an miR-1236 mimic for 48 h. The black arrow indicates the position of 402C>G mutation site.
- C A schematic diagram of the luciferase reporter constructs generated by inserting the predicted miR-1236-target sequences of WT and 402C>G *FOXL2* mRNAs in the 3'-UTR of *luciferase*.
- D Luciferase activities were measured in KGN cells, using the reporter constructs shown in (C), after transfection with a control miRNA or an miR-1236 mimic for 48 h.
- E, F miR-1236 mutants, in which the C that pairs with G402 of the *FOXL2* mutant was substituted with either G (miR-1236-G) (E) or U (miR-1236-U) (F), were cotransfected into KGN cells with one of the reporter constructs described above. Luciferase activities were subsequently determined. Arrows indicate the mismatched sites.
- G *In vitro* annealing kinetics of miR-1236 with 230 nt-long transcripts of WT or variant *FOXL2*. ³²P-labeled miR-1236 (0.5 nM) was incubated with increasing concentrations of synthetic *FOXL2* transcripts (0, 2.5, 12.5, 25, or 50 nM). *FOXL2* mRNA-miR-1236 complexes were resolved on a 6% native gel and detected by autoradiography (left). The predicted K_{dS} for the WT and 402C>G *FOXL2* transcripts are presented in the right graph.

Data information: The data are expressed as the mean \pm SEM from three independent experiments, performed in triplicate. The *P* values were analyzed by unpaired, two-tailed Student's *t*-test (***P* < 0.001).

Source data are available online for this figure.

speculate that formation of the miR-1236-AGO3-RISC on the variant *FOXL2* mRNA may induce translational repression, which leads to rapid degradation of the variant *FOXL2* mRNA via pathways that do not involve the slicer activity of Ago3 (e.g., decapping and exosome activities).

To determine whether miR-1236 mediates these effects, AGO-depleted KGN cells were transfected with miR-1236, and changes in *FOXL2* mRNA-expression levels were evaluated by allele-specific real-time RT-PCR. miR-1236 overexpression did not alter WT *FOXL2* mRNA expression in cells with AGO1, AGO2, AGO3, or AGO4 depletion (Fig 4C; left graph). In contrast, miR-1236 transfection did not significantly affect the variant *FOXL2* mRNA expression in AGO3-knockdown cells (Fig 4C; right graph). miR-1236 transfection was partially effective in AGO2-knockdown cells, but efficiently downregulated variant *FOXL2* mRNA levels in AGO1- or 4-depleted cells (Fig 4C; right graph). In addition, to ascertain the role of AGO3 in the activity of miR-1236 in promoting variant *FOXL2* mRNA decay, changes in *FOXL2* mRNA-expression levels were determined in miR-1236-KO cells after silencing each AGO. Our results were similar to those shown in Fig 4A when the control cell line expressing miR-1236 (miR-1236^{+/+}) was used in these experiments (Fig 4D; left graph). In sharp contrast, depletion of AGO1, 3, or 4 failed to increase the level of variant *FOXL2* mRNA in miR-1236-KO cells (miR-1236^{-/-}; Fig 4D; right graph). Consistent with the results shown in Fig 4A, AGO2 depletion increased both WT and variant *FOXL2* mRNA expression in miR-1236^{-/-} cells (Fig 4D; right graph). These results imply that AGO3 is the major AGO responsible for miR-1236-mediated decay of the variant *FOXL2* mRNA.

To assure miRISC formation of AGO3 with the variant *FOXL2* mRNA, the level of enriched variant *FOXL2* with AGO1-4 was determined following transfection of equally expressed FLAG-tagged AGOs, miR-1236, and a pGL3c-CDS-variant *FOXL2* construct (Fig 3A) in 293T cells, which lacked the variant *FOXL2* transcript. As shown in Fig 4E, variant *FOXL2* mRNA was highly enriched in the AGO3 immunoprecipitate compared with the other AGO immunoprecipitates. We further detected the *in vivo* formation of a miRISC comprising endogenously expressed variant *FOXL2* mRNA, miR-1236, and AGO3. The variant *FOXL2* transcript was highly incorporated into the AGO3 immunoprecipitate, whereas negligible WT *FOXL2* was incorporated (Fig 4F). Preferential incorporation of the variant *FOXL2* transcript with AGO3 was

reversed by transfection of the miR-1236-G mutant into KGN cells, in which predominant incorporation of WT *FOXL2* mRNA occurred with a concomitant decrease in the *FOXL2* protein level (Fig 4F). In addition, we confirmed the association of endogenous miR-1236 with AGO3 miRISC by analyzing AGO3-coupled with miR-1236 in KGN cells (Appendix Fig S6). Together, these results indicate that a functional miRISC consisting of AGO3, miR-1236, and the variant *FOXL2* mRNA preferentially formed in KGN cells.

KGN cells express all four AGOs, and a comparison of their relative abundances indicated that AGO3 was the most abundant (Appendix Fig S7A), whereas the level of AGO2 mRNA was the highest in 293T cells (Appendix Fig S7B). Further, we analyzed the relative abundances of AGO mRNAs in individual AGCT tissues from patients, using high-throughput deep sequencing. As shown in Fig 4G, we observed dramatically higher expression of AGO3 mRNA-expression levels compared with those of AGO1, AGO2, and AGO4 in 20 independent patients with AGCT, supporting a previously unidentified role of AGO3 as a functional miRISC component in AGCT cells.

Identification of DHX9 as a functional component for AGO3-miRISC regulation of variant *FOXL2* mRNA

We further investigated potential functional AGO3-miRISC components that could regulate expression of the variant *FOXL2* mRNA. Since most miRNAs target in CDS regions in plants (Iwakawa & Tomari, 2013), candidate human proteins were selected based on searches for homologous genes in plants that regulate the activities of miRISCs, AGO2-binding proteins, and RNA-binding proteins. Among them, silencing DHX9, an ATP-dependent RNA helicase A that has been reported to function as a siRNA-loading and-recognition factor for AGO2-siRISC assembly (Robb & Rana, 2007; Fu & Yuan, 2013), prominently affected the abundance of variant *FOXL2* mRNA (Fig 5A). In particular, DHX9 silencing restored the variant *FOXL2* mRNA level to that of the WT mRNA, involving a 6-fold increase of variant *FOXL2* mRNA compared with a 1.7-fold increase of the WT mRNA (Fig 5A). In contrast, depletion of GW182, a well-known miRISC component that associates with AGO2, increased the abundances of both the WT and variant *FOXL2* mRNAs (Fig 5A). Consistent with these effects on the mRNA level, DHX9- or GW182-depletion also increased expression of the *FOXL2* protein (Fig 5B).




Figure 4.

Figure 4. Identification of AGO3 as the major miRISC regulator for variant FOXL2 mRNA degradation.

- A, B Changes in WT and variant *FOXL2* mRNA-expression levels were assessed by real-time RT-PCR (A) or Western blot analysis (B) after transfecting KGN cells with siRNAs against AGO mRNAs for 48 h. The data (mean \pm SEM) are from three independent experiments, performed in triplicate.
- C The mRNA levels of WT (left) and variant *FOXL2* (right) were determined in KGN cells by real-time RT-PCR, after transfecting a control miRNA or miR-1236. The data (mean \pm SEM) are from three independent experiments, performed in triplicate.
- D The mRNA levels of WT and the variant *FOXL2* in control (left) and miR-1236^{-/-} KGN cells (right) after transfecting siRNAs against AGO mRNAs were determined by allele-specific real-time RT-PCR. The data (mean \pm SEM) are from three independent experiments, performed in triplicate.
- E 293T cells were transfected with an miR-1236 mimic (50 nM) for 24 h, followed by cotransfection with expression vectors encoding FLAG/HA-tagged variants of the indicated human AGOs and pGL3c-CDS-MT for 24 h. The empty p3XFLAG-CMV-10 vector was used as control. Co-immunoprecipitated mRNAs were reverse transcribed, and the cDNA products were used for allele-specific real-time PCR analysis of the *FOXL2* variant and *GAPDH* mRNAs (top). The level of variant *FOXL2* mRNA immunoprecipitated using FLAG-tagged AGO proteins was normalized using the level of *GAPDH* mRNA from the same lysates. The immunoprecipitated-AGO proteins were detected by Western blotting (bottom). The data (mean \pm SEM) are from three independent experiments.
- F *In vivo* association of AGO3-mediated miRISC formation with *FOXL2* mRNAs is shown. Following transfection of a control miRNA or the miR-1236-G mutant into KGN cells, AGO3-mediated RISC-associated RNAs were isolated by immunoprecipitation with an anti-AGO3 antibody. IgG was used as a control. The co-immunoprecipitated mRNAs were reverse transcribed using a *FOXL2*-430-R primer binding downstream of the 402C>G site. The cDNA products were used for *FOXL2* allele-specific PCR analysis with the *FOXL2*-279F primer (Appendix Fig S1), and a representative result (top left) is shown. Quantitative real-time RT-PCR results that examined *FOXL2* mRNAs, normalized using the level of *GAPDH* mRNA (bottom left), are also presented. Western blot analysis of immunoprecipitated AGO3 and inputs are shown in the right panel. The data are presented as the mean \pm SEM of two independent experiments.
- G RNA-seq analysis was performed to determine AGO-expression levels (transcripts per million) from the individual tissues from 20 independent AGCT patients. X-axis represents mRNAs of AGO1 to 4. The box plot represents the minimum value, first quartile, median, third quartile, and maximum value of a data set. The whiskers extend to the most extreme data points not considered outliers, and the outliers are represented as dots.

Data information: Different letters ($P < 0.05$; Student–Newman–Keuls test) or asterisks (***) ($P < 0.001$) denote statistically significant differences. Source data are available online for this figure.

Immunoprecipitation analysis of AGOs with DHX9 or GW182 showed that DHX9 exhibited stronger binding affinity to AGO3, whereas GW182 bound AGO2 more tightly (Fig 5C). Furthermore, to detect the miR-1236-loaded RISC in KGN cell lysates, miRISC pull-down experiments were performed using biotin-tagged 2'-O-methyl oligonucleotides complementary to miR-1236. Efficient pull-down of miR-1236 using the miR-1236-complementary biotin-tagged 2'-O-methyl oligonucleotides was confirmed as demonstrated via depletion of endogenous miR-1236 in the supernatant (Appendix Fig S8). Immunoblot analyses of the pull-down assays showed that miRISC associated with endogenous miR-1236 and with AGO3, DHX9, AGO2, and GW182 proteins, whereas no association was observed in the control (Fig 5D). Their specific interaction was assured as *GAPDH*, a non-miRISC component, was not copurified with the miR-1236-complementary biotin-tagged 2'-O-methyl oligonucleotides (Fig 5D). In particular, we observed an increase in the association with AGO3 and DHX9 to miR-1236 compared with AGO2 and GW182 based on the relative quantification of bound proteins compared to the input (Fig 5D). Thus, these results indicate that AGO3 and DHX9 are components of the miRISC with miR-1236.

In addition, we determined the functional role of DHX9 for *in vivo* AGO3–miRISC formation with the variant *FOXL2* mRNA. Consistent with the data shown in Fig 4F, a predominant incorporation of the variant *FOXL2* transcript over the WT transcript in AGO3 immunoprecipitates was observed (Fig 5E). However, DHX9 depletion significantly decreased the incorporation of variant *FOXL2* mRNA in AGO3 immunoprecipitates without affecting the incorporation of WT *FOXL2* mRNA (Fig 5E). To assess the effect of DHX9 on miR-1236 loading with AGOs, miRISCs were immunoprecipitated using anti-AGO3 or anti-AGO2 antibodies in control and DHX9-depleted KGN cells, and changes in associated miR-1236 were assessed using the TaqMan® microRNA assay. The amount of miR-1236 pulled down via anti-AGO3, but not with anti-AGO2, was significantly decreased when DHX9 was depleted (Fig 5F), which suggests that DHX9 plays a role in loading miR-1236 specifically with AGO3.

Furthermore, we investigated whether the alternative miR-1236-RISC, comprising AGO3 and DHX9, had a specific location preference for the miRNA-responsive element (MIRE) to either the CDS or 3'-UTR. Luciferase assays were performed in AGO3-, DHX9-, AGO2-, or GW182-depleted KGN cells using two independent reporters. These reporters contained a 231-bp segment of *FOXL2* that harbored the 402C>G mutated nucleotide, which were inserted into the CDS (pGL3c-*FOXL2*-CDS) or the 3'-UTR (pGL3c-*FOXL2*-UTR) of *luciferase* as described in Fig 3A and C. Interestingly, we found that the miR-1236-induced decrease of the luciferase activity from pGL3c-*FOXL2*-CDS was abolished in AGO3- or DHX9-silenced cells, but not in AGO2- or GW182-silenced cells (Fig 5G). Conversely, the miR-1236-induced decrease of the luciferase activity from pGL3c-*FOXL2*-UTR was abolished in AGO2- or GW182-silenced cells, but not in AGO3- or DHX9-silenced cells (Fig 5H). Taken together, these results suggest that the miR-1236-AGO3-RISC acts preferentially on the MIRE positioned within the CDS region, whereas the miR-1236-AGO2-RISC acts preferentially on the MIRE positioned within the 3'-UTR region.

Oncogenic function of miR-1236 in AGCT cells

Next, we investigated the oncogenic effects of miR-1236 in AGCT cells. We found that transfecting KGN cells with an miR-1236 mimic or anti-miR-1236 significantly decreased or increased the numbers of annexin V-positive apoptotic cells and increased or decreased the cellular viability, respectively (Fig 6A and B, Appendix Fig S9A and B). Overexpression of other anti-miRNAs, which showed no significant effects on *FOXL2* expression (Fig 2A and B), did not alter KGN cell viability (Appendix Fig S9C). Moreover, treatment with the miR-1236 mimic accelerated cell cycle progression to S phase, whereas anti-miR-1236 induced cell cycle arrest at G₀/G₁ phase (Fig 6C and D). Transfection of either miR-1236 or anti-miR-1236 in *FOXL2*-silenced cells failed to alter the numbers of apoptotic cells, cell cycle, and cell viability (Fig 6A–D, Appendix Fig S9D and E), indicating that the effects of miR-1236 on KGN cells were *FOXL2*-

dependent. The effect of *FOXL2* on GCT cell migration was also examined using Transwell chambers, and we found that miR-1236 or anti-miR-1236 transfection promoted or suppressed cell

migration, respectively, and these effects were abolished upon *FOXL2* depletion (Fig 6E and F). Transfecting miR-1236 into COV434 cells did not alter either cell viability or cell migration and




Figure 5.

Figure 5. DHX9 as a preferential interactor with AGO3 and an essential molecule for AGO3-mediated degradation of the variant FOXL2 mRNA.

- A Changes in WT and variant *FOXL2* mRNA levels in KGN cells were assessed by real-time RT-PCR after transfecting siRNAs against the indicated factors for 48 h. The data are presented as the mean \pm SEM from three independent experiments, performed in triplicate.
- B *FOXL2* protein-expression levels were determined by Western blotting after transfecting KGN cells with control, DHX9, or GW182 siRNAs for 48 h. Quantification of *FOXL2* protein expression is presented in the bottom panel. The data are presented as the mean \pm SEM from three independent experiments. The *P* values were analyzed by unpaired, two-tailed Student's *t*-test (***P* < 0.01, ****P* < 0.001).
- C Relative binding affinities of DHX9 and GW182 to AGOs. 293T cells were transfected with expression vectors encoding the indicated FLAG/HA-tagged AGOs, and cell extracts were prepared and immunoprecipitated with an anti-FLAG antibody, followed by immunoblot analyses (top). The empty p3XFLAG-CMV-10 vector was used as a control. The band intensities of immunoprecipitated DHX9 and GW182 were quantified and normalized following pulldown with the indicated AGOs (bottom). The data are presented as the mean \pm SEM from three independent experiments. * and # indicate statistically significant differences in the respective amounts of DHX9 or GW182 bound to AGO1. The *P* values were analyzed by unpaired, two-tailed Student's *t*-test (*P* < 0.05).
- D Association of endogenous miR-1236 with RISC components in KGN cells was determined via pulldown assays using immobilized 2'-*O*-methylated oligonucleotides (2'-*O*-Me oligos) complementary to miR-1236 followed with a pulldown using streptavidin-coupled Dynabeads and Western blot analyses (top). Relative quantification of bound proteins compared with proteins from the input is presented as fold enrichment (bottom). Efficient pulldown of endogenous miR-1236 using the 2'-*O*-Me oligos was confirmed with depleted miR-1236 in the discarded supernatant following the pulldown (Appendix Fig S8). As a control, 2'-*O*-Me oligos not complementary to miR-1236 were used. The data are the means \pm SEM from three independent experiments. The *P* values were analyzed by unpaired, two-tailed Student's *t*-test (**P* < 0.05, ***P* < 0.01).
- E Following transfection of control siRNA or siDHX9 into KGN cells, AGO3-mediated RISC-associated RNAs were immunoprecipitated using an anti-AGO3 antibody. IgG was used as a control. Co-immunoprecipitated mRNAs were reverse transcribed using a *FOXL2*-430-R primer binding downstream of the 402C>G site. The cDNA products were used for *FOXL2* allele-specific PCR analysis with a *FOXL2*-279F primer (Appendix Fig S1A), and a representative result obtained by RT-PCR (top) is shown. Quantitative real-time RT-PCR results (middle) are also presented as fold enrichment of *FOXL2* mRNAs normalized using the level of *GAPDH* mRNA. Western blots of immunoprecipitated AGO3 and the inputs are shown in the bottom panel. The data are presented as the mean \pm SEM from three independent experiments. Different letters denote statistically significant differences (*P* < 0.05; Student–Newman–Keuls test).
- F We examined whether DHX9 affected the association between miR-1236 and AGOs. After transfecting KGN cells with control siRNA or siDHX9, the total RNA and AGOs-mediated RISC-associated RNAs were isolated following immunoprecipitations using anti-AGO3 or anti-AGO2 antibodies. The AGOs-immunoprecipitated RNAs were extracted using an acidic phenol:chloroform mixture (5:1, pH 4.3) and precipitated with isopropanol using 10% of 3 M NaOAc (pH 5.2). The enrichment of miR-1236 within miRISCs was detected using the TaqMan[®] microRNA assay in the immunoprecipitated RNAs and normalized using the level of total miR-1236. The data (means \pm SEM) are presented as the fold enrichment calculated from three independent experiments. Different letters denote statistically significant differences (*P* < 0.05; Student–Newman–Keuls test).
- G, H Luciferase activities of the reporter constructs presented in Fig 3A and C were measured in KGN cells after transfecting the miR-1236 mimic, indicated siRNAs, and either pGL3c-CDS-*FOXL2* MT or pGL3c-UTR-*FOXL2* MT for 48 h. The data are expressed as the means \pm SEM from three independent experiments and were performed in triplicate. The *P* values were analyzed by unpaired, two-tailed Student's *t*-test (****P* < 0.001).

Source data are available online for this figure.

had no effect on *FOXL2* protein expression (Fig 6G and H), validating the 402C>G allele-specific oncogenic effects of miR-1236. Additionally, we confirmed the oncogenic properties of miR-1236 using miR-1236-KO KGN cells and obtained results that were consistent with the effects observed after anti-miR-1236 transfection. miR-1236^{-/-} KGN cells exhibited significantly reduced cell survival, proliferation, and migration compared with control KGN cells, and miR-1236^{-/+} KGN cells showed intermediate activities (Fig 6I–K). In sharp contrast, the oncogenic characteristics were not altered in miR-1236^{-/-} COV434 cells (Fig 6L).

In addition, we performed RNA-seq analysis to examine the transcriptomic profile in KGN cells following anti-miR-1236 transfection. Genes were considered differentially expressed if the adjusted *P* < 0.05 and if there was at least a 1.5-fold change in expression. Pairwise differential expression analysis identified 136 differentially expressed gene transcripts, including 65 genes downregulated via miR-1236 depletion. Gene ontology (GO) pathway enrichment analysis showed significant associations with pathways including those that mediate apoptosis, signal transduction, angiogenesis, and tumor necrosis factor-mediated signaling (Appendix Fig S10A). Heat maps and unsupervised hierarchical clustering are shown in Appendix Fig S10B. Upregulated *FOXL2* mRNA levels in anti-miR-1236 transfected cells were confirmed (log₂ fold change = 0.650365, *P* = 6.89e-07) (Appendix Fig S10C and D). Furthermore, changes in mRNA levels of reported, known targets of miR-1236 were analyzed from RNA-seq data sets and validated using real-time PCR analysis (Appendix Fig S10C and D). Except for *FOXL2* and *p21*,

whose transcript levels were increased, none of the other tested genes showed altered profiles upon miR-1236 inhibition (Appendix Fig S10D). In regard to the upregulated *p21* mRNA in anti-miR-1236-transfected cells, this change is likely associated with upregulated *FOXL2* expression since we previously demonstrated that *p21* is a transcriptional target gene of *FOXL2* (Kim *et al*, 2014). At the same time, this observation could be due to inhibition of miR-1236 using anti-miR-1236 as *p21* is also a target for miR-1236 (Wang *et al*, 2016). Regardless, since both *FOXL2* and *p21* inhibit cell proliferation, miR-1236-mediated downregulation of *FOXL2* and *p21* would likely promote oncogenesis in cells. In addition, the data in Fig 6 show that oncogenic effects of miR-1236 are directly associated with expression levels of *FOXL2*.

***In vivo* suppression of metastasis by miR-1236 KO and inverse expression between miR-1236 and the variant *FOXL2* mRNA in tissues from AGCT patients**

We further performed an *in vivo* xenograft mouse experiment using the miR-1236^{-/-} KGN cells and examined the effect of miR-1236 loss on AGCT metastasis. Xenografting control KGN cells resulted in AGCT metastasis to the intestine in mice (Fig 7A and B). In contrast, miR-1236^{-/-} cell-xenografted mice showed significantly fewer metastasized tumor nodules (Fig 7A and B). By analyzing the relative abundances of *FOXL2* transcripts in these tumors, we confirmed that the miR-1236-KO tumors expressed more variant *FOXL2* mRNA than WT mRNA (Fig 7C). These *in vivo* results provided further

support that miR-1236 acts as a critical molecule for AGCT metastasis by preferentially targeting the variant *FOXL2* mRNA.

Moreover, we examined whether the aggressiveness of AGCT was indeed linked to the ability of miR-1236 to downregulate mRNA expression of the 402C>G *FOXL2* variant in patients with AGCT. Clinical tissues from AGCT patients (exhibiting metastasis or no evidence of metastasis) were examined for miR-1236 and *FOXL2* mRNA expression. Notably, increased miR-1236 and decreased 402C>G *FOXL2* mRNA levels were evident in metastatic AGCT

tissues compared with non-metastatic AGCT tissues (Fig 7D and E). In contrast, the expression of WT *FOXL2* mRNA was similar in both AGCT groups (Fig 7F). Moreover, comparative analysis of the miR-1236 and *FOXL2* mRNA-expression levels in tissues from AGCT patients revealed a strong inverse correlation between miR-1236 and variant *FOXL2* mRNA expression (Fig 7G). Such a correlation was not observed between miR-1236 and WT *FOXL2* mRNA (Fig 7H). Together, these results provided further support that miR-1236 caused haploinsufficiency of the tumor-suppressor *FOXL2* by




Figure 6.

Figure 6. FOXL2-mediated oncogenic effects of miR-1236 on cell death, cell cycle progression, and cell migration.

- A–D KGN cells were transfected with 200 nM of scrambled control or FOXL2-specific siRNAs for 24 h. Then, KGN cells were further transfected with 20 nM of control miRNA, miR-1236, control anti-miRNA, or anti-miR-1236. The proportion of annexin V-positive apoptotic cells (A and B) and the population at S phase (C and D) were analyzed by flow cytometry. The data are presented as the mean \pm SEM of three independent experiments. Different letters denote statistically significant differences ($P < 0.0001$; Student–Newman–Keuls test). Efficient silencing of *FOXL2* using specific siRNAs was confirmed by Western blotting.
- E, F KGN cells were transfected with control siRNA or siFOXL2 for 24 h. Then, KGN cells were further transfected with control miRNA or miR-1236 (E) or with control anti-miRNA or anti-miR-1236 (F) for 48 h, and Transwell-migration assays were performed. The migrated cells were imaged under a bright-field microscope (100 \times magnification, scale bar = 100 μ m). The results are from three independent experiments and represent fold changes in the average number of cells/field (mean \pm SEM). Different letters denote statistically significant differences ($P < 0.01$; Student–Newman–Keuls test).
- G, H COV434 cells were transfected with miR-control or miR-1236 for 48 h, after which cell viabilities (G) and migration abilities (H) were measured. The data are presented as the mean \pm SEM of three independent experiments. Immunoblots showing no change in FOXL2 protein are presented in the top panel, and images of migrated cells are presented in the right panel. The migrated cells were imaged under a bright-field microscope (100 \times magnification, scale bar = 100 μ m).
- I–K The properties of miR-1236^{-/-} and miR-1236^{-/-} KGN cells versus control KGN cells were assessed by measuring cell viability (I), cell proliferation (J), and cell migration (K). The data are presented as the mean \pm SEM of three independent experiments. The migrated cells were imaged under a bright-field microscope (100 \times magnification; top of Figure 6K). Scale bar = 100 μ m.
- L No difference in the cell-migration activities of control and two independent miR-1236^{-/-} (#1 and #2) COV434 cell lines. The migrated cells were imaged under a bright-field microscope (100 \times magnification, scale bar = 100 μ m; top), and the results (bottom) represent fold changes in the average number of cells/field. The data are presented as the mean \pm SEM of three independent experiments.

Data information: Different letters denote statistically significant differences ($P < 0.0001$; Student–Newman–Keuls test).

Source data are available online for this figure.

inducing degradation of the variant *FOXL2* mRNA following recognition of the 402C>G locus as a target site, suggesting that miR-1236 plays an important etiological role in AGCT (Fig 7I).

Discussion

Data from recent studies have suggested that miRNAs can target CDS sites in human cells, and the presence of functional miRNA recognition elements in CDSs was addressed in a recent human study (Duursma *et al*, 2008; Forman *et al*, 2008; Elcheva *et al*, 2009; Schnall-Levin *et al*, 2011; Wang *et al*, 2012; Cai *et al*, 2015; Zhang *et al*, 2018). However, the regulatory mechanism underlying miRNA activity against the CDS of target genes is largely unknown. The general conceptual agreement that miRNAs target 3'-UTRs mostly arises from the idea that translating ribosomes occupy the CDS region and interfere with RISC formation. Thus, the binding of a miRISC to the CDS region of a transcript has been considered a less-efficient mechanism of action than binding to the 3'-UTR (Bartel, 2009; Gu *et al*, 2009). However, a recent report showed that RISC formation in the CDS region causes translation repression, but do not appear to alter the overall ribosome occupancy on the target mRNA (Zhang *et al*, 2018), indicating the existence of disparate mechanism for translation repression of target mRNA by CDS-targeted miRNAs. In addition, over 70% of mammalian genes generate mRNA transcripts with differing lengths and 3'-UTR sequences due to alternative cleavage and polyadenylation, which would abolish some miRNA-binding sites (Tian & Manley, 2017). Accordingly, variations in 3'-UTR sequences would decrease the efficiency of miRNA-mediated target mRNA degradation, which is often observed with oncogenes in cancer (Mayr *et al*, 2007; Mayr & Bartel, 2009). Therefore, targeting CDSs can assure effective downregulation of gene expression by miRNA, especially for disease-associated genes whose expression levels need to be tightly controlled.

miR-1236, the trans-acting RNA regulator identified in this study, is a mirtron located in an intron of the negative-elongation factor complex, member E (*NELFE*) gene (Okamura *et al*, 2007; Ruby *et al*, 2007; Wen *et al*, 2015). The genomic sequences of the miR-1236

region are well conserved from lower vertebrates including cavefish, lizards, and chickens to mammalian vertebrates (Appendix Fig S11). The steady-state level of miR-1236 expression seems to be more than 300 copies per KGN cell, based on northern blot analysis of small RNAs purified from 2×10^7 miR-1236^{+/+} KGN cells, which showed a radioactive intensity similar to 10 fmol of miR-1236 mimics (Appendix Fig S4I). According to a very recent miRNA profiling data by Kim *et al* (2019), in which they developed and adopted bias-minimized accurate quantification by sequencing (AQ-seq) technology, miR-1236 was fairly well expressed as being in the top 50% of abundance among all miRNAs detected in human cells. Cumulative recent data revealed altered various cellular responses after modulating miR-1236 expression levels in diverse cell types (Gao *et al*, 2015; Chen *et al*, 2016; Wang *et al*, 2016; Feng *et al*, 2019; Li *et al*, 2019; Wang *et al*, 2020). In this study, miR-1236 selectively targeted the mutated 402C>G locus in the *FOXL2* CDS, without affecting WT *FOXL2* mRNA expression. Preferential targeting of the mutated “G” allele by miR-1236 was also reported in recent single-nucleotide polymorphism (SNP) studies: specific miR-1236 binding to the G allele over the C allele (rs11536889) in the 3'-UTR of Toll-like receptor 4 (Zhao *et al*, 2019) and specific miR-1236 targeting of the G allele over the T allele (rs4246215) in the 3'-UTR of Flap endonuclease 1 (Nanda *et al*, 2018). Furthermore, our *in vivo* and *in vitro* analysis of the phenotypes of miR-1236 KO cells collectively showed that miR-1236 acted as an oncomiR in AGCT cells by mainly causing selective variant *FOXL2* mRNA degradation (Figs 6 and 7).

For miR-1236 activity against variant *FOXL2* mRNA, we identified AGO3 as a key AGO protein (Fig 5). For over two decades, AGO2 has been considered as the key AGO protein in miRISC formation. However, AGO3 may have a distinct, but overlapping functionality for mRNA degradation based on its ability to interact with a similar spectrum of RNA transcripts (Meister *et al*, 2004; Azuma-Mukai *et al*, 2008; Landthaler *et al*, 2008; Huang *et al*, 2014; Park *et al*, 2017). This study provides evidence supporting this notion, considering that AGO3 depletion completely restored the variant *FOXL2* mRNA-expression level to the WT level; that an *in vivo* miRISC comprised endogenously expressed variant *FOXL2* mRNA,




Figure 7.

Figure 7. Inhibition of *in vivo* metastasis by miR-1236 KO and inverse expression between miR-1236 and the variant *FOXL2* mRNA in tissues from AGCT patients.

- A–C The effect of miR-1236 KO on AGCT metastasis was assessed, using an *in vivo* xenograft mice model. (A) Representative images of tumor nodules (white arrows) formed in the intestines of nude mice xenografted with control or miR-1236^{-/-} KGN cells are shown (left and right; scale bar = 5 mm). Hematoxylin and eosin staining confirmed the pathological characteristics of the metastasized GCT nodules (middle, black arrows; scale bar = 50 μm). The black dashed circles are metastasized nodules of xenografted KGN cells found in mouse intestines. (B) The number of tumor nodules formed in the intestines was counted in control (*n* = 8) and miR-1236^{-/-} (*n* = 8) mice. (C) Allele-specific real-time RT–PCR analysis of the WT and 402C>G variant *FOXL2* mRNAs was performed using RNA extracted from tumor nodules from control or miR-1236^{-/-} mice. The *P* values were analyzed by unpaired, two-tailed Student's *t*-test (***P* < 0.01, ****P* < 0.001).
- D–F Box-and-whisker plots showing the relative expression of miR-1236 (D), variant *FOXL2* mRNA (E), and WT *FOXL2* mRNA (F), respectively, in 32 patients with non-metastasized AGCTs and 14 patients with metastasized AGCTs. X-axis indicates patient subgroups depending on whether they exhibit metastasized AGCTs (meta) or not (non-meta). The relative miR-1236 levels were measured using a TaqMan[®] microRNA RT–qPCR assay, with expression normalized to *RNU6B*. The levels of 402C>G or WT *FOXL2* mRNA were determined by real-time RT–PCR, and the data were normalized to paired-gDNA levels. The relative levels of miR-1236 and *FOXL2* mRNAs were quantified by setting the levels of AGCT #1 to 1. Real-time RT–PCR was performed in triplicate for each specimen. The box plot represents the minimum value, first quartile, median, third quartile, and maximum value of a data set. The whiskers extend to the most extreme data points not considered outliers, and the outliers are represented as dots. Comparisons between groups were performed using Student's *t*-test, and *P* values are presented.
- G, H The estimated regression line superimposed on the scatter plot of miR-1236 levels with 402C>G *FOXL2* mRNA (G) or WT *FOXL2* mRNA (H) in AGCT samples (*n* = 46) is shown, along with correlation coefficient (*r*) and *P* values.
- I The proposed model for *FOXL2* haploinsufficiency induced by the 402C>G mutation during AGCT development.

Source data are available online for this figure.

miR-1236, and AGO3; and that the abundance of AGO3 was highest among the AGOs in AGCT tissues (Fig 4). Thus, we conjecture that the association of AGO3–RISC with target RNAs could be stabilized or promoted by interactions with distinct binding partner proteins, thereby facilitating target mRNA decay in the CDS. We discovered DHX9 as one of such proteins, which interacted preferentially with AGO3 (Fig 5C). DHX9 depletion increased variant *FOXL2* mRNA expression to the WT level (Fig 5A), and AGO3 association with variant *FOXL2* decreased markedly in DHX9-depleted cells (Fig 5E). Compromised target mRNA incorporation and processing were likely due to the inefficient incorporation of miR-1236 into the AGO3–miRISC when DHX9 was depleted (Fig 5F). These findings suggest that DHX9 plays a role in loading miR-1236 specifically with AGO3. Indeed, the miRISC pulldown experiment using oligonucleotides that complemented miR-1236 provided evidence that supports the existence of an alternative miRISC comprising AGO3, DHX9, and miR-1236. Additionally, the miR-1236 within this alternative miRISC preferentially associated with AGO3 and DHX9 rather than AGO2 and GW182 in KGN cells (Fig 5D). Concurrently, when we compared the levels of mature miR-1236 in DHX9-depleted or DHX9-overexpressed cells, the total amount of mature miR-1236 was significantly increased in DHX9-depleted cells compared with the control, whereas miR-1236 levels decreased in DHX9-overexpressed cells (Appendix Fig S12A and B). As miR-1236 is the mirtron of the host gene *NELFE*, we examined whether modulating DHX9 affected *NELFE* expression levels, which was not the case (Appendix Fig S12C and D). To examine whether the increase in miR-1236 that was induced via DHX9-knockdown was also seen with other miRNAs, several other miRNAs were analyzed. We found that a subset of miRNAs also showed increased expression when DHX9 was knocked down (Appendix Fig S12E). We speculate that DHX9 may also play a role in the biogenesis of certain miRNAs, including miR-1236, although the underlying mechanism that accounts for this effect remains to be elucidated.

In contrast to effects of DHX9 depletion, depletion of GW182, of which association with AGO2 is essential for gene silencing by miRNAs in animals (Eulalio *et al*, 2009), increased the abundances of both the WT and variant *FOXL2* mRNAs (Fig 5A and B), implying that GW182 is required for miRISC action on the 3'-

UTR present in both *FOXL2* mRNAs. In addition, GW182 preferentially bound AGO2 among other AGOs (Fig 5C), suggesting the presence of preferential binding partners for each AGO. Furthermore, our luciferase data that used two independent reporters, which contained segments from *FOXL2* that harbored the 402C>G mutated site that were inserted into the CDS or the 3'-UTR of *luciferase*, demonstrated that the alternative miRISC containing AGO3 and DHX9 preferred the MIRE to be positioned at the CDS, whereas the conventional miRISC containing AGO2 and GW182 prefer the MIRE present at the 3'-UTR (Fig 5G and H). While the detailed regulatory mechanism by which this non-canonical miRISC involving AGO3 and DHX9 targets the variant *FOXL2* CDS needs to be studied further, it is worthwhile to note findings of a recent study showing that CDS-targeted miRNAs require extensive base-pairing at the 3' side rather than the 5' seed in a manner independent of GW182 expression (Zhang *et al*, 2018), which overlap with characteristics of miR-1236.

Previously, we showed that the 134C>W missense mutant protein of *FOXL2* caused by the 402C>G mutation in *FOXL2* induced serine 33 hyperphosphorylation by GSK3β, thereby accelerating ubiquitin-mediated proteasomal degradation of the mutant protein (Kim *et al*, 2014). Here, we demonstrated that the mutation dramatically destabilized variant *FOXL2* mRNA via the allele-specific activity of miR-1236. Our findings suggest an etiological model for *FOXL2*-mediated AGCT development; selective degradation of the variant *FOXL2* mRNA by an miRISC composed of AGO3, DHX9, and miR-1236 promotes a strong allelic imbalance of *FOXL2* expression, which is aggravated by accelerated degradation of the mutant *FOXL2* protein. These processes result in *FOXL2* haploinsufficiency and insufficient suppression of AGCT development (Fig 7I). In 2009, Shah *et al* reported RNA sequencing results of four AGCTs, in which they observed various ratios of the variant and WT *FOXL2* mRNA levels (9:1; 1:1, 4:6 and 6:4) (Shah *et al*, 2009). In the present study, we observed a dramatic decrease in the variant *FOXL2* mRNA levels compared with WT mRNA levels in 46 AGCT patients. Further studies are needed to investigate how this discrepancy arises from. One possibility is that the RNA-seq study by Shah *et al* focused on expression of both *FOXL2* mRNAs rather than measurement of relative abundance of WT versus variant mRNAs. In fact,

total sequencing and mapped reads in their study were not enough to accurately quantify gene expression. To obtain coverage over the full-sequence diversity of complex transcript libraries, including rare and low-expressed transcripts, up to 500 million reads are required (Fu *et al*, 2014). Our RNA-seq study generated, on average, 134,312,269 total sequence reads and approximately 71,531 Mbp of total mapped base using Illumina HiSeq 2500 for 20 AGCT samples. In addition, we employed three other methods to assess the relative abundance of the WT versus variant *FOXL2* mRNA and obtained a consistent allelic imbalance.

In conclusion, our study provides a pathogenic mechanism in which a disease-associated heterozygous missense mutation can induce strong degradation of a variant mRNA by an alternative miRISC, which largely contributes to the haploinsufficiency of the gene and, consequently, development of the associated disease. We identified miR-1236 as an oncogenic miRNA in AGCT, suggesting a promising therapeutic target and prognostic parameter for patients with AGCT. Our study also provides evidence of the importance of nucleotide substitutions, such as somatic mutations and SNPs, in the CDS in regulating gene expression at the level of mRNA stability. A research approach that diverges from focusing mainly on the effects of missense mutations on protein activity may uncover many previously unsuspected mechanisms involving the regulation of mRNA stability by phenotype-associated missense mutations. In addition, we identified miR-1236 as an oncogenic miRNA in AGCT, suggesting a promising therapeutic target and prognostic parameter for patients with AGCT.

Materials and Methods

Cells culture and reagents

Human AGCT-derived KGN cells (Riken, Tsukuba, Japan) and JGCT-derived COV434 cells (Sigma-Aldrich, St. Louis, MO, USA) were cultured in Dulbecco's modified Eagle's medium (DMEM)-Ham's F12 (Caisson, North Logan, UT, USA) and DMEM (Caisson), respectively. 293T cells (American Type Culture Collection) were cultured in DMEM. The media were supplemented with 10% fetal bovine serum (FBS) and 1% penicillin-streptomycin (Caisson). The cells were grown in 5% CO₂ at 37°C. Other reagents were purchased from Sigma-Aldrich, unless otherwise indicated.

Plasmids

Plasmids expressing Myc-tagged variants of WT *FOXL2* and the C134W variant were prepared as described previously (Park *et al*, 2010; Kim *et al*, 2011; Kim & Bae, 2014). A 231-bp coding region (including nucleotide residue 402) of *FOXL2* containing the putative miRNA recognition site was amplified from gDNA of KGN cells using the following primers (Cosmo Genetech, Seoul, Korea): CDS-F (5'-CATGCCATGGCAAAGGGCTGGCAAATAG-3'), CDS-R (5'-CATGCCATGGGCGCCTCCGGCCCCGAAGAG-3'), 3'-UTR-F (5'-GCTCTAGAAGGGCTGGCAAATAGCA-3'), and 3'-UTR-R (5'-GCTCTAGACGGCGCCTCCGGCCCCGA-3'). The amplified fragment was then

cloned into the pGL3-control vector (Promega, Madison, WI, USA) at the *NcoI* or *XbaI* sites (restriction enzymes from Takara Bio, Shiga, Japan) in the CDS or 3'-UTR of the luciferase gene, respectively, generating pGL3c-CDS-*FOXL2s* and pGL3c-UTR-*FOXL2s*. A slicer-incompetent E638A mutant of AGO3 was generated using the following primers: AGO3-F (5'-CGCTAGCGGCCCGCATGGAAATCGG-3') with E638A-R (5'-TCCTGGATGATCGCTGTCCGGGGTCTCTG-3') and E638A-F (5'-CAGAGACCCCGACAGGCGATCATCCAGGA-3') with Mul1-R (5'-TAGGTGCTGGAAGTACGCGTCCAGTTACATG-3'). The mammalian expression vectors, FLAG/HA-tagged AGO1 (Addgene plasmid #10820), AGO2 (Addgene plasmid #10822), AGO3 (Addgene plasmid #10823), and AGO4 (Addgene plasmid #10824) were gifts from Dr. Thomas Tuschl (Meister *et al*, 2004).

Human subjects and GCT tissues

Formalin-fixed paraffin-embedded (FFPE) block sections of AGCT ($n = 46$) specimens from patients who visited the Seoul Asan Medical Center and the Bundang CHA Women's Hospital were analyzed. The present study was reviewed and approved by the Seoul Asan Medical Center and the Bundang CHA Women's Hospital Institutional Review Board. Informed consent was obtained from all subjects who participated in this study. This study was carried out in compliance with approved guidelines.

gDNA and RNA extraction

Total RNA from FFPE sections was extracted using the PureLink™ FFPE Total RNA Isolation kit (Invitrogen, Carlsbad, CA, USA), following the manufacturer's protocol. Total RNA in cultured cells was isolated using the TRIzol Reagent (Invitrogen), according to the manufacturer's instructions. gDNA was extracted from the deparaffinized tissues and cells using an Intron G-DEX™ Genomic DNA Extraction Kit (Intron, Seongnam, Korea), according to the manufacturer's protocol.

RNA library construction and ultra-deep RNA sequencing

Total RNA concentrations were calculated using Quant-IT RiboGreen (Invitrogen). To determine the % of RNA fragments > 200 bp in length, samples are run on the TapeStation RNA ScreenTape (Agilent Technologies, Waldbronn, Germany). A sequencing library was constructed using 100 ng of total RNA and a TruSeq RNA Access Library Prep Kit (Illumina, Inc., San Diego, CA, USA), according to the manufacturer's protocols. Briefly, total RNA was first fragmented into small pieces using divalent cations, at an elevated temperature. The cleaved RNA fragments were copied into first strand cDNA using SuperScript II reverse transcriptase (Invitrogen, cat#18064014) and random primers. Second-strand cDNA synthesis was performed using DNA Polymerase I, RNase H, and dUTP. The cDNA fragments were subjected to an end-repair process, involving the addition of a single "A" base and then adapter ligation. The products were then purified and enriched by 15 cycles of PCR to create the cDNA library. All libraries were normalized, and six libraries were pooled into a single hybridization/capture reaction. The pooled libraries were incubated with a cocktail of biotinylated oligonucleotides corresponding to coding regions of the genome. Targeted library molecules were captured via hybridized

biotinylated oligonucleotide probes using streptavidin-conjugated beads. After two rounds of hybridization/capture reactions, the enriched library molecules were subjected to a second round of PCR amplification in 10 cycles. The captured libraries were quantified using KAPA Library Quantification Kits for Illumina Sequencing platforms according to the qPCR Quantification Protocol Guide (Kapa Biosystems, catalog number KK4854) and qualified using the TapeStation D1000 ScreenTape assay (Agilent Technologies, catalog number 5067-5582). The indexed libraries were then loaded into an Illumina HiSeq 2500 instrument (Illumina, Inc.), and paired-end (2 × 100 bp) sequencing was performed by Macrogen, Inc. (Seoul, Korea). We generated, on average, 134,312,269 total sequence reads and approximately 71,531 Mbp of total mapped base.

Pyrosequencing

The PyroMark PCR Kit (Qiagen, Hilden, Germany) was used for pyrosequencing with a forward primer (Pyro-FOXL2-F: 5'-AGAAGGGCTGGCAAATAGCATC-3') and reverse biotinylated primer (Pyro-FOXL2-R: 5'-CCGGAAGGGCCTTTCAT-3'), or a reverse primer (Pyro-FOXL2-R2: 5'-TAGTTGCCCTTCTCGAACATGTC-3') and a forward biotinylated primer (Pyro-FOXL2-F2: 5'-CATCGGAAGTTCCCGTTCTA-3'). The PCR products were purified using streptavidin Sepharose HP beads (GE Healthcare, Buckinghamshire, UK), followed by hybridization with the sequencing primers (FOXL2-seqF: 5'-CGCAAGGGCAACTACT-3' or FOXL2-seqR2: 5'-CCTTCTCGAACATGTC-3'), as described in the PyroMark Q48 vacuum workstation guide (Qiagen). The sequencing data were analyzed using PyroMark Q48 software (Qiagen). The Pyrosequencing was performed and analyzed by Macrogen, Inc.

Allele-specific PCR analysis

Allele-specific PCRs were performed based on a previous study (Song *et al*, 2019). To amplify each *FOXL2* allele, allele-specific primers were designed with a 3' mismatch at the variable nucleotide at 402. The PCR primers used for *FOXL2* amplification were the FOXL2-279F primer (5'-GAATAAGAAGGGCTGGCAAAT-3'), the WT-specific reverse primer (5'-CCTTCTCGAACATGTCCTCG-3'), and the 402C>G-specific reverse primer (5'-CCTTCTCGAACATGTCCTCC-3'). For *GAPDH* amplification, the GAPDH-F (5'-AGGGGCCATCCACAGTCTT-3') and GAPDH-R (5'-AGCCAAAAGGGTCATCATCTCT-3') primers were used. PCR was performed in 20- μ l reactions containing 50 ng gDNA or cDNA with DMSO (10%). PCR was performed for 34 cycles using 2 annealing temperatures: 50°C for cDNA from FFPE sections and 60°C for cDNA from cell lines. SP-Taq DNA polymerase and 2X HOT MasterMix with Dye were purchased from Cosmo Genetech and MGmed (Seoul, Korea), respectively. The PCR products were electrophoresed on a 2% agarose gel and visualized under ultraviolet (UV) light.

Reverse transcription-quantitative real-time PCR (RT-qPCR) analysis

Following confirmation of the quality and quantity of extracted total RNA samples with a Nanodrop 2000 (Thermo Scientific, Wilmington, DE, USA), cDNA was synthesized using an iScript cDNA Synthesis Kit (Bio-Rad Laboratories, Hercules, CA, USA). Samples for RT-qPCR were prepared and analyzed as previously described

(Lee *et al*, 2019). The same primers described above for the allele-specific PCR analysis were used for RT-qPCR analysis of *FOXL2* and *GAPDH* expression. Additional PCR primers used for RT-qPCR analysis were as follows: Dicer-F (5'-TCTCTTTCCCAACTGGCATC-3'), Dicer-R (5'-GGTGGTTTCGTTTTGATTTGC-3'), ZEB2-F (5'-CAAGAGCGCAAACAAGC-3'), ZEB2-R (5'-GGTTGGCAATACCGTCAT-3'), BCL-2-F (5'-GTGGAGGAGCTCTCAGGA-3'), BCL-2-R (5'-AGGTGCGGTTCCAGTACTC-3'), p21-F (5'-TGTCCGTCAGAACCCATGC-3'), p21-R (5'-AAAGTCGAAGTCCATCGCT-3'), FIS-1-F (5'-ACCTGGCCGTGGGGAACC-3'), FIS-1-R (5'-AGTTCCTTGGCCTGGTTGTCTGG-3'), AFP-F (5'-CACGATCCAACCTGAGGCTGTCATTGC-3'), AFP-R (5'-CGGAATTCGATAAGGAAATCTCACATAAAAGTC-3'), ZEB1-F (5'-ATGCAGCTGACTGTGAAGT-3'), ZEB1-R (5'-GAAAATGCATCTGGTGTCC-3'), FOXL2-F (5'-AAGTACCTGCAGTCTGGCTT-3'), FOXL2-R (5'-CAGATGGTGTGCGTGGGAT-3'), p21-F (5'-TGTCCGTCAGAACCCATGC-3'), p21-R (5'-AAAGTCGAAGTCCATCGCTC-3'), PTEN-F (5'-AGTGGCGGAACCTGCAATCCTCAGT-3'), PTEN-R (5'-ACACAGGTACGGCTGAGGGAAC-3'), FLT4-F (5'-CTGGACCGAGTTTGTGGAGG-3'), FLT4-R (5'-GTCACATAGAAGTAGATGAGCCG-3'), HDAK3-F (5'-GTCGATGTTATTTCCCAGC-3'), HDAK3-R (5'-CCGATTTGGTGTGGGTGTT-3'), MAT2-F (5'-ATCATTACAGCCACCA-3'), MAT2-R (5'-CGATTATCAGATTCTCCCTC-3'), KLF8-F (5'-TCTGCAGGGACTACAGCAAG-3'), KLF8-R (5'-TCACATTGGTGAATCCGCTC-3'), SENP1-F (5'-CCCCATCATCACCCTCTGT-3'), SENP1-R (5'-CTTTGATCCACAGCTCTGCC-3'), DCAF1-F (5'-CTGCTATCTTGTGTCATTG-3'), DCAF1-R (5'-CTCTTCATCCTCATCCTCTG-3'), NELFE-F (5'-CAAGGTGGTCAAACGCTC-3'), and NELFE-R (5'-CTTCCCCTCAAGGGTTCGAG-3'). Gene expression levels were quantified using the $\Delta\Delta$ Ct method. All primers were purchased from Cosmo Genetech.

For miRNA analysis, ~10 ng of total RNA was reverse transcribed using the TaqMan® microRNA Reverse Transcription Kit (Applied Biosystems, Foster City, CA, USA) and the reverse-transcription primers from the TaqMan® microRNA Assay Kit (Applied Biosystems, *hsa-miR-1236* [Assay ID: 002761]; *hsa-miR-10b* [Assay ID: 002218]; *hsa-miR-132* [Assay ID: 002132]; *hsa-miR-204* [Assay ID: 000508]; *hsa-miR-210* [Assay ID: 467276_mat]; *hsa-miR-365* [Assay ID: 001020]; *hsa-miR-483* [Assay ID: 002560]; *RNU6B* [Assay ID: 001093]). RT-qPCR was performed in a CFX-96 Thermal Cycler and Detection System, using the TaqMan® microRNA Universal PCR Master Mix (Applied Biosystems) and TaqMan probes from the TaqMan® microRNA Assay Kit, according to the manufacturer's instructions. miRNA-expression levels were normalized to endogenous *RNU6B* expression.

RNA decay rate

To assess mRNA turnover, RNA synthesis was terminated by adding 5 μ g/ml ActD (Sigma-Aldrich) to the cell culture medium. At different time points (0, 0.5, 1, 2, 4, and 8 h) after ActD addition, the cells were harvested, and total RNA was isolated using the TRIzol reagent following the manufacturer's protocol. mRNA levels were determined by real-time PCR, as described above.

RNA interference

The target sequences of the siRNAs used in this study were as follows: siFOXL2 (5'-GGCAUCUACCAGUACAUCAdTdT-3'), siAGO1

(5'-GCACGGAAGUCCAUCUGAAUU-3' and 5'-GAGAAGAGGUGCUC AAGAAUU-3'), siAGO2 (5'-GCACGGAAGUCCAUCUGAAUU-3' and 5'-GCAGGACAAAGAUGUAUUUdTdT-3'), siAGO3 (5'-GCAUCAU UAUGCAAUAUGAAUU-3', 5'-GAAUUAGCAGAUUGGUAUU-3', and 5'-CAAGAUACCUUACGCACAAUU-3'), siAGO4 (5'-GGCCAGAACU AAUAGCAAUUU3' and 5'-GGCCAGAACUAAUAGCAAUUU-3'), siDH X9 (5'-CCAGGCAGAAATTCATGTGTG-3' and 5'-CAAUAUCUGUU AAUUGUdTdT-3'), siGW182 (5'-UGAUUGUUAGGCAUCUGGcdTd T-3' and 5'-GCCAGAUGCCUACAAUCA-3'), siADARBP1 (5'-GAUAG ACACCCAAAUCGUAdTdT-3'), siDDX20 (5'-GGAAUAAGUCAUACU UGG-3'), siFKBP5 (5'-GGAGCAACAGUAGAAUCCdTdT-3'), siGEMI N4 (5'-GGCACUGGCAGAAUUAACA-3'), siGPKOW (5'-CUGUGUAUG UCGGACAGAUdTdT-3'), siHSP90AA1 (5'-AACAUAGAAACUAAAA GCAUdTdT-3'), siHSP90B1 (5'-GAAGAAGCAUCUGAUUACcdTd T-3'), siLIG4 (5'-GCUAGAUGGUGAACGUAUG-3'), siMSI1 (5'-GGA GAAAGUGUGUAAAUUdTdT-3'), siNOTCH2 (5'-GGAGGUCUCAGU GGAUUAUdTdT-3'), siPAFAH1B1 (5'-UUUAGUCUCAGAUCUGUU GCUUCAdTdT-3'), siPPID (5'-GCAGGGAGCAAUUGACAG UdTdT-3'), and siYTHDF1 (5'-CCGCGUCUAGUUGUCAUGAA-3'). The sequence of the control siRNA was 5'-CCUACGCCACCAAUUUCGU-3'. All siRNAs were purchased from Bioneer (Seoul, Korea). Sense and anti-sense oligonucleotides were annealed in the presence of annealing buffer (Bioneer).

Immunoblot analysis

KGN cells (1×10^6) and COV434 cells (0.3×10^6) were transfected with the indicated plasmids or oligonucleotides using Lipofectamine 2000 (Invitrogen), according to the manufacturer's instructions. Cell lysates were prepared and subjected to sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) for subsequent immunoblotting with the respective antibodies. The protein signals on the membranes were detected using a ChemiDoc XRS + System Imager (Bio-Rad Laboratories), and the intensity of each band was quantified using Quantity One software (Bio-Rad Laboratories). For all immunoblot images presented in this manuscript, the membrane was sectioned according to the estimated molecular weights of the proteins of interest and probed with the indicated antibodies. All cropped blots were processed under the same experimental conditions. The following antibodies were used in this study: rabbit anti-FOXO2 (Park *et al*, 2010), rabbit anti-AGO1 (5053S; Cell Signaling Technology, Danvers, MA, USA), rat anti-AGO2 (11A9; Helmholtz Zentrum München, Germany), rabbit anti-AGO3 (5054S; Cell Signaling Technology), rabbit anti-AGO4 (6913S; Cell Signaling Technology), mouse anti-FLAG (F1804; Sigma-Aldrich), rabbit anti-DHX9 (ab26271; Abcam, Cambridge, MA, USA), rabbit anti-GW182 (NBP1-28751; Novus Biologicals, Littleton, CO, USA), mouse anti-Myc (2276S; Cell Signaling Technology), rabbit anti-GAPDH (sc-25778; Santa Cruz Biotechnology, Santa Cruz, CA, USA), and rabbit control-IgG (sc-2027; Santa Cruz Biotechnology).

miRNAs and antisense miRNAs

Sense-strand RNA oligonucleotides for miR-1236 (5'-CCUCUCCU CUUGUCUCUCCAG-3'), miR-1236-G (5'-CCUCUCCGUUGUCU CUCCAG-3'), miR-1236-U (5'-CCUCUCCUUGUCUCUCCAG-3'), and the non-specific control RNA (5'-CCUCGUGCCGUCCAU

CAGGUAGUU-3'; Song *et al*, 2013) were supplied by Genolution Pharmaceuticals, Inc. (Seoul, Korea). Antisense miRNAs for mature miR-1236 (5'-CTGGAGAGACAAGGGGAAGAGG-3'), miR-145 (5'-AG GGATTCCTGGGAAAAGTGGAC-3'), miR-204 (5'-AGGCATAGGATG ACAAAGGGAA-3'), miR-423-3p (5'-ACTGAGGGGCTCAGACCGAG CT-3'), and miR-484 (5'-ATCGGGAGGGGACTGAGCCTGA-3'), and a negative control miRNA (22-mer scrambled probe with a random sequence of 5'-GTGTAACACGTCTATACGCCCA-3') (Nielsen *et al*, 2011), with no known complementary sequence target among human transcripts, were purchased from Cosmo Genetech.

Full-length cDNA preparation using the CAGE method

Full-length cDNA was generated by the CAGE method (Takahashi *et al*, 2012). Briefly, 5 μ g of total RNA was prepared from KGN cells. cDNA was reverse transcribed using reverse transcriptase (Takara), biotinylated with Biotin (Long Arm) Hydrazide (Vector Laboratories, Burlingame, CA, USA), and cap-trapped to capture 5'-completed cDNAs using Streptavidin C1 Dynabeads (Invitrogen).

Genomic engineering of CRISPR/Cas9-nickase-mediated miR-1236 KO cell lines

miR-1236 KO cells were generated based on a protocol described by Ran *et al* (2013). Briefly, to generate vectors encoding the Cas9 (D10A) nickases, pSpCas9n(BB)-2A-GFP (Addgene, Cambridge, MA, USA) was mutagenized by PCR amplification. The primers used for site-specific mutagenesis were D10A-F (5'-TAGAGGTACCCGTTACA TAAC-3'), D10A-R (5'-CTGAAGATCTCTTGCAGATAG-3'), Mut-D10A-F (5'-TACAGCATCGGCCTGGCCATCGGC-3'), and Mut-D10A-R (5'-GGTGCCGATGGCCAGGCCGATGCT-3'). Four guide RNAs were designed targeting the protospacer adjacent motif (PAM) sequence at the miR-1236* region (sgRNA-miR-1236-1 and -2) or miR-1236 region (sgRNA-miR-1236-3 and -4) (Appendix Fig S4A and S4E), using CRISPR DESIGN (<http://crispr.mit.edu/>). To generate plasmids targeting miR-1236* or miR-1236, the pX458(D10A)-miR-1236-5p or pX458(D10A)-miR-1236-3p dual-guide oligonucleotide primers, respectively, were cloned into the pX458(D10A) vector. KGN or COV434 cells (1×10^6) were cultured in 100-mm dishes and transfected with 5 μ g of each CRISPR plasmid. The transfected cells were cultured for 24 h, harvested, and resuspended in phosphate-buffered saline (PBS; Ca²⁺/Mg²⁺-free, 1 mM EDTA, 25 mM HEPES pH 7.0), containing 1% heat-inactivated FBS. They were then sorted by flow cytometry (based on green fluorescent protein signaling), using a BD FACSAria II cell sorter (BD Bioscience, San Jose, CA, USA). Cells were sorted into individual wells of 96-well plates and then further expanded.

T7 endonuclease I (T7E1) assays

A gDNA fragment (762-bp) harboring miR-1236 was amplified with the primers, 5'-GATGGATGAAGCTTCCA-3' and 5'-ACTCAGAATGG TACAGC-3'. The purified PCR products were denatured and reannealed in NEBuffer 2 (New England BioLabs, Ipswich, MA, USA), using a thermocycler. The hybridized PCR products were digested with T7E1 (New England BioLabs) for 2 h and resolved on 2% agarose gels.

Genotyping for miR-1236 KO

Allelic deletion was confirmed using the TOPcloner™ TA Core Kit (Enzymomics, Daejeon, Korea) and DNA sequencing (Cosmo Genetech).

Northern blot analysis

Total RNA (including miRNAs) was extracted from cultured cells using TRIzol (Invitrogen). Then, the small RNA species were isolated and concentrated using the mirVana PARIS Kit (Ambion). Small RNA fractions from cells (~1.5 µg) were separated on a 15 % polyacrylamide gel containing 8 M urea, visualized by ethidium bromide staining, and then transferred to a Hybond-XL membrane (Amersham Bioscience). Subsequently, the RNA was immobilized on the membrane with an ultraviolet cross-linker (UVP) and hybridized with a 5' end-labeled-oligonucleotide. Anti-miR-1236 (5'-CTGGAGAGACAAGGGGAAGAGG-3') and anti-miR-21 (5'-TCAA CATCAGTCTGATAAGCTA-3') were used as probes. The radioactive signals were analyzed using a Bio-Rad phosphorimager and the Quantity One image-analysis software package (Bio-Rad).

Luciferase reporter assays

Luciferase assays were performed as described previously (Kim *et al*, 2011). Briefly, KGN cells (4×10^5) were plated in 12-well plates and were transfected with 1 µg of the reporter construct (pGL3c-CDS-*FOXL2s* or pGL3c-UTR-*FOXL2s*), 200 ng of pCMV-β-galactosidase (Clontech, Palo Alto, CA, USA), 50 nM of either control or mimic miR-1236, and 200 nM of the indicated siRNAs using the Neon transfection system (Invitrogen). After 48 h, luciferase activities were measured with the FlexStation3 Microplate Reader (Molecular Devices, Sunnyvale, CA, USA).

In vitro transcription

In vitro transcription of 230-bp CDS sequences that included nucleotide 402 of WT and 402C>G *FOXL2* mRNA was performed using the MegashortScript™ T7 Kit (Thermo Scientific, Rockford, IL, USA). To generate the T7-*FOXL2* DNA template, pCMV-*FOXL2*-WT and pCMV-*FOXL2*-C134W (Kim *et al*, 2011) were used as templates and amplified by PCR, with a primer containing the T7 promoter sequence, T7-*FOXL2*-F (5'-TAATACGACTCACTATAGGGAGGGCTGGCAAATAGCA-3') and T7-*FOXL2*-R (5'-GGCGCTCCGGCCCGCA-3'). RNA was transcribed *in vitro* using T7 RNA polymerase according to the manufacturer's instructions.

In vitro annealing kinetics between miR-1236 and FOXL2 transcripts

We incubated 5'-labeled ³²P-miR-1236 (0.5 nM; PerkinElmer, Boston, MA, USA) with increasing concentrations of *in vitro*-transcribed WT or 402C>G *FOXL2* mRNAs (0, 2.5, 12.5, 25, and 50 nM) in hybridization buffer (100 mM NaCl, 20 mM Tris-HCl at pH 7.4, and 10 mM MgCl₂) at 65°C for 5 min. Then, the hybridization reactions were cooled slowly to room temperature. Aliquots were transferred into 1 volume (v) of stop buffer (20 mM Tris-HCl at pH 7.4, 10 mM EDTA, 2% [v/v] SDS, 8 M

urea, 0.025% [v/v] bromophenol blue) and analyzed by 6% native PAGE. Band intensities reflecting *FOXL2*-miRNA complex formation were detected and quantified using a phosphorimager and the OptiQuant software package (Packard, Meriden, CT, USA).

In vivo association of FOXL2 mRNAs or miR-1236 with RISC

293T cells were cotransfected with miR-1236-G mimic, siDHX9, or the indicated plasmids for 48 h. The cells were washed with cold 1× PBS, and RNA-protein complexes were cross-linked by UV-A irradiation (0.15 J/cm²). In each case, 10% of the cells was removed and kept as the input samples, and remaining cells were lysed in buffer containing 25 mM Tris-HCl at pH 7.4, 150 mM KCl, 0.5% NP-40, 2 mM EDTA, 1 mM NaF, 0.5 mM DTT, RNase inhibitor (Promega), 0.05% Tween 20, and protease inhibitors (GeneDEPOT, Barker, TX, USA), and centrifuged at 7,000 × g for 15 min at 4°C. Each lysate was pre-incubated with 2 µg IgG and 20 µl Dynabeads (Invitrogen) for 4 h at 4°C. Three milligrams of each lysate was precleared using a control rabbit IgG-Dynabeads complex for 30 min at 4°C. Then, the cell lysates were re-incubated with 5 µg FLAG antibody-coupled Dynabeads (50 µl) overnight on a rotator at 4°C. After incubation, the beads were washed five times with a buffer (300 mM KCl, 50 mM Tris-HCl pH 7.4, 1 mM MgCl₂, 0.1% NP-40, 0.05% Tween 20) and incubated with DNase I for 10 min at 37°C. The sample proteins were digested with proteinase K, and RNA was extracted using an acidic phenol: chloroform mixture (5: 1, pH 4.3) and precipitated with isopropanol in the presence of 10% of 3 M sodium acetate (NaOAc; pH 5.2). The purified RNAs were reverse transcribed to cDNA with the *FOXL2*-430-R primer (5'-GGTAGTTGCCCTTCTCGAAC-3'), which binds to a region downstream of the 402C>G site. Then, the cDNA products were used for *FOXL2* allele-specific RT-PCR or real-time RT-PCR analysis, with the *FOXL2*-279F primer. The relative abundance of miR-1236 was validated using TaqMan® microRNA assays. To detect endogenous association of AGO3-mediated RISCs and *FOXL2* mRNA, RNA-protein complexes formed in KGN cells (3×10^7) transfected with a control miRNA or miR-1236-G (10 nM), or a control siRNA or siDHX9 (200 nM), were cross-linked via UV irradiation (0.15 J/cm²). The lysates were incubated with 10 µl of AGO3 antibody-coupled Dynabeads (50 µl) overnight on a rotator at 4°C. The RNAs were purified as described above.

RISC capture using immobilized 2'-O-methyl oligonucleotides

RISC capture experiment was performed according to a previous study (Hutvagner *et al*, 2004). using biotin-tagged 2'-O-methyl oligonucleotides complementary to miR-1236. Biotin-tagged 2'-O-methyl oligonucleotides (100 pmol) complementary to miR-1236 (5'-bio-CUGGACGCUGGAGAGACAAGGGGAAGAGGUGUUCGA-3') and control (5'-bio-CUGGACGGUGUACACGUCUAUACGCCCAU GUUCGA-3') were incubated for 1 h at 4°C in PBST (PBS with 0.1% Tween-20) buffer containing 2.5% bovine serum albumin with 50 µl of streptavidin-coupled Dynabeads M280 (Invitrogen) to immobilize the oligonucleotides on the beads. Cell lysates (3 mg) were nutated with the immobilized 2'-O-methyl oligonucleotides for 1 h at room temperature. The bound miRISC was

analyzed using immunoblotting with the indicated antibodies. Relative binding was determined using ratios of protein from the input compared to bound RISC. The unbound supernatant was collected, and the total RNA was extracted to determine the capture efficiency of the mature miR-1236 using the TaqMan[®] microRNA assay.

Cell-viability assay

Cell-viability assays were performed as previously described (Kim *et al*, 2014).

5-bromo-2'-deoxyuridine cell-proliferation assay

Cell (1×10^4) proliferation was measured using the 5-Bromo-2'-deoxyuridine Labeling and Detection Kit III (Roche, Mannheim, Germany; Jin *et al*, 2016).

Cell-migration assay

A chemotaxis migration assay was performed using Transwell Permeable Supports (8- μ m pore size, 6.5-mm insert; Corning-Costar, Lowell, MA, USA). KGN cells (1×10^6) and COV434 cells (0.3×10^6) were transfected with either pcDNA3 or pcDNA3-Flag-FOXL2 (Park *et al*, 2010) or the indicated oligonucleotides. Cells were plated at a density of 2×10^5 cells/well in the top chamber containing DMEM/F12 and 1% FBS, whereas the bottom chamber contained DMEM/F12 and 10% FBS as a chemoattractant. After 2 to 6 h of incubation, the cells and medium were removed from the top chamber, and the migrated cells were fixed with 5% glutaraldehyde for 20 min and stained with 0.5% crystal violet for 20 min. Images of the migrated cells were taken at $\times 100$ magnification under a bright-field microscope. Quantification was performed under a light microscope with $\times 200$ magnification by counting the number of migrated cells in five random fields per chamber. The data are presented as the mean \pm SEM of three independent experiments and are shown as the fold change in the average number of cells.

Apoptosis and cell cycle analyses

Detection of Annexin V-positive apoptotic cells and cell cycle analysis was performed as previously reported (Kim *et al*, 2011; Jin *et al*, 2016).

RNA sequencing and transcriptome analysis of KGN cells

Next-generation RNA sequencing was performed using KGN cells (1×10^6) transfected with control oligonucleotides or anti-miR-1236 (50 nM) for 48 h with two biological replicates. Total RNA concentration was calculated by Quant-IT RiboGreen (Invitrogen, #R11490). To assess the integrity of the total RNA, samples are run on the TapeStation RNA screentape (Agilent Technologies). High-quality RNAs prepared, with RIN > 7.0, were used for RNA library construction. A library was independently prepared with 1 μ g of total RNA for each sample by Illumina TruSeq Stranded mRNA Sample Prep Kit (Illumina, #RS-122-2101). The first step

in the workflow involves purifying the poly-A containing mRNA molecules using poly-T-attached magnetic beads. Following purification, the mRNA is fragmented into small pieces using divalent cations under elevated temperature. The cleaved RNA fragments are copied into first-strand cDNA using SuperScript II reverse transcriptase (Invitrogen, #18064014) and random primers. This is followed by second-strand cDNA synthesis using DNA Polymerase I, RNase H and dUTP. These cDNA fragments then go through an end-repair process, the addition of a single "A" base, and then ligation of the adapters. The products are then purified and enriched with PCR to create the final cDNA library. The libraries were quantified using KAPA Library Quantification kits for Illumina Sequencing platforms according to the qPCR Quantification Protocol Guide (Kapa Biosystems, #KK4854) and qualified using the TapeStation D1000 ScreenTape (Agilent Technologies, # 5067-5582). Indexed libraries were then submitted to an Illumina NovaSeq (Illumina), and the paired-end (2×100 bp) sequencing was performed. Average sequencing depth across samples was 73.3 million reads. Data normalization and analyses were performed based on a previous study (Sindi *et al*, 2020). Genes were considered differentially expressed if the adjusted *P* value was < 0.05 and there was at least a 1.5-fold change in expression. GO (Gene Ontology) enrichment analysis was performed using DAVID v6.8 (<https://david.ncifcrf.gov/>), where significant enrichment was determined if the adjusted *P* value was < 0.05. Heat map displays were analyzed using MultiExperiment Viewer (MeV) (<https://sourceforge.net/projects/mev-tm4/>).

Tumor xenograft establishment and immunohistochemistry

Control or miR-1236-KO (miR-1236^{-/-}) KGN cells (5×10^6) were suspended in 0.1 ml Matrigel (1: 1, v/v, Corning, Tewksbury, MA) and injected subcutaneously into 7-week-old BALB/c nu/nu immunodeficient mice (weights ranged between 18 and 20 g) (ORIENT BIO, Seongnam, Korea). Mice were euthanized 2 months later, and the numbers of intraperitoneal nodules were counted. Half of the nodules were excised, fixed in 4% neutral buffered formalin, and embedded in paraffin before preparing 5- μ m sections that were stained with hematoxylin. The other half of the nodules were excised, and total RNA was isolated using the TRIzol reagent (Invitrogen), according to the manufacturer's instructions. The animal guidelines were approved by the Chung-Ang University Institutional Animal Care and Use Committee (IRB# CAU2012-0044), and the animals were treated as described in the protocol.

Statistical analysis

All details of the statistical analysis used in the experiments are included in the figure legends. Multiple-comparison analyses of values were performed using the Student–Newman–Keuls test, and Student's *t*-test was used for comparisons with control samples, using SAS version 9.2 (SAS Institute, Cary, NC, USA) and SigmaPlot (Systat Software, San Jose, CA, USA). The data are presented as mean \pm SEM, and *P* < 0.05 was considered to reflect as statistically significant difference.

Data Availability

Ultra-deep RNA Sequencing data have been deposited in the NCBI Sequence Read Archive (SRA) with the BioProject accession numbers of PRJNA555182 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA555182>) and PRJNA648830 (<https://www.ncbi.nlm.nih.gov/bioproject/PRJNA648830>).

Expanded View for this article is available online.

Acknowledgements

This research was supported by the National Research Foundation of Korea (NRF), which is funded by the Ministry of Science, ICT, and Future Planning (2019R1H1A2080015, 2019R1A2C2084563, NRF-2017R1A2B2011248, NRF-2015R1A5A1008958, and NRF-2018R1A5A1025077).

Author contributions

Conception and design: KL and JB Development of methodology: ES, HJ, SH, KL, and JB. Performance of experiments: ES, HJ, D-SS, YL, H-JH, and THK. Analysis and interpretation of data: YH, SH, KL, and JB. Writing and review of the manuscript: KL and JB Study supervision: KL and JB.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Azuma-Mukai A, Oguri H, Mituyama T, Qian ZR, Asai K, Siomi H, Siomi MC (2008) Characterization of endogenous human Argonautes and their miRNA partners in RNA silencing. *Proc Natl Acad Sci USA* 105: 7964–7969
- Bartel DP (2009) MicroRNAs: target recognition and regulatory functions. *Cell* 136: 215–233
- Broughton JP, Lovci MT, Huang JL, Yeo GW, Pasquinelli AE (2016) Pairing beyond the Seed Supports MicroRNA Targeting Specificity. *Mol Cell* 64: 320–333
- Cai Z, Cao R, Zhang K, Xue Y, Zhang C, Zhou Y, Zhou J, Sun H, Fu XD (2015) Oncogenic miR-17/20a Forms a Positive Feed-forward Loop with the p53 Kinase DAPK3 to Promote Tumorigenesis. *J Biol Chem* 290: 19967–19975
- Chen SY, Teng SC, Cheng TH, Wu KJ (2016) miR-1236 regulates hypoxia-induced epithelial-mesenchymal transition and cell migration/invasion through repressing SENP1 and HDAC3. *Cancer Lett* 378: 59–67
- Chi SW, Zang JB, Mele A, Darnell RB (2009) Argonaute HITS-CLIP decodes microRNA-mRNA interaction maps. *Nature* 460: 479–486
- Crisponi L, Deiana M, Loi A, Chiappe F, Uda M, Amati P, Bisceglia L, Zelante L, Nagaraja R, Porcu S et al (2001) The putative forkhead transcription factor FOXL2 is mutated in blepharophimosis/ptosis/epicanthus inversus syndrome. *Nat Genet* 27: 159–166
- Czech B, Hannon GJ (2011) Small RNA sorting: matchmaking for Argonautes. *Nat Rev Genet* 12: 19–31
- Dai A, Sun H, Fang T, Zhang Q, Wu S, Jiang Y, Ding L, Yan G, Hu Y (2013) MicroRNA-133b stimulates ovarian estradiol synthesis by targeting Foxl2. *FEBS Lett* 587: 2474–2482
- Duursma AM, Kedde M, Schrier M, le Sage C, Agami R (2008) miR-148 targets human DNMT3b protein coding region. *RNA* 14: 872–877
- Elcheva I, Goswami S, Noubissi FK, Spiegelman VS (2009) CRD-BP protects the coding region of betaTRCP1 mRNA from miR-183-mediated degradation. *Mol Cell* 35: 240–246
- Eulalio A, Tritschler F, Izaurralde E (2009) The GW182 protein family in animal cells: new insights into domains required for miRNA-mediated gene silencing. *RNA* 15: 1433–1442
- Feng W, Gong H, Wang Y, Zhu G, Xue T, Wang Y, Cui G (2019) circIFT80 functions as a ceRNA of miR-1236-3p to promote colorectal cancer progression. *Mol Ther Nucleic Acids* 18: 375–387
- Forman JJ, Legesse-Miller A, Collier HA (2008) A search for conserved sequences in coding regions reveals that the let-7 microRNA targets Dicer within its coding sequence. *Proc Natl Acad Sci USA* 105: 14879–14884
- Fu GK, Xu W, Wilhelmy J, Mindrinos MN, Davis RW, Xiao W, Fodor SP (2014) Molecular indexing enables quantitative targeted RNA sequencing and reveals poor efficiencies in standard library preparations. *Proc Natl Acad Sci USA* 111: 1891–1896
- Fu Q, Yuan YA (2013) Structural insights into RISC assembly facilitated by dsRNA-binding domains of human RNA helicase A (DHX9). *Nucleic Acids Res* 41: 3457–3470
- Gao R, Cai C, Gan J, Yang X, Shuang Z, Liu M, Li S, Tang H (2015) miR-1236 down-regulates alpha-fetoprotein, thus causing PTEN accumulation, which inhibits the PI3K/Akt pathway and malignant phenotype in hepatoma cells. *Oncotarget* 6: 6014–6028
- Gu S, Jin L, Zhang F, Sarnow P, Kay MA (2009) Biological basis for restriction of microRNA targets to the 3' untranslated region in mammalian mRNAs. *Nat Struct Mol Biol* 16: 144–150
- Hafner M, Landthaler M, Burger L, Khorshid M, Hausser J, Berninger P, Rothballer A, Ascano Jr M, Jungkamp AC, Munschauer M et al (2010) Transcriptome-wide identification of RNA-binding protein and microRNA target sites by PAR-CLIP. *Cell* 141: 129–141
- Hausser J, Syed AP, Bilén B, Zavolan M (2013) Analysis of CDS-located miRNA target sites suggests that they can effectively inhibit translation. *Genome Res* 23: 604–615
- Hock J, Meister G (2008) The Argonaute protein family. *Genome Biol* 9: 210
- Huang H, Li Y, Szulwach KE, Zhang G, Jin P, Chen D (2014) AGO3 Slicer activity regulates mitochondria-nuage localization of Armitage and piRNA amplification. *J Cell Biol* 206: 217–230
- Hutvagner G, Simard MJ, Mello CC, Zamore PD (2004) Sequence-specific inhibition of small RNA function. *PLoS Biol* 2: E98
- Iwakawa HO, Tomari Y (2013) Molecular insights into microRNA-mediated translational repression in plants. *Mol Cell* 52: 591–601
- Jamieson S, Butzow R, Andersson N, Alexiadis M, Unkila-Kallio L, Heikinheimo M, Fuller PJ, Anttonen M (2010) The FOXL2 C134W mutation is characteristic of adult granulosa cell tumors of the ovary. *Mod Pathol* 23: 1477–1485
- Jin H, Lee B, Luo Y, Choi Y, Choi EH, Jin H, Kim KB, Seo SB, Kim YH, Lee HH et al (2020) FOXL2 directs DNA double-strand break repair pathways by differentially interacting with Ku. *Nat Commun* 11: 2010
- Jin H, Lee K, Kim YH, Oh HK, Maeng YI, Kim TH, Suh DS, Bae J (2016) Scaffold protein FHL2 facilitates MDM2-mediated degradation of IER3 to regulate proliferation of cervical cancer cells. *Oncogene* 35: 5106–5118
- Kertesz M, Iovino N, Unnerstall U, Gaul U, Segal E (2007) The role of site accessibility in microRNA target recognition. *Nat Genet* 39: 1278–1284
- Kim H, Kim J, Kim K, Chang H, You K, Kim VN (2019) Bias-minimized quantification of microRNA reveals widespread alternative processing and 3' end modification. *Nucleic Acids Res* 47: 2630–2640
- Kim JH, Bae J (2014) Differential apoptotic and proliferative activities of wild-type FOXL2 and blepharophimosis-ptosis-epicanthus inversus syndrome (BPES)-associated mutant FOXL2 proteins. *J Reprod Dev* 60: 14–20

- Kim JH, Kim YH, Kim HM, Park HO, Ha NC, Kim TH, Park M, Lee K, Bae J (2014) FOXL2 posttranslational modifications mediated by GSK3beta determine the growth of granulosa cell tumours. *Nat Commun* 5: 2936
- Kim JH, Yoon S, Park M, Park HO, Ko JJ, Lee K, Bae J (2011) Differential apoptotic activities of wild-type FOXL2 and the adult-type granulosa cell tumor-associated mutant FOXL2 (C134W). *Oncogene* 30: 1653–1663
- Landthaler M, Gaidatzis D, Rothballer A, Chen PY, Soll SJ, Dinic L, Ojo T, Hafner M, Zavolan M, Tuschl T (2008) Molecular characterization of human Argonaute-containing ribonucleoprotein complexes and their bound target mRNAs. *RNA* 14: 2580–2596
- Lee J, Lee DH, Jeon CO, Lee K (2019) RNase G controls tpiA mRNA abundance in response to oxygen availability in *Escherichia coli*. *J Microbiol* 57: 910–917
- Li QH, Liu Y, Chen S, Zong ZH, Du YP, Sheng XJ, Zhao Y (2019) circ-CSPP1 promotes proliferation, invasion and migration of ovarian cancer cells by acting as a miR-1236-3p sponge. *Biomed Pharmacother* 114: 108832
- Liu J, Carmell MA, Rivas FV, Marsden CG, Thomson JM, Song JJ, Hammond SM, Joshua-Tor L, Hannon GJ (2004) Argonaute2 is the catalytic engine of mammalian RNAi. *Science* 305: 1437–1441
- Luo Y, Wu X, Ling Z, Yuan L, Cheng Y, Chen J, Xiang C (2015) microRNA133a targets Foxl2 and promotes differentiation of C2C12 into myogenic progenitor cells. *DNA Cell Biol* 34: 29–36
- Mayr C, Bartel DP (2009) Widespread shortening of 3'UTRs by alternative cleavage and polyadenylation activates oncogenes in cancer cells. *Cell* 138: 673–684
- Mayr C, Hemann MT, Bartel DP (2007) Disrupting the pairing between let-7 and Hmga2 enhances oncogenic transformation. *Science* 315: 1576–1579
- Meister G, Landthaler M, Patkaniowska A, Dorsett Y, Teng G, Tuschl T (2004) Human Argonaute2 mediates RNA cleavage targeted by miRNAs and siRNAs. *Mol Cell* 15: 185–197
- Nanda GG, Kumar MV, Pradhan L, Padhy B, Sundaray S, Das S, Alone DP (2018) rs4246215 is targeted by hsa-miR1236 to regulate FEN1 expression but is not associated with Fuchs' endothelial corneal dystrophy. *PLoS One* 13: e0204278
- Nielsen BS, Jorgensen S, Fog JU, Sokilde R, Christensen IJ, Hansen U, Brunner N, Baker A, Moller S, Nielsen HJ (2011) High levels of microRNA-21 in the stroma of colorectal cancers predict short disease-free survival in stage II colon cancer patients. *Clin Exp Metastasis* 28: 27–38
- Okamura K, Hagen JW, Duan H, Tyler DM, Lai EC (2007) The mirtron pathway generates microRNA-class regulatory RNAs in *Drosophila*. *Cell* 130: 89–100
- Park M, Shin E, Won M, Kim JH, Go H, Kim HL, Ko JJ, Lee K, Bae J (2010) FOXL2 interacts with steroidogenic factor-1 (SF-1) and represses SF-1-induced CYP17 transcription in granulosa cells. *Mol Endocrinol* 24: 1024–1036
- Park MS, Phan HD, Busch F, Hinckley SH, Brackbill JA, Wysocki VH, Nakanishi K (2017) Human Argonaute3 has slicer activity. *Nucleic Acids Res* 45: 11867–11877
- Ran FA, Hsu PD, Lin CY, Gootenberg JS, Konermann S, Trevino AE, Scott DA, Inoue A, Matoba S, Zhang Y et al (2013) Double nicking by RNA-guided CRISPR Cas9 for enhanced genome editing specificity. *Cell* 154: 1380–1389
- Robb GB, Rana TM (2007) RNA helicase A interacts with RISC in human cells and functions in RISC loading. *Mol Cell* 26: 523–537
- Ruby JG, Jan CH, Bartel DP (2007) Intronic microRNA precursors that bypass Drosha processing. *Nature* 448: 83–86
- Sasaki T, Shiohama A, Minoshima S, Shimizu N (2003) Identification of eight members of the Argonaute family in the human genome. *Genomics* 82: 323–330
- Schmidt D, Ovitt CE, Anlag K, Fehsenfeld S, Gredsted L, Treier AC, Treier M (2004) The murine winged-helix transcription factor Foxl2 is required for granulosa cell differentiation and ovary maintenance. *Development* 131: 933–942
- Schnall-Levin M, Rissland OS, Johnston WK, Perrimon N, Bartel DP, Berger B (2011) Unusually effective microRNA targeting within repeat-rich coding regions of mammalian mRNAs. *Genome Res* 21: 1395–1403
- Schnall-Levin M, Zhao Y, Perrimon N, Berger B (2010) Conserved microRNA targeting in *Drosophila* is as widespread in coding regions as in 3'UTRs. *Proc Natl Acad Sci USA* 107: 15751–15756
- Schumer ST, Cannistra SA (2003) Granulosa cell tumor of the ovary. *J Clin Oncol* 21: 1180–1189
- Shah SP, Kobel M, Senz J, Morin RD, Clarke BA, Wiegand KC, Leung G, Zayed A, Mehl E, Kalloger SE et al (2009) Mutation of FOXL2 in granulosa-cell tumors of the ovary. *N Engl J Med* 360: 2719–2729
- Sindi HA, Russomanno G, Satta S, Abdul-Salam VB, Jo KB, Qazi-Chaudhry B, Ainscough AJ, Szulcek R, Jan Bogaard H, Morgan CC et al (2020) Therapeutic potential of KLF2-induced exosomal microRNAs in pulmonary hypertension. *Nat Commun* 11: 1185
- Song NJ, Yoon HJ, Kim KH, Jung SR, Jang WS, Seo CR, Lee YM, Kweon DH, Hong JW, Lee JS et al (2013) Butein is a novel anti-adipogenic compound. *J Lipid Res* 54: 1385–1396
- Song W, Joo M, Yeom JH, Shin E, Lee M, Choi HK, Hwang J, Kim YI, Seo R, Lee JE et al (2019) Divergent rRNAs as regulators of gene expression at the ribosome level. *Nat Microbiol* 4: 515–526
- Su H, Trombly MI, Chen J, Wang X (2009) Essential and overlapping functions for mammalian Argonautes in microRNA silencing. *Genes Dev* 23: 304–317
- Takahashi H, Lassmann T, Murata M, Carninci P (2012) 5' end-centered expression profiling using cap-analysis gene expression and next-generation sequencing. *Nat Protoc* 7: 542–561
- Tay Y, Zhang J, Thomson AM, Lim B, Rigoutsos I (2008) MicroRNAs to Nanog, Oct4 and Sox2 coding regions modulate embryonic stem cell differentiation. *Nature* 455: 1124–1128
- Tian B, Manley JL (2017) Alternative polyadenylation of mRNA precursors. *Nat Rev Mol Cell Biol* 18: 18–30
- Uhlenhaut NH, Jakob S, Anlag K, Eisenberger T, Sekido R, Kress J, Treier AC, Klugmann C, Klasen C, Holter NI et al (2009) Somatic sex reprogramming of adult ovaries to testes by FOXL2 ablation. *Cell* 139: 1130–1142
- Wang C, Tang K, Li Z, Chen Z, Xu H, Ye Z (2016) Targeted p21(WAF1/CIP1) activation by miR-1236 inhibits cell proliferation and correlates with favorable survival in renal cell carcinoma. *Urol Oncol* 34: e23–e34
- Wang H, Guo M, Ding D, Yang F, Chen Z (2020) Long non-coding RNA NNT-AS1 contributes to cisplatin resistance via miR-1236-3p/ATG7 axis in lung cancer cells. *Oncotargets Ther* 13: 3641–3652
- Wang K, Long B, Jiao JQ, Wang JX, Liu JP, Li Q, Li PF (2012) miR-484 regulates mitochondrial network through targeting Fis1. *Nat Commun* 3: 781
- Wang T, Li F, Tang S (2015) MiR-30a upregulates BCL2A1, IER3 and cyclin D2 expression by targeting FOXL2. *Oncol Lett* 9: 967–971
- Wang Y, Yan S, Liu X, Zhang W, Li Y, Dong R, Zhang Q, Yang Q, Yuan C, Shen K et al (2014) miR-1236-3p represses the cell migration and invasion abilities by targeting ZEB1 in high-grade serous ovarian carcinoma. *Oncol Rep* 31: 1905–1910
- Wen J, Ladewig E, Shenker S, Mohammed J, Lai EC (2015) Analysis of nearly one thousand mammalian mirtrons reveals novel features of dicer substrates. *PLoS Comput Biol* 11: e1004441

Xue Y, Ouyang K, Huang J, Zhou Y, Ouyang H, Li H, Wang G, Wu Q, Wei C, Bi Y et al (2013) Direct conversion of fibroblasts to neurons by reprogramming PTB-regulated microRNA circuits. *Cell* 152: 82–96

Zhang K, Zhang X, Cai Z, Zhou J, Cao R, Zhao Y, Chen Z, Wang D, Ruan W, Zhao Q et al (2018) A novel class of microRNA-recognition elements that function only within open reading frames. *Nat Struct Mol Biol* 25: 1019–1027

Zhao X, Feng J, Zhang L, Zhao F, Li M, Du Y, Li Y, Wu Q, Li G (2019) One functional variant in the 3'-untranslated region of TLR4 is associated with the elevated risk of ventilator-associated pneumonia in the patients with chronic obstructive pulmonary disease. *J Cell Physiol* 234: 18879–18886

Zuker M (2003) Mfold web server for nucleic acid folding and hybridization prediction. *Nucleic Acids Res* 31: 3406–3415



License: This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDeriv 4.0 License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.