



HHS Public Access

Author manuscript

Curr Opin Psychiatry. Author manuscript; available in PMC 2022 January 01.

Published in final edited form as:

Curr Opin Psychiatry. 2021 January ; 34(1): 29–35. doi:10.1097/YCO.0000000000000665.

Addressing the Psychological Resilience During the Covid-19 Pandemic: A Rapid Review

Judite Blanc¹, Anthony Q. Briggs², Azizi A. Seixas¹, Marvin Reid³, Girardin Jean-Louis¹, Seithikurippu R. Pandi Perumal^{4,CA}

¹Department of Population Health, NYU Grossman School of Medicine, Center for Healthful Behavior Change, New York, USA.

²Department of Psychiatry, NYU Grossman School of Medicine, New York, USA.

³Tropical Metabolism Research Unit, Tropical Medicine Research Institute, the University of the West Indies, Kingston, Jamaica

⁴Somnogen Canada Inc., College Street, Toronto, ON, Canada.

Abstract

Purpose of Review: The mental health toll on populations exposed to COVID-19 is alarming and there is an urgent need to address with urgency. This current review provides insights on how individuals, communities, and specific populations such as healthcare workers and patients are leveraging pre and peri-COVID-19 factors to reinforce their psychological resilience during the global public health crisis.

Recent Findings: Examination of the extant literature indicated that populations around the world rely often on support from their loved-ones, closed significant others, outdoor and physical activities, and spirituality to cope with the COVID-19-related distress. Increased sense of meaning/purpose since the COVID-19 pandemic was also reported.

Summary: A portion of publications provided intervention models to reinforce resilience among specific populations during the COVID-19 pandemic. Nevertheless, it's not convincing that some

^{CA}**Corresponding author:** S.R. Pandi-Perumal, Somnogen Canada Inc., College Street, Toronto, ON, Canada, Pandiperumal2020@gmail.com.

Author's contributions: All authors met the ICMJE criteria for authorship. All authors contributed equally in the subsequent preparation of the manuscript. Conceived and designed the manuscript: GJL, SRP, JB; Wrote the initial draft: JB, ABQ, AAS; critically reviewed the manuscript: MR, GJL, SRP. Prior to submission, all authors (JB, ABQ, AAS, MR, GJL, and SRP) reviewed and accepted the final version of the manuscript.

Ethical approval: Not applicable.

Compliance with ethical standards: This article does not contain any studies with human participants or animals performed by any of the authors. No primary data have been reported.

Declaration of Competing Interests: The authors have read the journal's policy and have the following potential conflicts: SRP is a stockholder and the President and Chief Executive Officer of Somnogen Canada Inc., a Canadian Corporation. He declares that he has no competing interests that might be perceived to influence the content of this article. This does not alter the authors' adherence to all the journal policies.

All remaining authors declare that they have no proprietary, financial, professional, nor any other personal interest of any nature or kind in any product or services and/or company that could be construed or considered to be a potential conflict of interest that might have influenced the views expressed in this manuscript.

of these models can be applied universally. Additionally, it is important to note that in this category translational data was scarce.

Keywords

COVID-19; Infectious Disease; Mental health; Outbreak; Pandemic; Resilience

Introduction

1. Mental Health consequence of COVID

On March 11, the World Health Organization (WHO) declared the Novel Coronavirus (COVID-19) outbreak as a global pandemic. According to early October 2020 WHO's updates, a new surge of COVID-19 that has now spread to over 235 areas, countries, and territories, with over thirty-five million individuals contracting the virus. This has resulted in over 1 million deaths across the globe¹ and many more suffering from the economic, physical, and mental health consequences of its aftermath.¹ It is important to note that this COVID-19 translates to a broad range of somatic conditions ranging from fever or chills, cough, shortness of breath or difficulty breathing, fatigue, muscle or body aches, headache, the new loss of taste or smell, sore throat, congestion or runny nose, nausea or vomiting, diarrhea.^{2,3,6}

Studies reviewed by Vindegaard and Benros (2020) revealed that the vast majority of participants infected with COVID-19 are at high risk for post-traumatic stress (PTSS) (96.2%) and depressive symptoms. Individuals with preexisting psychiatric conditions reported a deterioration of psychiatric symptoms: 37.5% with an eating disorder and 56.2% reported additional anxiety symptoms.³ In the general population, the situation was no different, as studies found that individuals without a mental health history exhibited lower psychological well-being and greater levels of anxiety and depression in comparison to the period that preceded the COVID-19 pandemic.^{3,4} The burden of mental health was quite stark in the U.S, as the prevalence of depression symptoms in the U.S. was more than 3-fold higher during COVID-19 compared to the period before the COVID-19 pandemic.⁵⁻⁹ One group that appeared to be most affected was healthcare workers. According to Vindegaard and Benros (2020), frontline healthcare workers who treated COVID-19 patients were at increased risk of suffering from depression/depressive symptoms, anxiety, psychological distress, and poor sleep quality. During the peak of the pandemic in April 2020, 57% of New York healthcare workers surveyed suffered from acute stress, 48% and 33% endorsed depressive, and anxiety symptomatology respectively. The vast majority (61%) reported an increased sense of meaning/purpose since the COVID-19 pandemic.⁴ Furthermore, physical activity or exercise was the most frequent coping strategy (59%) besides access to an individual therapist with online self-guided counseling (33%) (Shechter et al. 2020). Moreover, U.S participants with lower socioeconomic status (SES), and greater exposure to stressors (e.g., loss of income) were more affected by depression symptoms.⁹

Our main goal in this review on “psychological resilience during COVID-19” is to summarize the main findings on the following aspects: mental health consequences of the current pandemic, factors that facilitate the relationship between COVID and mental health,

resilience during infectious diseases outbreak. Lastly, we describe lessons learned that can be used in other conditions.

2. Factors that facilitate the relationship between COVID-19 and mental health

The COVID-19 pandemic is associated with mortality, contamination rate, and strict biosecurity restrictions that are affecting many segments of a functional society. At the peak of the pandemic, the main objective was to mitigate the devastating impacts of populations around the globe that were struggling to cope with: neighborhood quarantines, social distancing, nighttime curfews, and states of emergency that have been imposed worldwide.^{10,11} The global economy receded to a precarious state with high levels of unemployment in major industries, and sectors coming to a screeching halt, geographic borders and markets closed for business, trade and agricultural production faltered, altogether creating a cascade of financial losses and psychological stress above and beyond the fear-provoking consequences of the virus itself.^{7,12–14} Most importantly, not all countries, societies, and communities were impacted equally. Many countries have no increased unemployment percentages.

This is not the first time the world has witnessed a global pandemic and catastrophe of epic proportions. The death toll of the influenza pandemic of 1918–1919 surpassed that of World War I (WWI). An estimate of 20 to 40 million people lost their lives. Also known as the Spanish Flu it has been cited as the most devastating epidemic in recorded world history.¹⁵

Less than one decade ago, another deadly epidemic emerged in Western Africa with Ebola virus disease (EVD) in 2013. It was the most widespread outbreak of Ebola virus disease (EVD) in history.¹⁶

Exposure to global epidemics such as the 2014–2015 Ebola and the 2003 SARS outbreaks has been associated with a high rate of common mental disorders among survivors in the short to long-term. The prevalence of any psychiatric disorder at 30 months post-SARS was 33.3%. One-fourth of the patients had post-traumatic stress disorder (PTSD), and 15.6% had depressive disorders. Additionally, there is growing evidence on the adverse psychological outcomes including PTSD symptoms, confusion, and anger that are associated with the isolation or quarantine experience imposed on suspected cases during those outbreaks.^{11, 17–21}

As of September 2020, the devastation caused by COVID-19 goes beyond that caused by the SARS epidemic and Ebola. Preliminary studies published on the mental health consequences of the current COVID-19 reveal an alarming rate of stress-related disorders among populations exposed to the pandemic. Symptoms of anxiety and depression (16–28%) and self-reported stress (8%) are common psychological reactions to the COVID-19 pandemic, and may be associated with disturbed sleep and may vary by level of exposure.²²

In a nationwide survey of 52,730 individuals from 36 provinces, autonomous regions, and municipalities, from Hong Kong, Macau, and Taiwan between January 31 and February 20, 2020, about 35% of the sample experienced psychological distress. Among this sample, in

line with previous findings, the COVID-19 Peri-traumatic Distress Index (CPDI) score was linked to gender, age, education, occupation, and region.²³

3. Psychological Resilience and Infectious Diseases Outbreak

Over the past decade, the human's ability to adapt in the face of tragedy, trauma, adversity, hardship, and ongoing significant life stressors has been studied under the rubric of "Psychological Resilience". Psychological resilience has garnered tremendous interest in mental health and has seized the attention of public health practitioners. As such, psychological resilience has emerged as a fundamental variable in reducing and preventing the negative psychological effects of the pandemic. Currently, PubMed contains more than 50,000 entries related to COVID-19; among them, more than 150 scientific publications focus on "Psychological resilience" and "COVID-19". Overall, publications studying psychological resilience during COVID-19 could be summarized in two categories.

- Psychiatric epidemiological research
- Report papers presenting models of interventions to reinforce Psychological resilience.

Psychiatric epidemiological research on resilience and mental health during COVID-19

Typically, those studies provide evidence on psychological resilience factors such as protective factors for anxiety, depression, post-traumatic stress, and other mental health disorders are included in this category. For instance, during the peak of the pandemic in China, Ran et al. (2020) found that psychological resilience (CD-RISC scores) was negatively associated with symptoms of depression, anxiety, and somatization. The authors concluded that participants with greater psychological resilience were less likely to exhibit emotional problems, while those with lower resilience factors were more likely to report negative emotional symptoms.¹² A similar finding was observed in Germany in an online survey which found that strategies such as maintaining a healthy lifestyle and social contacts, acceptance of anxiety and negative emotions, fostering self-efficacy, and information on where to get medical treatment when necessary was helpful. Nevertheless, the psychological burden was mostly linked to substance abuse and suppression of anxiety and negative emotions.¹¹

Key psychological resilience promoting factors during COVID-19 lockdown

At present, there is preliminary data on important key behaviors and other factors that may contribute to psychological resilience during the pandemic that originated from China in December 2019. For instance, among Chinese adolescents, a study explored the emotional resilience of middle school students learning at home between February and March 2020 as well as its effect on students' learning management ability. The results of that study indicated that emotional resilience, measured by the Chinese version of the "adolescent emotional resilience scale" (AERS), was lower among eighth-graders, compared to seventh-graders. Furthermore, negative emotional recovery worsened during that period, and

emotional resilience was positively related to learning management skills; positive emotional ability was a predictor of learning management skills.

In an online Turkish survey conducted among social media users using the “Short Psychological Resilience Scale” and the “Beck Depression Scale” (BDI), males, educators, university graduates and those without mental health problems presented stronger psychological resilience. Conversely, females, university students, high school and lower graduates, and individuals with mental health issues were at higher risk for depression.²⁵

One American study that focused on resilience (defined as the self-perceived trait-like ability to face threatening situations according to the Connor-Davidson Risk Scale-CD-RISC; Connor and Davidson, 2003) among 18–35 years Americans in early April 2020 found that greater resilience was predicted by a set of multiple modifiable factors including the number of days per week spent outdoor in the sunshine, daily exercise, perceived support from closed and loved-ones, sleep quality, perceived care and support from a significant closed-ones, and frequency of prayer. In summary, those who obtained higher scores on this combination of variables were more likely to exhibit stronger psychological resilience during the lockdown.^{32–35}

Bueanas et al. (2020) explored coping strategies using the “Temperament and character inventory-revised” (TCI-R) (Cloninger, Svrakic, Przybeck, & Whitehead, 1999) associated with symptomatic deterioration among people with eating disorders (ED). Accordingly, a worsening in ED symptomatology and other psychiatric symptoms (anxiety and depression) during the COVID-19 lockdown were linked to low self-directedness. Greater ED symptoms during confinement were associated with less adaptive coping strategies to adjust to recommended confinement leading to weight gain. This aligns well with other work in confinement as reported by Moccia et al. (2020). Additional variables such as age and ED severity were also associated with COVID-19 concerns.

Overall, resilience promoting factors related to the ongoing COVID-19 pandemic could be regrouped in a chronological perspective: 1) pre-COVID-19 factors, 2) peri-COVID-19, and lockdown factors, 3) Post-COVID-19 Lockdown factors (Figure 1).

4. Lessons learned that can be used in other conditions

These publications provide interventions models to reinforce resilience among specific populations during the COVID-19 pandemic. However, it is important to note that in this category translational data was scarce.

Individual-level—As shown in figure 1, modifiable peri-COVID-19 factors were leveraged frequently by individuals in their effort to cope with the pandemic. Accordingly, researchers have made many recommendations from the conceptual and methodological gap of psychological resilience perspective and programming standpoint such as learning stress management, social awareness and minimizing media consumption, increased knowledge on mental health education, continue to build support and bonding, finding ways to increase self-preservation through – physical fitness, reading, yoga, proper nutrition, laughter- finding creative outlets to social and interpersonal communication with individuals through small

groups and online communication^{3,5, 27, 28}. Virtual reality therapy to acclimatize individuals pre-travel or the growing interest in mental well-being health applications to be used as an informative tool for persons (i.e. health care workers) dealing with PTSD- whether it is working long hours in high-intensity environments^{27, 23, 12}, or individuals recovering from COVID-19-^{25,31}. It could also be the invention of technologically innovative solutions (i.e. mental well-being apps) as a mechanism to promote psychological resilience to foster a digitized community and reduce fear, stress, anxiety, and depression.^{20,21,22,35–38}

Institutional and community levels—Leveraging social and economic resources available from their individuals' immediate environment, emerged often in our rapid literature review on resilience-promoting factors during COVID-19. Blanc et al. (2020) in a qualitative study about coping lessons that high-income countries could learn from the Haitian resilience, revealed three main psychological resilient characteristics that may have helped Haitians to cope with the alarming predictions from international and national health authorities. These are- Hardiness, the ability to cope, move forward through traumatic events and large-scale disasters; With also the ability to make, imagine, and hope for new future possibilities through creative art. Finally, traditional, natural, and culturally specific interventions (alignment with spiritual beliefs and complementary healing practices) as a mechanism to restore harmony and balance.^{30, 40} These characteristics do not exist in social isolation but are reinforced through community and togetherness - These are acts that other western countries could leverage in their efforts to buster community resilience during this pandemic.

In the U.S., at the institutional level, for instance, Mount Sinai Health System (MSHS) has created an Employee, Faculty, and Trainee Crisis Support Taskforce (in Early March 2020) in New York City. The task force implemented a rapid needs assessment model guided by three priority areas that focus on addressing many challenges of the workforce and maintaining the wellbeing of the entire MSHS workforce concerning the COVID-19 pandemic.¹² The priority areas are (1) meeting the basic needs of the workforce throughout the crisis- this is executed through the Office of Well-being and Resilience (OBWR) human resources, recreation office, housing office, infection prevention, nursing - entails serving 5 categories- food (discounted or free), housing (onsite call room, on-campus option, local hotel at discounted rates), Transportation (free parking, discounted or free car rentals) (2) Enhancing Communications – consolidating system-wide messaging into a daily communication with links to a comprehensive website has helped streamline messaging and direct the workforce situated within multiple hospitals and numerous practice sites, to a single regularly updated resource (3) Developing Psychosocial and mental health supportive options- includes several options- self-care resource (i.e. mindfulness activities) to virtual support groups facilitated by social workers and psychologist and to one-on-one counseling sessions and 24/7 immediate crisis management, training mental health staff with exercising the flexibility to shift their responsibilities when needed, and creating new initiatives.¹³,

Conclusion

The mental health toll on populations exposed to COVID-19 is alarming. The current review provides insights on how individuals, communities, and specific populations such as

healthcare workers and patients are leveraging pre and peri-COVID-19 factors to reinforce their psychological resilience during the ongoing global public health crisis. Throughout the review, we noted that populations around the world rely often on support from their loved-ones, closed significant others, outdoor and physical activities, and spirituality to cope with the COVID-19 related distress. Increased sense of meaning/purpose since the COVID-19 pandemic was also reported. Concerning interventions, many scholars have recommended community and individual evidence-based models that could potentially reinforce psychological resilience during COVID-19. Nevertheless, we are skeptical about the universal character of such models and what is deemed legitimate hard evidence with the proper use of methods. Despite the weakness in their health infrastructures, some low-income countries seem to have been successful in coping with the pandemic, while in high-income countries, historically disadvantaged populations bear the greatest toll of the pandemic. Consequently, there is an emergency to fill the gap of scientifically-sounded and culturally-responsive data supporting the effectiveness of the aforementioned interventions, programs, or policies on psychological resilience during the pandemic.

Financial support and sponsorship:

Grant Number: T32HL1299953, United States

References

1. World Health Organization. WHO Director-General's opening remarks at the media briefing on COVID-19 (March 11 2020). <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>. Published March 11, 2020.
- 2**. Jean- Louis G, Turner A, Peng, Mengling L, Boutin-Foster C, McFarlane SI, Seixas A. Increased Metabolic Burden Among Blacks: A Putative Mechanism for Disparate Covid-19 Outcomes. *Diabetes Metab Syndr Obes* 2020; 13: 3471–3479. [PubMed: 33061507] Study demonstrates an overwhelming high prevalence of the metabolic factors related to COVID-19 among Blacks in New York, highlighting disparate metabolic burden among blacks as a possible mechanism conferring the greater burden of COVID-19 infection and mortality.
- 3**. Labrague LJ, De los Santos JAA. COVID-19 anxiety among front-line nurses: Predictive role of organizational support, personal resilience and social support. *J of nurs manag* 2020. This study examines the relative influence of personal resilience, social support and organisational support in reducing COVID-19 anxiety in front-line nurses
- 4*. Vindegaard N, Benros ME. COVID-19 pandemic and mental health consequences: systematic review of the current evidence. *Brai, Beha, and Immu* (2020). Study identified the indirect effects of the pandemic on general mental health are of increasing concern, particularly since the SARS-CoV-1 epidemic (2002–2003) was associated with psychiatric complications
5. Connor KM, Davidson JR. Development of a new resilience scale: The Connor-Davidson resilience scale (CDRISC). *Depr and anxie*, 2003, 18(2), 76–82. The study uses The Connor-Davidson Resilience scale (CD-RISC) to demonstrate is modifiable and can be improved with treatment.
6. Stewart DE, Yuen T. A systematic review of resilience in the physically ill. *Psychosomatics* 2011, 52, no. 3 : 199–209. [PubMed: 21565591] Developed a systematic review of resilience and related concepts in the physically ill to determine factors associated with predicting or promoting resilience.
- 7*. Farewell CV, Jewell J, Walls J, Leiferman JA. A mixed-methods pilot study of perinatal risk and resilience during COVID-19. *J of Pri Care & Comm Hea* 2020 11: 2150132720944074. The primary aim of this pilot study was to use mixed-methods to better understand the mental health and well-being effects of the coronavirus disease 2019 (COVID-19) pandemic, as well as sources of resilience, among women during the perinatal period.

8. Moccia L, Janiri D, Pepe M, Dattoli L, Molinaro M, De Martin V, Daniela Chieffo D, Fiorillo A, Sani G, Di Nicola M. Affective temperament, attachment style, and the psychological impact of the COVID-19 outbreak: an early report on the Italian general population. *Brain, behavior, and immunity* (2020). Study aims to investigate the psychological distress perceived by the Italian general population during the early phase of the COVID-19 pandemic, and to analyze affective temperament and adult attachment styles as potential mediators.
- 9**. Shechter A, Diaz F, Moise N, Anstey DE, Ye S, Agarwal S, Birk JL, Brodie D, Cannone DE, Chang B, Claassen J. Psychological distress, coping behaviors, and preferences for support among New York healthcare workers during the COVID-19 pandemic." *Gen hosp psyc* 2020, 66: 1–8. Study identifies characterized distress, coping, and preferences for support among NYC HCWs during the COVID-19 pandemic.
- 10**. Ettman CK, Abdalla SM, Cohen GH, Sampson L, Vivier PM, Galea S. Prevalence of Depression Symptoms in US Adults Before and During the COVID-19 Pandemic. *JAMA Network Open* 2020,3, no. 9 : e2019686–e2019686. [PubMed: 32876685] To estimate the prevalence of and risk factors associated with depression symptoms among US adults during vs before the COVID-19 pandemic.
- 11*. Petzold MB, Bendau A, Plag J, Pyrkosch L, Lea Mascarell Maricic, Betzler F, Rogoll J, Große J, Ströhle A. Risk, resilience, psychological distress, and anxiety at the beginning of the COVID-19 pandemic in Germany. *Bra and Behav* 2020: e01745. The aim of this study was to screen the population in Germany for negative impact on mental health in the current COVID-19 pandemic and to analyze possible risk and protective factors
12. Ran L, Wang W, Ai M, Kong Y, Chen J, Kuang L. Psychological resilience, depression, anxiety, and somatization symptoms in response to COVID-19: A study of the general population in China at the peak of its epidemic. *Soci Scie & Medi* 2020: 113261. This research investigated the relationship between psychological resilience and mental health (depression, anxiety, somatization symptoms) among the general population in China.
- 13**. Ripp J, Peccoraro L, Charney D. Attending to the emotional well-being of the health care workforce in a New York City health system during the COVID-19 pandemic. *Acad Medic* 2020. Utilization of Employee, Faculty, and Trainee Crisis Support Task force - rapid needs assessment model addressing challenges in the workforce and maintaining well-being.
14. Richard SA., Sugaya N, Simonsen L, Miller MA, Viboud C. A comparative study of the 1918–1920 influenza pandemic in Japan, USA and UK: mortality impact and implications for pandemic planning. *Epid & Infect* 2009 137, no. 8: 1062–1072. Identifies the mortality rate during the Spanish flu
- 15**. Shultz JM, Cooper JL, Baingana F, Oquendo MA, Espinel Z, Althouse BM, Marcelin LH, Towers S, Espinola M, McCoy CB, Mazurik L, "The role of fear-related behaviors in the 2013–2016 West Africa Ebola virus disease outbreak." *Curr psyc repo* 2016; 18, no. 11: 104.
16. Flaherty GT, Nasir N. Reiseangst: travel anxiety and psychological resilience during and beyond the COVID-19 pandemic. *J Travel Med.* 2020 9 1:taaa150. [PubMed: 32870275] The study uses virtual reality tools as a form of psychological resilience to address anxiety and stress in travelers.
17. Prime H, Wade M, Browne DT. Risk and resilience in family well-being during the COVID-19 pandemic. *Am Psychol* 2020; 75(5):631–643. [PubMed: 32437181] Study demonstrates how social support through family can be a mechanism of psychological resilience during COVID 19
18. Krešimir C, Popovi S, Šarlija M, Kesedži I. Impact of human disasters and Covid-19 pandemic on mental health: Potential of digital psychiatry *Psychiatria Danubina* 2020; 32, no. 1: 25–31. [PubMed: 32303026] The promotion for digital tools as a means in mitigation of stress-related disorders and assessment of stress resilience
19. Petzold MB, Bendau A, Plag J, Pyrkosch L, Mascarell Maricic L, Betzler F, Rogoll J, Große J, Ströhle A. Risk, resilience, psychological distress, and anxiety at the beginning of the COVID-19 pandemic in Germany. *Brain Behav* 2020; 7 7:e01745. doi: 10.1002/brb3.1745. Epub ahead of print. Authors aim of the study was to screen the population in Germany on the negative impact of COVID 19 pandemic and analyze possible risk and protective risk factors.
20. Alexopoulos AR, Hudson JG, Otenigbagbe O. The Use of Digital Applications and COVID-19. *Community Ment Health J.* 2020 10;56(7):1202–1203.. [PubMed: 32734311] This study promotes Resilience Among College Student Veterans Through an Acceptance-and-Commitment-Therapy

App: An Intervention Refinement Study”, 2020) the importance of repurposing their app to help these users to improve their emotional resilience and subsequently their ability to cope with the trauma

- 21*. Rajkumar RP. “COVID-19 and mental health: A review of the existing literature.” *Asian journal of psychiatry* (2020): 102066. [PubMed: 32302935] Study reveals such widespread outbreaks are associated with adverse mental health consequences.
22. Lupe SE, Keefer L, Szigethy E. Gaining resilience and reducing stress in the age of COVID-19. *Curr Opin Gastroenterol.* 2020; 36(4): 295–303. [PubMed: 32398567] The studies draw from the relevant literature on stress reactions, and offer some guidance on how to help patients mitigate the physiological and psychological impact of the pandemic through resilience-building techniques.
23. Qi H, Xiao S, Shi R, Ward MP, Chen Y, Tu W, Qing Q, Wang W, Wang X, and Zhang Z. COVID-19 transmission in Mainland China is associated with temperature and humidity: A time-series analysis. *Science of the Total Environment* (2020): 138778.
24. Farewell CV, Jewell J, Walls J, Leiferman JA. A mixed-methods pilot study of perinatal risk and resilience during COVID-19. *J of Prim Care & Comm Hea* 2020 11: 2150132720944074. The aim of this pilot study used mixed-methods to understand the mental health, well-being effects, and sources of resilience among women during the perinatal period and coronavirus disease 2019 (COVID-19) pandemic
25. Kara ar B, Canli D. Psychological Resilience and Depression during the Covid-19 Pandemic in Turkey. *Psychiatr Danub.* 2020; 32(2). 273–279. doi: 10.24869/psyd.2020.273. The purpose of this study is to investigate psychological resilience and depression in individuals during the period of COVID-19 outbreak in Turkey in relation to different variables.
- 26**. Labrague LJ, De Los Santos JAA. COVID-19 anxiety among front-line nurses: Predictive role of organisational support, personal resilience and social support. *J Nurs Manag* 2020 8:10.1111/jonm.13121. Examines the relative influence of personal resilience, social support and organisational support in reducing COVID-19 anxiety in front-line nurses.
27. Solano GA. RESILIENCE AND COVID-19. *Rev Colomb Obstet Ginecol.* 2020; 3;71(1):7–8. doi: 10.18597/rcog.3531. [PubMed: 32418371]
28. van Ageren J, Bartholomaeus J, Fassnacht DB, Iasiello M, Ali K, Lo L, Kyrios M. Using Internet-Based Psychological Measurement to Capture the Deteriorating Community Mental Health Profile During COVID-19: Observational Study. *JMIR Ment Health.* 2020 6 11;7(6) This study uses an internet-based self-assessment tool to collect data on psychological distress, mental well-being, and resilience in community cohorts during (n=673) and prior to the pandemic (two cohorts, n=1264)
- 29**. Choi S, Hong JY, Kim YJ, Park H. Predicting Psychological Distress Amid the COVID-19 Pandemic by Machine Learning: Discrimination and Coping Mechanisms of Korean Immigrants in the U.S. *Int J Environ Res Public Health.* 2020 8 20;17(17):E6057. [PubMed: 32825349] Study examines the predictive ability of discrimination-related variables, coping mechanisms, and sociodemographic factors on the psychological distress level of Korean immigrants in the US amid the COVID-19 pandemic.
- 30**. Blanc J, Elizabeth FL, Jolette J, Chimene C, Jean-Louis G. What the world could learn from the Haitian resilience while managing COVID-19. *Psychological Trauma: Theory, Research, Practice, and Policy* 2020, 12, no. 6: 569. Authors explain three concepts, hardiness, creativity, and traditional and complementary healings as a form of psychological resilience and the world should learn from these practices.
31. Stanton R, To QG, Khalesi S, Williams SL, Alley SJ, Thwaite TL, Fenning AS, Vandelanotte C. Depression, Anxiety and Stress during COVID-19: Associations with Changes in Physical Activity, Sleep, Tobacco and Alcohol Use in Australian Adults. *Int J Environ Res Public Health.* 2020 6 7;17(11):4065. Study examined the associations between psychological distress and changes in selected health behaviors since the onset of COVID-19 in Australia.
32. Tai DB, Shah A, Doubeni CA, Sia IG, Wieland ML. The disproportionate impact of COVID-19 on racial and ethnic minorities in the United States. *Clinical Infectious Diseases.* 2020 6 20. The authors provide insight into the disproportionate affect the pandemic has had on racial and ethnic groups.

- 33*. Abedi V, Olulana O, Avula V, Chaudhary D, Khan A, Shahjouei S, Li J, Zand R. Racial, Economic, and Health Inequality and COVID-19 Infection in the United States. *Journal of racial and ethnic health disparities*. 2020 9 1:1–1. [PubMed: 31410785] The authors study the association between race, health and socioeconomics with the virus.
34. Quandt SA, LaMonto NJ, Mora DC, Talton JW, Laurienti PJ, Arcury TA. COVID-19 Pandemic among Latinx Farmworker and Nonfarmworker Families in North Carolina: Knowledge, Risk Perceptions, and Preventive Behaviors. *Inter j of enviro rese and publ heal*. 2020 1;17(16):5786. Identify knowledge and preventive behaviors to reduce COVID-19 spread in the community.
35. Killgore WD, Taylor EC, Cloonan SA, Dailey NS. Psychological Resilience During the COVID-19 Lockdown. *Psychiatry Research*. 2020 6 9:113216. [PubMed: 32544705] The authors identify factors that may contribute psychological resilience.
- 36**. Raifman MA, Raifman JR. Disparities in the population at risk of severe illness from covid-19 by race/ethnicity and income. *American journal of preventive medicine* 2020. This study highlights the numbers of racial/ethnic and income disparities that should be considered in physical distancing and other protective measures.
37. LeRoy AS, Robles B, Kilpela LS, Garcini LM. Dying in the face of the COVID-19 pandemic: Contextual considerations and clinical recommendations. *Psychological Trauma: Theory, Research, Practice, and Policy*. 2020 6 11 *Psychological Trauma: Theory, Research, Practice, and Policy*. 2020 Jun 11. The authors discuss contextual and cultural considerations needed to facilitate healing and resiliency
- 38**. Chin-Hong P, Kevin MA, Haynes N, Albert MA. Pulling at the heart: COVID-19, race/ethnicity and ongoing disparities. *Nat Rev Cardiol* 2020; 17, 533–535. [PubMed: 32732989] Authors advocate for decisive attention to and action against ethnicity based inequities that undermine cardiovascular health.
- 39**. Wang Q, Xu R, Volkow DN. Increased risk of COVID –19 infection and mortality in people with mental disorders: analysis from electronic health records in the United States. *World Psych* 2020: 1–7. The authors examine the impact of a recent diagnosis of a mental disorder (i.e. bipolar, depression, ADHD) on the risk of for COVID-19 infection and related mortality and hospitalization rates.
40. Blanc J, Eugene D, Louis EF, Cadichon JM, Joseph J, Pierre A, Laine R, Alexandre M, Haung KY. Mental Health Among Children Older than 10 years Exposed to the Haiti 2010 Earthquake: a Critical Review. *Curr Psychiatry Rep* 2020 9 2: 22(11):57. [PubMed: 32876808] Study shows evidence that the exposure to major natural disaster such as the 2010 earthquake in Haiti may have devastating short- and long term effects on children’s mental health and overall development.

Key points

- Psychological resilience has emerged as a fundamental variable in reducing and preventing the negative psychological effects of the pandemic.
- Populations around the world rely often on support from their loved ones, closed significant others, outdoor and physical activities, and spirituality to cope with the COVID-19 related distress.
- Overall, resilience promoting factors related to the ongoing COVID-19 pandemic could be regrouped in a chronological perspective: 1) pre-COVID-19 factors, 2) peri-COVID-19 Lockdown factors, 3) Post-COVID-19 Lockdown factors




Figure 1:
Summary of Key psychological resilience promoting factors during COVID-19 lockdown