

## REVIEW

## Cancer Focus

# Lipid metabolism and cancer

Xueli Bian<sup>1\*</sup>, Rui Liu<sup>2\*</sup>, Ying Meng<sup>3\*</sup>, Dongming Xing<sup>1,4</sup>, Daqian Xu<sup>3</sup>, and Zhimin Lu<sup>3,5</sup>

**Dysregulation in lipid metabolism is among the most prominent metabolic alterations in cancer. Cancer cells harness lipid metabolism to obtain energy, components for biological membranes, and signaling molecules needed for proliferation, survival, invasion, metastasis, and response to the tumor microenvironment impact and cancer therapy. Here, we summarize and discuss current knowledge about the advances made in understanding the regulation of lipid metabolism in cancer cells and introduce different approaches that have been clinically used to disrupt lipid metabolism in cancer therapy.**

## Introduction

Lipids, together with proteins and nucleic acids, are essential components of biological membranes and building blocks that constitute cells. In addition, lipids are used in energy storage and metabolism and have important roles as signaling molecules for many cellular activities. The regulation of lipid metabolism, such as lipid uptake, synthesis, and hydrolysis, is essential for the maintenance of cellular homeostasis (Röhrig and Schulze, 2016). Cancer cells in tumor microenvironments, where nutrient availability is consistently changing during tumor progression, harness lipid metabolism to support their rapid proliferation, survival, migration, invasion, and metastasis.

Glycolipids and phospholipids (which are subcategorized into phosphoglycerides and sphingolipids), together with cholesterol, represent major components of biological membranes. Cholesterol is also a substrate for the synthesis of fat-soluble vitamins and steroid hormones (Luo et al., 2020). As major components of glycolipids and phospholipids, fatty acids (FAs) can be esterified with a glycerol moiety to form triglycerides, which are nonpolar lipids synthesized and stored in lipid droplets during high nutrient availability and hydrolyzed to generate ATP by FA oxidation (FAO, also called  $\beta$ -oxidation) under energy stress conditions. Aside from energy metabolism and membrane formation, lipids form second messengers derived from phospholipase-dependent hydrolyzation of membrane lipids and synthesis from essential FAs, whose availability is largely determined by lipids in the diet (Park et al., 2012).

Phospholipases (PLC, PLD, and PLA) can generate many bioactive second messengers, such as diacylglycerol, phosphatidic acid, lysophosphatidic acid, and arachidonic acid. These molecules trigger the activation of the RAS, phosphoinositide 3-kinases (PI3Ks), protein kinase C, RAC, RHO, and several other signaling axes that can promote tumorigenesis, which was intensively discussed previously and is not covered in this review (Moolenaar and Perrakis, 2011; Park et al., 2012). In addition, sterols, including oxysterol and cholesterol, are critical regulators of sterol regulatory element (SRE)-binding protein (SREBP) activation for downstream gene expression, and thus their levels affect lipogenesis in cancer. Cholesterol is a component of lipid rafts for signaling and can also covalently modify Hedgehog and Smoothened proteins for Hedgehog signaling activation (Porter et al., 1996; Xiao et al., 2017).

In this review, we summarize and discuss current knowledge about the advances in the regulation of lipid (FAs and cholesterol) uptake, lipogenesis, and FAO-dependent lipolysis in cancer cells. We dedicate a separate section at the end to discuss different approaches that might be used to disrupt lipid metabolism for cancer therapy.

## Lipid uptake

### FA uptake

Mammals produce only certain FAs, i.e., those carrying double bonds to the  $\Delta 9$  position of the hydrocarbon chain. Other FAs, particularly polyunsaturated FAs, are essential and obtained

<sup>1</sup>Cancer Institute of The Affiliated Hospital of Qingdao University and Qingdao Cancer Institute, Qingdao, China; <sup>2</sup>State Key Laboratory of Oral Diseases, National Clinical Research Center for Oral Diseases, Chinese Academy of Medical Sciences Research Unit of Oral Carcinogenesis and Management, West China Hospital of Stomatology, Sichuan University, Chengdu, China; <sup>3</sup>Zhejiang Provincial Key Laboratory of Pancreatic Disease, The First Affiliated Hospital, and Institute of Translational Medicine, Zhejiang University School of Medicine, Hangzhou, China; <sup>4</sup>School of Life Sciences, Tsinghua University, Beijing, China; <sup>5</sup>Zhejiang University Cancer Center, Hangzhou, China.

\*X. Bian, R. Liu, and Y. Meng contributed equally to this paper; Correspondence to Zhimin Lu: [zhiminlu@zju.edu.cn](mailto:zhiminlu@zju.edu.cn); Daqian Xu: [xudaqian@zju.edu.cn](mailto:xudaqian@zju.edu.cn); Dongming Xing: [xdm\\_tsinghua@163.com](mailto:xdm_tsinghua@163.com).

© 2020 Bian et al. This article is distributed under the terms of an Attribution–Noncommercial–Share Alike–No Mirror Sites license for the first six months after the publication date (see <http://www.rupress.org/terms/>). After six months it is available under a Creative Commons License (Attribution–Noncommercial–Share Alike 4.0 International license, as described at <https://creativecommons.org/licenses/by-nc-sa/4.0/>).

from the diet (Nakamura and Nara, 2004). The known FA protein transporters in the plasma membrane include cluster of differentiation 36 (CD36; also known as FA translocase), the family of FA transport proteins (also collectively known as SLC27), and plasma membrane FA-binding proteins (FABPs), all of which display increased gene and protein expression in tumors (Su and Abumrad, 2009).

High CD36 expression has been correlated with poor prognosis for patients with breast, ovarian, gastric, and prostate cancer (Koundouros and Poulogiannis, 2020). Deletion of *Cd36* in the prostate of cancer-susceptible *Pten*<sup>-/-</sup> mice or mammary tissues of MMTV-neu mice attenuated increased FA uptake in cancer and mitigated tumorigenesis (Feng et al., 2019; Watt et al., 2019). In breast cancer patients, CD36 expression increases following anti-HER2 therapy and correlates with poor survival. Correspondingly, the HER2 inhibitor lapatinib-resistant breast cancer cells increase FA uptake, and CD36 inhibition suppresses the growth of lapatinib-resistant, but not that of lapatinib-sensitive, tumor cells (Feng et al., 2019). High-fat diets induced NF- $\kappa$ B-dependent CD36 expression and elicited O-GlcNAcylation of CD36 at S468 and T470, which enhanced FA uptake and murine gastric cancer metastasis (Jiang et al., 2019). Hydrogen sulfide, which is implicated in cancer metastasis, induces CD36 expression and reduces C333-C272 disulfide bond formation in CD36, which activates the long-chain FA-binding conformation of CD36 to promote FA uptake and accelerate gastric cancer metastasis (Wang et al., 2019). Palmitic acid induces gastric cancer cell migration and invasion through CD36-dependent activation of the protein kinase B (PKB, also known as AKT)/ $\beta$ -catenin signaling pathway, whereas dietary oleic acid up-regulates CD36 expression and its expression-dependent activation of the Src-ERK1/2 pathway, which promotes cervical cancer cell growth (Pan et al., 2019; Yang et al., 2018). Palmitic acid or a high-fat diet can specifically boost the metastatic potential of metastasis-initiating oral cancer cells with high CD36 expression. Blockade of CD36 with neutralizing antibodies inhibits the metastasis of oral cancer cells in mice (Pascual et al., 2017), implying a role for dietary lipids and CD36 in tumor metastasis.

The ability of cancer cells to use exogenous lipids for the provision of unsaturated FAs appears to be dependent on oxygen levels and the type of oncogene expressed, and not all lipids are exploited equally (Snaebjornsson et al., 2020). Hypoxic tumor cells exhibit increased FA import dependent on hypoxia-inducible factor-1 $\alpha$  (HIF-1 $\alpha$ )-induced FABP3/7 expression, which accompanies reduced de novo FA synthesis because the catabolism of glucose to acetyl-coenzyme A (CoA) is decreased (Bensaad et al., 2014). Compared with myrAKT expression, which boosts de novo FA synthesis, oncogenic H-RasV12G or K-RasG12D expression under normoxic conditions recapitulates hypoxic conditions and re-establishes cell reliance on the uptake of lipids, among which lysophospholipids containing mono- or polyunsaturated acyl chains are internalized to a far greater extent than phospholipids and saturated lysophospholipids (Kamphorst et al., 2013). In addition to being blood resources, adipocytes in the tumor microenvironment provide extracellular FAs for tumor cells. FABP4 expression was detected in ovarian cancer cells at the adipocyte-

tumor cell interface. Omentum-metastatic ovarian cells activate the lipolysis in adipocytes to produce free FAs that can be subsequently secreted and taken up by cancer cells in a FABP4-dependent manner for increased FAO and rapid tumor growth (Gharpure et al., 2018; Nieman et al., 2011). In addition, omental adipocytes induce CD36 expression and associated lipid accumulation in ovarian cancer cells. Inhibition of CD36 attenuated adipocyte-induced cholesterol and lipid droplet accumulation in ovarian cancer cells, peritoneal dissemination of tumor cells, and tumor growth (Ladanyi et al., 2018). Similar to adipocytes, tissue-resident pancreatic stellate cell-derived cancer-associated fibroblasts in pancreatic ductal adenocarcinoma (PDAC) secrete lipids including lysophosphatidylcholines for cancer cell uptake and synthesis of phosphatidylcholines. In addition, PDAC cell-secreted enzyme autotaxin hydrolyzes lysophosphatidylcholines into lysophosphatidic acid as an extracellular signal to promote PDAC cell proliferation and migration (Auciello et al., 2019). Thus, hypoxia, overexpressed FA transporters in tumor cells, specific oncogene expression, and tumor cell-modulated stromal cells including adipocytes and fibroblasts can induce tumor cells to take up extracellular FAs and create mitogenic signals to sustain tumor cell proliferation.


#### Cholesterol uptake

Dietary cholesterol is absorbed by Niemann-Pick type C1-like 1 (NPC1L1) protein in the membrane of intestinal enterocytes (Altmann et al., 2004), where cholesterol is esterified by acyl-CoA:cholesterol acyltransferases (ACATs; also known as sterol O-acyltransferase) for uptake by the liver (Ko et al., 2020). The liver, the main cholesterol biosynthesis organ, delivers cholesterol as very-low-density lipoproteins to the bloodstream, where the very-low-density lipoproteins are processed into low-density lipoproteins (LDLs) for uptake by the LDL receptors (LDLRs) on peripheral cells. The cellularly absorbed cholesterol eventually reaches the ER for sensing, transport, or esterification (Goldstein and Brown, 2009).

LDLR expression was positively correlated with poor prognosis of the patients with small cell lung cancer (SCLC), breast cancers, and PDAC. LDLR depletion in HER2-overexpressing breast cancer cells and PDAC cells reduced cholesterol uptake and tumor growth in mice with hyperlipidemia (Gallagher et al., 2017; Guillaumond et al., 2015; Zhou et al., 2017). In addition, LDLR expression is transcriptionally up-regulated by epidermal growth factor receptor (EGFR) vIII mutant/PI3K-activated SREBP1 in glioblastoma (Guo et al., 2011). Loss of PTEN and subsequent PI3K/AKT activation in prostate cancer cells largely enhanced exogenous LDL uptake required for tumor growth (Yue et al., 2014). These findings underscore that the LDLR-mediated cholesterol uptake plays instrumental roles in the proliferation of some types of cancer cells.

#### Lipogenesis

Lipogenesis in normal tissue is primarily restricted to hepatocytes and adipocytes. Nevertheless, cancer cells activate lipogenesis in response to their high metabolic demand, even in the presence of exogenous lipid sources, or to serum-derived lipid



**Figure 1. Cytoplasmic acetyl-CoA production for lipid synthesis.** Cytoplasmic acetyl-CoA is produced from ACLY-catalyzed citrate and ACSs-catalyzed acetate. Glucose and glutamine contribute to citrate production from mitochondrial pyruvate oxidation in the TCA cycle and reductive carboxylation, respectively. PDH, pyruvate dehydrogenase;  $\alpha$ -KG,  $\alpha$ -ketoglutarate; GLS, glutaminase; PM, plasma membrane.

deficiency in the tumor microenvironment (Röhrig and Schulze, 2016). The main substrate for lipid synthesis is cytoplasmic acetyl-CoA, which can be derived from citrate by ATP-citrate lyase (ACLY) or acetate by acetyl-CoA synthetase (ACSs; Li et al., 2017a). Carbons from glucose and glutamine contribute to citrate production (Li et al., 2016b; Metallo et al., 2011; Fig. 1). Glucose-derived pyruvate is converted to acetyl-CoA by pyruvate dehydrogenase followed by citrate synthase-mediated production of citrate, which is then exported to cytosol from mitochondria by mitochondrial citrate transport proteins. Glutamine is converted to  $\alpha$ -ketoglutarate mediated by cytosolic glutaminase 1 (GLS1) or mitochondrial GLS2, which is followed by cytosolic isocitrate dehydrogenase 1 (IDH1)- and mitochondrial IDH2-dependent isocitrate and subsequent citrate production. Since both FA and cholesterol are synthesized from acetyl-CoA through a series of reactions, acetyl-CoA levels are a key element for lipid production (Fig. 2).


#### Acetyl-CoA-producing enzymes ACLY and ACSs

**ACLY.** ACLY catalyzes the conversion of citrate and CoA to oxaloacetate and acetyl-CoA. By inhibition of ACLY, bempedoic acid reduces patients' LDL cholesterol levels (Giral, 2020), displaying a critical role of ACLY in lipogenesis. The overexpression or enhanced activity of ACLY is correlated with tumor progression in glioblastoma, colorectal cancer, breast cancer, lung cancer, and hepatocellular carcinoma (HCC). Decreased ACLY expression reduces tumor cell viability and suppresses tumor cell proliferation, invasion, and metastasis in various cancer types (Hatzivassiliou et al., 2005; Khwairakpam et al., 2020).

ACLY, which is transcriptionally up-regulated by SREBP1, is allosterically activated by glycolytic intermediate fructose

6-phosphate and inhibited by citrate at high concentrations via homotropic allosteric regulation of ACLY. The citrate-dependent inhibition is abolished by ACLY S455 phosphorylation by AKT or protein kinase A (PKA; Berwick et al., 2002; Potapova et al., 2000). Cullin3-KLHL25 ubiquitin ligase targets ACLY for degradation and inhibits lipid synthesis and tumor progression (Zhang et al., 2016). Under high-glucose conditions, sirtuin-2 (SIRT2) deacetylase-promoted ACLY ubiquitylation and degradation are inhibited by lysine acetyltransferase 2B (KAT2B, also known as PCAF)-mediated acetylation at K540, K546, and K554. This acetylation is increased in human lung cancers and promotes lipogenesis and tumor growth (Lin et al., 2013). In addition, IKK $\beta$ -phosphorylated ACLY binds to IKK $\beta$ -phosphorylated and stabilized ubiquitin-specific protease-30 (USP30), which deubiquitinates and stabilizes ACLY in HCC cells (Gu et al., 2020). In diethylnitrosamine (DEN)/CCL<sub>4</sub>-induced liver tumors in mice, USP30 deletion attenuated lipogenesis, inflammation, and tumor growth (Gu et al., 2020).

In addition to participating in lipid metabolism in the cytosol, nucleus-translocated ACLY generates acetyl-CoA for histone acetylation and gene transcription regulation (Wellen et al., 2009). In melanoma cells, ACLY enhances acetyltransferase p300-dependent histone acetylation, thereby promoting PPAR $\gamma$  coactivator (PGC) 1 $\alpha$ -mediated mitochondrial biogenesis for cell proliferation (Guo et al., 2020). In PDAC, glioma, and prostate cancer cells, growth factors or oncogenic K-Ras expression promotes AKT-ACLY signaling, histone acetylation, cell proliferation, and tumor growth (Carrer et al., 2019; Lee et al., 2014). Following DNA damage, the ataxia telangiectasia mutated-AKT axis-induced nuclear ACLY S455 phosphorylation promotes histone acetylation at double-strand break sites, enabling BRCA1 recruitment and DNA repair by homologous recombination



**Figure 2. The FA and cholesterol biosynthesis.** FA biosynthesis starts with conversion of acetyl-CoA to malonyl-CoA by ACC. Acetyl-CoA and malonyl-CoA are then catalyzed into palmitic acid by FASN. Further elongation is mediated by elongases to form polysaturated FA. Palmitic acid and polysaturated FAs are desaturated into unsaturated FAs by SCD and other fatty acyl-CoA desaturases, respectively. Cholesterol biosynthesis starts with the condensation of two molecules of acetyl-CoA by ACAT to form acetoacetyl-CoA, which is further condensed with a third molecule of acetyl-CoA by HMG-CoA synthase (HMGCS) to form HMG-CoA. HMGCR then reduces HMG-CoA to mevalonate, which is converted to farnesyl pyrophosphate (FPP). Farnesyl pyrophosphate-converted squalene is oxidized by SM to produce 2,3-oxidosqualene, a precursor of cholesterol and sterols.

(Sivanand et al., 2017). Thus, ACLY-mediated acetyl-CoA production plays critical roles in multiple cellular activities in both the cytosol and nucleus.

**ACSS2.** ACSS produces acetyl-CoA via the ligation of acetate and CoA. ACSS1 and ACSS3 are mitochondrial proteins, and ACSS2 is localized in the cytoplasm and nucleus (Li et al., 2017b). ACSS2, which is transcriptionally up-regulated by SREBP, is expressed in a large proportion of human tumors and critical for acetate catalyzation to sustain cancer cell growth, especially under metabolic stress (Comerford et al., 2014; Luong et al., 2000). Acetate can be converted from fructose by the gut microbiota, and depletion of the microbiota or silencing of hepatic ACSS2 suppresses the conversion of bolus fructose into hepatic acetyl-CoA and FAs (Jang et al., 2018; Zhao et al., 2020). In addition, ACSS2 depletion or deficiency largely inhibits tumor cell growth and mouse tumor formation, highlighting a critical role of acetate consumption in the production of lipid biomass for tumor growth (Mashimo et al., 2014; Schug et al., 2015).

Limited amounts of oxygen, serum, or glucose increase the nuclear localization of ACSS2 (Bulusu et al., 2017; Li et al., 2017a, b). Glucose deprivation promotes 5' AMP-activated protein kinase (AMPK)-mediated ACSS2 S659 phosphorylation, resulting in exposure of its nuclear localization signal for nuclear translocation in glioblastoma cells. In complexes with transcription factor EB, ACSS2 activates lysosome- and autophagosome-related genes by locally producing acetyl-CoA for histone H3 acetylation at the promoter regions of these genes by using the acetate released from histone deacetylation, leading to lysosomal biogenesis and autophagy, cell survival, and brain tumor growth (Li et al., 2017a, b). In addition, ACSS2 enhances the expression and activation of liver X receptor (LXR)/retinoid X receptor transcription factors to promote lipogenesis during prolonged fasting (Huang et al., 2018). Analyses of human tumor specimens showed that ACSS2 S659 phosphorylation was positively correlated with glioma malignancy and poor survival of patients with non-SCLC (NSCLC; Li et al., 2017b; Yang et al., 2019), whereas high ACSS1/2 expression led to increased histone

H3 acetylation and FA synthase (FASN) expression in HCC patients (Bulusu et al., 2017). These findings reveal dual roles of ACSS2 in the direct involvement of de novo lipogenesis and the epigenetic regulation of gene expression for promoting lysosomal biogenesis, autophagy, and lipid metabolism.

#### FA biosynthesis enzymes

**Acetyl-CoA carboxylase (ACC).** ACC, the rate-limiting enzyme for FA synthesis, catalyzes the carboxylation of acetyl-CoA to malonyl-CoA (Fig. 2; Wei and Tong, 2015). Mammalian ACC has two tissue-specific isoforms: ACC1 (also known as ACC $\alpha$ , encoded by ACACA) and ACC2 (also known as ACC $\beta$ , encoded by ACACB). ACC1 is a cytosolic enzyme primarily expressed in lipogenic tissues (liver and adipose tissues) and is critical for FA synthesis. In contrast, ACC2 is embedded in mitochondrial outer membrane and is mainly expressed in oxidative tissues (heart and skeletal muscles), and ACC2-produced malonyl-CoA inhibits carnitine palmitoyltransferase I (CPT1, also known as palmitoyl-CoA transferase I, CAT1, or CCAT), which governs the rate-limiting step of FA uptake and FAO by mitochondria (Luo et al., 2012).

ACC1 is highly expressed in a variety of human cancers, including breast, prostate, liver, and gastric carcinoma, and the ACACA gene is present in recurrent amplicons associated with lowered survival of breast cancer patients (Chin et al., 2006; Luo et al., 2012). ACC1 depletion decreases FA synthesis and induces the apoptosis of prostate and breast tumor cells but not non-malignant cells (Brusselmans et al., 2005). ACC1 is regulated at the transcriptional level by SREBP and at the protein level by a complex interplay of phosphorylation, the binding of allosteric regulators, and protein-protein interactions (Koundouros and Poulogiannis, 2020). AMPK phosphorylates ACC1 S79 and ACC2 S212 and inhibits their activities, whereas the expression of ACC1/2 phosphorylation-dead mutants increases lipogenesis and lesions in mouse liver and the proliferation of human liver cancer cells (Ha et al., 1994; Lally et al., 2019). In addition, leptin and TGF $\beta$  signaling induce TGF $\beta$ -activated kinase (TAK)

1-AMPK-mediated ACC1 inhibition, resulting in increased cellular acetyl-CoA, Smad2 acetylation and activation, and breast cancer cell invasion and metastasis. These results suggest that fine-tuning ACC1 activity is critical for balanced FA synthesis and other cellular activities (Rios Garcia et al., 2017). ACC1 is allosterically activated by citrate binding and inactivated by the binding of palmitoyl-CoA and other fatty acyl-CoAs, as well as the BRCA1 C-terminal (BRCT) domains (Hunkeler et al., 2018). Mutated BRCA1, which is associated with the predisposition to inherited cancer, or insulin-like growth factor-1 (IGF-1) stimulation disrupts the interaction between BRCA1 and inactive and phosphorylated ACC. Consequently, ACC is dephosphorylated and activated for FA synthesis (Koobotse et al., 2018; Moreau et al., 2006). ACC stability can be increased by its interaction with peptidyl-prolyl cis-trans isomerase PIN1 to inhibit its lysosomal degradation in prostate cancer cells and with aldo-keto reductase family 1 B10 (AKR1B10) to inhibit its ubiquitylation-dependent degradation in breast cancer cells (Ma et al., 2008; Ueda et al., 2019). ACC1 expression and subsequent de novo FA synthesis can also be up-regulated by calcium/calmodulin-dependent protein kinase kinase 2 (CAMKK2) expression in prostate cancer cells, likely in a protein translation-dependent manner (Penfold et al., 2018). ACC1 also forms a complex with CPT1 and prevents its mitochondrial distribution under nutrient-sufficient conditions. During metabolic stress, phosphorylated ACC1 dissociates from CPT1, which translocates to the mitochondria for FAO, thus maintaining HCC cell survival (Wang et al., 2016).

ACC2 protein is highly expressed in laryngocarcinoma, and ACC2 expression level is positively associated with clinical cancer stage and a decreased 5-yr survival rate (Li et al., 2019a). During nutrient abundance, prolyl hydroxylase domain protein 3 (PHD3) hydroxylates and activates ACC2, thereby repressing FAO. In acute myeloid leukemia (AML), PHD3 expression is dramatically decreased, contributing to a boost of FAO that drives AML cell proliferation and disease severity (German et al., 2016).

**FASN.** FASN condenses one molecule of acetyl-CoA and seven malonyl-CoA molecules into the 16-carbon palmitate (C16:0; Fig. 2). FASN overexpression and hyperactivity commonly occur in many human epithelial cancers and their preneoplastic lesions and are correlated with a higher risk of both cancer recurrence and death (Menendez and Lupu, 2007). FASN expression has been shown to be stimulated by PTEN loss, steroid hormones, and the activation of EGFR and ERBB2 in a manner dependent on the activation of the PI3K-AKT and ERK1/2 signaling cascades (Menendez and Lupu, 2007; Yang et al., 2002b). Tumor-associated FASN overexpression is preferentially regulated at the transcription level by SREBP1 (Rawson, 2003). Posttranslational regulation of FASN was exhibited by its interaction with androgen-regulated and overexpressed USP2a in prostate cancer and USP30 in HCC cells, both of which deubiquitinate and stabilize FASN (Graner et al., 2004; Gu et al., 2020). FASN can be acetylated by KAT8 and deacetylated by HDAC3. FASN acetylation enhances its association with the ubiquitin ligase TRIM21 for degradation and is frequently reduced in human HCC samples with up-regulated HDAC3

expression (Lin et al., 2016). The tumor suppressor speckle-type POZ protein is another ubiquitin ligase for FASN ubiquitylation. The speckle-type POZ protein mutants commonly found in prostate cancer cannot bind to FASN, thereby up-regulating FASN expression and lipid accumulation in prostate cancer cells (Gang et al., 2019). FASN stability is also regulated by its SUMOylation, which protects FASN against proteasomal degradation in breast cancer cells (Floris et al., 2020).

FASN inhibition decreases FA synthesis and induces malonyl-CoA accumulation to inhibit CPT1-mediated FAO and cause subsequent cell-cycle arrest and apoptosis of tumor cells (Bandyopadhyay et al., 2006; Menendez and Lupu, 2007). FASN inhibition alters membrane phospholipid composition, structure, and fluidity and lipid raft formation, thereby impairing the correct localization and/or functioning of tyrosine kinase receptors, such as EGFR and ERBB2 (Menendez and Lupu, 2007). FASN inhibition also dampens palmitate-dependent palmitoylation and activation of plasma membrane- and mitochondria-associated EGFR, resulting in EGFR ubiquitylation, abrogation of EGFR activation-induced mitochondrial fusion, blunted tumor growth, and increased sensitivity of cancer cells to EGFR inhibitors (Ali et al., 2018; Bollu et al., 2014). Overexpression of FASN reduces the expression of the E-twenty six (ets)-DNA-binding protein PEA3, a negative regulator of HER2 gene transcription, and consequently increases HER2 mRNA expression (Menendez et al., 2004; Xing et al., 2000). Although intrahepatic cholangiocarcinoma showed insensitivity to FASN depletion, an outcome likely attributed to the high expression of FA uptake-related proteins and robust long-chain FA uptake in intrahepatic cholangiocarcinoma cells, ablation of FASN significantly delayed mouse HCC formation. Notably, FASN deficiency promoted nuclear translocation and activation of SREBP2 for cholesterologenesis. SREBP2 inhibition in combination with FASN ablation completely prevented PTEN deficiency/c-Met-driven mouse hepatocarcinogenesis (Che et al., 2020; Li et al., 2016a), highlighting the critical role of FASN-dependent FA synthesis and SREBP2-mediated cholesterologenesis in HCC development.

**Stearoyl-CoA desaturase (SCD).** SCD (SCD1 and SCD5/hSCD2 in humans) is a rate-limiting enzyme in the ER that catalyzes the formation of a double bond at position  $\Delta 9$  in stearic acid (C18:0) and, to a lesser extent, palmitic acid (C16:0), to generate the monounsaturated FA (MUFA) oleic acid (C18:1) and palmitoleic acid (C16:1), respectively (Snaebjornsson et al., 2020; Fig. 2).

SCD requires NADPH and oxygen to function, further explaining why, under hypoxic conditions, cancer cells rely more on the exogenous supply of unsaturated FA-containing lipids. When cells are deprived of exogenous lipids, the subsequent SCD1 inhibition induced both ferroptosis and apoptosis. Inhibition of SCD1 decreases CoQ10, an endogenous membrane antioxidant whose depletion has been linked to ferroptosis. A concomitant decrease in unsaturated fatty acyl chains in the phospholipids of membranes, including the ER membrane, and an increase in long-chain saturated ceramides leads to the unfolded protein response, ER stress, and apoptosis (Ackerman and Simon, 2014; Tesfay et al., 2019). Inhibited desaturation impairs mitochondrial respiration, which results in oxidative stress, and decreases MUFA incorporation into cardiolipins, a specific class

of lipids exclusive to the inner mitochondrial membrane that bind cytochrome c and prevent its release from the membrane. SCD inhibition coincides with increased cytochrome c release and apoptosis (Peck and Schulze, 2016; Potze et al., 2016).

Activated EGFR in lung cancer cells phosphorylates SCD1 at Y55 to stabilize SCD1 expression leading to increased MUFA levels and lung tumor growth in mice (Zhang et al., 2017a). In Burkitt's lymphoma cells, overexpressed Myc induces the activation of ER stress sensor inositol-requiring enzyme 1 (IRE1) and transcriptional factor X-box binding protein 1 (XBP1) and subsequent SCD1 expression for maintaining ER homeostasis and cell survival (Xie et al., 2018).

SCD1 overexpression was also found in HCC and predicts the clinical response of HCC patients to sorafenib treatment and shorter disease-free survival. Suppression of SCD1 forces liver tumor-initiating cells to differentiate via the ER stress-induced unfolded protein response, resulting in enhanced sensitivity to sorafenib (Ma et al., 2017). SCD1 silencing or inhibition in xenograft mouse models also reduced formation of tumors derived from human stomach, colon, lung, and prostate cancer cells (Snaebjornsson et al., 2020), underscoring critical roles of SCD1 in tumor development.

### Cholesterol biosynthesis enzymes

**Mammalian 3-hydroxy-3-methylglutaryl (HMG)-CoA reductase (HMGCR).** HMGCR, the rate-limiting enzyme of the mevalonate pathway for cholesterol biosynthesis, is an ER-localized glycoprotein and converts HMG-CoA to mevalonate (Fig. 2; Liscum et al., 1985). The regulation of HMGCR is achieved at transcriptional, posttranscriptional, translational, and posttranslational levels. HMGCR gene transcription is activated by SREBP2, whereas its mRNA translation can be blocked by unknown mevalonate-derived nonsterol metabolites (Luo et al., 2020; Nakanishi et al., 1988). In addition, HNRNPA1 regulates the alternative splicing of HMGCR to increase the expression of an alternatively spliced HMGCR transcript lacking exon 13, which exhibits diminished HMGCR activity (Yu et al., 2014). HMGCR degradation can be induced by sterols, mostly oxysterols and methylated sterols, and vitamin E family members  $\delta$ -tocotrienol and  $\gamma$ -tocotrienol. Accumulated sterols in cells stimulate insulin-induced gene (INSIG) protein binding to the sterol-sensing domain of HMGCR and subsequent HMGCR ubiquitylation and degradation involving multiple ubiquitin ligases, the GP78 complex, TRC8 (also known as RNF139), and RNF145 (Luo et al., 2020). Under hypoxic conditions, hypoxia-inducible factor-1 $\alpha$  activates INSIG2 transcription to trigger HMGCR degradation and inhibit cholesterol biosynthesis in the liver (Hwang et al., 2017). AMPK phosphorylates HMGCR S872 and inhibits HMGCR-dependent cholesterol biosynthesis (Clarke and Hardie, 1990; Sato et al., 1993). SIRT1, which can be down-regulated by miR-34a, a microRNA increased in nonalcoholic fatty liver disease, deacetylates and activates LKB1 and LKB1-AMPK signaling for HMGCR inhibition (Min et al., 2012; Ruderman et al., 2010).

HMGCR expression is up-regulated in multiple types of cancer, including gastric cancer, glioblastoma, and prostate cancer. The overexpression of HMGCR promoted the growth and migration of these cancer cells, while HMGCR knockdown


inhibited tumorigenesis. HMGCR inhibition has been targeted for treating solid and blood cancers and cancer with drug resistance (Kong et al., 2018; Lee et al., 2018; Yang et al., 2020). Intriguingly, lipophilic HMGCR inhibitor simvastatin treatment reduced the mevalonate pathway-produced geranylgeranyl diphosphate, which is required for protein prenylation, and the prenylation levels of the small guanosine triphosphatase Rab5 in antigen-presenting cells, resulting in arrested endosomal maturation, prolonged antigen retention, enhanced antigen presentation, and antigen-specific antitumor immunity. In addition, simvastatin treatment robustly enhances cancer vaccinations and synergizes with anti-PD-1 antibodies for tumor treatment (Xia et al., 2018). These results highlight the potential of the mevalonate pathway as a target for boosting cancer immune therapy.

**Squalene monoxygenase (SM).** SM (encoded by *SQLE*), the second rate-limiting ER-associated cholesterol biosynthesis enzyme downstream of HMGCR, converts nonsterol intermediate squalene to 2,3(S)-oxidosqualene (Fig. 2). Similar to HMGCR, SM is also a target of SREBP2 (Sharpe and Brown, 2013). SM has a cholesterol-sensing domain (the first N-terminal 100 amino acids, termed SM N100), which regulates the proteasomal SM degradation in a cholesterol, but not INSIG-, 24,25-dihydrolanosterol-, or side-chain oxysterol-dependent manner (Chua et al., 2017; Gill et al., 2011; Padyana et al., 2019). Cholesterol-accelerated SM degradation is mediated by E3 ligase membrane-associated RING finger 6 (MARCH6), which is concomitantly stabilized by the cholesterol-dependent inhibition of MARCH6 autoubiquitylation (Sharpe et al., 2019; Zelcer et al., 2014). Squalene, as a feedforward factor for cholesterol synthesis, directly binds to the N100 region, thereby reducing the interaction of SM with and ubiquitination by MARCH6 (Yoshioka et al., 2020). In contrast to cholesterol, unsaturated FAs, such as oleate, can stabilize SM by inhibiting MARCH6-mediated degradation (Stevenson et al., 2014), thereby maintaining cholesterol synthesis.

The *SQLE* locus has increased copy numbers in multiple cancers, and its overexpression was detected in nonalcoholic fatty liver disease-induced HCC and linked to radioresistance in pancreatic cancer and progression or poor prognosis of breast cancer, prostate cancer, colorectal cancer, and squamous lung cancer (Brown et al., 2016; Cirmena et al., 2018; Liu et al., 2018). In addition, the proliferation of SCLC cells is sensitive to SM inhibition, which results not from cholesterol biosynthesis inhibition but from the toxic accumulation of the SM substrate squalene (Mahoney et al., 2019). The loss of SM expression, which has been found in anaplastic lymphoma kinase-positive anaplastic large cell lymphoma cells, contributes to cholesterol auxotrophy and renders these cells dependent on LDLR for cholesterol uptake and tumor growth (Garcia-Bermudez et al., 2019). These findings reveal distinct features of cholesterol metabolism in different types of tumor.

### Transcriptional regulation of lipogenesis

Lipogenesis is transcriptionally governed by SREBPs (Fig. 3), a family of helix-loop-helix leucine zipper transcription factors consisting of three isoforms: SREBP1a and SREBP1c encoded by



**Figure 3. Regulation of SREBP1/2 in cancer cells.** SREBP activity can be regulated at multiple levels and at different subcellular localizations. In the ER, sterols bind to SCAP and disrupt the direct interaction between SCAP and COPII for the SREBP ER exit. When sterol level decreases, SCAP dissociates from INSIGs and facilitates the incorporation of SCAP/SREBP into COPII-coated vesicles. mTORC1 suppresses autophagy and subsequent cholesterol trafficking from the lysosome to the ER, leading to SREBP2 activation. Long-chain unsaturated FAs inhibit SREBP activation through inhibition of INSIG1. AKT-phosphorylated PCK1 phosphorylates INSIG1/2 and disrupts the bindings of oxysterols to INSIG1/2 for SREBP1/2 activation. In addition, activated AMPK can phosphorylate SREBP1/2 for their retention in the ER. EGFR activation enhances N-glycosylation of SCAP, triggering its dissociation from INSIG1. In the Golgi, SREBP1/2 are cleaved by S1P and S2P, releasing the transcriptionally active SREBP1/2. HSP90 facilitates the SREBP-SCAP complex transit from the ER to the Golgi. PAQR3 potentiates SREBP processing in the Golgi, whereas TAK1-mediated phosphorylation of SREBP1/2 inhibits SREBP. In the nucleus, truncated SREBP1/2 bind to SREs within the promoters of their target genes. GSK3-phosphorylated SREBP1/2 undergoes ubiquitylation and degradation, which can be counteracted by acetylation of the ubiquitylated Lys residues of SREBP1/2. LD, lipid droplet.

the *SREBF1* gene, and SREBP2 encoded by the *SREBF2* gene. SREBP-1c is ubiquitously expressed, whereas SREBP-1a is expressed in intestinal epithelium, the heart, and macrophages, and SREBP2 is expressed in the liver and adipose tissue. SREBP1 mainly regulates the expression of FA synthesis genes and *LDLR*, whereas SREBP2 preferentially controls cholesterol biosynthesis gene expression (Horton et al., 2003; Im et al., 2011).

Inactive SREBPs reside in the ER membrane, where their C-terminal domains interact with the WD-repeat domain in the C terminus of SREBP cleavage-activating protein (SCAP; Gong et al., 2015). The N-terminal domain of SCAP binds to INSIG1 and INSIG2, forming an INSIG/SCAP/SREBP complex that retains SREBPs in the ER (Yabe et al., 2002; Yang et al., 2002a). The binding of coatamer II (COPII) to SCAP leads to the translocation of the SCAP-SREBP complex from the ER to the Golgi, where SREBPs are sequentially cleaved by membrane-bound site-1 protease (S1P) and site-2 protease (S2P), releasing the transcriptionally active N-terminal domains. Mature SREBPs then translocate to the nucleus and bind as homodimers to SREs and E-boxes within their target gene promoters (Nohturfft and Zhang, 2009). SREBP activity can be regulated at multiple levels and in different subcellular localizations, such as the ER, Golgi, and nucleus.

**Regulation of SREBPs in the ER.** SREBP activation is regulated by sterol fluctuations in the ER. Cholesterol binds to SCAP, thereby disrupting the interaction between SCAP and COPII and retaining SREBP in the ER (Shimano and Sato, 2017). Oxysterols, such as 25-hydroxycholesterol, are much more potent

than cholesterol in the SREBP's ER retention by binding to INSIGs and promoting INSIG binding to SCAP (Eberlé et al., 2004). When the sterol level decreases, SCAP dissociates from INSIGs, which facilitates the incorporation of SCAP/SREBP into COPII-coated vesicles (Menendez and Lupu, 2007). p53 transcriptionally induces the ATP-binding cassette transporter (ABCA1), a cholesterol transporter gene. p53 loss or ABCA1 ablation decreases the retrograde transportation of cholesterol from the plasma membrane to the ER, thereby promoting SREBP2 maturation and mouse liver tumorigenesis (Moon et al., 2019). ER-residing ACAT esterifies ER cholesterol with FAs to form cholesteryl esters for the storage in lipid droplets, thereby controlling ER cholesterol levels. Inhibition of ACAT accumulates ER cholesterol, reduces lipid droplet formation, and suppresses glioblastoma growth and the aggressiveness of prostate cancer cells by blocking SREBP1-regulated gene expression (Geng et al., 2016; Yue et al., 2014).

mTORC1 suppresses autophagy and maintains endosomal recycling to the plasma membrane, thereby preventing membrane-derived cholesterol from reaching lysosomes and subsequent ER localization, leading to SREBP2 activation (Eid et al., 2017). Long-chain unsaturated FAs inhibit SREBP activation through interaction with ubiquitin regulatory X domain-containing protein 8 (UBXD8). UBXD8 binds to polyubiquitylated INSIG1, in turn recruiting the valosin-containing protein complex, which facilitates INSIG1 degradation. Long-chain unsaturated FAs induce the detachment of UBXD8 from INSIG1, thereby stabilizing INSIG1 (Lee et al., 2008; Lee et al., 2010).

The INSIG/SCAP/SREBP complex can also be regulated in lipid-independent manners. Receptor tyrosine kinase activation and active K-RAS mutations in cancer cells induce AKT-mediated phosphorylation of S90 of cytosolic phosphoenolpyruvate carboxykinase 1 (PCK1), resulting in inhibition of the gluconeogenic activity of PCK1 and its translocation to the ER, where it acts as a protein kinase phosphorylating INSIG1 S207 and INSIG2 S151. Phosphorylated INSIGs reduce their binding to oxysterols, thereby activating SREBP-dependent lipogenesis for liver tumor formation. In addition, PCK1-mediated INSIG phosphorylation is associated with poor HCC prognoses (Xu et al., 2020). EGFR activation also leads to enhanced SCAP N-glycosylation to trigger its dissociation from INSIG1 to induce lipogenesis and glioblastoma growth (Cheng et al., 2015). In addition to regulation by the AKT-PCK1 axis, SREBP1 activation by AKT- and mTORC1-dependent or mTORC1-independent mechanisms was revealed (Düvel et al., 2010; Guo et al., 2009; Yecies et al., 2011). CREB-regulated transcription coactivator 2 (CRTC2) competes with the COPII complex subunit Sec23A to interact with Sec31A, thereby disrupting SREBP1 transport, and this inhibitory effect was attenuated by mTOR-mediated CRTC2 phosphorylation (Han et al., 2015). It was shown that insulin signaling and AKT activation suppress INSIG2 expression by promoting its mRNA decay (Yecies et al., 2011). In addition, INSIG1, but not INSIG2, is ubiquitinated by GP78 and degraded upon sterol depletion (Lee et al., 2006). INSIG1 degradation, which is also induced in HCC cells in an AKT-mediated INSIG1 S207 phosphorylation-dependent manner, results in SREBP activation-increased transcription of downstream genes, including *INSIG1* itself (Shao and Espenshade, 2012; Xu et al., 2020). The newly synthesized INSIG1 continues to be degraded until a sufficient amount of cholesterol is produced to induce a conformational change of SCAP that enables INSIG1 binding in normal cells (Gong et al., 2006). However, switching off SREBP activation by the feedback-produced cholesterol and INSIG1 protein is disrupted by INSIG1 S207 phosphorylation in tumor cells, and thus a high level of lipogenesis is maintained (Xu et al., 2020). In contrast to AKT-mediated INSIG phosphorylation, activated AMPK phosphorylates SREBPs for their potential retention in the ER, thereby suppressing lipogenesis in HepG2 liver cancer cells (Li et al., 2011). In addition, the SCAP-SREBP2-INSIG complex can be also retained in the ER upon its binding to the LATS2 tumor suppressor, interaction with ERLINs and RNF139 independent of ubiquitylation, or the RNF145-dependent SCAP ubiquitylation that prevents COPII binding (Aylon et al., 2016; Huber et al., 2013; Irisawa et al., 2009; Zhang et al., 2017b). Mice harboring liver-specific *Lats2* knockout spontaneously developed fatty liver disease in association with impaired p53 activation (Aylon et al., 2016).

Thus, the INSIG/SCAP/SREBP complex in the ER can be regulated by sterol and FA fluctuations; posttranslational modifications of INSIG, SCAP, and SREBP; mRNA and protein stability of INSIG; and the protein interaction that affects the binding of SCAP to COPII.

**Regulation of SREBPs in the Golgi apparatus.** Heat shock protein 90 (HSP90) binds to and stabilizes the SREBP-SCAP complex in both the ER and the Golgi and facilitates its transit to


the Golgi, whereas HSP90 inhibition leads to proteasomal degradation of the complex (Kuan et al., 2017). The Golgi-anchoring protein progesterin and adipoQ receptor 3 (PAQR3), which is transcriptionally induced under cholesterol-depleting conditions, interacts with SCAP and SREBP and tethers them to the Golgi to potentiate SREBP processing and enhance lipid synthesis (Xu et al., 2015). ER- or Golgi-localized SREBPs can be phosphorylated and inhibited by TAK1 without blocking their nuclear localization, and hepatic *Tak1* deficiency causes steatosis (Morioka et al., 2016).

**Regulation of nuclear SREBPs.** mTORC1 promotes the nuclear localization of mature SREBP1 through the phosphorylation and cytoplasmic retention of the phosphatidate phosphatase LPIN1, which, in its unphosphorylated state, inhibits SREBP1 by sequestering it at the nuclear periphery (Peterson et al., 2011). Nuclear SREBP can be degraded by the SCF<sup>FBXW7</sup> ubiquitin ligase dependent on GSK3-mediated SREBP phosphorylation (Sundqvist et al., 2005). This phosphorylation is prevented by SREBP1a R321 dimethylation by protein arginine methyltransferase 5 (PRMT5), resulting in increased lipogenesis, accelerated tumor cell proliferation, and poor prognosis for HCC patients (Liu et al., 2016). The SREBP degradation can be also inhibited by the acetylation of the ubiquitylated Lys residue by p300 and its related protein CBP (Giandomenico et al., 2003), and this inhibition can be abrogated by SIRT1 deacetylase (Walker et al., 2010). In addition, ERK-mediated phosphorylation of nuclear SREBP2 at S432 and S455 enhances SREBP2 activity, whereas SUMO-1-conjugating enzyme Ubc9-dependent SREBP SUMOylation decreases its transcriptional activity (Hirano et al., 2003; Kotzka et al., 2004).

The vicinity of the SRE motif of the promoter regions of lipogenic genes, including *SREBF2*, contains binding sites for transcription factors SP1 and/or NFY (Shimano and Sato, 2017). SP1 or NFY interacts with SREBPs and coordinates the expression of a subset of SREBP target genes. Hepatocyte nuclear factor 4 (HNF4) and PGC1 $\beta$  up-regulate SREBP activity also via direct binding (Lin et al., 2005; Misawa et al., 2003). Upon ER stress or glucose deprivation, ATF6 interacts with SREBP2, resulting in the recruitment of HDAC1 to the ATF6-SREBP2 complex and the inhibition of SREBP2 (Zeng et al., 2004). Thus, the activity of nuclear SREBPs can be regulated by phosphorylation, acetylation, ubiquitylation, and protein-protein interactions.

**Transcriptional regulation of SREBPs and the roles of LXRs in lipid homeostasis.** The *SREBF1* and *SREBF2* promoters each contain an SRE for their own autoloop activation, which generates positive feedback self-regulation upon SREBP activation (Shimano and Sato, 2017). In addition, LXR $\alpha$  and LXR $\beta$  form heterodimers with retinoid X receptor and bind to conserved LXR response elements in the promoters of *SREBF1* in the presence of different nuclear oxysterols, which function as LXR ligands, to promote lipid synthesis (Lin and Gustafsson, 2015; Repa et al., 2000). The expression of SREBP2 can be inhibited by the binding of FOXO3 to the *SREBF2* promoter, leading to the recruitment of SIRT6 to deacetylate histone H3 (Tao et al., 2013). In addition to transcriptional regulation, the negative regulation of SREBPs at the mRNA level by miR-29, miR-185, and miR-342 has also been revealed (Cheng et al., 2018).





**Figure 4. The FAO pathway.** FAs are activated to fatty acyl-CoA by fatty acyl-CoA synthetase. On the outer mitochondrial membrane, fatty acyl-CoA is converted to fatty acylcarnitine by CPT1 and shuttled into the mitochondrial matrix. CPT2 on the matrix side of the inner membrane then reconverts acylcarnitine to acyl-CoA, which is cleaved into acetyl-CoA by a repeated four-step cycle catalyzed sequentially by the activity of acyl-CoA dehydrogenase, enoyl-CoA hydratase, 3-hydroxyacyl CoA dehydrogenase, and 3-ketoacyl-CoA thiolase, resulting in the shortening of FAs by two carbons in each cycle. The breakdown product acetyl-CoA enters the TCA cycle, and the produced NADH and FADH<sub>2</sub> are coenzymes used in the electron transport chain to generate ATP. CACT, carnitine/acylcarnitine translocase; IMM, inner mitochondrial membrane; OMM, outer mitochondrial membrane.

The role of LXRs in lipid metabolism in cancer cells is complicated by the expression of their multiple downstream genes, and LXR has been shown to exert positive or negative regulation on the growth and survival of different tumor cells (Flaveny et al., 2015; Guo et al., 2011). LXRs can reduce intracellular cholesterol levels by enhancing cholesterol efflux mediated by up-regulated ABCA1 expression, and decreasing cholesterol uptake mediated by increased expression of inducible degrader of the LDLR (Idol), a ubiquitin ligase for LDLR degradation (Lin and Gustafsson, 2015; Repa et al., 2000; Zelcer et al., 2009). In contrast, in addition to activating SREBF1 expression, LXRs can increase lipogenesis by inducing FASN and SCD1 expression and enhances glycolysis by inducing phosphofructokinase-2 (PFK2) and glucokinase (GCK1) expression (Kim et al., 2009; Tao et al., 2013; Zhao et al., 2012). Given that LXR activation also induces the expression of cell cycle regulators and genes involved in immune cell functions, studying LXR in cancer in intact tumor immune microenvironments may elucidate more insightful information.

### Lipolysis and FAO

FAO allows the mitochondrial conversion of long-chain FAs into acetyl-CoA, which enters the TCA cycle, and NADH and FADH<sub>2</sub>, which are coenzymes used in the electron transport chain (Fig. 4). FAO occurs in peroxisomes when the FA chains are too long to be oxidized in mitochondria. On the outer mitochondrial membrane, fatty acyl-CoA is converted to fatty acylcarnitine by CPT1, for which the isoforms CPT1A, CPT1B, and CPT1C are predominantly expressed in the liver, muscle, and brain, respectively. Carnitine/acylcarnitine translocase located on the inner mitochondrial membrane shuttles acylcarnitine into the mitochondrial matrix. CPT2 on the matrix side of the inner

membrane reconverts acylcarnitine to acyl-CoA for being cleaved into acetyl-CoA, which enters the TCA cycle to generate ATP and malic enzyme-dependent NADPH (Fig. 4). In addition, FAO-generated citrate can be exported to the cytoplasm for NADPH production by IDH-mediated isocitrate oxidation. NADPH is a reducing agent that supports biosynthesis and redox homeostasis (Carracedo et al., 2013).

Many types of cancer, such as lung cancer with K-Ras mutation, triple-negative breast cancer, and glioma, exhibit high FAO activity (Carracedo et al., 2013; Padanad et al., 2016). Consistently, the overexpression of various FAO pathway proteins, including CPT1A, CPT1B, CPT1C, CPT-2, carnitine transporter CT2, and acyl-CoA synthetase long-chain family member 3 (ACSL3), is found in multiple types of cancers, while some FAO enzymes, such as CPT1A, are highly correlated with poor outcomes patients with AML or ovarian cancer (Carracedo et al., 2013). CPT1C, expressed normally in the brain, is up-regulated in human lung tumors, depending on AMPK, and promotes FAO, ATP production, and the resistance of tumor cells to energy stress (Zaugg et al., 2011).

The most prominent transcriptional regulators of FAO are PPARs (Poulsen et al., 2012). Promyelocytic leukemia protein, which is overexpressed in triple-negative breast cancer, activates PPAR and FAO by reducing PGC-1 $\alpha$  acetylation (Carracedo et al., 2012). Jumonji D3 (JMJD3) histone demethylase, interacting with SIRT1 and PPAR $\alpha$ , decreases H3K27me3 at FAO genes and thereby epigenetically activates the expression of these genes (Seok et al., 2018). Acting as a transcription factor, HNF4 directly activates FAO genes and is required for the renewal of intestinal stem cells (Chen et al., 2020). In breast cancer cells, c-Myc/PGC-1 $\beta$ /ERR $\alpha$  signaling induces FAO enzyme expression (Yan et al., 2017), and FAO inhibition blocked MYC-driven

breast tumor growth in mice (Camarda et al., 2016). In addition, mammary adipocyte-derived leptin up-regulates STAT3-induced CPT1B expression and FAO activity in breast cancer stem cells, which promotes cancer cell stemness and chemoresistance (Wang et al., 2018). Similar to CPT1 overexpression, acyl-CoA-binding protein (also known as diazepam binding inhibitor [DBI]) is highly expressed in glioblastoma and binds to acyl-CoAs to promote mitochondrial long-chain fatty acyl-CoA accumulation, FAO, and tumor growth (Duman et al., 2019). As the first and rate-limiting enzyme in FAO and a major producer of H<sub>2</sub>O<sub>2</sub> in peroxisomes, acyl-CoA oxidase 1 (ACOX1) exhibits increased succinylation and activity due to the down-regulated expression of SIRT5 desuccinylase, which enhances DNA damage response to oxidative stress in HCC cells (Chen et al., 2018). FAO can also be stimulated by the AMPK activation, which reduces ACC-produced malonyl CoA, an allosteric CPT1 inhibitor (Jeon et al., 2012). CircACCI, an ACC1-derived circular RNA with elevated expression in human colon cancer, regulates the assembly and activation of the AMPK complex under metabolic stress, thereby facilitating FAO and xenograft tumor growth (Li et al., 2019b).

In addition to its role in ATP production and cell proliferation, FAO is critical for NADPH homeostasis, mitochondrial function, and cell survival. FAO inhibition decreases NADPH levels with correspondingly increased ROS levels and apoptosis rates of glioma cells (Pike et al., 2011). During energy stress, AMPK activation increases NADPH production through FAO and inhibits NADPH consumption in FA synthesis, thereby inhibiting lung cancer cell death (Jeon et al., 2012).

MCL-1, which is a BCL-2 family protein linked to a FAO signature in AML, interacts with very-long-chain acyl-CoA dehydrogenase and promotes long-chain FA  $\beta$ -oxidation (Escudero et al., 2018). Truncated BH3 interacting domain death agonist, a proapoptotic mitochondrial protein, decreases CPT-1 activity and inhibits FAO, while overexpressed CPT-1 interacts with Bcl-2 and counteracts the effects exerted by BH3 interacting domain death agonist on FAO. Inhibition of FAO promotes Bak and Bax oligomerization, mitochondrial permeability transition, cytotoxic accumulation of long-chain FAs, ER stress, and subsequent apoptosis (Giordano et al., 2005; Ma et al., 2018; Samudio et al., 2010).

However, overly active FAO can be cytotoxic. In glioblastoma cells, highly expressed diacylglycerol-acyltransferase 1 (DGAT1) converts diacylglycerol and excessive fatty acyl CoA to triglycerides, which are stored in lipid droplets. Inhibiting DGAT1 resulted in excessive FAO and ROS production and apoptosis (Cheng et al., 2020). DGAT1/2 depletion also increased saturated toxic FAs in clear cell renal cell carcinoma cells and impaired mouse tumor growth (Ackerman et al., 2018; Chitraju et al., 2017). In obesity- and nonalcoholic steatohepatitis-driven HCC, CPT2 down-regulation prevents lipotoxicity in HCC cells and promotes liver tumorigenesis (Fujiwara et al., 2018). Thus, dynamic and balanced control of FAO supports tumor cell proliferation and survival by providing the needed ATP and NADPH, eliminating potentially toxic lipids, inhibiting proapoptotic pathways, and producing the metabolic intermediates required for anabolism.

## Targeting lipid metabolism for cancer treatment

The integrated and mutual regulation between oncogenic signaling and lipid metabolism promotes cancer cell growth, survival, proliferation, migration, invasion, and metastasis. Lipid metabolism in cancer cells and other cells in the tumor microenvironment, including immune cells, adipocytes, endothelial cells, and fibroblasts, is dynamically regulated and interconnected, and great efforts have been made to develop therapeutic drugs to intervene in lipid metabolism at different levels. Numerous preclinical studies on many inhibitors of a variety of lipid metabolism enzymes have been reported (Koundouros and Pouligiannis, 2020; Ma et al., 2018; Röhrig and Schulze, 2016) and will not be covered in detail here. Instead, advanced studies, especially clinical trials, are discussed.

To target cholesterol synthesis, statin family drugs, such as HMGCR inhibitors, are currently being tested as anticancer agents in multiple clinical trials (Mullen et al., 2016). Some inconsistent results were reported. Retrospective studies showed that statin treatment prolonged the survival of patients with multiple myeloma, colorectal cancer, and metastatic pancreatic cancer cotreated with first-line chemotherapy (Abdel-Rahman, 2019; Brånvall et al., 2020; Cardwell et al., 2014). In contrast, randomized phase III trials in patients with SCLC, metastatic colorectal cancer, HCC, or gastric cancer showed that addition of pravastatin or simvastatin to standard chemotherapy offers no additional benefit (Jouve et al., 2019; Kim et al., 2014; Lim et al., 2015; Seckl et al., 2017). Notably, a phase II study revealed that simvastatin in combination with the EGFR inhibitor gefitinib, but not afatinib, resulted in higher tumor response rates and longer progression-free survival than did EGFR inhibitor alone in NSCLC patients (Han et al., 2011; Lee et al., 2017). It was suggested that the lipophilic statin drugs more readily enter extrahepatic cells, whereas hydrophilic statins are more hepatoselective (Duncan et al., 2006). In addition, clinical data indicate that the anticancer effects of statins are both dose and time dependent (Kim et al., 2014). Thus, identification of predictive biomarkers for patient stratification and selection of appropriate statin type, dosage, and treatment durations likely provide more definitive evaluations of statins as adjuvant treatments.

Nelfinavir, an S2P inhibitor that has been used in HIV treatment, was administered concurrently with chemoradiotherapy during early-phase clinical trials, and the results showed encouraging antitumor activity and acceptable safety in patients with NSCLC, inoperable pancreatic cancer, or multiple myeloma, although the exact contribution of lipid synthesis inhibition by nelfinavir to the anticancer effect warrants further study (Das, 2019; Driessen et al., 2016; Rengan et al., 2012; Wilson et al., 2016).

To target FA synthesis, the FASN inhibitor TVB-2640 is currently being evaluated in phase II clinical trials as a single agent in NSCLC with KRAS multination (NCT03808558), in combination with paclitaxel and trastuzumab in triple-negative breast cancer (NCT03179904), or with the anti-angiogenic drug bevacizumab in high-grade astrocytoma (NCT03032484). Preclinical animal studies showed that ACC inhibitors ND-646 and ND-654 markedly suppressed growth of mouse lung tumor and

rat HCC, respectively (Lally et al., 2019; Svensson et al., 2016). In addition, the ACC inhibitor ND-630, originally developed for the treatment of nonalcoholic steatohepatitis, is currently undergoing phase I testing as a cancer treatment (NCT02876796; Snaebjornsson et al., 2020).

Given that lipid uptake and synthesis both contribute to lipid resources in cancer cells, simple treatment strategies inhibiting FA or cholesterol biosynthesis may be less effective due to the compensation from dietary lipids. In addition, the dynamic FAO regulation exerts critical roles in cancer progression. Thus, more specific and combinational approaches for the intervention of lipid uptake, synthesis, and lipolysis, such as FAO, are needed. Tumor lipid metabolism in each patient can manifest unique features and render profiles showing specific regulation with distinct genetic fingerprints determined by genetic mutations and epigenetic gene regulation in each type and subtype of cancer. A rational combination of conventional chemotherapy and/or radiation treatment with immunotherapy and targeted therapies, including lipid metabolism-targeted approaches, should be investigated for developing more efficient cancer treatments that do not elicit drug resistance.

### Conclusion and perspectives

Our current understanding of the aberrant regulation of lipid metabolism in cancer indicates increased comprehension of the metabolic wiring of cancer cells. Lipid metabolism in cancer cells can be regulated not only through intracellular oncogenic signaling but also by the input from the tumor microenvironment composed of various types of cells, cytokines, growth factors, DNA, RNA, and nutrients, including lipids. In turn, aberrant lipid metabolism reroutes oncogenic signaling pathways in cancer cells and affects neighboring normal cell populations through secretory components, including lipids. This complexity highlights the need for studying not only the lipid metabolism network in cancer cells but also the interconnected pathways in the tumor microenvironment and the effects of interfering with lipid metabolism in the cells in the tumor microenvironment on tumor progression and treatment responses, especially the antitumor immune and anti-angiogenesis responses. In addition, structural elucidation of oncogenic signaling-induced and posttranslationally modified lipid enzymes, as well as lipid metabolism-regulating enzymes, will facilitate the identification of specific interventions that target aberrantly regulated, but not normal, lipid metabolism. Insight into the tumor-specific regulation of lipid metabolism will reveal new and exciting therapeutic opportunities for cancer elimination with minimal side effects.

### Acknowledgments

This study was supported by Ministry of Science and Technology of the People's Republic of China grants (2020YFA0803300 to Z. Lu), the Zhejiang University Research Fund (188020\*194221901/029 to Z. Lu), the Leading Innovative and Entrepreneur Team Introduction Program of Zhejiang (2019R01001 to Z. Lu), the National Natural Science Foundation of China (81902880 to X. Bian), the Qingdao Postdoctoral Application Research Project

(to X. Bian), and the China Postdoctoral Science Foundation (2019M660160 to X. Bian). Z. Lu is the Kuancheng Wang Distinguished Chair.

Author contributions: All authors researched data for the article, contributed to the discussion of the content, wrote the article, and reviewed and/or edited the manuscript before submission.

Disclosures: The authors declare no competing interests exist.

Submitted: 24 July 2020

Revised: 15 October 2020

Accepted: 26 October 2020

### References

- Abdel-Rahman, O. 2019. Statin treatment and outcomes of metastatic pancreatic cancer: a pooled analysis of two phase III studies. *Clin. Transl. Oncol.* 21:810–816. <https://doi.org/10.1007/s12094-018-1992-3>
- Ackerman, D., and M.C. Simon. 2014. Hypoxia, lipids, and cancer: surviving the harsh tumor microenvironment. *Trends Cell Biol.* 24:472–478. <https://doi.org/10.1016/j.tcb.2014.06.001>
- Ackerman, D., S. Tumanov, B. Qiu, E. Michalopoulou, M. Spata, A. Azzam, H. Xie, M.C. Simon, and J.J. Kamphorst. 2018. Triglycerides Promote Lipid Homeostasis during Hypoxic Stress by Balancing Fatty Acid Saturation. *Cell Rep.* 24:2596–2605.e5. <https://doi.org/10.1016/j.celrep.2018.08.015>
- Ali, A., E. Levantini, J.T. Teo, J. Goggi, J.G. Clohessy, C.S. Wu, L. Chen, H. Yang, I. Krishnan, O. Kocher, et al. 2018. Fatty acid synthase mediates EGFR palmitoylation in EGFR mutated non-small cell lung cancer. *EMBO Mol. Med.* 10:e8313. <https://doi.org/10.15252/emmm.201708313>
- Altmann, S.W., H.R. Davis Jr., L.J. Zhu, X. Yao, L.M. Hoos, G. Tetzloff, S.P. Iyer, M. Maguire, A. Golovko, M. Zeng, et al. 2004. Niemann-Pick C1 Like 1 protein is critical for intestinal cholesterol absorption. *Science.* 303:1201–1204. <https://doi.org/10.1126/science.1093131>
- Auciello, F.R., V. Bulusu, C. Oon, J. Tait-Mulder, M. Berry, S. Bhattacharyya, S. Tumanov, B.L. Allen-Petersen, J. Link, N.D. Kendersky, et al. 2019. A Stromal Lysolipid-Autotaxin Signaling Axis Promotes Pancreatic Tumor Progression. *Cancer Discov.* 9:617–627. <https://doi.org/10.1158/2159-8290.CD-18-1212>
- Aylon, Y., A. Gershoni, R. Rotkopf, I.E. Biton, Z. Porat, A.P. Koh, X. Sun, Y. Lee, M.I. Fiel, Y. Hoshida, et al. 2016. The LATS2 tumor suppressor inhibits SREBP and suppresses hepatic cholesterol accumulation. *Genes Dev.* 30:786–797. <https://doi.org/10.1101/gad.274167.115>
- Bandyopadhyay, S., R. Zhan, Y. Wang, S.K. Pai, S. Hirota, S. Hosobe, Y. Takano, K. Saito, E. Furuta, M. Iizumi, et al. 2006. Mechanism of apoptosis induced by the inhibition of fatty acid synthase in breast cancer cells. *Cancer Res.* 66:5934–5940. <https://doi.org/10.1158/0008-5472.CAN-05-3197>
- Bensaad, K., E. Favaro, C.A. Lewis, B. Peck, S. Lord, J.M. Collins, K.E. Pinnick, S. Wigfield, F.M. Buffa, J.L. Li, et al. 2014. Fatty acid uptake and lipid storage induced by HIF-1 $\alpha$  contribute to cell growth and survival after hypoxia-reoxygenation. *Cell Rep.* 9:349–365. <https://doi.org/10.1016/j.celrep.2014.08.056>
- Berwick, D.C., I. Hers, K.J. Heesom, S.K. Moule, and J.M. Tavare. 2002. The identification of ATP-citrate lyase as a protein kinase B (Akt) substrate in primary adipocytes. *J. Biol. Chem.* 277:33895–33900. <https://doi.org/10.1074/jbc.M204681200>
- Bollu, L.R., J. Ren, A.M. Blessing, R.R. Katreddy, G. Gao, L. Xu, J. Wang, F. Su, and Z. Weihua. 2014. Involvement of de novo synthesized palmitate and mitochondrial EGFR in EGF induced mitochondrial fusion of cancer cells. *Cell Cycle.* 13:2415–2430. <https://doi.org/10.4161/cc.29338>
- Brånvall, E., S. Ekberg, S. Eloranta, T. Wåsterlid, B.M. Birman, and K.E. Smedby. 2020. Statin use is associated with improved survival in multiple myeloma: A Swedish population-based study of 4315 patients. *Am. J. Hematol.* 95:652–661. <https://doi.org/10.1002/ajh.25778>
- Brown, D.N., I. Caffa, G. Cirmena, D. Piras, A. Garuti, M. Gallo, S. Alberti, A. Nencioni, A. Ballestrero, and G. Zoppi. 2016. Squalene epoxidase is a bona fide oncogene by amplification with clinical relevance in breast cancer. *Sci. Rep.* 6:19435. <https://doi.org/10.1038/srep19435>
- Brusselmans, K., E. De Schrijver, G. Verhoeven, and J.V. Swinnen. 2005. RNA interference-mediated silencing of the acetyl-CoA-carboxylase- $\alpha$

- gene induces growth inhibition and apoptosis of prostate cancer cells. *Cancer Res.* 65:6719–6725. <https://doi.org/10.1158/0008-5472.CAN-05-0571>
- Bulusu, V., S. Tumanov, E. Michalopoulou, N.J. van den Broek, G. MacKay, C. Nixon, S. Dhayade, Z.T. Schug, J. Vande Voorde, K. Blyth, et al. 2017. Acetate Recapturing by Nuclear Acetyl-CoA Synthetase 2 Prevents Loss of Histone Acetylation during Oxygen and Serum Limitation. *Cell Rep.* 18:647–658. <https://doi.org/10.1016/j.celrep.2016.12.055>
- Camarda, R., A.Y. Zhou, R.A. Kohnz, S. Balakrishnan, C. Mahieu, B. Anderton, H. Eyob, S. Kajimura, A. Tward, G. Krings, et al. 2016. Inhibition of fatty acid oxidation as a therapy for MYC-overexpressing triple-negative breast cancer. *Nat. Med.* 22:427–432. <https://doi.org/10.1038/nm.4055>
- Cardwell, C.R., B.M. Hicks, C. Hughes, and L.J. Murray. 2014. Statin use after colorectal cancer diagnosis and survival: a population-based cohort study. *J. Clin. Oncol.* 32:3177–3183. <https://doi.org/10.1200/JCO.2013.54.4569>
- Carracedo, A., D. Weiss, A.K. Leliaert, M. Bhasin, V.C. de Boer, G. Laurent, A.C. Adams, M. Sundvall, S.J. Song, K. Ito, et al. 2012. A metabolic prosurvival role for PML in breast cancer. *J. Clin. Invest.* 122:3088–3100. <https://doi.org/10.1172/JCI62129>
- Carracedo, A., L.C. Cantley, and P.P. Pandolfi. 2013. Cancer metabolism: fatty acid oxidation in the limelight. *Nat. Rev. Cancer.* 13:227–232. <https://doi.org/10.1038/nrc3483>
- Carrer, A., S. Trefely, S. Zhao, S.L. Campbell, R.J. Norgard, K.C. Schultz, S. Sidoli, J.L.D. Parris, H.C. Affronti, S. Sivanand, et al. 2019. Acetyl-CoA Metabolism Supports Multistep Pancreatic Tumorigenesis. *Cancer Discov.* 9:416–435. <https://doi.org/10.1158/2159-8290.CD-18-0567>
- Che, L., W. Chi, Y. Qiao, J. Zhang, X. Song, Y. Liu, L. Li, J. Jia, M.G. Pilo, J. Wang, et al. 2020. Cholesterol biosynthesis supports the growth of hepatocarcinoma lesions depleted of fatty acid synthase in mice and humans. *Gut.* 69:177–186. <https://doi.org/10.1136/gutjnl-2018-317581>
- Chen, X.F., M.X. Tian, R.Q. Sun, M.L. Zhang, L.S. Zhou, L. Jin, L.L. Chen, W.J. Zhou, K.L. Duan, Y.J. Chen, et al. 2018. SIRT5 inhibits peroxisomal ACOX1 to prevent oxidative damage and is downregulated in liver cancer. *EMBO Rep.* 19:e45124. <https://doi.org/10.15252/embr.201745124>
- Chen, L., R.P. Vasoya, N.H. Toke, A. Parthasarathy, S. Luo, E. Chiles, J. Flores, N. Gao, E.M. Bonder, X. Su, and M.P. Verzi. 2020. HNF4 Regulates Fatty Acid Oxidation and Is Required for Renewal of Intestinal Stem Cells in Mice. *Gastroenterology.* 158:985–999.e9. <https://doi.org/10.1053/j.gastro.2019.11.031>
- Cheng, C., P. Ru, F. Geng, J. Liu, J.Y. Yoo, X. Wu, X. Cheng, V. Euthine, P. Hu, J.Y. Guo, et al. 2015. Glucose-Mediated N-glycosylation of SCAP Is Essential for SREBP-1 Activation and Tumor Growth. *Cancer Cell.* 28:569–581. <https://doi.org/10.1016/j.ccell.2015.09.021>
- Cheng, X., J. Li, and D. Guo. 2018. SCAP/SREBPs are Central Players in Lipid Metabolism and Novel Metabolic Targets in Cancer Therapy. *Curr. Top. Med. Chem.* 18:484–493. <https://doi.org/10.2174/1568026618666180523104541>
- Cheng, X., F. Geng, M. Pan, X. Wu, Y. Zhong, C. Wang, Z. Tian, C. Cheng, R. Zhang, V. Puduvali, et al. 2020. Targeting DGAT1 Ameliorates Glioblastoma by Increasing Fat Catabolism and Oxidative Stress. *Cell Metab.* 32:229–242.e8. <https://doi.org/10.1016/j.cmet.2020.06.002>
- Chin, K., S. DeVries, J. Fridlyand, P.T. Spellman, R. Roydasgupta, W.L. Kuo, A. Lapuk, R.M. Neve, Z. Qian, T. Ryder, et al. 2006. Genomic and transcriptional aberrations linked to breast cancer pathophysiology. *Cancer Cell.* 10:529–541. <https://doi.org/10.1016/j.ccr.2006.10.009>
- Chittraju, C., N. Mejhert, J.T. Haas, L.G. Diaz-Ramirez, C.A. Grueter, J.E. Imbriglio, S. Pinto, S.K. Koliwad, T.C. Walther, and R.V. Farese Jr. 2017. Triglyceride Synthesis by DGAT1 Protects Adipocytes from Lipid-Induced ER Stress during Lipolysis. *Cell Metab.* 26:407–418.e3. <https://doi.org/10.1016/j.cmet.2017.07.012>
- Chua, N.K., V. Howe, N. Jatana, L. Thukral, and A.J. Brown. 2017. A conserved degron containing an amphipathic helix regulates the cholesterol-mediated turnover of human squalene monooxygenase, a rate-limiting enzyme in cholesterol synthesis. *J. Biol. Chem.* 292:19959–19973. <https://doi.org/10.1074/jbc.M117.794230>
- Cirmena, G., P. Franceschelli, E. Isnaldi, L. Ferrando, M. De Mariano, A. Ballestrero, and G. Zoppi. 2018. Squalene epoxidase as a promising metabolic target in cancer treatment. *Cancer Lett.* 425:13–20. <https://doi.org/10.1016/j.canlet.2018.03.034>
- Clarke, P.R., and D.G. Hardie. 1990. Regulation of HMG-CoA reductase: identification of the site phosphorylated by the AMP-activated protein kinase in vitro and in intact rat liver. *EMBO J.* 9:2439–2446. <https://doi.org/10.1002/j.1460-2075.1990.tb07420.x>
- Comerford, S.A., Z. Huang, X. Du, Y. Wang, L. Cai, A.K. Witkiewicz, H. Walters, M.N. Tantawy, A. Fu, H.C. Manning, et al. 2014. Acetate dependence of tumors. *Cell.* 159:1591–1602. <https://doi.org/10.1016/j.cell.2014.11.020>
- Das, M. 2019. Nelfinavir with concurrent chemoradiotherapy in NSCLC. *Lancet Oncol.* 20:e561. [https://doi.org/10.1016/S1470-2045\(19\)30567-4](https://doi.org/10.1016/S1470-2045(19)30567-4)
- Driessen, C., M. Kraus, M. Joerger, H. Rosing, J. Bader, F. Hitz, C. Berset, A. Xyrafas, H. Hawle, G. Berthod, et al. 2016. Treatment with the HIV protease inhibitor nelfinavir triggers the unfolded protein response and may overcome proteasome inhibitor resistance of multiple myeloma in combination with bortezomib: a phase I trial (SAKK 65/08). *Haematologica.* 101:346–355. <https://doi.org/10.3324/haematol.2015.135780>
- Duman, C., K. Yaqubi, A. Hoffmann, A.A. Acikgöz, A. Korshunov, M. Bendzsus, C. Herold-Mende, H.K. Liu, and J. Alfonso. 2019. Acyl-CoA-Binding Protein Drives Glioblastoma Tumorigenesis by Sustaining Fatty Acid Oxidation. *Cell Metab.* 30:274–289.e5. <https://doi.org/10.1016/j.cmet.2019.04.004>
- Duncan, R.E., A. El-Soheby, and M.C. Archer. 2006. Statins and the risk of cancer. *JAMA.* 295:2720, author reply:2721–2722. <https://doi.org/10.1001/jama.295.23.2720-a>
- Düvel, K., J.L. Yecies, S. Menon, P. Raman, A.I. Lipovsky, A.L. Souza, E. Triantafellow, Q. Ma, R. Gorski, S. Cleaver, et al. 2010. Activation of a metabolic gene regulatory network downstream of mTOR complex 1. *Mol. Cell.* 39:171–183. <https://doi.org/10.1016/j.molcel.2010.06.022>
- Eberlé, D., B. Hegarty, P. Bossard, P. Ferré, and F. Foulfelle. 2004. SREBP transcription factors: master regulators of lipid homeostasis. *Biochimie.* 86:839–848. <https://doi.org/10.1016/j.biochi.2004.09.018>
- Eid, W., K. Dauner, K.C. Courtney, A. Gagnon, R.J. Parks, A. Sorisky, and X. Zha. 2017. mTORC1 activates SREBP-2 by suppressing cholesterol trafficking to lysosomes in mammalian cells. *Proc. Natl. Acad. Sci. USA.* 114:7999–8004. <https://doi.org/10.1073/pnas.1705304114>
- Escudero, S., E. Zaganjor, S. Lee, C.P. Mill, A.M. Morgan, E.B. Crawford, J. Chen, T.E. Wales, R. Mourtada, J. Luccarelli, et al. 2018. Dynamic Regulation of Long-Chain Fatty Acid Oxidation by a Noncanonical Interaction between the MCL-1 BH3 Helix and VLCAD. *Mol. Cell.* 69:729–743.e7. <https://doi.org/10.1016/j.molcel.2018.02.005>
- Feng, W.W., O. Wilkins, S. Bang, M. Ung, J. Li, J. An, C. Del Genio, K. Canfield, J. DiRenzo, W. Wells, et al. 2019. CD36-Mediated Metabolic Rewiring of Breast Cancer Cells Promotes Resistance to HER2-Targeted Therapies. *Cell Rep.* 29:3405–3420.e5. <https://doi.org/10.1016/j.celrep.2019.11.008>
- Flaveny, C.A., K. Griffett, B.-D. El-Gendy, M. Kazantzis, M. Sengupta, A.L. Amelio, A. Chatterjee, J. Walker, L.A. Solt, T.M. Kamenecka, and T.P. Burris. 2015. Broad Anti-tumor Activity of a Small Molecule that Selectively Targets the Warburg Effect and Lipogenesis. *Cancer Cell.* 28:42–56. <https://doi.org/10.1016/j.ccell.2015.05.007>
- Floris, A., M. Mazarei, X. Yang, A.E. Robinson, J. Zhou, A. Barberis, G. D'hallewin, E. Azara, Y. Spissu, A. Iglesias-Ara, et al. 2020. SUMOylation Protects FASN Against Proteasomal Degradation in Breast Cancer Cells Treated with Grape Leaf Extract. *Biomolecules.* 10:529. <https://doi.org/10.3390/biom10040529>
- Fujiwara, N., H. Nakagawa, K. Enooku, Y. Kudo, Y. Hayata, T. Nakatsuka, Y. Tanaka, R. Tateishi, Y. Hikiba, K. Misumi, et al. 2018. CPT2 down-regulation adapts HCC to lipid-rich environment and promotes carcinogenesis via acylcarnitine accumulation in obesity. *Gut.* 67:1493–1504. <https://doi.org/10.1136/gutjnl-2017-315193>
- Gallagher, E.J., Z. Zelenko, B.A. Neel, I.M. Antoniou, L. Rajan, N. Kase, and D. LeRoith. 2017. Elevated tumor LDLR expression accelerates LDL cholesterol-mediated breast cancer growth in mouse models of hyperlipidemia. *Oncogene.* 36:6462–6471. <https://doi.org/10.1038/ncr.2017.247>
- Gang, X., L. Xuan, X. Zhao, Y. Lv, F. Li, Y. Wang, and G. Wang. 2019. Speckle-type POZ protein suppresses lipid accumulation and prostate cancer growth by stabilizing fatty acid synthase. *Prostate.* 79:864–871. <https://doi.org/10.1002/pros.23793>
- Garcia-Bermudez, J., L. Baudrier, E.C. Bayraktar, Y. Shen, K. La, R. Guarecuco, B. Yucel, D. Fiore, B. Tavora, E. Freinkman, et al. 2019. Squalene accumulation in cholesterol auxotrophic lymphomas prevents oxidative cell death. *Nature.* 567:118–122. <https://doi.org/10.1038/s41586-019-0945-5>
- Geng, F., X. Cheng, X. Wu, J.Y. Yoo, C. Cheng, J.Y. Guo, X. Mo, P. Ru, B. Hurwitz, S.H. Kim, et al. 2016. Inhibition of SOAT1 Suppresses Glioblastoma Growth via Blocking SREBP-1-Mediated Lipogenesis. *Clin. Cancer Res.* 22:5337–5348. <https://doi.org/10.1158/1078-0432.CCR-15-2973>
- German, N.J., H. Yoon, R.Z. Yusuf, J.P. Murphy, L.W. Finley, G. Laurent, W. Haas, F.K. Satterstrom, J. Guarnerio, E. Zaganjor, et al. 2016. PHD3 Loss in Cancer Enables Metabolic Reliance on Fatty Acid Oxidation via Deactivation of ACC2. *Mol. Cell.* 63:1006–1020. <https://doi.org/10.1016/j.molcel.2016.08.014>

- Gharpure, K.M., S. Pradeep, M. Sans, R. Rupaimoole, C. Ivan, S.Y. Wu, E. Bayraktar, A.S. Nagaraja, L.S. Mangala, X. Zhang, et al. 2018. FABP4 as a key determinant of metastatic potential of ovarian cancer. *Nat. Commun.* 9:2923. <https://doi.org/10.1038/s41467-018-04987-y>
- Giandomenico, V., M. Simonsson, E. Grönroos, and J. Ericsson. 2003. Coactivator-dependent acetylation stabilizes members of the SREBP family of transcription factors. *Mol. Cell. Biol.* 23:2587–2599. <https://doi.org/10.1128/MCB.23.7.2587-2599.2003>
- Gill, S., J. Stevenson, I. Kristiana, and A.J. Brown. 2011. Cholesterol-dependent degradation of squalene monooxygenase, a control point in cholesterol synthesis beyond HMG-CoA reductase. *Cell Metab.* 13:260–273. <https://doi.org/10.1016/j.cmet.2011.01.015>
- Giordano, A., M. Calvani, O. Petillo, P. Grippo, F. Tuccillo, M.A. Melone, P. Bonelli, A. Calarco, and G. Peluso. 2005. tBid induces alterations of mitochondrial fatty acid oxidation flux by malonyl-CoA-independent inhibition of carnitine palmitoyltransferase-1. *Cell Death Differ.* 12: 603–613. <https://doi.org/10.1038/sj.cdd.4401636>
- Giral, P. 2020. Bempedoic Acid to Lower LDL Cholesterol - Safety and Efficacy. *N. Engl. J. Med.* 383:e49. <https://doi.org/10.1056/NEJMci1908495>
- Goldstein, J.L., and M.S. Brown. 2009. The LDL receptor. *Arterioscler. Thromb. Vasc. Biol.* 29:431–438. <https://doi.org/10.1161/ATVBAHA.108.179564>
- Gong, Y., J.N. Lee, P.C. Lee, J.L. Goldstein, M.S. Brown, and J. Ye. 2006. Sterol-regulated ubiquitination and degradation of Insig-1 creates a convergent mechanism for feedback control of cholesterol synthesis and uptake. *Cell Metab.* 3:15–24. <https://doi.org/10.1016/j.cmet.2005.11.014>
- Gong, X., J. Li, W. Shao, J. Wu, H. Qian, R. Ren, P. Espenshade, and N. Yan. 2015. Structure of the WD40 domain of SCAP from fission yeast reveals the molecular basis for SREBP recognition. *Cell Res.* 25:401–411. <https://doi.org/10.1038/cr.2015.32>
- Graner, E., D. Tang, S. Rossi, A. Baron, T. Migita, L.J. Weinstein, M. Lechpammer, D. Huesken, J. Zimmermann, S. Signoretti, and M. Loda. 2004. The isopeptidase USP2a regulates the stability of fatty acid synthase in prostate cancer. *Cancer Cell.* 5:253–261. [https://doi.org/10.1016/S1535-6108\(04\)00055-8](https://doi.org/10.1016/S1535-6108(04)00055-8)
- Gu, L., Y. Zhu, X. Lin, B. Lu, X. Zhou, F. Zhou, Q. Zhao, E.V. Prochownik, and Y. Li. 2020. The IKK $\beta$ -USP30-ACLY Axis Controls Lipogenesis and Tumorigenesis. *Hepatology*.:hep.31249. <https://doi.org/10.1002/hep.31249>
- Guillaumont, F., G. Bidaut, M. Ouassii, S. Servais, V. Gouirand, O. Olivares, S. Lac, L. Borge, J. Roques, O. Gayet, et al. 2015. Cholesterol uptake disruption, in association with chemotherapy, is a promising combined metabolic therapy for pancreatic adenocarcinoma. *Proc. Natl. Acad. Sci. USA.* 112:2473–2478. <https://doi.org/10.1073/pnas.1421601112>
- Guo, D., R.M. Prins, J. Dang, D. Kuga, A. Iwanami, H. Soto, K.Y. Lin, T.T. Huang, D. Akhavan, M.B. Hock, et al. 2009. EGFR signaling through an Akt-SREBP-1-dependent, rapamycin-resistant pathway sensitizes glioblastomas to antilipogenic therapy. *Sci. Signal.* 2:ra82. <https://doi.org/10.1126/scisignal.2000446>
- Guo, D., F. Reinitz, M. Youssef, C. Hong, D. Nathanson, D. Akhavan, D. Kuga, A.N. Amzajerd, H. Soto, S. Zhu, et al. 2011. An LXR agonist promotes glioblastoma cell death through inhibition of an EGFR/AKT/SREBP-1/LDLR-dependent pathway. *Cancer Discov.* 1:442–456. <https://doi.org/10.1158/2159-8290.CD-11-0102>
- Guo, W., J. Ma, Y. Yang, S. Guo, W. Zhang, T. Zhao, X. Yi, H. Wang, S. Wang, Y. Liu, et al. 2020. ATP-Citrate Lyase Epigenetically Potentiates Oxidative Phosphorylation to Promote Melanoma Growth and Adaptive Resistance to MAPK Inhibition. *Clin. Cancer Res.* 26:2725–2739. <https://doi.org/10.1158/1078-0432.CCR-19-1359>
- Ha, J., S. Daniel, S.S. Broyles, and K.H. Kim. 1994. Critical phosphorylation sites for acetyl-CoA carboxylase activity. *J. Biol. Chem.* 269:22162–22168.
- Han, J.Y., S.H. Lee, N.J. Yoo, L.S. Hyung, Y.J. Moon, T. Yun, H.T. Kim, and J.S. Lee. 2011. A randomized phase II study of gefitinib plus simvastatin versus gefitinib alone in previously treated patients with advanced non-small cell lung cancer. *Clin. Cancer Res.* 17:1553–1560. <https://doi.org/10.1158/1078-0432.CCR-10-2525>
- Han, J., E. Li, L. Chen, Y. Zhang, F. Wei, J. Liu, H. Deng, and Y. Wang. 2015. The CREB coactivator CRTC2 controls hepatic lipid metabolism by regulating SREBP1. *Nature.* 524:243–246. <https://doi.org/10.1038/nature14557>
- Hatzivassiliou, G., F. Zhao, D.E. Bauer, C. Andreadis, A.N. Shaw, D. Dhanak, S.R. Hingorani, D.A. Tuveson, and C.B. Thompson. 2005. ATP citrate lyase inhibition can suppress tumor cell growth. *Cancer Cell.* 8:311–321. <https://doi.org/10.1016/j.ccr.2005.09.008>
- Hirano, Y., S. Murata, K. Tanaka, M. Shimizu, and R. Sato. 2003. Sterol regulatory element-binding proteins are negatively regulated through SUMO-1 modification independent of the ubiquitin/26 S proteasome pathway. *J. Biol. Chem.* 278:16809–16819. <https://doi.org/10.1074/jbc.M212448200>
- Horton, J.D., N.A. Shah, J.A. Warrington, N.N. Anderson, S.W. Park, M.S. Brown, and J.L. Goldstein. 2003. Combined analysis of oligonucleotide microarray data from transgenic and knockout mice identifies direct SREBP target genes. *Proc. Natl. Acad. Sci. USA.* 100:12027–12032. <https://doi.org/10.1073/pnas.1534923100>
- Huang, Z., M. Zhang, A.A. Plec, S.J. Estill, L. Cai, J.J. Repa, S.L. McKnight, and B.P. Tu. 2018. ACS2 promotes systemic fat storage and utilization through selective regulation of genes involved in lipid metabolism. *Proc. Natl. Acad. Sci. USA.* 115:E9499–E9506. <https://doi.org/10.1073/pnas.1806635115>
- Huber, M.D., P.W. Vesely, K. Datta, and L. Gerace. 2013. Erlins restrict SREBP activation in the ER and regulate cellular cholesterol homeostasis. *J. Cell Biol.* 203:427–436. <https://doi.org/10.1083/jcb.201305076>
- Hunkeler, M., A. Hagmann, E. Stüttfeld, M. Chami, Y. Guri, H. Stahlberg, and T. Maier. 2018. Structural basis for regulation of human acetyl-CoA carboxylase. *Nature.* 558:470–474. <https://doi.org/10.1038/s41586-018-0201-4>
- Hwang, S., A.D. Nguyen, Y. Jo, L.J. Engelking, J. Brugarolas, and R.A. DeBose-Boyd. 2017. Hypoxia-inducible factor 1 $\alpha$  activates insulin-induced gene 2 (Insig-2) transcription for degradation of 3-hydroxy-3-methylglutaryl (HMG)-CoA reductase in the liver. *J. Biol. Chem.* 292:9382–9393. <https://doi.org/10.1074/jbc.M117.788562>
- Im, S.S., L. Yousef, C. Blaschitz, J.Z. Liu, R.A. Edwards, S.G. Young, M. Raffatellu, and T.F. Osborne. 2011. Linking lipid metabolism to the innate immune response in macrophages through sterol regulatory element binding protein-1a. *Cell Metab.* 13:540–549. <https://doi.org/10.1016/j.cmet.2011.04.001>
- Irisawa, M., J. Inoue, N. Ozawa, K. Mori, and R. Sato. 2009. The sterol-sensing endoplasmic reticulum (ER) membrane protein TRC8 hampers ER to Golgi transport of sterol regulatory element-binding protein-2 (SREBP-2)/SREBP cleavage-activated protein and reduces SREBP-2 cleavage. *J. Biol. Chem.* 284:28995–29004. <https://doi.org/10.1074/jbc.M109.041376>
- Jang, C., S. Hui, W. Lu, A.J. Cowan, R.J. Morscher, G. Lee, W. Liu, G.J. Tesz, M.J. Birnbaum, and J.D. Rabinowitz. 2018. The Small Intestine Converts Dietary Fructose into Glucose and Organic Acids. *Cell Metab.* 27: 351–361.e3. <https://doi.org/10.1016/j.cmet.2017.12.016>
- Jeon, S.M., N.S. Chandel, and N. Hay. 2012. AMPK regulates NADPH homeostasis to promote tumour cell survival during energy stress. *Nature.* 485:661–665. <https://doi.org/10.1038/nature11066>
- Jiang, M., N. Wu, B. Xu, Y. Chu, X. Li, S. Su, D. Chen, W. Li, Y. Shi, X. Gao, et al. 2019. Fatty acid-induced CD36 expression via O-GlcNAcylation drives gastric cancer metastasis. *Theranostics.* 9:5359–5373. <https://doi.org/10.7150/thno.34024>
- Jouve, J.L., T. Lecomte, O. Bouché, E. Barbier, F. Khemissa Akouz, G. Riachi, E. Nguyen Khac, I. Ollivier-Hourmand, M. Debette-Gratien, R. Faroux, et al. PRODIGE-11 investigators/collaborators. 2019. Pravastatin combination with sorafenib does not improve survival in advanced hepatocellular carcinoma. *J. Hepatol.* 71:516–522. <https://doi.org/10.1016/j.jhep.2019.04.021>
- Kamphorst, J.J., J.R. Cross, J. Fan, E. de Stanchina, R. Mathew, E.P. White, C.B. Thompson, and J.D. Rabinowitz. 2013. Hypoxic and Ras-transformed cells support growth by scavenging unsaturated fatty acids from lysophospholipids. *Proc. Natl. Acad. Sci. USA.* 110:8882–8887. <https://doi.org/10.1073/pnas.1307237110>
- Khwairakpam, A.D., K. Banik, S. Girisa, B. Shabnam, M. Shakibaei, L. Fan, F. Arfuso, J. Monisha, H. Wang, X. Mao, et al. 2020. The vital role of ATP citrate lyase in chronic diseases. *J. Mol. Med. (Berl.)*. 98:71–95. <https://doi.org/10.1007/s00109-019-01863-0>
- Kim, T.H., H. Kim, J.M. Park, S.S. Im, J.S. Bae, M.Y. Kim, H.G. Yoon, J.Y. Cha, K.S. Kim, and Y.H. Ahn. 2009. Interrelationship between liver X receptor alpha, sterol regulatory element-binding protein-1c, peroxisome proliferator-activated receptor gamma, and small heterodimer partner in the transcriptional regulation of glucokinase gene expression in liver. *J. Biol. Chem.* 284:15071–15083. <https://doi.org/10.1074/jbc.M109.006742>
- Kim, S.T., J.H. Kang, J. Lee, S.H. Park, J.O. Park, Y.S. Park, H.Y. Lim, I.G. Hwang, S.C. Lee, K.W. Park, et al. 2014. Simvastatin plus capecitabine-cisplatin versus placebo plus capecitabine-cisplatin in patients with previously untreated advanced gastric cancer: a double-blind randomised phase 3 study. *Eur. J. Cancer.* 50:2822–2830. <https://doi.org/10.1016/j.ejca.2014.08.005>

- Ko, C.W., J. Qu, D.D. Black, and P. Tso. 2020. Regulation of intestinal lipid metabolism: current concepts and relevance to disease. *Nat. Rev. Gastroenterol. Hepatol.* 17:169–183. <https://doi.org/10.1038/s41575-019-0250-7>
- Kong, Y., L. Cheng, F. Mao, Z. Zhang, Y. Zhang, E. Farah, J. Bosler, Y. Bai, N. Ahmad, S. Kuang, et al. 2018. Inhibition of cholesterol biosynthesis overcomes enzalutamide resistance in castration-resistant prostate cancer (CRPC). *J. Biol. Chem.* 293:14328–14341. <https://doi.org/10.1074/jbc.RA118.004442>
- Koobotse, M., J. Holly, and C. Perks. 2018. Elucidating the novel BRCA1 function as a non-genomic metabolic restraint in ER-positive breast cancer cell lines. *Oncotarget.* 9:33562–33576. <https://doi.org/10.18632/oncotarget.26093>
- Kotzka, J., S. Lehr, G. Roth, H. Avci, B. Knebel, and D. Muller-Wieland. 2004. Insulin-activated Erk-mitogen-activated protein kinases phosphorylate sterol regulatory element-binding Protein-2 at serine residues 432 and 455 in vivo. *J. Biol. Chem.* 279:22404–22411. <https://doi.org/10.1074/jbc.M401198200>
- Koundouros, N., and G. Poulgiannis. 2020. Reprogramming of fatty acid metabolism in cancer. *Br. J. Cancer.* 122:4–22. <https://doi.org/10.1038/s41416-019-0650-z>
- Kuan, Y.C., T. Hashidume, T. Shibata, K. Uchida, M. Shimizu, J. Inoue, and R. Sato. 2017. Heat Shock Protein 90 Modulates Lipid Homeostasis by Regulating the Stability and Function of Sterol Regulatory Element-binding Protein (SREBP) and SREBP Cleavage-activating Protein. *J. Biol. Chem.* 292:3016–3028. <https://doi.org/10.1074/jbc.M116.767277>
- Ladanyi, A., A. Mukherjee, H.A. Kenny, A. Johnson, A.K. Mitra, S. Sundarasan, K.M. Nieman, G. Pascual, S.A. Benitah, A. Montag, et al. 2018. Adipocyte-induced CD36 expression drives ovarian cancer progression and metastasis. *Oncogene.* 37:2285–2301. <https://doi.org/10.1038/s41388-017-0093-z>
- Lally, J.S.V., S. Ghoshal, D.K. DePeralta, O. Moaven, L. Wei, R. Masia, D.J. Erstad, N. Fujiwara, V. Leong, V.P. Houde, et al. 2019. Inhibition of Acetyl-CoA Carboxylase by Phosphorylation or the Inhibitor ND-654 Suppresses Lipogenesis and Hepatocellular Carcinoma. *Cell Metab.* 29:174–182.e5. <https://doi.org/10.1016/j.cmet.2018.08.020>
- Lee, J.N., B. Song, R.A. DeBose-Boyd, and J. Ye. 2006. Sterol-regulated degradation of Insig-1 mediated by the membrane-bound ubiquitin ligase gp78. *J. Biol. Chem.* 281:39308–39315. <https://doi.org/10.1074/jbc.M608999200>
- Lee, J.N., X. Zhang, J.D. Feramisco, Y. Gong, and J. Ye. 2008. Unsaturated fatty acids inhibit proteasomal degradation of Insig-1 at a postubiquitination step. *J. Biol. Chem.* 283:33772–33783. <https://doi.org/10.1074/jbc.M806108200>
- Lee, J.N., H. Kim, H. Yao, Y. Chen, K. Weng, and J. Ye. 2010. Identification of Ubx28 protein as a sensor for unsaturated fatty acids and regulator of triglyceride synthesis. *Proc. Natl. Acad. Sci. USA.* 107:21424–21429. <https://doi.org/10.1073/pnas.1011859107>
- Lee, J.V., A. Carrer, S. Shah, N.W. Snyder, S. Wei, S. Venneti, A.J. Worth, Z.F. Yuan, H.W. Lim, S. Liu, et al. 2014. Akt-dependent metabolic reprogramming regulates tumor cell histone acetylation. *Cell Metab.* 20:306–319. <https://doi.org/10.1016/j.cmet.2014.06.004>
- Lee, Y., K.H. Lee, G.K. Lee, S.H. Lee, K.Y. Lim, J. Joo, Y.J. Go, J.S. Lee, and J.Y. Han. 2017. Randomized Phase II Study of Afatinib Plus Simvastatin Versus Afatinib Alone in Previously Treated Patients with Advanced Nonadenocarcinomatous Non-small Cell Lung Cancer. *Cancer Res. Treat.* 49:1001–1011. <https://doi.org/10.4143/crt.2016.546>
- Lee, J.S., A. Roberts, D. Juarez, T.T. Vo, S. Bhatt, L.O. Herzog, S. Mallya, R.J. Bellin, S.K. Agarwal, A.H. Salem, et al. 2018. Statins enhance efficacy of venetoclax in blood cancers. *Sci. Transl. Med.* 10:eaq1240. <https://doi.org/10.1126/scitranslmed.aag1240>
- Li, Y., S. Xu, M.M. Mihaylova, B. Zheng, X. Hou, B. Jiang, O. Park, Z. Luo, E. Lefai, J.Y. Shyy, et al. 2011. AMPK phosphorylates and inhibits SREBP activity to attenuate hepatic steatosis and atherosclerosis in diet-induced insulin-resistant mice. *Cell Metab.* 13:376–388. <https://doi.org/10.1016/j.cmet.2011.03.009>
- Li, L., L. Che, K.M. Tharp, H.M. Park, M.G. Pilo, D. Cao, A. Cigliano, G. Latte, Z. Xu, S. Ribback, et al. 2016a. Differential requirement for de novo lipogenesis in cholangiocarcinoma and hepatocellular carcinoma of mice and humans. *Hepatology.* 63:1900–1913. <https://doi.org/10.1002/hep.28508>
- Li, X., Y. Jiang, J. Meisenhelder, W. Yang, D.H. Hawke, Y. Zheng, Y. Xia, K. Aldape, J. He, T. Hunter, et al. 2016b. Mitochondria-Translocated PGK1 Functions as a Protein Kinase to Coordinate Glycolysis and the TCA Cycle in Tumorigenesis. *Mol. Cell.* 61:705–719. <https://doi.org/10.1016/j.molcel.2016.02.009>
- Li, X., X. Qian, and Z. Lu. 2017a. Local histone acetylation by ACS2 promotes gene transcription for lysosomal biogenesis and autophagy. *Autophagy.* 13:1790–1791. <https://doi.org/10.1080/15548627.2017.1349581>
- Li, X., W. Yu, X. Qian, Y. Xia, Y. Zheng, J.H. Lee, W. Li, J. Lyu, G. Rao, X. Zhang, et al. 2017b. Nucleus-Translocated ACS2 Promotes Gene Transcription for Lysosomal Biogenesis and Autophagy. *Mol. Cell.* 66:684–697.e9. <https://doi.org/10.1016/j.molcel.2017.04.026>
- Li, K., C. Zhang, L. Chen, P. Wang, Y. Fang, J. Zhu, S. Chen, J. Du, B. Shen, K. Wu, and Y. Liu. 2019a. The role of acetyl-coA carboxylase2 in head and neck squamous cell carcinoma. *PeerJ.* 7:e7037. <https://doi.org/10.7717/peerj.7037>
- Li, Q., Y. Wang, S. Wu, Z. Zhou, X. Ding, R. Shi, R.F. Thorne, X.D. Zhang, W. Hu, and M. Wu. 2019b. CircACC1 Regulates Assembly and Activation of AMPK Complex under Metabolic Stress. *Cell Metab.* 30:157–173.e7. <https://doi.org/10.1016/j.cmet.2019.05.009>
- Lim, S.H., T.W. Kim, Y.S. Hong, S.W. Han, K.H. Lee, H.J. Kang, I.G. Hwang, J.Y. Lee, H.S. Kim, S.T. Kim, et al. 2015. A randomised, double-blind, placebo-controlled multi-centre phase III trial of XELIRI/FOLFIRI plus simvastatin for patients with metastatic colorectal cancer. *Br. J. Cancer.* 113:1421–1426. <https://doi.org/10.1038/bjc.2015.371>
- Lin, C.Y., and J.A. Gustafsson. 2015. Targeting liver X receptors in cancer therapeutics. *Nat. Rev. Cancer.* 15:216–224. <https://doi.org/10.1038/nrc3912>
- Lin, J., R. Yang, P.T. Tarr, P.H. Wu, C. Handschin, S. Li, W. Yang, L. Pei, M. Uldry, P. Tontonoz, et al. 2005. Hyperlipidemic effects of dietary saturated fats mediated through PGC-1beta coactivation of SREBP. *Cell.* 120:261–273. <https://doi.org/10.1016/j.cell.2004.11.043>
- Lin, R., R. Tao, X. Gao, T. Li, X. Zhou, K.L. Guan, Y. Xiong, and Q.Y. Lei. 2013. Acetylation stabilizes ATP-citrate lyase to promote lipid biosynthesis and tumor growth. *Mol. Cell.* 51:506–518. <https://doi.org/10.1016/j.molcel.2013.07.002>
- Lin, H.P., Z.L. Cheng, R.Y. He, L. Song, M.X. Tian, L.S. Zhou, B.S. Groh, W.R. Liu, M.B. Ji, C. Ding, et al. 2016. Destabilization of Fatty Acid Synthase by Acetylation Inhibits De Novo Lipogenesis and Tumor Cell Growth. *Cancer Res.* 76:6924–6936. <https://doi.org/10.1158/0008-5472.CAN-16-1597>
- Liscum, L., J. Finer-Moore, R.M. Stroud, K.L. Luskey, M.S. Brown, and J.L. Goldstein. 1985. Domain structure of 3-hydroxy-3-methylglutaryl coenzyme A reductase, a glycoprotein of the endoplasmic reticulum. *J. Biol. Chem.* 260:522–530.
- Liu, L., X. Zhao, L. Zhao, J. Li, H. Yang, Z. Zhu, J. Liu, and G. Huang. 2016. Arginine Methylation of SREBP1a via PRMT5 Promotes De Novo Lipogenesis and Tumor Growth. *Cancer Res.* 76:1260–1272. <https://doi.org/10.1158/0008-5472.CAN-15-1766>
- Liu, D., C.C. Wong, L. Fu, H. Chen, L. Zhao, C. Li, Y. Zhou, Y. Zhang, W. Xu, Y. Yang, et al. 2018. Squalene epoxidase drives NAFLD-induced hepatocellular carcinoma and is a pharmaceutical target. *Sci. Transl. Med.* 10:eaap9840. <https://doi.org/10.1126/scitranslmed.aap9840>
- Luo, D.X., D.J. Tong, S. Rajput, C. Wang, D.F. Liao, D. Cao, and E. Maser. 2012. Targeting acetyl-CoA carboxylases: small molecular inhibitors and their therapeutic potential. *Recent Patents Anticancer Drug Discov.* 7:168–184. <https://doi.org/10.2174/157489212799972918>
- Luo, J., H. Yang, and B.L. Song. 2020. Mechanisms and regulation of cholesterol homeostasis. *Nat. Rev. Mol. Cell Biol.* 21:225–245. <https://doi.org/10.1038/s41580-019-0190-7>
- Luong, A., V.C. Hannah, M.S. Brown, and J.L. Goldstein. 2000. Molecular characterization of human acetyl-CoA synthetase, an enzyme regulated by sterol regulatory element-binding proteins. *J. Biol. Chem.* 275:26458–26466. <https://doi.org/10.1074/jbc.M004160200>
- Ma, J., R. Yan, X. Zu, J.M. Cheng, K. Rao, D.F. Liao, and D. Cao. 2008. Aldoketo reductase family 1 B10 affects fatty acid synthesis by regulating the stability of acetyl-CoA carboxylase-alpha in breast cancer cells. *J. Biol. Chem.* 283:3418–3423. <https://doi.org/10.1074/jbc.M707650200>
- Ma, M.K.F., E.Y.T. Lau, D.H.W. Leung, J. Lo, N.P.Y. Ho, L.K.W. Cheng, S. Ma, C.H. Lin, J.A. Copland, J. Ding, et al. 2017. Stearoyl-CoA desaturase regulates sorafenib resistance via modulation of ER stress-induced differentiation. *J. Hepatol.* 67:979–990. <https://doi.org/10.1016/j.jhep.2017.06.015>
- Ma, Y., S.M. Temkin, A.M. Hawkrigde, C. Guo, W. Wang, X.Y. Wang, and X. Fang. 2018. Fatty acid oxidation: An emerging facet of metabolic transformation in cancer. *Cancer Lett.* 435:92–100. <https://doi.org/10.1016/j.canlet.2018.08.006>
- Mahoney, C.E., D. Pirman, V. Chubukov, T. Sleger, S. Hayes, Z.P. Fan, E.L. Allen, Y. Chen, L. Huang, M. Liu, et al. 2019. A chemical biology screen identifies a vulnerability of neuroendocrine cancer cells to SQLE

- inhibition. *Nat. Commun.* 10:96. <https://doi.org/10.1038/s41467-018-07959-4>
- Mashimo, T., K. Pichumani, V. Vemireddy, K.J. Hatanpaa, D.K. Singh, S. Sirasanagandla, S. Nannepaga, S.G. Piccirillo, Z. Kovacs, C. Foong, et al. 2014. Acetate is a bioenergetic substrate for human glioblastoma and brain metastases. *Cell*. 159:1603–1614. <https://doi.org/10.1016/j.cell.2014.11.025>
- Menendez, J.A., and R. Lupu. 2007. Fatty acid synthase and the lipogenic phenotype in cancer pathogenesis. *Nat. Rev. Cancer*. 7:763–777. <https://doi.org/10.1038/nrc2222>
- Menendez, J.A., L. Vellon, I. Mehmi, B.P. Oza, S. Ropero, R. Colomer, and R. Lupu. 2004. Inhibition of fatty acid synthase (FAS) suppresses HER2/neu (erbB-2) oncogene overexpression in cancer cells. *Proc. Natl. Acad. Sci. USA*. 101:10715–10720. <https://doi.org/10.1073/pnas.0403390101>
- Metallo, C.M., P.A. Gameiro, E.L. Bell, K.R. Mattaini, J. Yang, K. Hiller, C.M. Jewell, Z.R. Johnson, D.J. Irvine, L. Guarente, et al. 2011. Reductive glutamine metabolism by IDH1 mediates lipogenesis under hypoxia. *Nature*. 481:380–384. <https://doi.org/10.1038/nature10602>
- Min, H.K., A. Kapoor, M. Fuchs, F. Mirshahi, H. Zhou, J. Maher, J. Kellum, R. Warnick, M.J. Contos, and A.J. Sanyal. 2012. Increased hepatic synthesis and dysregulation of cholesterol metabolism is associated with the severity of nonalcoholic fatty liver disease. *Cell Metab.* 15:665–674. <https://doi.org/10.1016/j.cmet.2012.04.004>
- Misawa, K., T. Horiba, N. Arimura, Y. Hirano, J. Inoue, N. Emoto, H. Shimano, M. Shimizu, and R. Sato. 2003. Sterol regulatory element-binding protein-2 interacts with hepatocyte nuclear factor-4 to enhance sterol isomerase gene expression in hepatocytes. *J. Biol. Chem.* 278:36176–36182. <https://doi.org/10.1074/jbc.M302387200>
- Moolenaar, W.H., and A. Perrakis. 2011. Insights into autotaxin: how to produce and present a lipid mediator. *Nat. Rev. Mol. Cell Biol.* 12:674–679. <https://doi.org/10.1038/nrm3188>
- Moon, S.H., C.H. Huang, S.L. Houlihan, K. Regunath, W.A. Freed-Pastor, J.P. Morris IV, D.F. Tschaharganeh, E.R. Kastenhuber, A.M. Barsotti, R. Culp-Hill, et al. 2019. p53 Represses the Mevalonate Pathway to Mediate Tumor Suppression. *Cell*. 176:564–580.e19. <https://doi.org/10.1016/j.cell.2018.11.011>
- Moreau, K., E. Dizin, H. Ray, C. Luquin, E. Lefai, F. Foufelle, M. Billaud, G.M. Lenoir, and N.D. Venezia. 2006. BRCAL affects lipid synthesis through its interaction with acetyl-CoA carboxylase. *J. Biol. Chem.* 281:3172–3181. <https://doi.org/10.1074/jbc.M504652200>
- Morioka, S., K. Sai, E. Omori, Y. Ikeda, K. Matsumoto, and J. Ninomiya-Tsuji. 2016. TAK1 regulates hepatic lipid homeostasis through SREBP. *Oncogene*. 35:3829–3838. <https://doi.org/10.1038/ncr.2015.453>
- Mullen, P.J., R. Yu, J. Longo, M.C. Archer, and L.Z. Penn. 2016. The interplay between cell signalling and the mevalonate pathway in cancer. *Nat. Rev. Cancer*. 16:718–731. <https://doi.org/10.1038/nrc.2016.76>
- Nakamura, M.T., and T.Y. Nara. 2004. Structure, function, and dietary regulation of delta6, delta5, and delta9 desaturases. *Annu. Rev. Nutr.* 24:345–376. <https://doi.org/10.1146/annurev.nutr.24.121803.063211>
- Nakanishi, M., J.L. Goldstein, and M.S. Brown. 1988. Multivalent control of 3-hydroxy-3-methylglutaryl coenzyme A reductase. Mevalonate-derived product inhibits translation of mRNA and accelerates degradation of enzyme. *J. Biol. Chem.* 263:8929–8937.
- Nieman, K.M., H.A. Kenny, C.V. Penicka, A. Ladanyi, R. Buell-Gutbrod, M.R. Zillhardt, I.L. Romero, M.S. Carey, G.B. Mills, G.S. Hotamisligil, et al. 2011. Adipocytes promote ovarian cancer metastasis and provide energy for rapid tumor growth. *Nat. Med.* 17:1498–1503. <https://doi.org/10.1038/nm.2492>
- Nohturfft, A., and S.C. Zhang. 2009. Coordination of lipid metabolism in membrane biogenesis. *Annu. Rev. Cell Dev. Biol.* 25:539–566. <https://doi.org/10.1146/annurev.cellbio.24.110707.175344>
- Padanad, M.S., G. Konstantinidou, N. Venkateswaran, M. Melegari, S. Rindhe, M. Mitsche, C. Yang, K. Batten, K.E. Huffman, J. Liu, et al. 2016. Fatty Acid Oxidation Mediated by Acyl-CoA Synthetase Long Chain 3 Is Required for Mutant KRAS Lung Tumorigenesis. *Cell Rep.* 16:1614–1628. <https://doi.org/10.1016/j.celrep.2016.07.009>
- Padyana, A.K., S. Gross, L. Jin, G. Cianchetta, R. Narayanaswamy, F. Wang, R. Wang, C. Fang, X. Lv, S.A. Biller, et al. 2019. Structure and inhibition mechanism of the catalytic domain of human squalene epoxidase. *Nat. Commun.* 10:97. <https://doi.org/10.1038/s41467-018-07928-x>
- Pan, J., Z. Fan, Z. Wang, Q. Dai, Z. Xiang, F. Yuan, M. Yan, Z. Zhu, B. Liu, and C. Li. 2019. CD36 mediates palmitate acid-induced metastasis of gastric cancer via AKT/GSK-3 $\beta$ / $\beta$ -catenin pathway. *J. Exp. Clin. Cancer Res.* 38:52. <https://doi.org/10.1186/s13046-019-1049-7>
- Park, J.B., C.S. Lee, J.H. Jang, J. Ghim, Y.J. Kim, S. You, D. Hwang, P.G. Suh, and S.H. Ryu. 2012. Phospholipase signalling networks in cancer. *Nat. Rev. Cancer*. 12:782–792. <https://doi.org/10.1038/nrc3379>
- Pascual, G., A. Avgustinova, S. Mejetta, M. Martín, A. Castellanos, C.S. Atoloni, A. Berenguer, N. Prats, A. Toll, J.A. Hueto, et al. 2017. Targeting metastasis-initiating cells through the fatty acid receptor CD36. *Nature*. 541:41–45. <https://doi.org/10.1038/nature20791>
- Peck, B., and A. Schulze. 2016. Lipid desaturation - the next step in targeting lipogenesis in cancer? *FEBS J.* 283:2767–2778. <https://doi.org/10.1111/febs.13681>
- Penfold, L., A. Woods, P. Muckett, A.Y. Nikitin, T.R. Kent, S. Zhang, R. Graham, A. Pollard, and D. Carling. 2018. CAMKK2 Promotes Prostate Cancer Independently of AMPK via Increased Lipogenesis. *Cancer Res.* 78:6747–6761. <https://doi.org/10.1158/0008-5472.CAN-18-0585>
- Peterson, T.R., S.S. Sengupta, T.E. Harris, A.E. Carmack, S.A. Kang, E. Balderas, D.A. Guertin, K.L. Madden, A.E. Carpenter, B.N. Finck, and D.M. Sabatini. 2011. mTOR complex 1 regulates lipin 1 localization to control the SREBP pathway. *Cell*. 146:408–420. <https://doi.org/10.1016/j.cell.2011.06.034>
- Pike, L.S., A.L. Smift, N.J. Croteau, D.A. Ferrick, and M. Wu. 2011. Inhibition of fatty acid oxidation by etomoxir impairs NADPH production and increases reactive oxygen species resulting in ATP depletion and cell death in human glioblastoma cells. *Biochim. Biophys. Acta*. 1807:726–734. <https://doi.org/10.1016/j.bbabi.2010.10.022>
- Porter, J.A., K.E. Young, and P.A. Beachy. 1996. Cholesterol modification of hedgehog signaling proteins in animal development. *Science*. 274:255–259. <https://doi.org/10.1126/science.274.5285.255>
- Potapova, I.A., M.R. El-Maghrabi, S.V. Doronin, and W.B. Benjamin. 2000. Phosphorylation of recombinant human ATP:citrate lyase by cAMP-dependent protein kinase abolishes homotropic allosteric regulation of the enzyme by citrate and increases the enzyme activity. Allosteric activation of ATP:citrate lyase by phosphorylated sugars. *Biochemistry*. 39:1169–1179. <https://doi.org/10.1021/bi992159y>
- Potze, L., S. Di Franco, C. Grandela, M.L. Pras-Raves, D.I. Picavet, H.A. van Veen, H. van Lenthe, F.B. Mullauer, N.N. van der Wel, A. Luyf, et al. 2016. Betulinic acid induces a novel cell death pathway that depends on cardiolipin modification. *Oncogene*. 35:427–437. <https://doi.org/10.1038/ncr.2015.102>
- Poulsen, L.L., M. Siersbæk, and S. Mandrup. 2012. PPARs: fatty acid sensors controlling metabolism. *Semin. Cell Dev. Biol.* 23:631–639. <https://doi.org/10.1016/j.semcdb.2012.01.003>
- Rawson, R.B. 2003. The SREBP pathway--insights from Insigs and insects. *Nat. Rev. Mol. Cell Biol.* 4:631–640. <https://doi.org/10.1038/nrm1174>
- Rengan, R., R. Mick, D. Pryma, M.A. Rosen, L.L. Lin, A.M. Maity, T.L. Evans, J.P. Stevenson, C.J. Langer, J. Kucharczuk, et al. 2012. A phase I trial of the HIV protease inhibitor nelfinavir with concurrent chemoradiotherapy for unresectable stage IIIA/IIIB non-small cell lung cancer: a report of toxicities and clinical response. *J. Thorac. Oncol.* 7:709–715. <https://doi.org/10.1097/JTO.0b013e3182435aa6>
- Repa, J.J., G. Liang, J. Ou, Y. Bashmakov, J.M. Lobaccaro, I. Shimomura, B. Shan, M.S. Brown, J.L. Goldstein, and D.J. Mangelsdorf. 2000. Regulation of mouse sterol regulatory element-binding protein-1c gene (SREBP-1c) by oxysterol receptors, LXRalpha and LXRbeta. *Genes Dev.* 14:2819–2830. <https://doi.org/10.1101/gad.844900>
- Rios Garcia, M., B. Steinbauer, K. Srivastava, M. Singhal, F. Mattijssen, A. Maida, S. Christian, H. Hess-Stumpp, H.G. Augustin, K. Müller-Decker, et al. 2017. Acetyl-CoA Carboxylase 1-Dependent Protein Acetylation Controls Breast Cancer Metastasis and Recurrence. *Cell Metab.* 26:842–855.e5. <https://doi.org/10.1016/j.cmet.2017.09.018>
- Röhrig, F., and A. Schulze. 2016. The multifaceted roles of fatty acid synthesis in cancer. *Nat. Rev. Cancer*. 16:732–749. <https://doi.org/10.1038/nrc.2016.89>
- Ruderman, N.B., X.J. Xu, L. Nelson, J.M. Cacicedo, A.K. Saha, F. Lan, and Y. Ido. 2010. AMPK and SIRT1: a long-standing partnership? *Am. J. Physiol. Endocrinol. Metab.* 298:E751–E760. <https://doi.org/10.1152/ajpendo.00745.2009>
- Samudio, I., R. Harmancey, M. Fiegl, H. Kantarjian, M. Konopleva, B. Korchin, K. Kaluarachchi, W. Bornmann, S. Duvvuri, H. Taegtmeier, and M. Andreeff. 2010. Pharmacologic inhibition of fatty acid oxidation sensitizes human leukemia cells to apoptosis induction. *J. Clin. Invest.* 120:142–156. <https://doi.org/10.1172/JCI38942>
- Sato, R., J.L. Goldstein, and M.S. Brown. 1993. Replacement of serine-871 of hamster 3-hydroxy-3-methylglutaryl-CoA reductase prevents phosphorylation by AMP-activated kinase and blocks inhibition of sterol synthesis induced by ATP depletion. *Proc. Natl. Acad. Sci. USA*. 90:9261–9265. <https://doi.org/10.1073/pnas.90.20.9261>
- Schug, Z.T., B. Peck, D.T. Jones, Q. Zhang, S. Grosskurth, I.S. Alam, L.M. Goodwin, E. Smethurst, S. Mason, K. Blyth, et al. 2015. Acetyl-CoA

- synthetase 2 promotes acetate utilization and maintains cancer cell growth under metabolic stress. *Cancer Cell*. 27:57–71. <https://doi.org/10.1016/j.ccell.2014.12.002>
- Seckl, M.J., C.H. Ottensmeier, M. Cullen, P. Schmid, Y. Ngai, D. Muthukumar, J. Thompson, S. Harden, G. Middleton, K.M. Fife, et al. 2017. Multi-center, Phase III, Randomized, Double-Blind, Placebo-Controlled Trial of Pravastatin Added to First-Line Standard Chemotherapy in Small-Cell Lung Cancer (LUNGSTAR). *J. Clin. Oncol.* 35:1506–1514. <https://doi.org/10.1200/JCO.2016.69.7391>
- Seok, S., Y.C. Kim, S. Byun, S. Choi, Z. Xiao, N. Iwamori, Y. Zhang, C. Wang, J. Ma, K. Ge, et al. 2018. Fasting-induced JMJD3 histone demethylase epigenetically activates mitochondrial fatty acid  $\beta$ -oxidation. *J. Clin. Invest.* 128:3144–3159. <https://doi.org/10.1172/JCI97736>
- Shao, W., and P.J. Espenshade. 2012. Expanding roles for SREBP in metabolism. *Cell Metab.* 16:414–419. <https://doi.org/10.1016/j.cmet.2012.09.002>
- Sharpe, L.J., and A.J. Brown. 2013. Controlling cholesterol synthesis beyond 3-hydroxy-3-methylglutaryl-CoA reductase (HMGCR). *J. Biol. Chem.* 288:18707–18715. <https://doi.org/10.1074/jbc.R113.479808>
- Sharpe, L.J., V. Howe, N.A. Scott, W. Luu, L. Phan, J.M. Berk, M. Hochstrasser, and A.J. Brown. 2019. Cholesterol increases protein levels of the E3 ligase MARCH6 and thereby stimulates protein degradation. *J. Biol. Chem.* 294:2436–2448. <https://doi.org/10.1074/jbc.RA118.005069>
- Shimano, H., and R. Sato. 2017. SREBP-regulated lipid metabolism: convergent physiology - divergent pathophysiology. *Nat. Rev. Endocrinol.* 13:710–730. <https://doi.org/10.1038/nrendo.2017.91>
- Sivanand, S., S. Rhoades, Q. Jiang, J.V. Lee, J. Benci, J. Zhang, S. Yuan, I. Viney, S. Zhao, A. Carrer, et al. 2017. Nuclear Acetyl-CoA Production by ACLY Promotes Homologous Recombination. *Mol. Cell.* 67:252–265.e6. <https://doi.org/10.1016/j.molcel.2017.06.008>
- Snaebjornsson, M.T., S. Janaki-Raman, and A. Schulze. 2020. Greasing the Wheels of the Cancer Machine: The Role of Lipid Metabolism in Cancer. *Cell Metab.* 31:62–76. <https://doi.org/10.1016/j.cmet.2019.11.010>
- Stevenson, J., W. Luu, I. Kristiana, and A.J. Brown. 2014. Squalene monooxygenase, a key enzyme in cholesterol synthesis, is stabilized by unsaturated fatty acids. *Biochem. J.* 461:435–442. <https://doi.org/10.1042/BJ20131404>
- Su, X., and N.A. Abumrad. 2009. Cellular fatty acid uptake: a pathway under construction. *Trends Endocrinol. Metab.* 20:72–77. <https://doi.org/10.1016/j.tem.2008.11.001>
- Sundqvist, A., M.T. Bengochea-Alonso, X. Ye, V. Lukiyanchuk, J. Jin, J.W. Harper, and J. Ericsson. 2005. Control of lipid metabolism by phosphorylation-dependent degradation of the SREBP family of transcription factors by SCF(Fbw7). *Cell Metab.* 1:379–391. <https://doi.org/10.1016/j.cmet.2005.04.010>
- Svensson, R.U., S.J. Parker, L.J. Eichner, M.J. Kolar, M. Wallace, S.N. Brun, P.S. Lombardo, J.L. Van Nostrand, A. Hutchins, L. Vera, et al. 2016. Inhibition of acetyl-CoA carboxylase suppresses fatty acid synthesis and tumor growth of non-small-cell lung cancer in preclinical models. *Nat. Med.* 22:1108–1119. <https://doi.org/10.1038/nm.4181>
- Tao, R., X. Xiong, R.A. DePinho, C.X. Deng, and X.C. Dong. 2013. Hepatic SREBP-2 and cholesterol biosynthesis are regulated by FoxO3 and Sirt6. *J. Lipid Res.* 54:2745–2753. <https://doi.org/10.1194/jlr.M039339>
- Tesfay, L., B.T. Paul, A. Konstorum, Z. Deng, A.O. Cox, J. Lee, C.M. Furdul, P. Hegde, F.M. Torti, and S.V. Torti. 2019. Stearoyl-CoA Desaturase 1 Protects Ovarian Cancer Cells from Ferroptotic Cell Death. *Cancer Res.* 79:5355–5366. <https://doi.org/10.1158/0008-5472.CAN-19-0369>
- Ueda, K., Y. Nakatsu, T. Yamamotoya, H. Ono, Y. Inoue, M.K. Inoue, Y. Mizuno, Y. Matsunaga, A. Kushiyama, H. Sakoda, et al. 2019. Prolyl isomerase Pin1 binds to and stabilizes acetyl CoA carboxylase 1 protein, thereby supporting cancer cell proliferation. *Oncotarget.* 10:1637–1648. <https://doi.org/10.18632/oncotarget.26691>
- Walker, A.K., F. Yang, K. Jiang, J.Y. Ji, J.L. Watts, A. Purushotham, O. Boss, M.L. Hirsch, S. Ribich, J.J. Smith, et al. 2010. Conserved role of SIRT1 orthologs in fasting-dependent inhibition of the lipid/cholesterol regulator SREBP. *Genes Dev.* 24:1403–1417. <https://doi.org/10.1101/gad.1901210>
- Wang, M.D., H. Wu, G.B. Fu, H.L. Zhang, X. Zhou, L. Tang, L.W. Dong, C.J. Qin, S. Huang, L.H. Zhao, et al. 2016. Acetyl-coenzyme A carboxylase alpha promotion of glucose-mediated fatty acid synthesis enhances survival of hepatocellular carcinoma in mice and patients. *Hepatology.* 63:1272–1286. <https://doi.org/10.1002/hep.28415>
- Wang, T., J.F. Fahrman, H. Lee, Y.J. Li, S.C. Tripathi, C. Yue, C. Zhang, V. Lifshitz, J. Song, Y. Yuan, et al. 2018. JAK/STAT3-Regulated Fatty Acid  $\beta$ -Oxidation Is Critical for Breast Cancer Stem Cell Self-Renewal and Chemoresistance. *Cell Metab.* 27:136–150.e5. <https://doi.org/10.1016/j.cmet.2017.11.001>
- Wang, R., B. Tao, Q. Fan, S. Wang, L. Chen, J. Zhang, Y. Hao, S. Dong, Z. Wang, W. Wang, et al. 2019. Fatty-acid receptor CD36 functions as a hydrogen sulfide-targeted receptor with its Cys333-Cys272 disulfide bond serving as a specific molecular switch to accelerate gastric cancer metastasis. *EBioMedicine.* 45:108–123. <https://doi.org/10.1016/j.ebiom.2019.06.037>
- Watt, M.J., A.K. Clark, L.A. Selth, V.R. Haynes, N. Lister, R. Rebello, L.H. Porter, B. Niranjani, S.T. Whitby, J. Lo, et al. 2019. Suppressing fatty acid uptake has therapeutic effects in preclinical models of prostate cancer. *Sci. Transl. Med.* 11:eaaau5758. <https://doi.org/10.1126/scitranslmed.aau5758>
- Wei, J., and L. Tong. 2015. Crystal structure of the 500-kDa yeast acetyl-CoA carboxylase holoenzyme dimer. *Nature.* 526:723–727. <https://doi.org/10.1038/nature15375>
- Wellen, K.E., G. Hatzivassiliou, U.M. Sachdeva, T.V. Bui, J.R. Cross, and C.B. Thompson. 2009. ATP-citrate lyase links cellular metabolism to histone acetylation. *Science.* 324:1076–1080. <https://doi.org/10.1126/science.1164097>
- Wilson, J.M., E. Fokas, S.J. Dutton, N. Patel, M.A. Hawkins, C. Eccles, K.Y. Chu, L. Durrant, A.G. Abraham, M. Partridge, et al. 2016. ARCI: A phase II trial of the HIV protease inhibitor Nelfinavir in combination with chemoradiation for locally advanced inoperable pancreatic cancer. *Radiother. Oncol.* 119:306–311. <https://doi.org/10.1016/j.radonc.2016.03.021>
- Xia, Y., Y. Xie, Z. Yu, H. Xiao, G. Jiang, X. Zhou, Y. Yang, X. Li, M. Zhao, L. Li, et al. 2018. The Mevalonate Pathway Is a Druggable Target for Vaccine Adjuvant Discovery. *Cell.* 175:1059–1073.e21. <https://doi.org/10.1016/j.cell.2018.08.070>
- Xiao, X., J.J. Tang, C. Peng, Y. Wang, L. Fu, Z.P. Qiu, Y. Xiong, L.F. Yang, H.W. Cui, X.L. He, et al. 2017. Cholesterol Modification of Smoothed Is Required for Hedgehog Signaling. *Mol. Cell.* 66:154–162.e10. <https://doi.org/10.1016/j.molcel.2017.02.015>
- Xie, H., C.H. Tang, J.H. Song, A. Mancuso, J.R. Del Valle, J. Cao, Y. Xiang, C.V. Dang, R. Lan, D.J. Sanchez, et al. 2018. IRE1 $\alpha$  RNase-dependent lipid homeostasis promotes survival in Myc-transformed cancers. *J. Clin. Invest.* 128:1300–1316. <https://doi.org/10.1172/JCI95864>
- Xing, X., S.C. Wang, W. Xia, Y. Zou, R. Shao, K.Y. Kwong, Z. Yu, S. Zhang, S. Miller, L. Huang, and M.C. Hung. 2000. The ets protein PEA3 suppresses HER-2/neu overexpression and inhibits tumorigenesis. *Nat. Med.* 6:189–195. <https://doi.org/10.1038/72294>
- Xu, D., Z. Wang, Y. Zhang, W. Jiang, Y. Pan, B.L. Song, and Y. Chen. 2015. PAQR3 modulates cholesterol homeostasis by anchoring Scap/SREBP complex to the Golgi apparatus. *Nat. Commun.* 6:8100. <https://doi.org/10.1038/ncomms9100>
- Xu, D., Z. Wang, Y. Xia, F. Shao, W. Xia, Y. Wei, X. Li, X. Qian, J.H. Lee, L. Du, et al. 2020. The gluconeogenic enzyme PCK1 phosphorylates INSIG1/2 for lipogenesis. *Nature.* 580:530–535. <https://doi.org/10.1038/s41586-020-2183-2>
- Yabe, D., M.S. Brown, and J.L. Goldstein. 2002. Insig-2, a second endoplasmic reticulum protein that binds SCAP and blocks export of sterol regulatory element-binding proteins. *Proc. Natl. Acad. Sci. USA.* 99:12753–12758. <https://doi.org/10.1073/pnas.162488899>
- Yan, X., G. Zhang, F. Bie, Y. Lv, Y. Ma, M. Ma, Y. Wang, X. Hao, N. Yuan, and X. Jiang. 2017. Eugenol inhibits oxidative phosphorylation and fatty acid oxidation via downregulation of c-Myc/PGC-1 $\beta$ /ERR $\alpha$  signaling pathway in MCF10A-ras cells. *Sci. Rep.* 7:12920. <https://doi.org/10.1038/s41598-017-13505-x>
- Yang, T., P.J. Espenshade, M.E. Wright, D. Yabe, Y. Gong, R. Aebersold, J.L. Goldstein, and M.S. Brown. 2002a. Crucial step in cholesterol homeostasis: sterols promote binding of SCAP to INSIG-1, a membrane protein that facilitates retention of SREBPs in ER. *Cell.* 110:489–500. [https://doi.org/10.1016/S0092-8674\(02\)00872-3](https://doi.org/10.1016/S0092-8674(02)00872-3)
- Yang, Y.A., W.F. Han, P.J. Morin, F.J. Chrest, and E.S. Pizer. 2002b. Activation of fatty acid synthesis during neoplastic transformation: role of mitogen-activated protein kinase and phosphatidylinositol 3-kinase. *Exp. Cell Res.* 279:80–90. <https://doi.org/10.1006/excr.2002.5600>
- Yang, P., C. Su, X. Luo, H. Zeng, L. Zhao, L. Wei, X. Zhang, Z. Varghese, J.F. Moorhead, Y. Chen, and X.Z. Ruan. 2018. Dietary oleic acid-induced CD36 promotes cervical cancer cell growth and metastasis via up-regulation Src/ERK pathway. *Cancer Lett.* 438:76–85. <https://doi.org/10.1016/j.canlet.2018.09.006>
- Yang, X., F. Shao, S. Shi, X. Feng, W. Wang, Y. Wang, W. Guo, J. Wang, S. Gao, Y. Gao, et al. 2019. Prognostic Impact of Metabolism Reprogramming Markers Acetyl-CoA Synthetase 2 Phosphorylation and Ketohexokinase-A Expression in Non-Small-Cell Lung Carcinoma. *Front. Oncol.* 9:1123. <https://doi.org/10.3389/fonc.2019.01123>



- Yang, J., L. Wang, and R. Jia. 2020. Role of *de novo* cholesterol synthesis enzymes in cancer. *J. Cancer*. 11:1761–1767. <https://doi.org/10.7150/jca.38598>
- Yecies, J.L., H.H. Zhang, S. Menon, S. Liu, D. Yecies, A.I. Lipovsky, C. Gorgun, D.J. Kwiatkowski, G.S. Hotamisligil, C.H. Lee, and B.D. Manning. 2011. Akt stimulates hepatic SREBP1c and lipogenesis through parallel mTORC1-dependent and independent pathways. *Cell Metab*. 14:21–32. <https://doi.org/10.1016/j.cmet.2011.06.002>
- Yoshioka, H., H.W. Coates, N.K. Chua, Y. Hashimoto, A.J. Brown, and K. Ohgane. 2020. A key mammalian cholesterol synthesis enzyme, squalene monooxygenase, is allosterically stabilized by its substrate. *Proc. Natl. Acad. Sci. USA*. 117:7150–7158. <https://doi.org/10.1073/pnas.1915923117>
- Yu, C.Y., E. Theusch, K. Lo, L.M. Mangravite, D. Naidoo, M. Kutilova, and M.W. Medina. 2014. HNRNPA1 regulates HMGCR alternative splicing and modulates cellular cholesterol metabolism. *Hum. Mol. Genet*. 23: 319–332. <https://doi.org/10.1093/hmg/ddt422>
- Yue, S., J. Li, S.Y. Lee, H.J. Lee, T. Shao, B. Song, L. Cheng, T.A. Masterson, X. Liu, T.L. Ratliff, and J.X. Cheng. 2014. Cholesteryl ester accumulation induced by PTEN loss and PI3K/AKT activation underlies human prostate cancer aggressiveness. *Cell Metab*. 19:393–406. <https://doi.org/10.1016/j.cmet.2014.01.019>
- Zaugg, K., Y. Yao, P.T. Reilly, K. Kannan, R. Kiarash, J. Mason, P. Huang, S.K. Sawyer, B. Fuerth, B. Faubert, et al. 2011. Carnitine palmitoyltransferase 1C promotes cell survival and tumor growth under conditions of metabolic stress. *Genes Dev*. 25:1041–1051. <https://doi.org/10.1101/gad.198721>
- Zelcer, N., C. Hong, R. Boyadjan, and P. Tontonoz. 2009. LXR regulates cholesterol uptake through Idol-dependent ubiquitination of the LDL receptor. *Science*. 325:100–104. <https://doi.org/10.1126/science.1168974>
- Zelcer, N., L.J. Sharpe, A. Loregger, I. Kristiana, E.C. Cook, L. Phan, J. Stevenson, and A.J. Brown. 2014. The E3 ubiquitin ligase MARCH6 degrades squalene monooxygenase and affects 3-hydroxy-3-methylglutaryl coenzyme A reductase and the cholesterol synthesis pathway. *Mol. Cell. Biol*. 34:1262–1270. <https://doi.org/10.1128/MCB.01140-13>
- Zeng, L., M. Lu, K. Mori, S. Luo, A.S. Lee, Y. Zhu, and J.Y. Shyy. 2004. ATF6 modulates SREBP2-mediated lipogenesis. *EMBO J*. 23:950–958. <https://doi.org/10.1038/sj.emboj.7600106>
- Zhang, C., J. Liu, G. Huang, Y. Zhao, X. Yue, H. Wu, J. Li, J. Zhu, Z. Shen, B.G. Haffty, et al. 2016. Cullin3-KLHL25 ubiquitin ligase targets ACLY for degradation to inhibit lipid synthesis and tumor progression. *Genes Dev*. 30:1956–1970. <https://doi.org/10.1101/gad.283283.116>
- Zhang, J., F. Song, X. Zhao, H. Jiang, X. Wu, B. Wang, M. Zhou, M. Tian, B. Shi, H. Wang, et al. 2017a. EGFR modulates monounsaturated fatty acid synthesis through phosphorylation of SCD1 in lung cancer. *Mol. Cancer*. 16:127. <https://doi.org/10.1186/s12943-017-0704-x>
- Zhang, L., P. Rajbhandari, C. Priest, J. Sandhu, X. Wu, R. Temel, A. Castrillo, T.Q. de Aguiar Vallim, T. Sallam, and P. Tontonoz. 2017b. Inhibition of cholesterol biosynthesis through RNF145-dependent ubiquitination of SCAP. *eLife*. 6:e28766. <https://doi.org/10.7554/eLife.28766>
- Zhao, L.F., Y. Iwasaki, M. Nishiyama, T. Taguchi, M. Tsugita, M. Okazaki, S. Nakayama, M. Kambayashi, S. Fujimoto, K. Hashimoto, et al. 2012. Liver X receptor  $\alpha$  is involved in the transcriptional regulation of the 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene. *Diabetes*. 61: 1062–1071. <https://doi.org/10.2337/db11-1255>
- Zhao, S., C. Jang, J. Liu, K. Uehara, M. Gilbert, L. Izzo, X. Zeng, S. Trefely, S. Fernandez, A. Carrer, et al. 2020. Dietary fructose feeds hepatic lipogenesis via microbiota-derived acetate. *Nature*. 579:586–591. <https://doi.org/10.1038/s41586-020-2101-7>
- Zhou, T., J. Zhan, W. Fang, Y. Zhao, Y. Yang, X. Hou, Z. Zhang, X. He, Y. Zhang, Y. Huang, and L. Zhang. 2017. Serum low-density lipoprotein and low-density lipoprotein expression level at diagnosis are favorable prognostic factors in patients with small-cell lung cancer (SCLC). *BMC Cancer*. 17:269. <https://doi.org/10.1186/s12885-017-3239-z>