


ARTICLE

Open Access

Transcriptomic and genetic approaches reveal an essential role of the NAC transcription factor SINAP1 in the growth and defense response of tomato

Jiao Wang¹, Chenfei Zheng¹, Xiangqi Shao¹, Zhangjian Hu¹, Jianxin Li¹, Ping Wang¹, Anran Wang¹, Jingquan Yu¹ and Kai Shi¹ 

Abstract

With global climate change, plants are frequently being exposed to various stresses, such as pathogen attack, drought, and extreme temperatures. Transcription factors (TFs) play crucial roles in numerous plant biological processes; however, the functions of many tomato (*Solanum lycopersicum* L.) TFs that regulate plant responses to multiple stresses are largely unknown. Here, using an RNA-seq approach, we identified *SINAP1*, a NAC TF-encoding gene, which was strongly induced by various stresses. By generating *SINAP1* transgenic lines and evaluating their responses to biotic and abiotic stresses in tomato, we found that *SINAP1*-overexpressing plants showed significantly enhanced defense against two widespread bacterial diseases, leaf speck disease, caused by *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000, and root-borne bacterial wilt disease, caused by *Ralstonia solanacearum*. In addition, *SINAP1* overexpression dramatically improved drought tolerance in tomato. Although the *SINAP1*-overexpressing plants were shorter than the wild-type plants during the early vegetative stage, eventually, their fruit yield increased by 10.7%. Analysis of different hormone contents revealed a reduced level of physiologically active gibberellins (GAs) and an increased level of salicylic acid (SA) and abscisic acid (ABA) in the *SINAP1*-overexpressing plants. Moreover, EMSAs and CHIP-qPCR assays showed that *SINAP1* directly activated the transcription of multiple genes involved in GA deactivation and both SA and ABA biosynthesis. Our findings reveal that *SINAP1* is a positive regulator of the tomato defense response against multiple stresses and thus may be a potential breeding target for improving crop yield and stress resistance.

Introduction

With global climate change, various stresses, such as pathogen attack, drought, and extreme temperatures, are occurring more frequently across the world. As sessile organisms, plants are faced with such biotic and abiotic stresses in situ, which often drastically affect their growth and cause tremendous economic losses. Plants have evolved sophisticated defense mechanisms to precisely

sense and respond to pathogen attack and abiotic stresses. Briefly, plant cells detect environmental stimuli via specific sensors or receptors, which trigger downstream responses. These responses occur via the reactive oxygen species (ROS) burst; phosphorylation of mitogen-activated protein (MAP) kinases; transcriptional reprogramming; and metabolic changes at physiological, biochemical, cellular and molecular levels.

Transcription factors (TFs) play crucial roles in the transcriptional reprogramming of stress-related genes^{1–3}. In particular, NAC proteins such as NAM, ATAF1/2, and CUC2 compose a large plant-specific TF family and have

Correspondence: Kai Shi (kaishi@zju.edu.cn)

¹Department of Horticulture, Zhejiang University, 866 Yuhangtang Road, 310058 Hangzhou, People's Republic of China

These authors contributed equally: Jiao Wang, Chenfei Zheng

© The Author(s) 2020



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

a highly conserved N-terminal domain that functions as a DNA-binding domain (known as the NAC domain) and a variable C-terminal domain that is responsible for transcriptional regulation. Extensive studies have revealed that NAC TFs are involved in a wide variety of plant biological processes, including plant growth and development, leaf senescence, secondary wall formation, and responses to biotic and abiotic stresses^{4–11}. Nonetheless, the functions of NAC TFs are highly species specific. For instance, transgenic plants overexpressing *ATAF1* show reduced resistance to *Pseudomonas syringae* pv. *tomato* (*Pst*) DC3000, *Botrytis cinerea* (*B. cinerea*) and *Alternaria brassicicola* in *Arabidopsis*^{12,13}, while the *ATAF1* homolog in rice, *OsNAC6*, plays a positive role in plant defense, as transgenic plants overexpressing *OsNAC6* show enhanced resistance to *Magnaporthe grisea*¹⁴. Similarly, two closely related NAC TFs in tomato, jasmonic acid 2 (*JA2*) and *JA2*-like (*JA2L*), play opposite roles in immunity against *Pst* DC3000 by differentially regulating stomatal closure and reopening¹⁵. *Solanum lycopersicum* stress-related NAC1 (*SISR1*) functions positively in the defense against *Pst* DC3000 and *B. cinerea*, while *SISR1* acts as a negative regulator of drought tolerance in tomato¹⁶. Thus, the functions of NAC proteins in response to different stresses are complex and still obscure.

Many previous studies have demonstrated that some NAC TFs, such as *SINAC35* and *SISR1*, induce plant defense responses via salicylic acid (SA) or jasmonic acid (JA) (and their crosstalk) signaling pathways during pathogen attack. Under abiotic stress conditions, *SINAC5*, *SINAM1*, and other NAC TFs activate the transcription of defense-related genes mainly via the JA or abscisic acid (ABA) signaling pathways^{17,18}. It therefore seems that plant hormones play critical roles in the functions of NAC TFs, which is worthy of further investigation.

Tomato (*S. lycopersicum* L.) is one of the most popular vegetable species in the world. However, many tomato cultivars are susceptible to pathogen infections and abiotic stresses, which cause billions of dollars of crop yield losses every year^{19,20}. In particular, *Pst* DC3000, which is an aboveground bacterial pathogen that causes leaf speck disease, resulting in a large yield penalty of vegetable crops, has been used as a model pathogen for understanding plant–bacterial interactions since the early 1990s²¹. *Ralstonia solanacearum*, a belowground bacterial pathogen that causes root-borne bacterial wilt disease, is one of the most aggressive and destructive pathogens worldwide due to the high mortality rates of diseased plants and the lack of effective control measures²². Moreover, drought is considered one of the most significant abiotic stresses, severely limiting plant growth and crop productivity⁹. There are 101 NAC TFs in tomato, but only ~10–20 NAC TFs have been functionally

characterized¹⁸. Here, using transcriptomic and genetic approaches, we identified a NAC TF, *SINAP1*, and characterized its functions in defense against *Pst* DC3000 and *R. solanacearum*, as well as in drought tolerance. This study highlights the positive role of *SINAP1* TFs in the defense against multiple stresses in tomato, suggesting that *SINAP1* can be a compelling biotechnological target to improve crop resilience.

Results

Changes in the transcript levels of tomato NAC genes in response to *Pst* DC3000 infection


To analyze the expression of tomato NAC genes in response to *Pst* DC3000 infection, we acquired the expression ratio (*Pst* DC3000/mock) data from two transcriptome profiles generated at 6 and 12 h post-inoculation (hpi) with *Pst* DC3000 in tomato leaves. We screened and obtained a list of 101 NAC TFs and their gene expression ratios after *Pst* DC3000 infection (Fig. 1a and Supplementary Table S2). Of these, 11 NAC genes were significantly upregulated by *Pst* DC3000 infection (expression ratio >1.5, and RPKM > 0) (Supplementary Table S2). We further analyzed the expression of these 11 NAC genes at 12 hpi via qRT-PCR. As shown in Fig. 1b, the expression of Solyc05g007770.2.1 (hereafter, *SINAP1*) was the greatest after 12 h of *Pst* DC3000 infection, which was more than 23-fold that in the control plants.

To determine the subcellular localization of tomato *SINAP1*, a 35S:*SINAP1*-GFP construct was transiently expressed in *Nicotiana benthamiana* (*N. benthamiana*) leaves. We found that *SINAP1*-GFP localized to the plasma membrane and nucleus (Fig. 1c).

SINAP1 inhibits vegetative growth but increases fruit yield of tomato plants

To explore the function of *SINAP1* in tomato, we generated *SINAP1*-overexpressing plants. The *SINAP1*-HA fusion protein was detected in two independent homozygous F2 progeny lines, OE-*SINAP1*-1 and OE-*SINAP1*-2, by western blot analysis (Fig. 2a). The *SINAP1*-overexpressing plants presented reduced plant height and leaf area compared with those of the WT plants up to 75 days after sowing (Fig. 2b, c), without a significant difference in their stem diameters (Fig. 2d). The net photosynthesis rate (Pn) of leaves significantly decreased in the *SINAP1*-overexpressing plants compared with the WT plants (Supplementary Fig. S1). *SINAP1* overexpression initially inhibited vegetative growth, although the height of *SINAP1*-overexpressing plants was close to that of the WT plants at 75 days after sowing (Fig. 2b).

Compared to those of the WT plants, the size of the fruits of both transgenic *SINAP1*-overexpressing plants was smaller (Fig. 2e, g), and the fresh weight per fruit of



the OE-*SINAP1*-1 and OE-*SINAP1*-2 plants decreased by 41.0% and 34.1%, respectively (Fig. 2f). However, compared with those of the WT plants, the fruit number per plant and the fruit yield of the *SINAP1*-overexpressing plants increased significantly. The fruit number per plant of the OE-*SINAP1*-1 and OE-*SINAP1*-2 plants increased by 92.6% and 81.5%, respectively (Fig. 2h), and the fruit yield increased by 10.7% and 11.0%, respectively (Fig. 2j). The dry matter contents of the roots and stems were nearly the same, whereas the dry matter contents of the leaves and fruits showed significant differences between the *SINAP1*-overexpressing plants and WT plants (Fig. 2i). Compared to those of the WT plants, the dry matter contents of leaves of the OE-*SINAP1*-1 and OE-*SINAP1*-2 plants decreased by 12.9% and 17.1%, respectively, while the dry matter contents of the fruits increased by 11.6% and 14.1%, respectively (Fig. 2i).

SINAP1 positively regulates disease resistance and drought tolerance

To determine the role of *SINAP1* in the defense against multiple stresses, first, WT and *SINAP1*-overexpressing plants were inoculated with *Pst* DC3000. We found that the *SINAP1*-overexpressing plants showed significantly enhanced resistance to *Pst* DC3000, as reflected by reduced cell death and bacterial populations in tomato leaves compared to those in the WT plants (Fig. 3a, b). Moreover, we inoculated the WT and *SINAP1*-overexpressing plants with *R. solanacearum*. In this case, the *SINAP1*-overexpressing plants also showed increased resistance to *R. solanacearum*, as evidenced by a relatively healthy appearance and reduced bacterial populations (Fig. 3c–d). At 12 dpi, the WT plants showed severe wilting, while the *SINAP1*-overexpressing plants exhibited minor symptoms (Fig. 3c). Taken together, these results




Fig. 2 Effects of *SINAP1* overexpression on plant growth and fruit phenotype. **a** Detection of the *SINAP1*-HA fusion protein in two independent homozygous *SINAP1*-overexpressing lines (OE-*SINAP1*-1 and -2) by western blot analysis using an anti-HA (Pierce) monoclonal antibody and the large subunit of Rubisco (RBC; as a loading control) in this experiment. **b**, **c**, **d** Plant phenotype in the early growth stage and growth indexes. The images were taken at 40 days (**c**) after sowing. Bar = 5 cm. Plant height and width were determined every 5 days after 25 days of sowing. **e** Plant phenotype in the advanced reproductive stage. Bar = 10 cm. **f**, **g** Fresh weight per fruit (**g**) and ripened fruit phenotype. Bar = 1 cm. **h**, **j** Fruit number per plant and yield (kg/m²). **i** Dry matter contents of tomato plant roots, stems, leaves, and fruits. The results are presented as the averages \pm SDs of five plants, which were grown in a greenhouse. The different letters indicate significant differences between treatments ($P < 0.05$, Tukey's test)

indicate that, in tomato, *SINAP1* plays a positive role in the defense against both above- and belowground bacterial pathogens.

Furthermore, the WT and *SINAP1*-overexpressing plants were subjected to water deprivation for 14 days. Compared with the WT plants, the *SINAP1*-overexpressing plants showed enhanced tolerance to drought, as reflected by relatively minor wilting and significantly reduced ion leakage in the leaves under drought (Fig. 3e–f). These results demonstrate that *SINAP1* also

functions as a positive regulator of drought tolerance in tomato.

***SINAP1* suppresses the accumulation of GAs and promotes the biosynthesis of SA and ABA**

Plant hormones play critical roles in the regulation of plant growth and defense responses to biotic and abiotic stresses. To determine whether *SINAP1* participates in the regulation of plant hormone homeostasis and subsequent plant growth and defense, we first measured the





Fig. 3 Effects of *SINAP1* overexpression on tomato disease resistance and drought tolerance. **a** Trypan blue staining for cell death in tomato leaves sampled at 2 dpi with *Pst* DC3000. Bar = 500 μ m. **b** *Pst* DC3000 bacterial population at 2 dpi. **c** Phenotypes of plants at 12 dpi with *R. solanacearum*. Bar = 5 cm. **d** *R. solanacearum* bacterial population at 12 dpi. **e** Phenotypes of plants after the imposition of drought for 7 days. Bar = 5 cm. **f** Ion leakage of plant leaves under drought for 7 days. The results represent the averages \pm SDs, $n = 3$. The different letters indicate significant differences between treatments ($P < 0.05$, Tukey's test). The experiments described above were performed three times, each yielding similar results

contents of growth- and stress-related plant hormones, such as GA, SA and ABA, in WT and *SINAP1*-overexpressing plants. As shown in Fig. 4a, the levels of GA₄, an important biologically active GA in plants, were lower in the *SINAP1*-overexpressing plants than in the WT plants, while the contents of SA and ABA were higher. Moreover, we analyzed the expression of genes involved in GA, SA and ABA metabolism, including *GIBBERELLIN 2-OXIDASE 3* (*SIGA2ox3*), which encodes a major GA deactivation enzyme, *PHENYLALANINE AMMONIA-LYASE 3* (*SIPAL3*), which encodes a major SA biosynthesis-related enzyme, and *9-CIS-EPOXYCAROTENOID DIOXYGENASE 1* (*SINCED1*), which encodes a rate-limiting enzyme involved in ABA biosynthesis. The transcript levels of *SIGA2ox3*, *SIPAL3* and *SINCED1* all significantly increased in the *SINAP1*-overexpressing plants compared with the WT plants (Fig. 4b). These results imply that

SINAP1 suppresses GA accumulation and promotes SA and ABA biosynthesis by regulating the transcription of *SIGA2ox3*, *SIPAL3* and *SINCED1* in tomato.

SINAP1* directly targets *SIGA2ox3*, *SIPAL3*, and *SINCED1

To further confirm whether *SINAP1*, a NAC TF, can regulate *SIGA2ox3*, *SIPAL3*, and *SINCED1* transcription through direct interaction between *SINAP1* and the promoter regions of target genes, we conducted an EMSA in vitro. Typically, NAC TFs preferentially bind to CACG, which is referred to as the NAC core-binding site, of their target promoters²³. Sequence analysis showed that the promoters of *SIGA2ox3*, *SIPAL3*, and *SINCED1* all had a NAC core-binding site. We then designed DNA probes that encompassed the NAC core-binding site and generated *SINAP1*-His fusion proteins. The results showed that the *SINAP1*-His fusion protein could bind to the




promoters of *SIGA2ox3*, *SIPAL3* and *SINCED1* and cause a mobility shift (Fig. 5a). When the NAC core binding sequences in the *SIGA2ox3*, *SIPAL3*, and *SINCED1* probes were mutated (-mut), the binding ability of SINAP1 to the probes was lost (Fig. 5a). We then conducted ChIP assays to determine the direct interaction between SINAP1 and the promoters of the three genes in vivo. After ChIP-qPCR analysis with an anti-HA antibody, the *SIGA2ox3*, *SIPAL3*, and *SINCED1* promoters were significantly enriched in the OE-SINAP1-HA samples compared with those in the WT control sample, while the IgG control was not enriched (Fig. 5b). These results suggest that SINAP1 could directly target *SIGA2ox3*, *SIPAL3*, and *SINCED1* to regulate the metabolism of relevant hormones in tomato plants.

Discussion

Extreme weather events are occurring more frequently with global climate change. Thus, plants are faced with more adverse environmental conditions imposed by multiple stresses at the same time. TF-based genetic engineering is a powerful tool improving crop tolerance to stress. However, very few TFs that can regulate plant

responses to multiple stresses have been functionally characterized in tomato. In this study, we screened *SINAP1* from among all the members of the whole tomato NAC TF family (Fig. 1) and found that *SINAP1* can significantly enhance defense against multiple stresses and increase fruit yields (Figs. 2 and 3). In-depth molecular investigations revealed that *SINAP1* directly activated the transcription of genes involved in GA deactivation, as well as SA and ABA biosynthesis (Figs. 4 and 5). Our findings suggest that the tomato NAC TF *SINAP1* is a potentially valuable candidate for genetic engineering programs aiming to generate high-yielding crops that are tolerant to multiple stresses.

NAC TFs have been shown to play dual roles (positive and negative) in the plant defense response, depending on the stresses and plant species. Here, we found that *SINAP1* can significantly increase defense against multiple stresses, including defense against both above- and belowground bacterial pathogens, namely, *Pst* DC3000 and *R. solanacearum*, and tolerance to drought (Fig. 3). Notably, the tomato *SINAP2* and Arabidopsis AtNAP (Arabidopsis ortholog of *SINAP1*) *SINAP1* homologs have been identified as central positive regulators of leaf



senescence^{6,7}. There are several other functionally identified NAC TFs that are involved in the defense response to stresses in tomato. For example, JA2 and JA2L regulate immunity against bacterial pathogens¹⁵. Another NAC TF, *S. lycopersicum* JUNGBRUNNEN 1 (*SIJUB1*), plays a positive role in tomato tolerance to drought⁹. Based on these studies, it appears that these NAC TFs regulate only a single stress response in tomato, while SINAP1 regulates the response to multiple stresses. Additionally, we found that SINAP1 inhibited the growth but increased the fruit yield of tomato plants. Compared with the WT plants, *SINAP1*-overexpressing plants were shorter and had concomitantly smaller leaves (Fig. 2b–c), the phenotype of which was similar to that of transgenic tomato lines overexpressing *AtJUB1*⁹. SINAP1 significantly increased the fruit number per plant and fruit yield, possibly by allocating more photoassimilates from the leaves to the fruits (Fig. 2h–j), implying that there may be a trade-off between vegetative and reproductive development. Similarly, life history theory indicates that the vegetative and reproductive functions of plants compete for common limited resources. An increased carbon allocation to reproductive development results in a relative reduction


in vegetative growth^{24,25}. Overall, SINAP1 seems to be an essential NAC TF that plays critical roles in the defense against multiple stresses and the regulation of plant growth and fruit productivity.

During plant-pathogen interactions, SA is recognized as a major plant defense hormone against biotrophic and hemibiotrophic pathogen infection²⁶. We found that SINAP1 played a positive role in the defense against bacterial pathogens by increasing the SA contents and by selectively activating the expression of *SIPAL3*, which is involved in SA biosynthesis (Figs. 4 and 5). Compared with the WT plants, the *SINAP1*-overexpressing plants had higher SA contents and *SIPAL3* transcript levels (Fig. 4a–b), and the SINAP1 protein could bind to the promoter regions of *SIPAL3* (Fig. 5a–b). Similarly, a previous study revealed that ectopic overexpression of *SINAC35* in tobacco increased resistance to *R. solanacearum* by up-regulating the expression of the SA-responsive defense genes *PR1a*, *NPR1*, *PR2*, and *PR5*²⁷. Conversely, JA2 and JA2L play opposite roles in the defense against bacterial pathogens by regulating pathogen-induced stomatal closure and reopening, respectively¹⁵. Nonetheless, these NAC TFs are involved in the defense against pathogens via different

mechanisms, which may be related to their localization. JA2 and JA2L are expressed predominantly in guard cells¹⁵; however, SINAP1 was localized to the mesophyll cells of tomato leaves (Fig. 1c).

In addition, SINAP1 is involved in drought tolerance. Previous studies demonstrated that a reduced GA level promoted plant tolerance to drought, which was based on the fact that tomato plants overexpressing *AtGAMT1* (which encodes an enzyme that can catalyze the methylation of active GAs to generate inactive GA methyl esters) exhibited typical GA-deficiency phenotypes and increased tolerance to drought²⁸. The plant hormone ABA plays a crucial role in the defense against drought stress¹. Typically, plants synthesize ABA in different tissues to initiate defense responses under stress conditions²⁹. Therefore, we analyzed the contents of GAs and ABA in both *SINAP1*-overexpressing and WT plants. We found that the active GA₄ content was reduced in the *SINAP1*-overexpressing plants compared to the WT plants, while the ABA content increased (Fig. 4a). The transcription of *SIGA2ox3* and *SINCE1* was significantly induced by *SINAP1* overexpression (Fig. 4b), and the SINAP1 protein could bind to the promoter regions of *SIGA2ox3* and *SINCE1* (Fig. 5a–b). These results imply that, as a positive regulator, SINAP1 could induce drought tolerance, possibly by activating the expression of *SIGA2ox3* and *SINCE1*, which are involved in GA deactivation and ABA biosynthesis in tomato. Similar to SINAP1, SIJUB1 also plays a positive role in drought tolerance. SIJUB1 targets *SIDREB1*, *SIDREB2*, and *SIDELLA* to transcriptionally regulate GA signaling pathways⁹. In addition, GAs play a critical role in the regulation of plant growth and act as negative regulators of growth cessation³⁰. Transgenic tomato lines overexpressing *AtJUB1* show a stunted phenotype due to GA and BR deficiencies⁹. Thus, we speculate that SINAP1 directly activates *SIGA2ox3* transcription to inhibit plant growth via GA deactivation in tomato.

Additionally, emerging evidence has indicated the presence of complex crosstalk between different hormones to coordinate plant growth and development, as well as responses to stress. For example, many constitutive defense mutants with elevated SA levels exhibit dwarf plant phenotypes, implying that SA may function antagonistically to GA to modulate plant growth³¹. In contrast, GA synergistically functions with SA in regulating disease resistance³². The crosstalk between SA and ABA is mainly antagonistic, and many studies show that ABA can attenuate plant defense responses³³. Moreover, GA and ABA play antagonistic roles in the regulation of seed germination, root growth, fruit set and responses to abiotic stresses^{34,35}. In our study, we found that there were significant differences in the contents of GA, SA and ABA hormones together with altered plant morphology, fruit size, and stress



tolerance between the WT and *SINAP1*-overexpressing plants (Figs. 2–4). We speculate that crosstalk among the three hormones, rather than the activity of a single hormone, might regulate these various processes in *SINAP1*-overexpressing plants, which needs further investigation. Moreover, SINAP1 directly activated the transcription of genes involved in the metabolism of these multiple stress-related hormones (Figs. 4 and 5), which appears to be a possible unique feature of SINAP1 as a positive regulator of both biotic and abiotic stress responses.

In conclusion, we identified a vital tomato NAC TF, SINAP1, which positively regulates the defense against multiple stresses by promoting GA deactivation and by stimulating SA and ABA biosynthesis (Fig. 6), further indicating the importance of NAC TFs in crop improvement.

Materials and methods

Plant materials and growth conditions

All the tomato (*S. lycopersicum* L.) lines generated in the present study were in the Condine Red (CR) wild-type (WT) background. To construct transgenic plants overexpressing *SINAP1*, the coding DNA sequence (CDS) of *SINAP1* was obtained from tomato cDNA via PCR using specific primers (Supplementary Table S1). The PCR product was subsequently inserted into a pFGC1008-3HA binary plasmid vector under the control of the 35S CaMV promoter. The resulting OE-SINAP1-3HA plasmid was transformed into tomato (cultivar Condine Red),

mediated by *Agrobacterium tumefaciens* strain GV3101. The transgenic plants were further verified by western blotting using anti-HA (Pierce) monoclonal antibodies (Fig. 2a). Two independent F2 progeny homozygous lines (OE-SINAP1-1 and -2) were used for phenotypic and molecular characterization.

Tomato seeds were germinated in 72-well trays in growth medium consisting of a mixture of peat and vermiculite (3:1, v/v). The tomato seedlings were transferred to plastic pots when two true leaves had fully expanded. The growth conditions were as follows: 500 $\mu\text{mol m}^{-2} \text{s}^{-1}$ of photosynthetic photon flux density (PPFD), a 13/11 hours (h) (day/night) photoperiod, a 25/21 °C (day/night) air temperature and 80% relative humidity. Approximately 4 weeks after sowing, tomato seedlings at the ~5-leaf stage were used for each treatment.

Pathogen inoculation and disease symptom assays

The bacterium *Pst* DC3000 was cultured in King's B solid media consisting of 25 mg mL⁻¹ rifampicin at 28 °C overnight and resuspended in 10 mM MgCl₂. Whole plants were inoculated with the bacterial suspension at a concentration of 10⁷ colony-forming units (CFU) mL⁻¹ together with 0.02% Silwet L-77 by spraying³⁶. Trypan blue staining and bacterial population counting (CFU) were conducted at 2 days post-inoculation (dpi) to assess disease symptoms^{37,38}.

The bacterium *R. solanacearum* was cultured on caseino peptone agar (CPG) supplemented with 1% triphenyltetrazolium chloride (TZC) at 28 °C for two days. The bacteria were resuspended in sterile water at 10⁹ CFU mL⁻¹. Bacterial suspensions (~50 mL) were applied to each plastic pot of tomato plants using the root inoculation method³⁹. CFU were measured to calculate the bacterial populations using stem slices at 12 dpi³⁸.

Drought treatments and ion leakage assays

For drought treatments, a water-withholding experiment was performed for ~14 days, while control plants were maintained under well-watered conditions by regular watering. For ion leakage measurements, initial electrical conductivity was measured using a conductometer (SI Analytics, Mainz, Germany) after immersing ~0.2 g of tomato leaves in 20 mL of deionized water and shaking them at 28 °C for 2 h. Total conductivity was measured after the leaves were boiled at 95 °C for half an hour and returned to 25 °C. The electrolyte leakage is presented as the percentage of the initial conductivity of the total conductivity.

Gene expression analysis

RNA was isolated using an RNA extraction kit (Easy-Do Biotech, Zhejiang, China) and reverse transcribed using a ReverTra Ace qPCR RT Kit (Toyobo, Tokyo, Japan) according to the manufacturers' instructions. Quantitative

real-time PCR (qRT-PCR) was conducted using a Light Cycler 480 II real-time PCR system (Roche, Basel, Switzerland). Each reaction buffer (20 μl) consisted of 10 μl of SYBR Green PCR Master Mix (Takara), 7.2 μl of water, 0.4 μl each of forward and reverse primers, and 2 μl of cDNA. The PCR conditions were as follows: 3 min at 94 °C, followed by 40 cycles of 30 s at 95 °C, 30 s at 58 °C and 1 min at 72 °C. The specific primers used for the target genes and internal control actin genes are listed in Supplementary Table S1.

Subcellular localization

SINAP1 was cloned into a pCambia2300 (Cambia) vector with a GFP tag at the C-terminus under the control of the 35S CaMV promoter. Transgenic tobacco leaves were infiltrated with the resulting 35S:SINAP1-GFP construct mediated by *Agrobacterium tumefaciens* strain GV3101, according to the methods of Liao et al.⁴⁰. The transgenic tobacco used in this experiment contained a nuclear localization protein that could emit red fluorescent signal (NLS-mCherry)⁴¹. At 48 h after infiltration, the fluorescence of the leaves was observed and recorded with a Zeiss LSM 780 confocal microscope; the excitation/emission wavelengths for GFP were 488 nm/500–530 nm and 561 nm/580–620 nm for NLS-mCherry. The primers used in this experiment are presented in Supplementary Table S1.

Electrophoretic mobility shift assays and chromatin immunoprecipitation

To generate pET-32a-His-SINAP1 constructs, the full-length CDS of SINAP1 was cloned via PCR using specific primers (Supplementary Table S1). The PCR product was then inserted into a pET-32a vector between the Sac I and Hind III sites. The recombinant SINAP1-His protein was expressed in *Escherichia coli* BL 21 (DE3) and purified following the instructions provided by the Novagen pET purification system. The ends of the oligonucleotide probes were labeled with biotin according to the instructions from a Biotin 3' End DNA Labeling Kit (89818, Pierce) and annealed to double-stranded probe DNA. An electrophoretic mobility shift assay (EMSA) was conducted according to the instructions of a Light Shift Chemiluminescent EMSA Kit (20148, Thermo Fisher Scientific). The DNA probes used in the EMSA are shown in Supplementary Table S1.

Chromatin immunoprecipitation (ChIP) experiments were conducted with lateral leaflets from the upper leaves of mature 35S:SINAP1-HA and WT plants following the instructions of a EpiQuik Plant ChIP Kit (P-2003-2, Epigentek). Approximately 1.5 g of leaf tissue was used and immunoprecipitated with anti-HA antibodies (26183, Pierce), with goat anti-mouse IgG antibodies (Millipore, AP124P) used as negative controls in this

experiment. The primers used for ChIP-qPCR are listed in Supplementary Table S1.

Measurement of phytohormones

Endogenous gibberellins (GAs) levels in tomato leaves were measured using a derivation approach coupled with nano-LC-electrospray ionization-quadrupole-time-of-flight-MS analysis^{42,43}. SA and ABA in the tomato leaves were extracted and analyzed by light chromatography (LC)/tandem mass spectrometry (MS/MS) on a 1290 Infinity HPLC system coupled to a 6460 Triple Quad LC-MS device (Agilent Technologies), according to the methods of Wang et al.⁴⁴.

RNA-seq library preparation and sequencing

To extract RNA for sequencing, tomato leaves were collected at 12 h after *Pst* DC3000 infection and then immediately frozen in liquid nitrogen. Each treatment consisted of three biological replicates, and the samples of each biological replicate were taken from at least three plants. RNA-seq was conducted by staff at LC-Bio Technologies (Hangzhou, China).

Another RNA-seq dataset (6 h post-inoculation with *Pst* DC3000) was obtained from the “Tomato functional genomics database” (TFGD), and the transcriptome of tomato leaves treated with different bacteria and PAMPs was sequenced by BTI at Cornell University.

Statistical analysis

At least three independent biological replicates sampled from different plants were included in each experiment. The data obtained were subjected to analysis of variance by SAS software, version 8 (SAS Institute), and the averages were compared using Tukey’s test ($P < 0.05$).

Accession numbers

Sequence data from this article can be found in the Sol Genomics Network (<http://solgenomics.net/>) database under the following accession numbers: Solyc02g069960, Solyc03g083880, Solyc03g115850, Solyc04g009440, *SINAP1* (Solyc05g007770), Solyc05g021090, Solyc05g055470, Solyc06g069710, Solyc06g074170, Solyc06g083850, Solyc07g063410, *SIACTIN* (Solyc03g078400), *SIGA2ox3* (Solyc01g079200), *SIPAL3* (Solyc09g007920), and *SINCED1* (Solyc07g056570).

Acknowledgements

This work was supported by the National Key Research and Development Program of China (2018YFD1000800), the National Natural Science Foundation of China (31822046, 31772355), the Natural Science Foundation of Zhejiang Province for Distinguished Young Scholar (LR19C150001), and the Fundamental Research Funds for the Central Universities.

Author contributions

K.S. conceived the research; J.W., C.Z., and K.S. designed the experiments; J.W., C.Z., X.S., Z.H., J.L., P.W., and A.W. performed the research and analyzed the data;

J.Y. provided technical/intellectual support for the research; and J.W. and K.S. wrote the article, with contributions from other authors.

Conflict of interest

The authors declare that they have no conflict of interest.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41438-020-00442-6>).

Received: 20 May 2020 Revised: 1 November 2020 Accepted: 5 November 2020

Published online: 25 December 2020

References

1. Ahammed, G. J. et al. Tomato *WRKY81* acts as a negative regulator for drought tolerance by modulating guard cell H₂O₂-mediated stomatal closure. *Environ. Exp. Bot.* **171**, 103960 (2020).
2. Ganie, S. A., Ahammed, G. J. & Wani, S. H. Vascular plant one zinc-finger (VOZ) transcription factors: novel regulators of abiotic stress tolerance in rice (*Oryza sativa* L.). *Genet. Resour. Crop Ev.* **67**, 799–807 (2020).
3. Hénanff, G. L. et al. Grapevine NAC1 transcription factor as a convergent node in developmental processes, abiotic stresses, and necrotrophic/biotrophic pathogen tolerance. *J. Exp. Bot.* **64**, 4877–4893 (2013).
4. Chai, M. F. et al. The NAC transcription factor OsSWN1 regulates secondary cell wall development in *Oryza sativa*. *J. Plant Biol.* **58**, 44–51 (2015).
5. Christianson, J. A., Dennis, E. S., Llewellyn, D. J. & Wilson, I. W. ATAF NAC transcription factors: regulators of plant stress signaling. *Plant Signal. Behav.* **5**, 428–432 (2010).
6. Guo, Y. F. & Gan, S. S. AtNAP, a NAC family transcription factor, has an important role in leaf senescence. *Plant J.* **46**, 601–612 (2006).
7. Ma, X. et al. The NAC transcription factor SINAP2 regulates leaf senescence and fruit yield in tomato. *Plant Physiol.* **177**, 1286–1302 (2018).
8. Olsen, A. N., Ernst, H. A., Leggio, L. L. & Skriver, K. NAC transcription factors: structurally distinct, functionally diverse. *Trends Plant Sci.* **10**, 79–87 (2005).
9. Thirumalaikumar, V. P. et al. NAC transcription factor JUNGBRUNNEN1 enhances drought tolerance in tomato. *Plant Biotechnol. J.* **16**, 354–366 (2018).
10. Tran, L. S. P., Nishiyama, R., Yamaguchi-Shinozaki, K. & Shinozaki, K. Potential utilization of NAC transcription factors to enhance abiotic stress tolerance in plants by biotechnological approach. *GM Crops* **1**, 32–39 (2010).
11. Zhong, R. Q., Lee, C. H. & Ye, Z. H. Global analysis of direct targets of secondary wall NAC master switches in *Arabidopsis*. *Mol. Plant* **3**, 1087–1103 (2010).
12. Wang, X. et al. The *Arabidopsis* ATAF1, a NAC transcription factor, is a negative regulator of defense responses against necrotrophic fungal and bacterial pathogens. *Mol. Plant Microbe Interact.* **22**, 1227–1238 (2009).
13. Wu, Y. et al. Dual function of *Arabidopsis* ATAF1 in abiotic and biotic stress responses. *Cell Res.* **19**, 1279–1290 (2009).
14. Nakashima, K. et al. Functional analysis of a NAC-type transcription factor OsNAC6 involved in abiotic and biotic stress-responsive gene expression in rice. *Plant J.* **51**, 617–630 (2007).
15. Du, M. M. et al. Closely related NAC transcription factors of tomato differentially regulate stomatal closure and reopening during pathogen attack. *Plant Cell* **26**, 3167–3184 (2014).
16. Liu, B. et al. Tomato NAC transcription factor SISR1 positively regulates defense response against biotic stress but negatively regulates abiotic stress response. *PLoS ONE* **9**, e102067 (2014).
17. Nakashima, K., Takasaki, H., Mizoi, J., Shinozaki, K. & Yamaguchi-Shinozaki, K. NAC transcription factors in plant abiotic stress responses. *Biochim. Biophys. Acta* **1819**, 97–103 (2012).
18. Tweneboah, S. & Oh, S.-K. Biological roles of NAC transcription factors in the regulation of biotic and abiotic stress responses in solanaceous crops. *J. Plant Biotechnol.* **44**, 1–11 (2017).
19. Nuruddin, M., Madramootoo, C. A. & Dodds, G. T. Effects of water stress at different growth stages on greenhouse tomato yield and quality. *Hortic. Sci.* **38**, 1389–1393 (2003).
20. Qi, S. et al. The RING finger E3 Ligase SpRING is a positive regulator of salt stress signaling in salt-tolerant wild tomato species. *Plant Cell Physiol.* **57**, 528–539 (2016).

21. Xin, X. F. & He, S. Y. *Pseudomonas syringae* pv. *tomato* DC3000: a model pathogen for probing disease susceptibility and hormone signaling in plants. *Annu. Rev. Phytopathol.* **51**, 473–498 (2013).
22. French, E., Kim, B. S., Rivera-Zuluaga, K. & Iyer-Pascuzzi, A. S. Whole root transcriptomic analysis suggests a role for auxin pathways in resistance to *Ralstonia solanacearum* in tomato. *Mol. Plant Microbe Interact.* **31**, 432–444 (2018).
23. Tran, L. S. et al. Isolation and functional analysis of Arabidopsis stress-inducible NAC transcription factors that bind to a drought-responsive cis-element in the early responsive to dehydration stress 1 promoter. *Plant Cell* **16**, 2481–2498 (2004).
24. Obeso, J. R. The costs of reproduction in plants. *N. Phytol.* **155**, 321–348 (2002).
25. Reekie, E. G. & Avila-Sakar, G. The shape of the trade-off function between reproduction and growth. In *Reproductive Allocation in Plants*. 189–214 <https://doi.org/10.1016/B978-012088386-8/50007-7> (2005).
26. Vlot, A. C., Dempsey, D. A. & Klessig, D. F. Salicylic acid, a multifaceted hormone to combat disease. *Annu. Rev. Phytopathol.* **47**, 177–206 (2009).
27. Wang, G. et al. A stress-associated NAC transcription factor (SINAC35) from tomato plays a positive role in biotic and abiotic stresses. *Physiol. Plant.* **158**, 45–64 (2016).
28. Nir, I., Moshelion, M. & Weiss, D. The Arabidopsis gibberellin methyl transferase 1 suppresses gibberellin activity, reduces whole-plant transpiration and promotes drought tolerance in transgenic tomato. *Plant Cell Environ.* **37**, 113–123 (2014).
29. Chae, L., Woonhee, B., Jangho, J., Jung-Hyun, K. & Sung, L. Function of ABA in stomatal defense against biotic and drought stresses. *Int. J. Mol. Sci.* **16**, 15251–15270 (2015).
30. Sun, T. P. The molecular mechanism and evolution of the GA-GID1-DELLA signaling module in plants. *Curr. Biol.* **21**, 338–345 (2011).
31. Huot, B., Yao, J., Montgomery, B. L. & He, S. Y. Growth-defense tradeoffs in plants: a balancing act to optimize fitness. *Mol. Plant* **7**, 1267–1287 (2014).
32. Robert-Seilaniantz, A., Grant, M. & Jones, J. D. G. Hormone crosstalk in plant disease and defense: more than just jasmonate-salicylate antagonism. *Annu. Rev. Phytopathol.* **49**, 317–343 (2011).
33. Cao, F. Y., Yoshioka, K. & Desveaux, D. The roles of ABA in plant–pathogen interactions. *J. Plant Res.* **124**, 489–499 (2011).
34. Vishal, B. & Kumar, P. P. Regulation of seed germination and abiotic stresses by gibberellins and abscisic acid. *Front. Plant Sci.* **9**, 838 (2018).
35. Kumar, R., Khurana, A. & Sharma, A. K. Role of plant hormones and their interplay in development and ripening of fleshy fruits. *J. Exp. Bot.* **16**, 4561–4575 (2014).
36. Li, X. et al. Tomato–*Pseudomonas syringae* interactions under elevated CO₂ concentration: the role of stomata. *J. Exp. Bot.* **66**, 307–316 (2015).
37. Bai, S. et al. Structure-function analysis of barley NLR immune receptor MLA10 reveals its cell compartment specific activity in cell death and disease resistance. *PLoS Pathog.* **8**, e1002752 (2012).
38. Wolfe, J., Hutcheon, C. J., Higgins, V. J. & Cameron, R. K. A functional gene-for-gene interaction is required for the production of an oxidative burst in response to infection with avirulent *Pseudomonas syringae* pv. *tomato* in Arabidopsis thaliana. *Physiol. Mol. Plant P.* **56**, 253–261 (2000).
39. Caldwell, D., Kim, B.-S. & Iyer-Pascuzzi, A. S. *Ralstonia solanacearum* differentially colonizes roots of resistant and susceptible tomato plants. *Phytopathology* **107**, 528–536 (2017).
40. Liao, Y. W. K. et al. Salicylic acid binding of mitochondrial alpha-ketoglutarate dehydrogenase E2 affects mitochondrial oxidative phosphorylation and electron transport chain components and plays a role in basal defense against tobacco mosaic virus in tomato. *N. Phytol.* **205**, 1296–1307 (2015).
41. Yang, Q. S. et al. PpHB22, a member of HD-Zip proteins, activates PpDAM1 to regulate bud dormancy transition in ‘Suli’ pear (Pyrus pyrifolia White Pear Group). *Plant Physiol. Biochem.* **127**, 355–365 (2018).
42. Chen, M. L. et al. Highly sensitive and quantitative profiling of acidic phytohormones using derivatization approach coupled with nano-LCESI-Q-TOF-MS analysis. *J. Chromatogr. B. Anal. Technol. Biomed. Life Sci.* **905**, 67–74 (2012).
43. Li, X. J. et al. DWARF overexpression induces alteration in phytohormone homeostasis, development, architecture and carotenoid accumulation in tomato. *Plant Biotechnol. J.* **14**, 1021–1033 (2016).
44. Wang, F. et al. Phytochrome A and B function antagonistically to regulate cold tolerance via abscisic acid-dependent jasmonate signaling. *Plant Physiol.* **170**, 459–471 (2016).