

Office of Research and Development, and Biomedical Laboratory Research and Development (Washington, DC, USA) (grant no. BX-000282-05). The other authors have no conflicts to declare.

References

1. https://www.who.int/gho/ncd/risk_factors/blood_pressure_prevalence_text/en/.
2. <https://www.cdc.gov/nchs/products/databriefs/db289.htm#ref1>.
3. Hu G, Jousilahti P, Tuomilehto J. Joint effects of history of hypertension at baseline and type 2 diabetes at baseline and during follow-up on the risk of coronary heart disease. *Eur Heart J* 2007;**28**:3059–3066.
4. Zanchetti A, Thomopoulos C, Parati G. Randomized controlled trials of blood pressure lowering in hypertension: a critical reappraisal. *Circ Res* 2015;**116**:1058–1073.
5. Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, Chen J, He J. Global disparities of hypertension prevalence and control: a systematic analysis of population-based studies from 90 countries. *Circulation* 2016;**134**:441–450.
6. Pickering TG. The clinical significance of diurnal blood pressure variations. Dippers and nondippers. *Circulation* 1990;**81**:700–702.
7. Kimura G. Kidney and circadian blood pressure rhythm. *Hypertension* 2008;**51**:827–828.
8. Zhang LL, Zhang Y, Cheng YQ, Zhang JM, Liu HQ, Wang WZ, Mehta JL, Xiong ZG, Su DF, Liu AJ. Metabolic syndrome emerges after artificial selection for low baroreflex sensitivity. *CNS Neurosci Ther* 2018;**24**:828–836.
9. Kario K. Nocturnal hypertension: new technology and evidence. *Hypertension* 2018;**71**:997–1009.
10. Boggia J, Li Y, Thijss L, Hansen TW, Kikuya M, Björklund-Bodegård K, Richart T, Ohkubo T, Kuznetsova T, Torp-Pedersen C, Lind L, Ibsen H, Imai Y, Wang J, Sandoya E, O'Brien E, Staessen JA; International Database on Ambulatory blood pressure monitoring in relation to Cardiovascular Outcomes (IDACO) investigators. Prognostic accuracy of day versus night ambulatory blood pressure: a cohort study. *Lancet* 2007;**370**:1219–1229.
11. Salles GF, Reboldi G, Fagard RH, Cardoso CR, Pierdomenico SD, Verdecchia P, Eguchi K, Kario K, Hoshide S, Polonia J, de la Sierra A, Hermida RC, Dolan E, O'Brien E, Roush GC; Investigators ABC-H. Prognostic effect of the nocturnal blood pressure fall in hypertensive patients: the Ambulatory Blood Pressure Collaboration in Patients With Hypertension (ABC-H) meta-analysis. *Hypertension* 2016;**67**:693–700.
12. Hermida RC, Ayala DE, Mojón A, Fernández JR. Bedtime ingestion of hypertension medications reduces the risk of new-onset type 2 diabetes: a randomised controlled trial. *Diabetologia* 2016;**59**:255–265.
13. Hermida RC, Ayala DE, Mojón A, Fernández JR. Risk of incident chronic kidney disease is better reduced by bedtime than upon-awakening ingestion of hypertension medications. *Hypertens Res* 2018;**41**:342–353.
14. Wang C, Ye Y, Liu C, Zhou Y, Lv L, Cheng C, Li S, Lou T, Liu X. Evening versus morning dosing regimen drug therapy for chronic kidney disease patients with hypertension in blood pressure patterns: a systematic review and meta-analysis. *Intern Med J* 2017;**47**:900–906.
15. Bowles NP, Thosar SS, Herzog MX, Shea SA. Chronotherapy for hypertension. *Curr Hypertens Rep* 2018;**20**:97.
16. Ishikawa J, Shimizu M, Sugiyama Edison E, Yano Y, Hoshide S, Eguchi K, Kario K; J-TOP (Japan Morning Surge-Target Organ Protection) Study Investigators Group. Assessment of the reductions in night-time blood pressure and dipping induced by antihypertensive medication using a home blood pressure monitor. *J Hypertens* 2014;**32**:82–89.
17. Kario K, Hoshide S, Shimizu M, Yano Y, Eguchi K, Ishikawa J, Ishikawa S, Shimada K. Effect of dosing time of angiotensin II receptor blockade titrated by self-measured blood pressure recordings on cardiorenal protection in hypertensives: the Japan Morning Surge-Target Organ Protection (J-TOP) study. *J Hypertens* 2010;**28**:1574–1583.
18. Hermida RC, Crespo JJ, Domínguez-Sardiña M, Otero A, Moyá A, Ríos MT, Sineiro E, Castiñeira MC, Callejas PA, Pousa L, Salgado JL, Durán C, Sánchez JJ, Fernández JR, Mojón A, Ayala DE; Hygia Project Investigators. Bedtime hypertension treatment improves cardiovascular risk reduction: the Hygia Chronotherapy Trial. *Eur Heart J* 2019;doi:10.1093/eurheartj/ehz754.

Corrigendum

doi:10.1093/eurheartj/ehaa467
Online publish-ahead-of-print 25 May 2020

Corrigendum to: Prediction of individualized lifetime benefit from cholesterol lowering, blood pressure lowering, antithrombotic therapy, and smoking cessation in apparently healthy people [Eur Heart J 2020;41:1190–1199]

The article describes the development of a prediction algorithm. The formula of this algorithm, presented in the supplementary material, contains logarithmic transformations. These transformations were erroneously presented as ‘log’, but should have been written as ‘ln’. These errors have been corrected in the supplementary material.

Published on behalf of the European Society of Cardiology. All rights reserved. © The Author(s) 2020. For permissions, please email: journals.permissions@oup.com.