Published in final edited form as:

Am J Otolaryngol. 2021; 42(1): 102823. doi:10.1016/j.amjoto.2020.102823.

Tolerability and effectiveness of povidone-iodine or mupirocin versus saline sinus irrigations for chronic rhinosinusitis

Victoria S. Lee^{a,*}, Paul S. Pottinger^b, Greg E. Davis^a

^aDepartment of Otolaryngology – Head and Neck Surgery, University of Washington, Box 356515, Seattle, WA 98195, United States of America

^bDepartment of Medicine, Division of Allergy & Infectious Diseases, University of Washington, 1959 NE Pacific St., Courier BB-302, P.O. Box 356130, Seattle, WA 98195, United States of America

The authors regret there is an error in the published manuscript that has come to their attention. The concentration of the povidone-iodine irrigation used in this study was 0.1%. This concentration is still within the bactericidal range. The published concentration of 1% resulted from an inaccurate calculation. This error is present in the following sections: (1) Abstract, Results section, line 3; (2) Abstract, Conclusions section, line 3; (3) Table 1, row 2/column 1; (4) Materials and Methods, Section 2.2, lines 2 and 3; (5) Results, Section 3.3.2, line 4; and (6) Conclusions, line 6.

The authors would like to apologise for any inconvenience caused.

^{*}Corresponding author at: Department of Otolaryngology – Head and Neck Surgery, University of Illinois at Chicago, 1855 W. Taylor St., Room 2.42 (MC 648), Chicago, IL 60612, United States of America. vlee39@uic.edu, (V.S. Lee).