



Published in final edited form as:

Mol Cell. 2016 December 01; 64(5): 888–899. doi:10.1016/j.molcel.2016.09.040.

Hsp90 and p23 Molecular Chaperones Control Chromatin Architecture by Maintaining the Functional Pool of the RSC Chromatin Remodeler

Frank J. Echtenkamp^{1,*}, Zlata Gvozdenov^{1,2,*}, Nicholas L. Adkins³, Yang Zhang¹, Melinda Lynch-Day¹, Shinya Watanabe³, Craig L. Peterson³, Brian C. Freeman^{1,+}

¹University of Illinois, Urbana-Champaign, Department of Cell and Developmental Biology, 601 S. Goodwin Avenue, Urbana, IL 61801 USA

²Technische Universität München, Department Chemie, Garching, 85748 Germany

³University of Massachusetts Medical School, Program of Molecular Medicine, 373 Plantation Street, Worcester, MA 01605 USA

Abstract

Molecular chaperones govern protein homeostasis being allied to the beginning (folding) and ending (degradation) of the protein life cycle. Yet, the Hsp90 system primarily associates with native factors including fully assembled complexes. The significance of these connections is poorly understood. To delineate why Hsp90 and its cochaperone p23 interact with a mature structure we focused on the RSC chromatin remodeler. Both Hsp90 and p23 triggered the release of RSC from DNA or a nucleosome. While Hsp90 only freed bound RSC, p23 enhanced nucleosome remodeling prior to discharging the complex. In vivo, RSC mobility and remodeling function were chaperone-dependent. Our results suggest Hsp90 and p23 contribute to proteostasis by chaperoning mature factors through energetically unfavorable events thereby maintaining the cellular pool of active native proteins. In the case of RSC, p23 and Hsp90 promote a dynamic action allowing a limited number of remodelers to effectively maintain chromatin in a pliable state.

Graphical Abstract

address correspondence to: Dr. Brian C. Freeman, 217-244-2662, bfree@illinois.edu.


*authors contributed equally

+lead contact

AUTHOR CONTRIBUTIONS

FJE, ZG, NLA, MLD, SW and BCF conducted the experiments, ZG and YZ analyzed genomic data, BCF designed the experiments and wrote the paper.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



INTRODUCTION

The protein homeostasis or proteostasis process safeguards a cell's polypeptide population by continuously scrutinizing the conformational status of each protein. A persistent monitoring facilitates the forward flow of biological pathways including the folding of nascent polypeptide chains and removal of damaged proteins (Hartl *et al.*, 2011). This high volume work is predominately carried out by members of the diverse molecular chaperone family and disruptions in the process correlate with the onset of a variety of diseases including neurodegeneration, type II diabetes, and heart failure (Balch *et al.*, 2008; Powers *et al.*, 2009). The promiscuous, transient binding activities of the abundant chaperones are well suited to perform such high-volume tasks, which are further challenged by the vast sequence and structural variety of a proteome. Adding to the complexity is the range of conformations any protein can assume and still be within the scope of a native factor (Bartlett and Radford, 2009). Likely, different structural forms distinguish distinct functional states of each factor. Here, we suggest that the Hsp90 and p23 chaperones contribute to proteostasis by manipulating fully assembled protein structures thereby fostering the available pool of functional complexes (*i.e.*, facilitating specific activities).

In eukaryotes the central molecular chaperones are Hsp90 and Hsp70 along with their respective cochaperones (Röhl *et al.*, 2013). While the Hsp70 hub manages early and late events (*i.e.*, folding and degradation), the Hsp90 system preferentially associates with near native or native proteins (DeZwaan and Freeman, 2008). Although Hsp90 can form long-lived interactions with metastable proteins (*e.g.*, unactivated kinases and steroid hormone receptors), Hsp90 components generally associate weakly with their protein targets (Zhao *et al.*, 2005; Millson *et al.*, 2005; McClellan *et al.*, 2007; Sharma *et al.*, 2012; Taipale *et al.*,

2012; Taipale *et al.*, 2014). However, the relevance of transient chaperone interactions with seemingly native clients is yet to be fully understood.

As numerous Hsp90-targets are part of large multi-subunit structures, an influence in the assembly and/or disassembly of protein machines has been proposed (Makhnevych and Houry, 2012). In addition to Hsp90, its cochaperones often connect to the same complexes, albeit through distinct subunits, suggesting a joint chaperone action with protein structures (Echtenkamp and Freeman, 2012). Yet, studies outlining the roles of Hsp90 and any of its cochaperones on the same client are limited. Here, we investigated the impact of the p23 cochaperone and Hsp90 on the Remodel the Structure of Chromatin (RSC) complex, which is an ~1.5 mDa protein structure composed of 17 subunits (Cairns *et al.*, 1996). Our prior work genetically connected the yeast p23 gene with *RSC7* and two different studies linked *HSP90* with *RSC14* (Zhao *et al.*, 2005; McClellan *et al.*, 2007; Echtenkamp *et al.*, 2011).

RSC is one of eight chromatin remodelers in budding yeast, which are conserved through humans, responsible for depositing histones and positioning nucleosomes across the genome (Clapier and Cairns, 2009). RSC, however, is the only chromatin remodeler essential for growth (Cairns *et al.*, 1996). In brief, remodelers provide a critical means to regulate genomic activities as these complexes alter nucleosome spacing to control access to the underlying DNA (Venkatesh and Workman, 2015). A key feature of the chromatin process is the fast kinetics of histone repositioning/removal since many events (*e.g.*, DNA repair or transcription) must be undertaken promptly to maintain cellular homeostasis.

Challenging the effectiveness of remodeling pathways is the overabundance of nucleosomes relative to the total number of remodelers (Ghaemmaghami *et al.*, 2003; Brogaard *et al.*, 2012; Chong *et al.*, 2015). Although a division of labor amongst the remodelers may alleviate some of the burden, it likely doesn't account for the efficiency of the remodeling system. For instance, RSC favors nucleosomes at RNAP II gene promoters helping to maintain promoter-associated Nucleosome Depleted Regions (NDRs) (Parnell *et al.*, 2008; Yen *et al.*, 2012). Yet, within minutes of a signaling event RSC can redistribute to hundreds of new gene promoters to create NDRs at these loci (Damelin *et al.*, 2002; Ng *et al.*, 2002). While nucleation of RSC to certain promoters is mediated by select transcription factors along with the RSC DNA binding subunits Rsc3 and Rsc30 (Angus-Hill *et al.*, 2001; Badis *et al.*, 2008), the events triggering the release of RSC to allow the rapid relocation are not understood. Given the genetic links between both chaperones and RSC subunits along with the functional dependence of chromatin on p23 (Wilson *et al.*, 2006; Zelin *et al.*, 2012), we felt that the RSC complex served as an ideal model to understand how molecular chaperones modulate an assembled multi-subunit protein structure.

RESULTS

P23 is not a classic proteostasis factor of RSC proteins and complex assembly

The connection between p23 and RSC was initially discovered in a Synthetic Genetic Array (SGA) screen (Echtenkamp *et al.*, 2011). We validated the genetic interaction between *SBA1* (yeast p23 gene) and the non-essential RSC subunit gene *RSC7* using a plasmid dropout spot-test assay (Figure 1A). The only known role for Rsc7 is in the assembly of the RSC

structure, as incorporation of Rsc3 and Rsc30 is impaired in *rsc7* yeast (Wilson *et al.*, 2006). As construction of protein complexes is a common molecular chaperone function, we checked whether p23 mediates classic RSC proteostasis events. We found that steady state RSC protein levels and assimilation of the subunits into the RSC structure were similar in parental and *p23* yeast (Figure 1B and 1C). Thus, p23 is not a typical proteostasis factor of individual RSC proteins or RSC subunit assembly.

Next, we checked whether the p23 link might entail an undetermined, specialized role of Rsc7 or if it involved the RSC chromatin remodeler. Typically, remodeling function is tied to the catalytic motor of each remodeler (Clapier and Cairns, 2009). Complicating our analysis is the essential nature of the RSC ATPase encoding gene *STH1* (Cairns *et al.*, 1996). To circumvent this difficulty we took advantage of the *sth1* temperature sensitive derivative Sth1-F793S. Unlike other *sth1* allelic protein products, Sth1-F793S degrades at restrictive temperatures (>37°C) and therefore serves as an effective null (Titus *et al.*, 2010). Since *STH1* is essential, the final growth test is not performed at the restrictive temperature but rather the cells are exposed and then grown at a permissive temperature. In brief, exponentially growing parental, *p23*, *sth1-F793S*, and *p23 sth1-F793S* yeast were shifted to 37°C for 24 h, cells were then serially diluted, spotted onto media, and incubated at 30°C. The double mutant *p23 sth1-F793S* survived poorly relative to the other strains indicating a possible physiological connection between p23 and the remodeling function of RSC (Figure 1D).

P23 triggers release of RSC from promoter DNA

We have shown that the maintenance of Nucleosome Depleted Regions (NDRs) is contingent upon p23 (Zelin *et al.*, 2012). We had postulated that p23's ability to regulate the DNA binding activities of numerous transcription factors accounted for the NDR effects; however, transcription factors do not directly move nucleosomes. Besides transcription factors an increase in the DNA occupancy of sites associated with chromatin modifiers, including Rsc3 and Rsc30 elements, was apparent (Zelin *et al.*, 2012). Of note, the majority of the genetic links between p23 and the various chromatin modifiers involve DNA binding subunits, suggesting a common basis for the connections (Echtenkamp *et al.*, 2011). Hence, we checked the influence of p23 on the DNA binding activity of RSC.

RSC uses multiple subunits to bind comparably to naked or nucleosomal DNA with the Rsc3/30 heterodimer providing sequence selectivity (Angus-Hill *et al.*, 2001; Sengupta *et al.*, 2001; Damelin *et al.*, 2002; Titus *et al.*, 2010). To monitor RSC DNA interactions we used a probe representing a section of the *PHO8* promoter that has two Rsc3/30 binding sites. RSC positions nucleosomes at the *PHO8* promoter in vivo and in vitro (Wippo *et al.*, 2011). Purified RSC bound the PHO8 probe and the addition of recombinant p23 dissociated the RSC-DNA structure (Figure 2A). Dissolution of the complex was dependent upon p23 chaperone activity since the defective mutant p23⁸⁴ had no apparent impact on RSC binding (Figure 2A) (Toogun *et al.*, 2007). While the effective p23 amounts were seemingly high, the applied levels are comparable to the nuclear concentration of p23 (~4 μM) (Figure S1A). Of note, p23 does not itself interact with DNA (Figure S1B).

To further mimic in vivo settings where numerous RSC binding sites are available, we challenged the radiolabeled-DNA RSC complex with unlabeled PHO8 probe and observed an additive effect of p23 and competitor DNA on loss of the radiolabeled RSC structure (Figure 2B). To further understand the impact of p23 on RSC we utilized fluorescence anisotropy. We found that p23 accelerates the off-rate of RSC from DNA alone or in the presence of competitor DNA (Figure S2). Hence, p23 disassembles RSC-DNA complexes thereby fostering the transition of RSC to new sites.

The nucleosome remodeling activity of RSC is advanced by p23

Although RSC binds naked and nucleosomal DNA with comparable affinities, the target of interest is a nucleosome (Sengupta *et al.*, 2001). To test whether p23 influences RSC-nucleosome interactions we built a mononucleosome with a centrally localized histone octamer using a 601 nucleosome-positioning DNA as the probe in an EMSA-based assay (Sengupta *et al.*, 2001). RSC bound the nucleosome independent of ATP (Figure 3A Complex 1) and formed an activated nucleosome structure upon ATP addition (Complex 2) (Lorch *et al.*, 1998). In the presence of p23 the remodeling reaction was driven to completion as evidenced by the release of terminally localized mononucleosomes and/or free DNA (Figure 3A lanes 10–13). Of note, p23 promoted the release of slid nucleosomes or free DNA even in the absence of competitor DNA, which is typically added to remove RSC from its modified target (Figure S3A). In the absence of ATP, p23 triggered dissociation of RSC from the unremodeled mononucleosomes (Figure 3A lane 14). Thus, p23 differentially impacts RSC in an ATP-dependent manner (*i.e.*, in the presence of ATP p23 fosters the remodeling reaction but in the absence of ATP p23 dissociates the idle complex).

Comparable to the EMSA results, p23 enhanced RSC-mediated histone repositioning and eviction using a mononucleosome sliding assay (Figure S3B and S3C). Importantly, the chaperone activity of p23 was required to induce RSC-dependent mononucleosome sliding and histone eviction (Figure 3B). Besides RSC we assessed the chaperone impact on ISW2, which is one of two remodelers without an apparent genetic connection to p23 (Echtenkamp *et al.*, 2011). P23 did not alter ISW2 remodeling activity (Figure S3D). Overall, p23 appears to selectively modulate RSC by targeting the remodeler-nucleosome complexes at potentially unproductive stages (*i.e.*, post-remodeling or in the absence of ATP) and either drives the reaction to completion (+ATP) or releases the stalled complex (–ATP). In both cases p23 contributes to RSC action by recycling otherwise inert RSC-nucleosome assemblies.

Rsc3 is released from DNA by p23

To identify the RSC components influenced by p23 we screened the subunits by yeast 2-hybrid. Of the 17, 3 displayed a relationship with p23 (Figure 4A). Rsc30 had a standard positive 2-hybrid result, as the *HIS3* reporter was active upon co-expression of the bait (DBD-p23) and prey (AD-Rsc30). On the other hand, AD-Rsc3 and AD-Sfh1 both autoactivated the *GAL1-HIS3* reporter yet coexpression of DBD-p23 abrogated the autoactivation. As we used a standard Gal4-based 2-hybrid strategy and RSC is known to modulate the endogenous *GAL1* promoter in an Rsc3-dependent manner (James *et al.*, 1996;

Floer *et al.*, 2010), we suspected that the DNA binding activity of Rsc3 drove the autoactivation. By increasing the local concentration of p23 through the expression of DBD-p23, AD-Rsc3 is dissociated from the *GAL1* promoter. As Sfh1 and Rsc3 have been directly linked it is possible that AD-Sfh1 joins the endogenous Rsc3 to activate the reporter or Sfh1 itself might bind the promoter DNA (Sengupta *et al.*, 2001; Campsteijn *et al.*, 2007).

We checked our premise that p23 targets Rsc3 to disassemble protein-DNA complexes by EMSA using purified recombinant p23, Rsc3, and Rsc30. While p23 removed Rsc3 from DNA it did not affect Rsc30 (Figure 4B and 4C). By anisotropy it was apparent that p23 accelerated the off-rate of Rsc3 but does not appreciably impact Rsc30 (Figure S4). Significantly, p23 dissociated the Rsc3/Rsc30 heterodimer from DNA indicating that the presence of Rsc3 is sufficient for p23 to act on a DNA-bound protein complex (Figure 4D).

Hsp90 disassembles RSC-nucleosome structures and impedes remodeling

Besides p23, Hsp90 shares a genetic connection with RSC through the *RSC14/LDB7* gene (Zhao *et al.*, 2005; McClellan *et al.*, 2007). Rsc14 works in conjunction with Rsc7 to assimilate the Rsc3/30 heterodimer into the RSC complex (Wilson *et al.*, 2006). As Hsp90 and p23 are also considered partners, we checked whether yeast Hsp90 affects the DNA binding activity of RSC and found that Hsp90 effectively disengaged RSC from naked DNA (Figure 5A). Of note, Hsp90 and p23 were only additive in their abilities to dislodge RSC from DNA and not synergistic (data not shown). In examining RSC nucleosome binding and remodeling activities we discovered a difference in chaperone effects on RSC. While p23 fostered completion of the remodeling reaction (Figure 2B), Hsp90 simply freed RSC from unremodeled nucleosomes even in the presence of ATP (Figure 5B lanes 10–13). By 2-hybrid, Hsp90 relieved the reporter auto-activation by AD-Rsc3 and AD-Sfh1 but Hsp90 did not interact with AD-Rsc3 (Figure S5). Hence, Hsp90 can serve as a general release factor of RSC-DNA/nucleosome complexes but unlike p23 it does not impart a stimulatory influence on RSC remodeling activity.

To investigate the influence of Hsp90 on chromatin architecture *in vivo* we used the DNase-Seq assay to map DNase I Hypersensitive Sites (DHSs) across the genome in yeast expressing either wild type (WT) Hsp90 or the temperature sensitive mutant G170D, which inactivates in ~5 min at temperatures >37°C (Nathan and Lindquist, 1995). DHSs mark areas of open chromatin including active gene promoters, terminators, and enhancers (Bell *et al.*, 2011). Analysis of the DHS patterns following a 15 min incubation at 37°C revealed a dependence on Hsp90 since ~27% of the DHSs in the WT background were lost in G170D and 585 (~10%) novel sites appear (Figure 5C). For comparison, ~50% of the DHSs are reduced in the p23 null cells with a gain of ~25% unique sites (Zelin *et al.*, 2012). Additionally, the average DHS width doubled in *p23* relative to parental cells yet the DHS lengths in G170D and WT were comparable (data not shown). Likely, the differences are attributable to the ability of p23 to modulate both RSC mobility and remodeling whereas Hsp90 only controls the release of RSC from a target.

Besides detecting open chromatin, DNase-Seq can gauge the relative DNA occupancy of select proteins since cognate DNA elements are protected from DNase I cleavage (Hesselberth *et al.*, 2009). In cells expressing WT Hsp82 136 Rsc3 and 137 Rsc30 consensus

sites were bound whereas in G170D the numbers rose to 275 and 272, respectively. We believe that in the absence of Hsp90 RSC associates more stably with DNA. A remarkable aspect of the DNase-Seq data is how quickly the DHS and footprint changes were detected (15 min) indicating that the chromatin landscape is actively maintained by an Hsp90-dependent mechanism.

At DHSs with Rsc3/30 elements occupied in both yeast backgrounds the DNase I cleavage levels were elevated in G170D suggesting RSC continued remodeling at the sites thus lowering nucleosome levels and increasing access to the DNA (Figure 5D). Supporting this contention was the finding that RSC DNA occupancy increased at these DHSs following Hsp90 inactivation (Figure 5E). As p23 is present in both strains, chaperone assistance for the remodeling reaction is available. Reinforcing a continued activity of RSC in the absence of Hsp90 is the finding that few DHSs with Rsc3/30 footprints were lost in the G170D background (3 of the 1876; Figure 5C). Overall, the DNase-Seq data support our contention that Hsp90 dissociates RSC from DNA and that RSC nucleosome remodeling activity is not Hsp90-dependent. Perhaps the ability of Hsp90 to mobilize remodelers accounts for the previous observation that Hsp90 generally localizes to NDRs (Sawarkar *et al.*, 2012).

RSC and p23 colocalize along the genome

Given the dual impact of p23 on RSC in vitro, we focused on p23 to better understand how a chaperone might influence a remodeler in vivo. To start, we checked whether p23 is in position to regulate RSC across the yeast genome. We mapped the genomic sites associated with p23 by ChIP-Seq and found 572 high quality peaks (P values <0.001). We compared the p23 pattern to a recent Sth1 map (Parnell *et al.*, 2015) and found 149 sites in common or ~40% of the Sth1 peaks (Figure 6A). Comparison to another RSC map, which was made by targeting 5 different RSC subunits along with microarrays comprised of intergenic DNA (Ng *et al.*, 2002), showed that p23 intersected with ~36% of the RSC sites (Figure S6A). Although Fisher exact tests using 2×2 contingency tables indicated high correlations between the p23 and RSC peaks (P values $<1e^{-104}$ or $<1e^{-71}$, respectively), the overlap between the RSC sites was limited to 80 (Figure S6B). While the larger overlap between both RSC maps and p23 relative to the two RSC patterns was unexpected, we suspect it reflects the dynamic nature of the RSC complex. Likely, RSC repositions in response to the growth conditions of a given experiment. As we believe p23 transiently but repeatedly interacts with RSC to promote a rapid exchange with chromatin, the p23-RSC association would be independent of the experimental status.

Regardless of the analyzed RSC dataset, commonalities in the shared p23-RSC peaks were apparent. For example, 46 of the 149 shared sites between p23 and Sth1 were at tRNA genes where RSC continuously depletes nucleosomes to maintain some of the lowest histone occupancies along the genome (Rao *et al.*, 2005; Parnell *et al.*, 2008). Recently it was shown that RSC also is associated with fragile nucleosomes, which are found at highly expressed genes (Kubik *et al.*, 2015). Of the Sth1 peaks 284 overlapped with a fragile nucleosome and p23 was at 307 in total with 147 being co-associated with Sth1 (Figure 6B); comparable pattern was found with the other RSC genomic dataset (Figure S5C). Relative to other DNA binding protein or DNA elements known to be associated with p23 (Figure S5D), RSC-areas

displayed a greater overlap with p23 sites. Taken together, our genomic data demonstrate that p23 localizes across the genome at sites of RSC action.

RSC activities are p23-dependent in vivo

To address whether p23 impacts the cellular function of RSC we exploited the common use of RSC within signal transduction pathways (Clapier and Cairns, 2009). Prompt reactions to diverse physiological cues necessitate the rapid mobilization of RSC to chromatin areas in need of remodeling in order to support growth (Damelin *et al.*, 2002; Ng *et al.*, 2002). For example, mutations in the RSC motor Sth1 sensitize cells to biological stressors including ethanol contact (Du *et al.*, 1998). Significantly, loss of *p23* in the *sth1-2* allelic background resulted in yeast hyper-sensitized to ethanol exposure (Figure 7A). Given the capacity of p23 to dissociate RSC from DNA/nucleosomes in vitro, we checked whether p23 altered RSC mobility in vivo. As RSC does not appear to have a set chromatin localization pattern under normal conditions (Ng *et al.*, 2002; Parnell *et al.*, 2015), we focused on monitoring the arrival kinetics of RSC at ethanol-regulated genes.

RSC nucleation at ethanol-responsive loci occurred in ~1 minute in parental cells but slowed to ~15 minutes in *p23* (Figure 7B; Figure S7A) and gene activation was delayed even further in the absence of p23 (Figure 7C; Figure S7B). We assessed the local chromatin structure using a Micrococcal nuclease mapping assay (Bai *et al.*, 2011). In parental yeast, NDRs near the transcription start sites were apparent along with well-positioned -1/+1 nucleosomes whose occupancies declined after ethanol addition (Figure 7D; Figure S7C). In contrast to parental cells, nucleosome levels at the promoters reduced slowly correlating with the kinetics of gene activation (Figure 7D; Figure S7C). Hence, in the absence of p23 there is a sequential lag in the nucleation of RSC at target promoters followed by a belated remodeling of nearby nucleosomes. Thus, the in vivo data correlate well with our biochemical studies showing a dual role of p23 in promoting the transition of RSC between target sites and in fostering nucleosome remodeling. Overall, p23 associates with RSC across the genome where it fosters a dynamic action critical for manipulating RSC conformations during remodeling and for rapidly relocating RSC in response to fluctuating physiological conditions.

DISCUSSION

Proteins are core units for generating and supporting life. As such, significant resources are directed towards supporting protein health including an integrated network of pathways mediating polypeptide synthesis, folding, assembly, and degradation that collectively form the proteostasis process (Hartl *et al.*, 2011). Proficiency of the system is reliant on a continuous sampling of each factor with variances in conformational states dictating protein fates. Here, we suggest that the persistent activities of p23 and Hsp90 elevate the functionality of assembled protein complexes by exploiting the RSC chromatin remodeler as a molecular model. In addition to the proteostatic contributions, our studies reveal a significant dependency of chromatin architecture on Hsp90.

While the use of molecular chaperones in building protein structures has long been appreciated (Ellis, 1987), a post-assembly role typically is not considered. Yet, the overall

goal of proteostasis is the upkeep of a healthy proteome (Balch *et al.*, 2008). As the maintenance of any system is more efficient than de novo synthesis and destruction, a continuous chaperone action with mature factors would be beneficial for perpetuating an active native protein landscape. In this scenario, the binding energy of a chaperone-client interaction dissuades non-native forms thereby retaining the target within its ensemble of native configurations (Bartlett and Radford, 2009). Hence, the sizable Hsp90 chaperone system provides a low energy mechanism to habilitate a healthy proteome.

Besides proteostatic contributions, we believe Hsp90 and p23 are further used to meet a specialized need of the nucleus (Sawarkar and Paro, 2013; Echtenkamp and Freeman, 2014). The nuclear compartment usually is not associated with proteostasis since conventional protein life cycle events happen in the cytosol (*e.g.*, translation and degradation). Yet, the upkeep of polypeptides in the nucleus is essential for vitality. Highlighting this point is the occurrence of nuclear protein inclusions in about a third of the known aggregation diseases (Woulfe, 2007). Hence, maintaining properly functioning nuclear factors has significant long-term implications in addition to the immediate actions of these proteins (*e.g.*, remodel chromatin).

In housing a genome, the nucleus encounters unique challenges centering on the need to package chromosomes into a confined space while allowing ‘on call’ access to the entire length of the DNA. The basic process of condensing chromosomes with histones while controlling access to the underlying DNA by altering nucleosome spacing using remodelers is straightforward (Clapier and Cairns, 2009). Yet, the typical chromatin-model does not take into account the disparity in the copy numbers of nucleosomes and remodelers. For example, yeast use ~70,000 nucleosomes to condense their genomes yet there are only ~200–2000 RSC particles (Ghaemmaghami *et al.*, 2003; Brogaard *et al.*, 2012; Chong *et al.*, 2015). Even though 7 additional remodelers exist, the sum of the assembled particles is still substoichiometric totaling ~1,300–14,000. Hence, remodelers must be able to rapidly transition between targets in order to effectively manage the overall chromatin architecture in response to fluctuating physiological conditions.

RSC mobility is vital since RSC fosters many genomic pathways requiring quick access to the underlying DNA including DNA repair, stress and cell cycle transcription programs (Clapier and Cairns, 2009; Sinha and Peterson, 2009). For example, exposure to ethanol, rapamycin, or hydrogen peroxide triggers the redistribution of RSC to hundreds of new sites including promoter nucleosomes that are remodeled to allow gene expression within minutes (Damelin *et al.*, 2002; Ng *et al.*, 2002). In the absence of p23 the ethanol-induced translocation timing of RSC was impaired (Figure 7A). Even after RSC associated with a promoter the kinetics of nucleosome remodeling were further delayed in *p23* yeast (Figure 7D; Figure S5C). Hence, by modulating multiple steps in the remodeling pathway the contribution of p23 is considerable.

Why might a chromatin remodeler depend on molecular chaperones? Standard remodeling models advocate that the energy of ATP-hydrolysis drives the various structural changes occurring in either the remodeler complex or nucleosome during the course of the reaction (Clapier and Cairns, 2009). Yet, sizeable conformational shifts follow the initial binding

event between RSC and a nucleosome. Electron micrograph studies revealed a bowl-like shape of RSC with a central depression holding the nucleosome (Asturias *et al.*, 2002; Leschziner *et al.*, 2007). In the EM structures the nucleosomal DNA is disordered since RSC binding energy, independent of ATP, separates a large section of the DNA from the histones (Asturias *et al.*, 2002; Lorch *et al.*, 2010). Crosslinking studies identified 4 RSC subunits in close proximity to the DNA including Sth1, Rsc2, Rsc3, and Rsc4 (Sengupta *et al.*, 2001). We suggest that the binding energy of chaperone interactions shifts the conformational status of the RSC proteins bound to DNA, which triggers the release of the complex. In a manner comparable to the dissociation of the DNA-Rsc3/Rsc30 heterodimer by p23, we favor a model in which select RSC subunits are targeted by the chaperones to trigger release of the entire complex.

The capacity of p23 and Hsp90 to dissociate RSC from either free or nucleosomal DNA might rationalize why the timing and efficiency of remodeling is significantly different *in vitro* vs. *in vivo*. For example, RSC redistributes across a genome within minutes and remodels numerous nucleosomes (Damelin *et al.*, 2002; Ng *et al.*, 2002), but under conditions mimicking the cellular state where nucleosomal arrays are in excess of RSC only ~2 arrays are remodeled every 120 min *in vitro* (Logie *et al.*, 1999). Attempts to enhance the efficiency with hyperacetylated histones resulted in a declined rate of ~0.6 array/120 min. Likely, the acetylation stabilizes the RSC-nucleosome interaction thereby further slowing the reaction. The longevity of RSC binding is even sufficient to block ligation of nucleosomal DNA *in vitro* (Lorch *et al.*, 2001). *In vivo*, however, chromatin remodelers dynamically associate with nucleosomes having half-lives in the range of seconds even at gene promoters being actively remodeled (Johnson *et al.*, 2008). We suspect the presence of p23 and Hsp90 account for the kinetic differences.

In general, genomic processes function through cooperative mechanisms to rapidly nucleate multi-component structures at precise locations. While high-affinity interactions produce robust and select actions, the inherent stability of such assembled complexes would impede the timing of biological events. We believe Hsp90 and p23 limit the timing of protein-DNA interactions thereby creating a highly dynamic nuclear environment to accelerate DNA-associated pathways (Echtenkamp and Freeman, 2014). Of note, an early discovery of yeast p23 showed that loss or overexpression of this chaperone results in chromosome instability, which led to its naming as Chromosome Stability 18 (*CST18*) (Ouspenski *et al.*, 1999). Significantly, the combination of Hsp90 and histone deacetylase inhibitors has proven to be synergistic in clinical trials (Yang *et al.*, 2008). While a mechanism involving the acetylation status of Hsp90 had been suggested, our studies open an unexplored therapeutic possibility by showing that proper chromatin architecture relies on Hsp90. Perhaps in a manner analogous to the more common role of molecular chaperones in protein folding, Hsp90 and p23 contribute to nuclear function by circumventing otherwise long-lived, unproductive structures (*i.e.*, pathway intermediates) thereby ensuring an effective forward flow of biological events including pathways organizing chromatin.

EXPERIMENTAL PROCEDURES

Yeast

The utilized *Saccharomyces cerevisiae* strains are described in Table S1. BY4741 *rsc7::KanR p23::His3 pRS316 p23*-locus was produced through tetrad dissection. All p23 strains were created using a PCR based method (Longtine *et al.*, 1998).

Genetic interaction assays

A 5-fold serial dilution spot test assay determined the viability of single (*p23*, *rsc7*, *sth1-F793S*) and double (*p23 rsc7*, *p23 sth1-F793S*) mutants. The Rsc7-associated experiments used plates lacking uracil or containing 5-FOA and incubations at 30°C. The Sth1-F793S related tests incubated cells at the non-permissive temperature (37°C) for 24 h prior to spotting and incubation at 22°C. A 3-fold serial dilution spot test checked the viability of single (*p23* and *sth1-2*) and double (*p23 sth1-2*) mutants along with growth on ethanol/glycerol or glucose containing media.

Protein purification

TAP-tagged RSC complex was isolated as described (Sinha *et al.*, 2009). Recombinant *Xenopus laevis* histones, p23, p23⁵⁰, and p23⁸⁴ were purified as described (Luger *et al.*, 1999; Toogun *et al.*, 2007). Recombinant Rsc3 and Rsc30 were purified as His₆-SUMO fusion proteins (Butt *et al.*, 2005).

Nucleosome reconstitution

The 200 bp DNA fragment of the 601 positioning sequence was generated by PCR from pGem-3Z 601. Nucleosomes were assembled using purified recombinant histone octamers by salt dialysis (Luger *et al.*, 1999).

EMSAs

RSC DNA binding assays used a radiolabeled *PHO8* probe (7 nM). Reactions were resolved on pre-chilled 3.2 % native polyacrylamide GTG. For competition assays, the reactions were incubated with 2, 4, 8, or 16-fold unlabeled *PHO8* promoter DNA for 5 min prior to resolution. Nucleosomes, RSC and p23 were incubated for 10 min at r.t. in binding buffer (10 mM TRIS, 40 mM NaCl, 0.1 mM EDTA, 1 mM DTT, 5% glycerol, 50 µg/ml BSA) along with 1 mM ATP, as marked.

Nucleosome sliding assay

Nucleosomes (5 nM) were incubated with RSC (1 nM), p23 (as marked) in sliding buffer (7 mM HEPES, 0.1 mM EDTA, 0.06% Tween, 55 mM NaCl, 1 mM DTT, 3 mM MgCl₂, 1% glycerol, 100 µg/ml BSA) at 30°C for 20 min.

Yeast 2-hybrid

Screen was completed using the Matchmaker GAL4 2-hybrid system (Clontech). RSC subunits and p23 were cloned into pGAD-T7 and pGBK-T7 and expressed in AH109 as

indicated. A 10-fold serial dilution spot test assay was used. Cells were plated on SD +His or SD-His plates and incubated at 30°C.

Chromatin immunoprecipitation

ChIP assay was performed as previously described (Kuras and Struhl, 1999). Briefly, Rsc3-TAP and *p23* Rsc3-TAP strains were growing logarithmically in YPD before ethanol (7.5%) treatment. Isolated DNA was analyzed by qPCR using primers select to the ethanol responsive loci and the control locus *POL1* (Ng *et al.*, 2002). RSC localization was measured by comparing the ratio of each locus over that of *POL1* for each time point, then comparing the fold increase of each time point to the 0 minute time point.

Quantitative PCR

Rsc3-TAP and *p23*Rsc3-TAP strains were grown as in the ChIP experiment. Cells were collected at the indicated time points and flash frozen. RNA was isolated using hot phenol extraction and treated with Turbo DNase (Life Technologies). cDNA was made using the SuperScript III First-Strand Synthesis Supermix for qRT-PCR kit. All time points were normalized to ACT1 RNA levels.

Nucleosome mapping

MNase digestion of chromatin was performed on spheroplasts as described (Bai *et al.*, 2011). Digested DNA and genomic DNA were analyzed with overlapping primer pairs for each gene promoter generating ~100 bp PCR products. Nucleosomal occupancy levels were calculated by the fold change between genomic DNA and MNase digested DNA normalized to the *PHO5*TATA (-1) nucleosome.

DNase-Seq analysis

The DNase-Seq protocol was adapted from an established method (Hesselberth *et al.*, 2009) as described (Zelin *et al.*, 2012).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGEMENTS

Support by the Public Service grants CA155333 and GM118306. The authors have no conflicts of interest. We'd like to thank Brian Nguyen, Janhavi Kolhe, and Anqi Wang for their helpful comments.

REFERENCES

- Angus-Hill ML, Schlichter A, Roberts D, Erdjument-Bromage H, Tempst P, and Cairns BR (2001) The Rsc3/Rsc30 zinc cluster dimer reveals novel roles for the chromatin remodeler RSC in gene expression and cell cycle control. *Mol. Cell* 7, 741–751. [PubMed: 11336698]
- Asturias FJ, Chung WH, Kornberg RD, and Lorch Y (2002). Structural analysis of the RSC chromatin-remodeling complex. *Proc. Natl. Acad. Sci. USA* 99, 13477–13480. [PubMed: 12368485]
- Badis G, Chan ET, van Bake H, Pena-Castillo L, Tillo D, Tsui K, Carlson CD, Gossett AJ, Hasinoff MJ, Warren CL, et al. (2008). A library of yeast transcription factor motifs reveals a widespread

function for Rsc3 in targeting nucleosome exclusion at promoters. *Mol. Cell.* 32, 878–887. [PubMed: 19111667]

Bai L, Ondracka A, and Cross FR (2011). Multiple sequence-specific factors generate the nucleosome-depleted region on CLN2 promoter. *Mol. Cell* 42, 465–476. [PubMed: 21596311]

Balch WE, Morimoto RI, Dillin, and A., Kelly JW (2008). Adapting proteostasis for disease intervention. *Science* 319, 916–919. [PubMed: 18276881]

Bartlett AI, and Radford SE (2009). An expanding arsenal of experimental methods yields an explosion of insights into protein folding mechanisms. *Nat. Struct. Mol. Biol.* 16, 582–588. [PubMed: 19491935]

Bell O, Tiwari VK, Thomä NH, and Schübeler D (2011). Determinants and dynamics of genome accessibility. *Nat. Rev. Genet.* 12, 554–564. [PubMed: 21747402]

Brogaard K, Xi L, Wang J-P, and Widom J (2012). A map of nucleosome positions in yeast at base-pair resolution. *Nature* 486, 496–501. [PubMed: 22722846]

Butt TR, Edavettal SC, Hall JP, and Mattern MR (2005) SUMO fusion technology for difficult-to-express proteins. *Protein Expr. Purif.* 43, 1–9. [PubMed: 16084395]

Cairns BR, Lorch Y, Li Y, Zhang M, Lacomis L, Erdjument-Bromage H, Tempst P, Du J, Laurent B, and Kornberg RD (1996). RSC, an essential abundant chromatin-remodeling complex. *Cell* 87, 1249–1260. [PubMed: 8980231]

Campsteijn C, Wijnands-Collin A-MJ, and Logie C (2007). Reverse genetic analysis of the yeast chromatin remodeler reveals a role for RSC3 and SNF5 homolog 1 in ploidy maintenance. *PLoS Genet.* 3, e92. [PubMed: 17542652]

Chong YT, Koh JLY, Friesen H, Duffy SK, Cox MJ, Moses A, Moffat J, Boone C, and Andrews BJ (2015). Yeast proteome dynamics from single cell imaging and automated analysis. *Cell* 161, 1413–1424. [PubMed: 26046442]

Clapier CR, and Cairns BR (2009). The biology of chromatin remodeling complexes. *Annu. Rev. Biochem.* 78, 273–304. [PubMed: 19355820]

Damelin M, Simon I, Moy TI, Wilson B, Komili S, Tempst P, Roth FP, Young RA, Cairns BR, and Silver PA (2002). The genome-wide localization of Rsc9, a component of the RSC chromatin-remodeling complex, changes in response to stress. *Mol. Cell* 9, 563–573. [PubMed: 11931764]

DeZwaan DC, and Freeman BC (2008). HSP90: The Rosetta stone for cellular protein dynamics? *Cell Cycle* 7, 1006–1012. [PubMed: 18414022]

Du J, Nasir I, Benton BK, Kladde MP, and Laurent BC (1998). Sth1p, a *Saccharomyces cerevisiae* Snf2p/Swi2p homolog, is an essential ATPase in RSC and differs from Snf/Swi in its interactions with histones and chromatin-associated proteins. *Genetics* 150, 987–1005. [PubMed: 9799253]

Echtenkamp FJ, Zelin E, Oxelmark E, Woo JI, Andrews BJ, Garabedian M, and Freeman BC (2011) Global functional map of the p23 molecular chaperone reveals an extensive cellular network. *Mol. Cell* 43, 229–241. [PubMed: 21777812]

Echtenkamp FJ, and Freeman BC (2012). Expanding the cellular molecular chaperone network through the ubiquitous cochaperones. *Biochim. Biophys. Acta* 1823, 668–673. [PubMed: 21889547]

Echtenkamp FJ, and Freeman BC (2014). Molecular chaperone-mediated nuclear protein dynamics. *Curr. Prot. Pept. Sci.* 15, 216–224.

Ellis J, (1987). Proteins as molecular chaperones. *Nature* 328, 378–379. [PubMed: 3112578]

Floer M, Wang X, Prabhu V, Berrozpe G, Narayan S, Spagna D, Alvarez D, Kendall J, Krasnitz A, Stepansky A, et al. (2010). A RSC/nucleosome complex determines chromatin architecture and facilitates activator binding. *Mol. Cell* 141, 407–418.

Ghaemmaghami S, Huh W-K, Bower K, Howson RW, Belle A, Dephoure N, O’Shea EK, and Weissman JS (2003). Global analysis of protein expression in yeast. *Nature* 425, 737–741. [PubMed: 14562106]

Hartl FU, Bracher A, and Hayer-Hartl M (2011). Molecular chaperones in protein folding and proteostasis. *Nature* 475, 324–332. [PubMed: 21776078]

Hesselberth JR, Chen X, Zhang Z, Sabo PJ, Sandstrom R, Reynolds AP, Thurman RE, Neph S, Kuehn MS, Noble WS, et al. (2009). Global mapping of protein-DNA interactions in vivo by digital genomic footprinting. *Nat. Methods* 6, 283–289. [PubMed: 19305407]

- James P, Halladay J, and Craig EA (1996). Genomic libraries and a host strain designed for highly efficient two-hybrid selection in yeast. *Genetics* 144, 1425–1436. [PubMed: 8978031]
- Johnson TA, Elbi C, Parekh BS, Hager GL and John S (2008). Chromatin remodeling complexes interact dynamically with a glucocorticoid receptor–regulated promoter. *Mol. Biol. Cell* 19, 3308–3322. [PubMed: 18508913]
- Kubik S, Bruzzone MJ, Jacquet P, Falcone J-L, Rougemont J, and Shore D (2015). Nucleosome stability distinguishes two different promoter types at all protein-coding genes in yeast. *Mol. Cell* 60, 422–434. [PubMed: 26545077]
- Kuras L, and Struhl K (1999). Binding of TBP to promoters in vivo is stimulated by activators and requires Pol II holoenzyme. *Nature* 399, 609–13. [PubMed: 10376605]
- Leschziner AE, Saha A, Wittmeyer J, Zhang Y, Bustamante C, Cairns BR, and Nogales E (2007). Conformational flexibility in the chromatin remodeler RSC observed by electron microscopy and the orthogonal tilt reconstruction method. *PNAS* 104, 4913–4918. [PubMed: 17360331]
- Logie C, Tse C, Hansen JC, and Peterson CL (1999). The core histone N-terminal domains are required for multiple rounds of catalytic chromatin remodeling by the SWI/SNF and RSC complexes. *Biochemistry* 38, 2514–22. [PubMed: 10029546]
- Longtine MS, McKenzie A III, Demarini DJ, Shah NG, Wach A, Brachet A, Philippsen P, and Pringle JR (1998). Additional modules for versatile and economical PCR-based gene deletion and modification in *Saccharomyces cerevisiae*. *Yeast* 14, 953–961. [PubMed: 9717241]
- Lorch Y, Cairns BR, Zhang M, and Kornberg RD (1998). Activated RSC-nucleosome complex and persistently altered form of the nucleosome. *Cell* 94, 29–34. [PubMed: 9674424]
- Lorch Y, Zhang M, Kornberg RD (2001). RSC unravels the nucleosome. *Mol. Cell.* 7, 89–95. [PubMed: 11172714]
- Lorch Y, Maier-Davis B, and Kornberg RD (2010). Mechanism of chromatin remodeling. *Proc. Natl. Acad. Sci. U S A* 107, 3458–3462. [PubMed: 20142505]
- Luger K, Rechsteiner TJ, and Richmond TJ (1999). Preparation of nucleosome core particle from recombinant histones. *Methods Enzymol.* 304, 3–19. [PubMed: 10372352]
- Makhnevych T, and Houry WA (2012). The role of Hsp90 in protein complex assembly. *Biochim. Biophys. Acta* 1823, 674–682. [PubMed: 21945180]
- McClellan AJ, Xia Y, Deutschbauer AM, Davis RW, Gerstein M, and Frydman J (2007). Diverse cellular functions of the Hsp90 molecular chaperone uncovered using systems approaches. *Cell* 131, 121–135. [PubMed: 17923092]
- Millson SH, Truman AW, King V, Prodromou C, Pearl LH, and Piper PW (2005). A two-hybrid screen of the yeast proteome for Hsp90 interactors uncovers a novel Hsp90 chaperone requirement in the activity of a stress-activated mitogen-activated protein kinase, Slf2p (Mpk1p). *Eukaryot. Cell* 4, 849–860. [PubMed: 15879519]
- Nathan DF, and Lindquist S (1995). Mutational analysis of Hsp90 function: interactions with a steroid receptor and a protein kinase. *Mol. Cell. Biol.* 15, 3917–3925. [PubMed: 7791797]
- Ng HH, Robert F, Young RA, and Struhl K (2002). Genome-wide location and regulated recruitment of the RSC nucleosome-remodeling complex. *Genes Dev.* 16, 806–819. [PubMed: 11937489]
- Ouspenski II, Elledge SJ, and Brinkley BR (1999). New yeast genes important for chromosome integrity and segregation identified by dosage effects on genome stability. *Nucleic Acids Res.* 27, 3001–3008. [PubMed: 10454593]
- Parnell TJ, Huff JT, and Cairns BR (2008). RSC regulates nucleosome positioning at Pol II genes and density at Pol III genes. *EMBO J.* 27, 100–110. [PubMed: 18059476]
- Parnell TJ, Schlichter A, Wilson BG, and Cairns BR (2015). The chromatin remodelers RSC and ISW1 display functional and chromatin-based promoter antagonism. *eLife* 4, e06073. [PubMed: 25821983]
- Powers ET, Morimoto RI, Dillin A, Kelly JW, and Balch WE (2009). Biological and chemical approaches to diseases of proteostasis deficiency. *Annu. Rev. Biochem.* 78, 959–991. [PubMed: 19298183]
- Rao B, Shibata Y, Strahl BD, and Lieb JD (2005). Dimethylation of histone H3 at lysine 36 demarcates regulatory and nonregulatory chromatin genome-wide. *Mol. Cell. Biol.* 25, 9447–9459. [PubMed: 16227595]

- Röhl A, Rohrberg, and J., Buchner J (2013). The chaperone Hsp90: changing partners for demanding clients. *Trends Biochem. Sci.* 38, 253–262. [PubMed: 23507089]
- Sawarkar R, Sievers C, and Paro R (2012). Hsp90 globally targets paused RNA polymerase to regulate gene expression in response to environmental stimuli. *Cell* 149, 807–818. [PubMed: 22579285]
- Sawarkar R, and Paro R (2013) Hsp90@chromatin.nucleus: an emerging hub of a networker. *Trends Cell Biol.* 23, 193–201. [PubMed: 23286900]
- Sengupta SM, VanKanegan M, Persinger J, Logie C, Cairns BR, Peterson CL, and Bartholomew B (2001). The interactions of yeast SWI/SNF and RSC with the nucleosome before and after chromatin remodeling. *J. Biol. Chem.* 276, 12636–12644. [PubMed: 11304548]
- Sharma K, Vabulas RM, Macek B, Pinkert S, Cox J, Mann M, and Hartl FU (2012). Quantitative proteomics reveals that Hsp90 inhibition preferentially targets kinases and the DNA damage response. *Mol. Cell Proteomics* 11, M111.014654.
- Sinha M, and Peterson CL (2009). Chromatin dynamics during repair of chromosomal DNA double-strand breaks. *Epigenomics* 1, 371–385. [PubMed: 20495614]
- Taipale M, Krykbaeva I, Koeva M, Kayatekin C, Westover KD, Karras GI, and Lindquist S (2012). Quantitative analysis of Hsp90-client interactions reveals principles of substrate recognition. *Cell* 150, 987–1001. [PubMed: 22939624]
- Taipale M, Tucker G, Peng J, Krykbaeva I, Lin Z-Y, Larsen B, Choi H, Berger B, Gingras A-C, Lindquist S (2014). A quantitative chaperone interaction network reveals the architecture of cellular protein homeostasis pathways. *Cell* 158, 434–448. [PubMed: 25036637]
- Titus LC, Dawson TR, Rexer DJ, Ryan KJ, and Wentz SR (2010). Members of the RSC chromatin-remodeling complex are required for maintaining proper nuclear envelope structure and pore complex localization. *Mol. Biol. Cell* 21, 1072–1087. [PubMed: 20110349]
- Toogun OA, Zeiger W, and Freeman BC (2007). The p23 molecular chaperone promotes functional telomerase complexes through DNA dissociation. *Proc. Natl. Acad. Sci. USA* 104, 5765–5770. [PubMed: 17389357]
- Venkatesh S, and Workman JL (2015). Histone exchange, chromatin structure and the regulation of transcription. *Nat. Rev.* 16, 178–189.
- Wilson B, Erdjument-Bromage H, Tempst P, and Cairns BR (2006). The RSC chromatin remodeling complex bears an essential fungal-specific protein module with broad functional roles. *Genetics* 172, 795–809. [PubMed: 16204215]
- Wippo CJ, Israel L, Watanabe S, Hochheimer A, Peterson CL and Korber P (2011). The RSC chromatin remodeling enzyme has a unique role in directing the accurate positioning of nucleosomes. *EMBO J.* 30, 1277–1288. [PubMed: 21343911]
- Woulfe JM (2007). Abnormalities of the nucleus and nuclear inclusions in neurodegenerative disease: a work in progress. *Neuropathol. Appl. Neurobiol.* 33, 2–42. [PubMed: 17239006]
- Yen K, Vinayachandran V, Batta K, Koerber RT, and Pugh BF (2012). Genome-wide nucleosome specificity and directionality of chromatin remodelers. *Cell* 149, 1461–1473. [PubMed: 22726434]
- Yang Y, Rao R, Shen J, Tang Y, Fiskus W, Nechtman J, Atadja P, and Bhalla K (2008). Role of Acetylation and Extracellular Location of Heat Shock Protein 90 α in Tumor Cell Invasion. *Cancer Res.* 68, 4833–4842. [PubMed: 18559531]
- Zelin E, Zhang Y, Toogun OA, Zhong S, and Freeman BC (2012). The p23 molecular chaperone and GCN5 acetylase jointly modulate protein-DNA dynamics and open chromatin status. *Mol. Cell* 48, 459–470. [PubMed: 23022381]
- Zhao R, Davey M, Hsu Y-C, Kaplanek P, Tong A, Parsons AB, Krogan N, Cagney G, Mai D, Greenblatt J, et al. (2005). Navigating the chaperone network: an integrative map of physical and genetic interactions mediated by the Hsp90 chaperone. *Cell* 120, 715–727. [PubMed: 15766533]
- Zhu C, Byers KJRP, McCord RP, Shi Z, Berger MF, Newburger DE, Saulrieta K, Smith Z, Shah MV, and Radhakrishnan M (2009). High-resolution DNA-binding specificity analysis of yeast transcription factors. *Genome Res.* 19, 556–566. [PubMed: 19158363]




Figure 1. The p23 chaperone is physiologically linked to RSC but not as a quality control factor. (A) The viability of single (*p23* and *rsc7*) or double (*p23 rsc7*) gene knockout yeast were evaluated with a plasmid dropout test. The yeast carry an *URA3*-marked covering plasmid with the endogenous *P23* locus and the vector was either positively (SD-Ura) or negatively (SD+5FOA) selected. (B) Steady-state protein levels of the indicated RSC subunits were assayed by immunoblot analysis using whole cell extracts prepared from logarithmically growing parental or *p23* cells. (C) Assembly of the RSC protein complex is p23-independent. Using the standard tandem affinity pull down assay RSC complexes were isolated from extracts prepared from *TAP-RSC2* parental or *p23 TAP-RSC2* yeast and the relative levels of the marked RSC subunits was checked. (D) Cell viability of a temperature sensitive *sth1* allele is p23-dependent. Parental, *p23*, *sth1-F793S*, and *p23 sth1-F793S* yeast were grown at 30°C (permissive) or exposed to 37°C (non-permissive) for 24 h and viability was then checked using a spot test assay and incubation at 30°C.




Figure 3. P23 promotes RSC-mediated nucleosome remodeling in vitro.

(A) RSC nucleosome binding and remodeling activity was checked by EMSA using nucleosomes prepared with radiolabeled 601 positioning DNA and titrations of purified RSC (1, 5, 10 nM). The influence of ATP (1 mM) and p23 (1, 3, 5, 15 μ M) were assessed as indicated. The migration positions of free DNA, nucleosomes with a centrally or terminally positioned histone, and RSC-nucleosome structures (Complex 1 and 2) are marked. (B) A nucleosome sliding assay was used to confirm the influence of p23 on RSC remodeling activity and to check if p23 chaperone function was required using the chaperone-deficient mutant p23⁸⁴.




Figure 4. Rsc3 is directly targeted for dissociation by p23.

(A) Interactions between p23 and each RSC subunit were monitored by yeast 2-hybrid. The influence of a p23 titration (0, 4, 12, 32 μ M) on the DNA binding activities of purified Rsc3 (90 nM) (B) or Rsc30 (90 nM) (C) was checked by EMSA using a radiolabeled Rsc3/30 element from the *ACCI/TIM23* loci as a probe (Zhu *et al.*, 2009). (D) The DNA binding capacity of the Rsc3/30 heterodimer (50 nM) in the presence or absence of p23 (6 μ M) was checked by EMSA using radiolabeled *PHO8* promoter DNA as a probe.




Figure 5. Hsp90 dissociates RSC from DNA and/or nucleosomes but does not enhance remodeling activity.

(A) RSC DNA binding activity was monitored by EMSA using purified RSC and radiolabeled *PHO8* promoter DNA. The influence of Hsp90 on RSC-DNA structures was determined by titrating in Hsp90 (0, 2, or 20 μ M). (B) RSC nucleosome binding and remodeling activity was checked by EMSA using nucleosomes prepared with radiolabeled 601 positioning DNA and titrations of purified RSC (1, 5, 10 nM). The influence of ATP (1 mM) and Hsp90 (4 μ M) were assessed as indicated. The migration positions of free DNA, nucleosomes with a centrally or terminally positioned histone, and RSC-nucleosome structures (Complex 1 and 2) are marked. (C) DHSs were identified by deep-sequencing samples following limited DNase I digestion of chromatin within nuclei prepared from yeast expressing either wild type Hsp82 or the G170D *ts* mutant as the sole source of Hsp90. Prior to collecting the yeast the cells were incubated at 37°C for 15 min, which inactivate s G170D (Nathan and Lindquist, 1995). The green bar represents DHSs unique to wild type, red is DHSs specific to G170D, and grey bars are the remaining sites with the number of overlaps indicated in parentheses. (D) At shared DHSs containing either an occupied Rsc30 (DHS 1) or Rsc3 (DHS 2) consensus element an increase in the levels of DNase I cleavage events was observed. A representative example of the DNA cleavage density taken from chromosome XI (60,938–109,058) is shown with the positions of the occupied Rsc30 or Rsc3 sites marked. (E) RSC DNA occupancy increased at the DHSs with enhanced nuclease sensitivity. ChIP was used to measure the relative association of RSC at DHS1 and DHS2 in

the parental and G170D yeast incubated at 37°C for 15 min. The ChIP data represent the averages of 3 independent replicas and error bars are SEM.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 6. The genomic localization profiles of RSC and p23 overlap.

The genomic association pattern of yeast p23 was determined using α -p23 antibody, chromatin immunoprecipitation, high-throughput sequencing, and bioinformatics. **(A)** The identified genomic sites bound by p23 were compared to an established map of the RSC motor subunit Sth1 (Parnell *et al.*, 2015). Of the 149 overlapping sites, 60 contain Rsc3/30 consensus elements. **(B)** The overlap between p23 and the known patterns of Sth1 and fragile nucleosomes across the yeast genome was determined (Kubik *et al.*, 2015).




Figure 7. RSC promoter recruitment and remodeling activity is compromised in *p23* cells. (A) Loss of *p23* increases the temperature sensitivity of *sth1-2* yeast grown on ethanol/glycerol (EtOH/Gly) but not on glucose. Logarithmic phase yeast were serially diluted, spotted onto the indicated media, and grown at 30°C or 37°C. (B) The temporal occupancy of RSC at the *HXK1* ethanol-responsive promoter was determined by ChIP using parental *TAP-RSC3* (solid gray) or *p23 TAP-RSC3* (dashed black) yeast. The time course was initiated by the addition of ethanol (7.5% final) to logarithmically growing cells. (C) The ethanol-induced RNA levels of *HXK1* was monitored by RT-qPCR in parental *TAP-RSC3* (solid gray) and *p23 TAP-RSC3* (dashed black) at the noted times following addition of ethanol (7.5%). *ACT1* RNA levels were used for normalization. (D) The relative nucleosome positions and densities at the *HXK1* gene were determined in parental (gray) and *p23* (black) yeast without (open) or with (closed) exposure to ethanol (7.5%) for 15 or 120 min. The ChIP and RT-qPCR data represent the averages of 7 independent replicas, the MNase-mapping is the average of 3 replicas, and the error bars are SEM.