

Review Article

Non-coding RNAs: emerging regulators of glucose metabolism in hepatocellular carcinoma

Yongting Lai^{1*}, Hairong Huang^{2*}, Mubalake Abudoureyimu³, Xinrong Lin³, Chuan Tian³, Ting Wang³, Xiaoyuan Chu^{1,3}, Rui Wang^{1,3}

¹Department of Medical Oncology, Jinling Hospital, The First School of Clinical Medicine, Southern Medical University, Nanjing, China; ²Department of Cardiothoracic Surgery, Jinling Hospital, Nanjing, China; ³Department of Medical Oncology, School of Medicine, Jinling Hospital, Nanjing University, Nanjing, China. *Equal contributors.

Received August 20, 2020; Accepted November 21, 2020; Epub December 1, 2020; Published December 15, 2020

Abstract: Reprogramming of metabolism is one of the hallmarks of cancer, among which glucose metabolism dysfunction is the most prominent feature. The glucose metabolism of tumor cells is significantly different from that of normal cells. Glucose metabolism reprogramming of hepatocellular carcinoma (HCC) has become an important research hotspot in the field of HCC, a variety of tumor metabolic interventions have been applied clinically. Moreover, various Non-coding RNAs (ncRNAs) including microRNAs (miRNAs), long non-coding (lncRNAs) as well as circular RNAs (circRNAs), have recently been proved to play potential roles in glucose metabolism. This review summarizes the effects of ncRNAs on HCC that participate in glucose metabolism and discuss the related mechanisms to find potential and effective targeted treatments for HCC.

Keywords: Hepatocellular carcinoma, glucose metabolism, glycolysis, non-coding RNA, microRNAs, long non-coding RNAs, circular RNAs

Introduction

Liver cancer is the sixth highest-incidence cancer and it is also the 4th most deadly cancer with easy metastasis and poor prognosis. There were approximately 850,000 new cases and 780,000 deaths of liver cancer occurred in 2018. HCC accounts for 75% to 85% of primary liver cancer and more than half of the world's HCC cases occur in China [1].

For early-stage HCC, a comprehensive treatment plan is mainly based on surgery, combined with transcatheter arterial chemoembolization (TACE) and radiofrequency ablation [2]. However, it is regrettable that most patients have reached an advanced stage or distant metastasis at the first diagnosis, that losing the opportunity for surgery. As for advanced HCC, there is no standard treatment, and the 5-year survival rate of HCC is 3% to 5% [3].

Despite the unremitting efforts of researchers, the key molecular mechanism of HCC development remains inconclusive, limiting the progression of therapeutic regimens. Recently,

multiple lines of evidence have shown that metabolic reprogramming is closely related to the occurrence and development of HCC, among which glucose metabolism reprogramming is one of the most prominent features [4]. The mechanisms of glucose metabolic reprogramming often involve gene mutations, especially C-MYC and P53 [5, 6]. While changes in the expression or activity of glucose metabolism genes and related glucose metabolism enzymes also have global effects [7].

Non-coding RNAs (ncRNAs) are RNA transcribed from DNA and unable to encode proteins, which account for the majority of RNAs (**Figure 1**). With the improvement and maturity of ncRNAs identification technology, the crucial role of ncRNAs in tumorigenesis has been increasingly recognized. They participate in almost all biological functions, including proliferation, apoptosis, migration, invasion, EMT, cancer stem cells and drug resistance [8, 9].

Multiple lines of evidence have manifested that ncRNAs, mainly miRNAs, lncRNAs and circular RNAs (circRNAs) may play pivotal roles in repro-

NcRNAs orchestrates glucose metabolism in HCC




Figure 1. A. The biogenesis of ncRNAs. miRNAs are transcribed into pri-miRNA under the help of RNA Polymerase II, pri-miRNA is processed into a pre-miRNA by Drosha/DGCR8 in the nucleus, and then transported to the cytoplasm via Exportin-5, pre-miRNA is further cleaved by Dicer to form mature miRNA. lncRNAs are transcribed by RNA Polymerase II or III. B. CircRNAs formed by exon circularization.

gramming glucose metabolism, including glycolytic pathway, pentose phosphate pathway (PPP) and gluconeogenesis via altering the expression or activity of the glucose metabolism genes and related glucose metabolism enzymes (**Figure 2**) and regulating the activation of various signaling pathways (**Figure 3**).

MiRNAs

MiRNAs are a class of ncRNAs that about 22 nt in length. MiRNAs completely or incompletely combine with the 3' untranslated regions (3'-UTR) of the target genes, leading to direct degradation or blocked translation of the target genes, thereby decreasing the expression of

these specific mRNAs by either mRNA cleavage or translational repression [10-12]. In recent years, more and more studies have found that miRNAs are involved in the regulation of glucose metabolism, including glycolytic pathway, pentose phosphate pathway (PPP) and gluconeogenesis. Here we discuss the role of miRNAs on glucose metabolism of HCC. The related studies' contents are summarized in **Table 1**.

MiRNAs involved in aerobic glycolysis

Aerobic glycolysis also called the "Warburg effect", which preference for tumor cells acquire energy through glycolysis rather than oxidative phosphorylation even in the sufficient aerobic and mitochondrial function [13]. Glycolysis also provides raw materials for other anabolic [13]. Furthermore, aerobic glycolysis can produce a large amount of lactic acid and creating an acidic microenvironment, which is conducive to tumor cell invasion and metastasis [15]. Therefore, inhibition of aerobic glycolysis may be a promising anti-tumor therapy [16-19].

The regulation of glycolysis is mainly through manipulating the activity of the rate-controlling enzymes (key enzymes), including hexokinase (HK), 6-phosphate fructokinase (PFK) and pyruvate kinase (PK). Hypoxia-inducible factors (HIFs), particularly HIF-1 α is a crucial mediator of hypoxia response, promoting glycolysis process to adapt cancer cells to hypoxic environment [20]. On the one hand, emerging studies demonstrated that alters miRNAs expression profiles may orchestrate the glycolytic pathway of HCC cells by regulating the expression of glycolysis-related enzymes, including HK, PFK, PK, HIFs. On the other hand, different glucose levels also alter the level of miRNAs. For example, the expression of miR-483-3p is low under

NcRNAs orchestrates glucose metabolism in HCC




Figure 2. Glycolysis pathway, PPP and gluconeogenesis pathway of HCC, and the role of ncRNAs in glucose metabolism via regulating rate-controlling enzyme and GLUT.

low glucose, while under high glucose conditions, the expression of miR-483-3p increases and inhibits apoptosis [21].

MiRNAs and HK: HK is the first rate-limiting enzymes of aerobic glycolysis, which catalysis phosphorylation of glucose to glucose 6-phosphate [22]. The overexpression of HK has been highlighted in HCC cells. On the contrary, sup-

pression of HK expression may induce HCC cells apoptosis. Therefore, it is considered to be a vital molecule in the glycolysis pathway and has been proposed as a therapeutic target for cancer.

In mammals, there are four hexokinase isoforms, HKI, HKII, HKIII, and HKIV. On the one hand, miRNAs weaken glycolysis of HCC by

NcRNAs orchestrates glucose metabolism in HCC




Figure 3. NcRNAs may play an important role in regulating glucose metabolism of HCC through different signal pathways and mechanisms.

directly targeting HKII expressions. For instance, miR-885-5p, miR-143 and miR-125b have proved plays a decisive role in limiting glycolysis in HCC cells by targeting HKII, thereby inhibiting the HCC growth [23-25]. On the other hand, miRNAs elevate glycolysis of HCC by stabilizing HKII expression. For example, miR-455-3p induces cell proliferation, metastasis and glycolysis by increasing HKII expression, which is achieved by stabilizing HKII protein through proteasome. Mir-455-3-p directly targets the 3'UTR of AMP-activated protein kinase subunit beta 2 (AMPK β 2), an important role in the AMPK pathway. Suppress AMPK β 2 expression has been shown to induced p-mTOR, Snail and HKII expressions, leading to enhanced cell glycolysis [26].

Furthermore, HKI has also been found to be involved in the regulation of glycolysis. MiR-139-5p negative regulate HKI and 6-phospho-fructo-2-kinase/fructose-2,6-biphosphatase 3 (PFKFB3) expression by directly targeting the ETS1, which is a transcription factor and bound

to the promoters of the HKI and PFKFB3 genes [27].

MiRNAs and PFK: PFK, the second rate-limiting enzyme in glycolysis that catalyzes fructose-6-phosphate to fructose-1,6-bisphosphate [28]. Mammals have three PFK isoforms: liver (PFKL), muscle (PFKM) and platelet (PFKP). Expression of PFKP was significantly up-regulated in a variety of cancers, including brain cancer, pancreatic cancer, breast cancer and HCC [29-31].

Recently, several miRNAs have been shown to be involved in regulating glycolysis of HCC by altering the expression of PFK. For instance, miR-520 family, including miR-520a-3p, miR-520b, and miR-520e are indicated to inhibited glycolysis by target the 3'UTR of PFKP in HCC. On the contrary, inhibit the expression of miR-520a/b/e was notably potentiated the rate of glycolysis [32]. MiR-338-3p dampens glycolysis by directly interacted with PFKL. MiR-338-3p act as a tumor suppressor was down-regulated

NcRNAs orchestrates glucose metabolism in HCC

Table 1. MiRNAs involved in glucose metabolism in HCC

Glucose metabolism	microRNA	Effect	Target	Mechanism	Reference
glycolytic pathway	miR-885-5p	Decrease	HKII	-	[23]
	miR-143	Decrease	HKII	-	[24]
	miR-125b	Decrease	-	Inhibit HKII	[25]
	miR-455-3p	Increase	AMPK β 2	stabilizes HKII protein active AMPK pathway	[26]
	miR-139-5p	Decrease	ETS1	Inhibit HKI and PFKFB3 expression	[27]
	miR-520 family	Decrease	PFKP	-	[32]
	miR-338-3p	Decrease	PFKL	-	[33]
	miR-139-5p	Decrease	PKM2	-	[27]
	miR-199a	Decrease	PKM2	-	[36]
	miR-491-5p	Decrease	PKM2	-	[37]
	miR-338-3p	Decrease	PKM2	-	[38]
	miR-122	Decrease	PKM2	-	[39]
	miR-4417	Increase	-	Promote PKM2 phosphorylation	[40]
	miR-365a-3p	Increase	-	Inhibit PKM2 degradation	[41]
	miR-374b	Decrease	hnRNPA1	Inhibit PKM2 expression	[42]
	miR-142-3p	Decrease	LDHA	-	[44]
	miR-383	Decrease	LDHA	-	[45]
	miR-34a	Decrease	LDHA	-	[46]
	miR-100-5p	Decrease	LDHA	-	[47]
	miR-592	Decrease	WSB1	Disrupting HIF-1 α stabilization	[51]
	miR-199a-5p	Decrease	HIF-1 α	-	[52]
	miR-3662	Decrease	HIF-1 α	Inactivate ERK and JNK	[53]
	miR-145	Decrease	-	Inhibit HIF-1 α and PDK1 expression	[54]
	miR-873	Increase	NDFIP1	Active AKT/mTOR	[56]
	miR-199b-5p	Decrease	-	Up-regulates HIF-1 α transcription through overexpression of NPAS2	[57]
	mitomiR-181a-5p	Decrease	-	Impair electron transport chain	[59]
	miR-342-3p	Decrease	IGF-1R	Downregulate GLUT1 Inactivate PI3K/AKT	[61]
	miR-455-5p	Decrease	IGF-1R	Downregulate GLUT1 Inactivate PI3K/AK	[62]
	miR-129-5p	Decrease	PDK4	-	[55]
	pentose phosphate pathway	miR-122	Decrease	G6PD	-
miR-4641		Decrease	PCK1	-	[69]
gluconeogenesis	miR-517a	Decrease	FBP1	-	[72]
	miR-23a	Decrease	G6PC PGC-1 α	-	[74]
	miR-96	Increase	IRS-1	-	[76]
	miR-122	Increase	-	Promoted glutaminolysis	[78]

“-”: unknown. Abbreviations: HCC: Hepatocellular carcinoma; HK: Hexokinase; PFK: 6-phosphate fructokinase; PK: Pyruvate kinase; HIFs: Hypoxia-inducible factors; AMPK β 2: AMP-activated protein kinase subunit beta 2; PFKFB3: 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3; PKM2: Pyruvate kinase M2; LDHA: Lactate dehydrogenase A; IGF-1R: Insulin-like growth factor-1 receptor; GLUT: Glucose transporters; PDK4: Pyruvate dehydrogenase kinase 4; PEPCK: Phosphoenolpyruvate carboxylase; FBP1: Fructose 1,6-bisphosphatase; G6PC: Glucose 6-phosphatase; PGC-1 α : Peroxisome proliferator-activated receptor gamma, coactivator 1 alpha; IRS-1: Insulin receptor substrate 1.

in HCC, while, the expression of miR-338-3p was increased after HCC cells treatment with 125I irradiation, which was considered as a potential strategy for HCC. The up-regulated miR-338-3p elevates the suppression of PFKL expression, thereby inhibiting glycolysis of HCC [33].

Moreover, miRNAs can regulate PFKFB to inhibit glycolysis. For instance, miR-139-5p reduced the expression of HKI and PFKFB3, thereby inhibiting aerobic glycolysis [27]. PFKFB is an allosteric activator of PFK1 and an effective stimulator of glycolysis. PFKFB3 is related to many aspects of tumorigenesis and development. And recent studies also found that PFKFB3 regulates immune response and the sensitivity of sorafenib in HCC [17, 34].

MiRNAs and PK: PK is the last rate-limiting enzyme in glycolysis, which transfers phosphate group from phosphoenolpyruvate to adenosine diphosphate (ADP) to produce pyruvate ATP [35]. So it is not surprising that miRNAs can regulate glycolysis and the progression of HCC through manipulating PKM2 expression. For example, miR-199a, miR-491-5p, miR-338-3p, and miR-122 act as a tumor suppressor that induces apoptosis, growth arrest and suppresses glycolysis of HCC cells by direct binding to 3'UTR of PKM2, which is one of isozymes of pyruvate kinase PK, and is universally expressed in embryonic development, tissue repair, and tumors [36-39].

Several miRNAs act as cancer-promoting factors by promoting PKM2 activation. For example, miR-4417 could promote the phosphorylation of PKM2 and facilitates the proliferation and glycolysis of HCC cells [40]. miR-365a-3p dampens PKM2 degradation and provokes Akt/mTOR signaling pathway activation via targeting linc01554, and thereby accelerates glycolytic of HCC cells [41].

MiRNAs not only manipulates the proliferation, metastasis and invasion of HCC through regulating PKM2, but also regulates the drug sensitivity of HCC. MiR-374b reverses the sensitivity of sorafenib-resistant HCC cells to sorafenib by counteracting PKM2-mediated glycolysis pathway via binding to the 3'-UTR of hnRNPA1, which is a RNA-binding protein [42]. It provides new ideas for the treatment of sorafenib resistance.

MiRNAs and lactate dehydrogenase: Lactate dehydrogenase A (LDHA) catalyzes the last key step of glycolysis, catalyzing lactate dehydrogenation to pyruvate. Down-regulation of LDHA expression notably inhibits the proliferation, invasion and migration of HCC cells [43]. Several miRNAs act as tumor suppressors, and decrease the expression of LDHA by binding to the 3'-UTR of LDHA mRNA, thereby inhibiting aerobic glycolysis. For instance, miR-142-3p, miR-383, miR-34a and miR-100-5p inhibit aerobic glycolysis and cell proliferation of HCC by targeting the 3'-UTR of LDHA [44-47].

MiRNAs and HIF-1 α : Emerging data have indicated that hypoxia of cancer cells is the initial factor for malignant transformation and even metastasis of tumors, and also one of the key factors that lead to the resistance of tumor cells to radiochemotherapy [48]. Among them, hypoxia-inducible factor-1 alpha (HIF-1 α) is an important transcriptional regulator under hypoxia, which has the function of promoting tumor angiogenesis and glucose metabolism, affecting tumor cell proliferation [49]. Under the induction of HIF-1 α , glycolytic-related enzyme expression increased, which further increase glycolytic activity, thus improve the imbalance between energy supply and energy consumption caused by hypoxia in tumors [50].

Several miRNAs have been demonstrated to decreased the efficacy of glycolysis by suppressing HIF-1 α expression. For instance, miR-592 disrupts HIF-1 α protein stabilization and inhibited glycolysis in HCC cells via targeting WSB1 mRNA [51]. Moreover, WSB1 negatively regulates JAK-STAT signaling pathway. MiR-199a-5p inhibits glycolysis by directly targets HIF-1 α [52]. Interestingly, HIF-1 α overexpression can in turn inhibit the abundance of miR-199a-5p under hypoxic environment. miR-3662 directly targets HIF-1 α , and negatively regulates the activation of ERK and JNK signaling pathways in HCC, thereby dampened glycolysis [53]. miR-145 attenuates the expression of HIF-1 α and PDK1, opposing glycolysis and suppress cell survival of HCC cells [54].

Other mechanisms: MiR-129-5p targets the mitochondrial matrix protein pyruvate dehydrogenase kinase 4 (PDK4), which diminished phosphorylation of the E1 α subunit of pyruvate dehydrogenase (PDH) complex and hinders glycolysis [55].

NcRNAs orchestrates glucose metabolism in HCC

MiR-873 promotes the Warburg effect through activating AKT/mTOR signaling pathway via targeting NDFIP1, which triggers metabolic shift and NDFIP1 was shown to suppress the PTEN/AKT signaling pathway activation [56].

MiRNAs also involved in glycolysis of HCC by regulating circadian gene expression. For example, miR-199b-5p prevents glycolysis via inhibiting NPAS2 expression, which is a circadian gene and notable boosts glycolysis through elevating the transcription of HIF-1 α and downregulated the expression of PGC-1 α [57]. Numerous studies have indicated that the consequences of circadian rhythm disturbances are related to many diseases, including obesity, type 2 diabetes and cancer [58]. Additionally, NPAS2 promoted glycolysis by heterodimeric with BMAL1, another core circadian rhythm factor, which regulates the expression of a variety of target genes including glycolysis in HCC cells [57].

MitomiR-181a-5p damages mitochondrial function and accelerates glycolysis in HCC by regulating the electron transport chain (ETC) [59]. MitomiRs refers to miRNAs located in the mitochondria. It is well known that mitochondria are the site of oxidative phosphorylation and adenosine triphosphate (ATP) production, and the explanation of the mechanism of oxidative phosphorylation is mainly based on the electron transport chain [60]. So it is not surprising that dampened ECT can inhibit oxidative phosphorylation and promotes glycolysis in HCC.

miR-342-3p and miR-455-5p attenuate glycolysis of HCC cells by target insulin-like growth factor-1 receptor (IGF-1R), which has been verified to activates the intracellular AKT signaling pathway, then up-regulates the expression of GLUT1 on the plasma membrane and enhances glycolysis in HCC cells [61, 62].

MiRNAs involved in PPP

The PPP is also known as hexose phosphate bypass, which is a glucose catabolic pathway commonly found in animals, plants and microorganisms. In addition to providing energy, the PPP provides a variety of raw materials for anabolic metabolism, for example, NADPH and ribose-5-phosphate. Therefore, the PPP is an important multifunctional metabolic pathway [63, 64].

Multiple lines of evidence indicated that miRNAs participate in PPP by altering G6PD expression. G6PD is a rate-limiting enzyme of the PPP, and its expression is significantly up-regulated in HCC. MiR-122 plays a tumor suppressive role in HCC and dampens PPP process by targeting G6PD [65].

MiRNAs involved in gluconeogenesis

Gluconeogenesis is the process by which an organism converts a variety of non-carbohydrate carbon substrates into free glucose. In mammals, the liver is the main organ of gluconeogenesis that ensures the blood sugar levels are normal. It has been reported that gluconeogenesis pathway is reduced in HCC. Additionally, Metformin was originally considered as an oral hypoglycemic agent. Recently, metformin has attracted people's attention due to its anti-tumor therapeutic effect in inhibiting liver gluconeogenesis [22].

Gluconeogenesis and glycolysis are coordinated, glycolysis is extremely active in HCC cells, and thereby the activity of gluconeogenesis is inhibited accordingly. Gluconeogenesis appears to be the reverse reaction of glycolysis, because the seven steps of gluconeogenesis are all reverse reactions of glycolysis and are catalyzed by the same enzymes. But there are three steps in glycolysis, which are irreversible reactions, and these three steps must be bypassed during gluconeogenesis [66].

These three steps are bypassed by the following three rate-controlling enzymes: phosphoenolpyruvate carboxykinase (PEPCK), Fructose 1,6-bisphosphatase (FBP1), and glucose 6-phosphatase (G6Pase) [67]. They can not only affect the overall speed of the entire metabolic pathway but also change the direction of metabolism.

Emerging evident verified that miRNAs play critical roles in gluconeogenesis of HCC by influencing the expression of the rate-controlling enzyme, including PEPCK, FBP1 and G6PC.

MiRNAs and PEPCK: PEPCK is the first rate-controlling enzyme of gluconeogenesis, which catalyzes the irreversible reaction of phosphoenolpyruvate (PEP) and HCO₃⁻ to oxaloacetate (OAA) and inorganic phosphoric acid. It is the first and most important reaction in the pro-

NcRNAs orchestrates glucose metabolism in HCC

cess of gluconeogenesis [68]. Accumulating evidence has indicated that PCK1 overexpression could block the glycolysis process and initiate the gluconeogenesis process by potentiating PEPCK expression.

MiR-4641 attenuates the expression of PEPCK by targeting PCK1, thereby inhibiting gluconeogenesis and promoting the growth and migration of HCC cells [69]. PCK1 is the coding gene of PEPCK and is widely involved in metabolic and biological processes such as glucose metabolism, lipid metabolism, diabetes, and tumor cell proliferation and apoptosis [70].

MiRNAs and FBP1: FBP1 is the second rate-controlling enzyme of the gluconeogenesis process. It hydrolyzes fructose 1,6-diphosphate (FDP) to phosphoric acid and fructose 6-diphosphate (F6P) [67]. The expression of FBP1 was suppressed in various cancers. In addition, HCC patients with low FBP1 expression have a higher malignant classification, including tumor enlargement, poor differentiation, impaired gluconeogenesis and enhanced glycolysis [71]. Therefore, FBP1 is expected to become a reliable prognostic marker for HCC patients.

MiR-517a inhibits gluconeogenesis, promotes glycolysis of HCC via directly targeting FBP1 [72]. MiR-517a was dominantly overexpressed and FBP1 expression was significantly lower in HCC cells and tissues. Ectopic expression of FBP1 upregulates gluconeogenesis and weakens miR-517a induced cell proliferation.

MiRNAs and G6Pase: G6Pase is the last rate-controlling enzyme of the gluconeogenesis process, which catalyzes glucose 6 phosphates to glucose [67]. G6Pase expression is significantly decreased in HCC cell lines and clinical tissue, G6Pase expression also correlated with tumor grade in HCCs. Moreover, G6Pase deficiency leads to glycogen storage disease type-1a (GSD-1a), while HCC is a long-term complication of GSD-1a. Restore G6Pase expression normalizes glucose homeostasis and prevents the development of HCC in the initial stage [73].

Recently, several miRNAs have been proved to reduce gluconeogenesis of HCC by suppressing the expression of G6Pase. Wang et al. showed that miR-23a inhibits gluconeogenesis and promotes HCC progression by targeting G6PC, which is encoding the key gluconeogenic enzymes

G6Pase, thereby suppressing the expression of G6Pase [74]. Moreover, miR-23a inhibits gluconeogenesis via targeting peroxisome proliferator-activated receptor gamma, coactivator 1 alpha (PGC-1 α), which has been demonstrated to accelerate hepatic gluconeogenesis in previous studies [75].

Other mechanisms: Jeong et al. found that miR-96 is up-regulated in mitochondrial dysfunction HCC cells and exhibit insulin resistance. Furthermore, mitochondrial dysfunction induced miR-96 overexpression, increase the level of gluconeogenesis in HCC cells via targeting the 3'UTR of Insulin receptor substrate 1 (IRS-1), which resulted in inhibition of gluconeogenesis in HCC cells [76].

It has been affirmed that mammalian cells utilize glutamine (Gln) as an alternative energy source of glucose and as an anaplerotic source for biomass generation. Glutamine-derived oxaloacetate is converted to PEP by PEPCK2, and then participates in the gluconeogenesis and other biosynthesis pathways [77]. Some miRNA can regulate gluconeogenesis by altering glutamine metabolism. For example, the level of miR-122 is reduced in HCC, and its expression is negative correlates with malignant classification. Silence miR-122 expression promoted glutaminolysis but suppressed gluconeogenesis in the mouse model. In contrast, ectopic expression of miR-122 promotes gluconeogenesis [78].

LncRNAs

LncRNAs refers to transcripts longer than 200 nucleotide units and is not involved in protein-coding [79]. LncRNAs regulations are diverse, which have been shown to regulate almost every step of gene expression. LncRNAs may serve as signals, decoys, guides or scaffolds. They also act as "sponge" or competing endogenous RNAs (ceRNAs) through the combination of their complementary miRNA response elements (MREs) and the primary miRNAs, playing a positive or negative role in the processing and expression of mature mRNAs, thereby indirectly participating in a variety of physiological process [80].

Growing researches have reported that lncRNAs play an important role in various tumors, including HCC. Although lncRNAs have been exten-

NcRNAs orchestrates glucose metabolism in HCC

Table 2. LncRNAs and circRNAs involved in glucose metabolism in HCC

Glucose metabolism	LncRNA/circRNA	Effect	Target	Mechanism	Signaling pathway	Reference
glycolysis	linc01554	Decrease	-	Promote PKM2 ubiquitination	Inactive Akt/mTOR	[42]
	lncRNA Ftx	Increase	-	Promote GLUT	active PPAR γ	[85]
	lncRNA WFDC21P	Decrease	-	Suppress PFKP and PKM2 transcription	-	[86]
	linc-RoR	Decrease	miR-145	Inhibit HIF-1 α and PDK1 expression	-	[54]
	lncRNAIDH1-AS1	Decrease	-	Inhibit HIF-1 α expression	-	[89]
	lncRNA RAET1K	Increase	miR-100-5p	Increase LDHA expression	-	[47]
gluconeogenesis	MALAT1	Decrease	-	Enhance TCF7L2 translation	Active Wnt and mTOR pathway	[90]
glycolysis	circMAT2B	Increase	-	Promote PKM2 expression	-	[39]
gluconeogenesis	circC3P1	Increase	miR-4641	Promote PCK1 expression	-	[69]

"-": unknown. Abbreviations: HCC: Hepatocellular carcinoma; PKM2: Pyruvate kinase M2; HIFs: Hypoxia-inducible factors; GLUT: Glucose transporters; PPAR γ : Peroxisome proliferator-activated receptor γ ; PFKP: the platelet isoform of phosphofructokinase; PDK1: Pyruvate dehydrogenase kinase; LDHA: Lactate dehydrogenase A; MALAT1: Metastasis-associated lung adenocarcinoma transcript 1; TCF7L2: Transcription factor 7 like 2.

sively studied in regulating gene expression, the role of lncRNAs in glucose metabolism is not yet clear. The related research contents are listed in **Table 2**.

LncRNAs involved in glycolytic pathway

An increasing number of studies have verified that lncRNAs could influence the glycolysis pathway in a variety of tumors by regulating the expression and activation of glycolytic enzymes. We summarize the influence of lncRNAs on glycolytic pathway and related mechanisms, including PI3K/Akt/mTOR pathway, PPAR γ , PFK/PKM2, and HIF-1 α , so as to find potential and effective targeted therapies for HCC.

LncRNAs and PI3K/Akt/mTOR pathway: The PI3K/AKT/mTOR pathway is an intracellular signaling pathway that is often activated in various types of cancer cells, and its regulatory role in the process of glycolysis has been emphasized in HCC [81].

linc01554 diminished the rate of glycolytic by accelerating PKM2 degradation and suppresses Akt/mTOR signaling pathway activation. linc01554 is highly expressed in liver, while it is frequently down-regulated in HCC in the level of protein mRNA and DNA contrast with adjacent normal tissues. Moreover, silence linc01554 was significantly associated with aggressive clinicopathological features [41]. Furthermore, overexpression of linc01554 in combination with Akt inhibitor MK2206 exhibits a synergistic effect compared with used each alone, which shed new light on introducing Akt inhibitor to HCC treatment.

LncRNAs and PPAR γ : Peroxisome proliferator-activated receptor γ (PPAR γ) belongs to the family of PPARs, which plays a crucial regulatory role in cell differentiation, proliferation, metabolism and tumorigenesis [82]. Currently, PPAR γ is the most extensively researched subtype. It has been demonstrated that ectopic expression of PPAR γ inhibits WNT/ β -catenin pathway and then downregulates PDK1, thus suppressed glycolysis [83].

lncRNA Ftx facilitates glucose consumption through promotes GLUT, including GUL1 and GUL4, and inhibits tumor necrosis factor (TNF) α and leptin expression via targeting PPAR γ in HCC cells [84]. Furthermore, lncRNA Ftx potentiates glycolysis of HCC via directly targeting PPAR γ , which elevating the activity and expression of glycolytic enzymes (LDH and PFKL) and decreases the activity of Krebs-cycle-associated molecules (TNF α , leptin and PDK1).

LncRNAs and PFK/PKM2: lncRNA WFDC21P diminished glycolysis via decreasing the expression and activity of PFKP and PKM2 [85]. Moreover, lncRNA WFDC21P is positively regulated by Nur77, which is a member of the orphan nuclear receptor NR4A family. Nur77 is down-regulated in HCC and shows the ability to inhibit glycolysis and promotes gluconeogenesis by stabilizing PEPCK1 [86]. Additionally, linc01554 diminished the rate of glycolytic by accelerating PKM2 degradation [41]. lncRNA Ftx contributes to glycolysis of HCC via enhancing the activity and expression of PFKL [84].

LncRNA and HIF-1 α : Recently, Takahashi et al. uncover linc-RoR knockdown significantly decreased HIF-1 α expression as well as PDK1

NcRNAs orchestrates glucose metabolism in HCC

expression, especially under hypoxia stress. Linc-RoR is a hypoxia-responsive lncRNAs, and thereby cancer cells release a large amount of linc-RoR under hypoxia context, facilitating cell survival in recipient cells by promoting glycolysis. In detail, Linc-RoR serves as a miRNA “sponge” to limit miR-145, which attenuates the expression of HIF-1 α and PDK1, opposing glycolysis and suppress cell survival of HCC cells [54].

LncRNAIDH1-AS1 potentiates the activity of isocitrate dehydrogenase 1 and augments the production of α -ketoglutarate under normoxia, attenuating the expression of HIF-1 α and inhibits glycolysis. Furthermore, MYC-dependent inhibition of lncRNAIDH1-AS1, induction of the Warburg effect by HIF1 α [87].

Additionally, HIF1 α boosting lncRNA RAET1K expression and facilitates glycolysis by binding to the promoter region of lncRNA RAET1K, and lncRNA RAET1K sponge miR-100-5p, which directly binding to LDHA and significantly inhibits glycolysis [47].

LncRNAs involved in gluconeogenesis

LncRNA metastasis-associated lung adenocarcinoma transcript 1 (MALAT1) acts as an oncogene in HCC. MALAT1 potentiated glycolytic and attenuated gluconeogenesis by enhancing the expression of Transcription factor 7 like 2 (TCF7L2) [88]. TCF7L2 is one of the earliest genes to be found and deeply studied and TCF7L2 transcription factor strongly activates the Wnt signaling pathway. And gluconeogenesis has been shown to be negatively regulated by TCF7L2 [89].

Moreover, MALAT1 activates the mTORC1 pathway by increasing phosphorylation of eIF4E binding protein (4EBP1) and enhancing the expression and function of the splicing oncoprotein SRSF1. 4EBP1 is an important downstream effector of mTORC1 pathway, and SRSF1 has been reported to activate mTOR and protein translation. Both the Wnt and mTOR signaling pathways have been suggested to play a negative regulator role in gluconeogenesis program of HCC [90].

CircRNAs

CircRNAs, a group of endogenous ncRNAs with covalently closed continuous circular structure

formed by exon circularization [91]. However, circRNAs were considered to be caused by splicing errors and without function in the previous. Due to the rapid development of high-throughput sequencing, increasingly circRNAs have been discovered and proved to be involved in a variety of biological processes.

The potential functions of circRNAs include: a) served as “miRNA sponge”, inhibit the function of the target miRNAs, one circRNAs may “sponge” multiple mRNAs; b) regulate the splicing of pre-mRNA, thereby affecting protein production; c) interact with proteins; d) translated into protein or polypeptide; e) Regulate the expression of parental genes [92].

Emerging evidence indicates that changes in circRNA expression profiles play pivotal roles in the initiation and development of various cancers, including breast cancer [93], colon cancer [94], gastric cancer [95] and HCC [96]. Even though various studies have highlight miRNAs and lncRNAs partly account for glucose reprogram of HCC, it was very limited research about circRNAs was involved in metabolic regulation in HCC. Recently, Li et al. found that circRNA MAT2B promotes glycolysis and endows HCC cells with clinical aggressiveness under hypoxic [38]. Mechanistically, circMAT2B promotes glycolysis and HCC progression via increasing the abundance of the miR-338-3p, which subsequently blocking PKM2. Moreover, circRNA circC3P1 has been proved to promote the gluconeogenesis process and suppress HCC growth and metastasis through miR-4641/PCK1 pathway [69]. CircC3P1 enhancing the expression of PCK1 by sponging miR-4641 in HCC. PCK1 is the coding gene of PEPCK, which is a rate-controlling enzyme of gluconeogenesis.

Conclusion and future directions

Glucose is the main nutritional component of the animal body, and a unique source of fuel for some organizations to generate and sustain biological function. The reprogramming of glucose metabolism is one of the hallmarks of HCC. This reprogramming is caused by various factors and is closely related to the initiation, development and poor prognosis of HCC. The glucose metabolic differences between HCC and normal cells may become potential new targets. Some related drugs are already undergoing clinical trials and are expected to be used in clinical later (**Table 3**).

NcRNAs orchestrates glucose metabolism in HCC

Table 3. Glucose metabolism targets and drugs which are in pre-clinical and clinical development for anti-tumor therapy

Target	Drug	Status	References
GLUTs	Phloretin	Preclinical	[99, 100]
	Fasentin	Preclinical	[101]
	STF-31	Preclinical	[102]
	WZB117	Preclinical	[103]
	Ritonavir	Phase III	[104]
	Silybin	Phase I	[105]
HKII	2-Deoxy-D-glucose	Phase II	[106, 107]
	Lonidamine	Phase II	[108]
	Genistein-27	Preclinical	[109]
	Benserazide	Preclinical	[110]
	Resveratrol	Phase I	[111]
	Astragalin	Preclinical	[25]
PDK	Chrysin	Preclinical	[112]
	Dichloroacetate	Phase I	[113-115]
LDHA	Oxamate	Preclinical	[116]
	FX11	Preclinical	[117]
	Quinoline-3-sulfonamide	Preclinical	[118]
	GNE-140	Preclinical	[119]
	PSTMB	Preclinical	[120]
PKM2	Shikonin	Preclinical	[121]
	Benserazide	Preclinical	[122]
PFKFB	3PO	Preclinical	[123]
	PFK158	Preclinical	[124]
GAPDH	Bromopyruvate	Preclinical	[125]
IDH	Enasidenib	Approved	[126]
	Ivosidenib	Approved	[127]
	DS-1001b	Preclinical	[128]
	Olutasidenib	Preclinical	[129]
	GSK864	Preclinical	[130]
	BAY1436032	Preclinical	[131]
	HMS-101	Preclinical	[132]
I-8	Preclinical	[133]	

Abbreviations: GLUT: Glucose transporters; HK: Hexokinase; PDK: Pyruvate dehydrogenase kinase; LDHA: Lactate dehydrogenase A; PKM2: Pyruvate kinase M2; PFKFB3: 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3; GAPDH: glyceraldehyde-3-phosphate dehydrogenase; IDH: Isocitrate dehydrogenase.

Extensive studies have demonstrated the functions of ncRNAs in glucose reprogramming. It provides us a novel perspective of tumorigenesis and potential therapeutic targets. In addition, the aberrant expression of glucose metabolism related ncRNAs may be serving as biomarkers for the diagnosis or prognosis of HCC. Moreover, some ncRNA-based therapeutics for other diseases has been used in clinical treatment. Although these ncRNAs may only serve as a fine-tuning mechanism, the synergistic

effect of multiple ncRNAs may lead to specific major metabolic changes in glucose metabolism in HCC. For example, Tang et al. synthesizes an artificial lncRNA (AlncRNA) that could target multiple sorafenib-resistance-related miRNAs simultaneously, including miR-21, miR-153, miR-216a, miR-217, and miR-494, restore the sensitivity of drug-resistant HCC cells to sorafenib again [97]. It brings a bright research prospect that ncRNAs combined with the glucose-metabolism-related-enzyme inhibitors would be a better choice than utilized inhibitors alone in the battle against HCC.

However, there are still some difficulties remain to be overcome. First, researches on glucose-metabolism-related ncRNAs are still very limited, especially in lncRNAs and circRNAs, which urgently needed to be explored. Second, how to efficiently deliver ncRNA molecules to the target is the biggest problem facing in their clinical application. There are several major problems with ncRNA compounds delivery: 1) Naked single-stranded RNA molecules are easily degraded by nucleases in the physiological environment; 2) RNA molecules are immunogenic and activate the immune system;

3) ncRNAs are biological macromolecules, and they are negatively charged, making it difficult to cross the cell membrane into cells; 4) The toxic effects of ncRNAs are unknown and may overlap the toxicity of existing chemotherapy drugs. 5) Liver is the site where the drug is acting and the site where the drug is metabolized. The amount of medicine, adverse reactions, and treatment of adverse reactions in patients with HCC need attention [98]. Therefore, it is necessary to design a suitable ncRNA de-

livery method or delivery vector to deliver ncRNAs to the target site to fully realize its huge disease treatment potential.

After decades of searching for ncRNA-based therapeutics, some ncRNA-based therapeutics has been approved for disease treatment. The development of ncRNA-based therapeutics in the future will focus on three aspects: 1) explore more ncRNA molecules that are critical for different glucose metabolism steps. 2) develop chemical modification technology for nucleic acid therapeutics to further improve the efficiency of ncRNA-based therapeutics; 3) develop diverse delivery systems for different types of ncRNA-based therapeutics based on the size and mechanism of action of ncRNAs; 4) combining ncRNA-based therapeutics with a variety of other drugs, such as combining ncRNAs with gene-editing tools, including CRISPR/Cas9-gRNA, antibodies, small molecules, or chemotherapeutics to maximize the effect of HCC treatment; 5) design individualized ncRNA-based therapeutics according to the etiology classification of patients by using gene sequencing technology.

Taken together, ncRNA-based therapies in orchestrates glucose metabolism of HCC have promising prospects. However, the evidence for the practical clinical application of ncRNAs is still very limited and desirable for further investigation.

Acknowledgements

The work was supported by grants from the National Natural Science Foundation of China (No.81772995 and 81472266). The Excellent Youth Foundation of Jiangsu Province, China (BK20140032); Jiangsu Province's Key Provincial Talents Program (No.ZDRCA2016090).

Consent for publication was obtained from all participants.

Disclosure of conflict of interest

None.

Abbreviations

HCC, Hepatocellular carcinoma; ncRNA, Non-coding RNA; miRNA, MicroRNA; lncRNA, Long noncoding RNA; circRNA, Circular RNA; TACE, Transcatheter arterial chemoembolization; GL-

UT, Glucose transporters; PPP, Pentose phosphate pathway; ATP, Adenosine triphosphate; HBP, Hexosamine biosynthetic pathway; HK, Hexokinase; PFK, 6-phosphate fructokinase; PK, Pyruvate kinase; HIFs, Hypoxia-inducible factors; VDAC, Voltage-dependent anion channel 1; 3'-UTR, 3'-untranslated region; AMPK β 2, AMP-activated protein kinase subunit beta 2; PFKFB3, 6-phosphofructo-2-kinase/fructose-2,6-biphosphatase 3; ADP, Adenosine diphosphate; PKM2, Pyruvate kinase M2; LDHA, Lactate dehydrogenase A; SOCS, Suppressor of cytokine signaling; ETC, Electron transport chain; IGF-1R, Insulin-like growth factor-1 receptor; PDK4, Pyruvate dehydrogenase kinase 4; PDH, Pyruvate dehydrogenase; PEPCK, Phosphoenolpyruvate carboxylase; FBP1, Fructose 1,6-bisphosphatase; G6PC, Glucose 6-phosphatase; GSD-1a, Glycogen storage disease type-1a; PGC-1 α , Peroxisome proliferator-activated receptor gamma, coactivator 1 alpha; FDP, Fructose 1,6-diphosphate; PEP, Phosphoenolpyruvate; OAA, Oxaloacetate; TCA, Tricarboxylic acid; IRS-1, Insulin receptor substrate 1; Gln, Glutamine; PPAR γ , Peroxisome proliferator-activated receptor γ ; TNF, Tumor necrosis factor; MALAT1, Metastasis-associated lung adenocarcinoma transcript 1; TCF7L2, Transcription factor 7 like 2; 4EBP1, eIF4E binding protein; gRNA, Guide RNA.

Address correspondence to: Rui Wang, Department of Medical Oncology, Jinling Hospital, The First School of Clinical Medicine, Southern Medical University, Nanjing, China. E-mail: wangrui218@163.com

References

- [1] Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA and Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; 68: 394-424.
- [2] Maluccio M and Covey A. Recent progress in understanding, diagnosing, and treating hepatocellular carcinoma. *CA Cancer J Clin* 2012; 62: 394-399.
- [3] Heimbach JK, Kulik LM, Finn RS, Sirlin CB, Abecassis MM, Roberts LR, Zhu AX, Murad MH and Marrero JA. AASLD guidelines for the treatment of hepatocellular carcinoma. *Hepatology* 2018; 67: 358-380.
- [4] Faubert B, Solmonson A and DeBerardinis RJ. Metabolic reprogramming and cancer progression. *Science* 2020; 368: w5473.

NcRNAs orchestrates glucose metabolism in HCC

- [5] David CJ, Chen M, Assanah M, Canoll P and Manley JL. HnRNP proteins controlled by c-Myc deregulate pyruvate kinase mRNA splicing in cancer. *Nature* 2010; 463: 364-368.
- [6] Huang Q, Li J, Xing J, Li W, Li H, Ke X, Zhang J, Ren T, Shang Y, Yang H, Jiang J and Chen Z. CD147 promotes reprogramming of glucose metabolism and cell proliferation in HCC cells by inhibiting the p53-dependent signaling pathway. *J Hepatol* 2014; 61: 859-866.
- [7] Baig MH, Adil M, Khan R, Dhadi S, Ahmad K, Rabbani G, Bashir T, Imran MA, Husain FM, Lee EJ, Kamal MA and Choi I. Enzyme targeting strategies for prevention and treatment of cancer: implications for cancer therapy. *Semin Cancer Biol* 2019; 56: 1-11.
- [8] Beermann J, Piccoli M, Viereck J and Thum T. Non-coding RNAs in development and disease: background, mechanisms, and therapeutic approaches. *Physiol Rev* 2016; 96: 1297-1325.
- [9] Fan C, Tang Y, Wang J, Xiong F, Guo C, Wang Y, Zhang S, Gong Z, Wei F, Yang L, He Y, Zhou M, Li X, Li G, Xiong W and Zeng Z. Role of long non-coding RNAs in glucose metabolism in cancer. *Mol Cancer* 2017; 16: 130.
- [10] Pasquinelli AE. MicroRNAs and their targets: recognition, regulation and an emerging reciprocal relationship. *Nat Rev Genet* 2012; 13: 271-282.
- [11] Sayed D and Abdellatif M. MicroRNAs in development and disease. *Physiol Rev* 2011; 91: 827-887.
- [12] Lu J and Clark AG. Impact of microRNA regulation on variation in human gene expression. *Genome Res* 2012; 22: 1243-1254.
- [13] Warburg OH. The classic: the chemical constitution of respiration ferment. *Clin Orthop Relat Res* 2010; 468: 2833-2839.
- [14] Li J, Liang N, Long X, Zhao J, Yang J, Du X, Yang T, Yuan P, Huang X, Zhang J, He X and Xing J. SDHC-related deficiency of SDH complex activity promotes growth and metastasis of hepatocellular carcinoma via ROS/NFκB signaling. *Cancer Lett* 2019; 461: 44-55.
- [15] Peppicelli S, Bianchini F and Calorini L. Extracellular acidity, a "reappreciated" trait of tumor environment driving malignancy: perspectives in diagnosis and therapy. *Cancer Metast Rev* 2014; 33: 823-832.
- [16] Liu J, Pan C, Guo L, Wu M, Guo J, Peng S, Wu Q and Zuo Q. A new mechanism of trastuzumab resistance in gastric cancer: MACC1 promotes the Warburg effect via activation of the PI3K/AKT signaling pathway. *J Hematol Oncol* 2016; 9: 76.
- [17] Chen D, Ning W, Jiang Z, Peng Z, Zhu L, Zhuang S, Kuang D, Zheng L and Wu Y. Glycolytic activation of peritumoral monocytes fosters immune privilege via the PFKFB3-PD-L1 axis in human hepatocellular carcinoma. *J Hepatol* 2019; 71: 333-343.
- [18] Mori Y, Yamawaki K, Ishiguro T, Yoshihara K, Ueda H, Sato A, Ohata H, Yoshida Y, Minamino T, Okamoto K and Enomoto T. ALDH-dependent glycolytic activation mediates stemness and paclitaxel resistance in patient-derived spheroid models of uterine endometrial cancer. *Stem Cell Rep* 2019; 13: 730-746.
- [19] Hong W, Cai P, Xu C, Cao D, Yu W, Zhao Z, Huang M and Jin J. Inhibition of glucose-6-phosphate dehydrogenase reverses cisplatin resistance in lung cancer cells via the redox system. *Front Pharmacol* 2018; 9: 43.
- [20] Wang H, Zhao L, Zhu L, Wang Y, Pan D, Yao J, You Q and Guo Q. Wogonin reverses hypoxia resistance of human colon cancer HCT116 cells via downregulation of HIF-1α and glycolysis, by inhibiting PI3K/Akt signaling pathway. *Mol Carcinogen* 2014; 53: E107-E118.
- [21] Li X, Cheng T, He Y, Zhou S, Wang Y, Zhang K and Yu P. High glucose regulates ERp29 in hepatocellular carcinoma by LncRNA MEG3-miRNA 483-3p pathway. *Life Sci* 2019; 232: 116602.
- [22] DeWaal D, Nogueira V, Terry AR, Patra KC, Jeon S, Guzman G, Au J, Long CP, Antoniewicz MR and Hay N. Hexokinase-2 depletion inhibits glycolysis and induces oxidative phosphorylation in hepatocellular carcinoma and sensitizes to metformin. *Nat Commun* 2018; 9: 446.
- [23] Xu F, Yan JJ, Gan Y, Chang Y, Wang HL, He XX and Zhao Q. miR-885-5p negatively regulates Warburg effect by silencing hexokinase 2 in liver cancer. *Mol Ther Nucleic Acids* 2019; 18: 308-319.
- [24] Fang R, Xiao T, Fang Z, Sun Y, Li F, Gao Y, Feng Y, Li L, Wang Y, Liu X, Chen H, Liu X and Ji H. MicroRNA-143 (miR-143) regulates cancer glycolysis via targeting hexokinase 2 gene. *J Biol Chem* 2012; 287: 23227-23235.
- [25] Li W, Hao J, Zhang L, Cheng Z, Deng X and Shu G. Astragaloside reduces hexokinase 2 through increasing mir-125b to inhibit the proliferation of hepatocellular carcinoma cells in vitro and in vivo. *J Agr Food Chem* 2017; 65: 5961-5972.
- [26] Lin Y, Wu M, Huang Y, Yeh C, Cheng M, Chi H, Tsai C, Chung I, Chen C and Lin K. Taurine up-regulated gene 1 functions as a master regulator to coordinate glycolysis and metastasis in hepatocellular carcinoma. *Hepatology* 2018; 67: 188-203.
- [27] Hua S, Lei L, Deng L, Weng X, Liu C, Qi X, Wang S, Zhang D, Zou X, Cao C, Liu L and Wu D. miR-139-5p inhibits aerobic glycolysis, cell proliferation, migration, and invasion in hepatocellular carcinoma via a reciprocal regulatory interaction with ETS1. *Oncogene* 2018; 37: 1624-1636.

NcRNAs orchestrates glucose metabolism in HCC

- [28] Lee J, Liu R, Li J, Zhang C, Wang Y, Cai Q, Qian X, Xia Y, Zheng Y, Piao Y, Chen Q, de Groot JF, Jiang T and Lu Z. Stabilization of phosphofructokinase 1 platelet isoform by AKT promotes tumorigenesis. *Nat Commun* 2017; 8: 949.
- [29] Yang Y, Ishak Gabra MB, Hanse EA, Lowman XH, Tran TQ, Li H, Milman N, Liu J, Reid MA, Locasale JW, Gil Z and Kong M. MiR-135 suppresses glycolysis and promotes pancreatic cancer cell adaptation to metabolic stress by targeting phosphofructokinase-1. *Nat Commun* 2019; 10: 809.
- [30] Lee J, Liu R, Li J, Wang Y, Tan L, Li X, Qian X, Zhang C, Xia Y, Xu D, Guo W, Ding Z, Du L, Zheng Y, Chen Q, Lorenzi PL, Mills GB, Jiang T and Lu Z. EGFR-phosphorylated platelet isoform of phosphofructokinase 1 promotes PI3K activation. *Mol Cell* 2018; 70: 197-210.
- [31] Peng M, Yang D, Hou Y, Liu S, Zhao M, Qin Y, Chen R, Teng Y and Liu M. Intracellular citrate accumulation by oxidized ATM-mediated metabolism reprogramming via PFKP and CS enhances hypoxic breast cancer cell invasion and metastasis. *Cell Death Dis* 2019; 10: 228.
- [32] Park Y, Kim S, Han HD, Sohn BH, Kim JH, Liang J, Lu Y, Rodriguez-Aguayo C, Lopez-Berestein G, Mills GB, Sood AK and Lee J. Tat-activating regulatory DNA-binding protein regulates glycolysis in hepatocellular carcinoma by regulating the platelet isoform of phosphofructokinase through microRNA 520. *Hepatology* 2013; 58: 182-191.
- [33] Zheng J, Luo J, Zeng H, Guo L and Shao G. 125I suppressed the Warburg effect via regulating miR-338/PFKL axis in hepatocellular carcinoma. *Biomed Pharmacother* 2019; 119: 109402.
- [34] Li S, Dai W, Mo W, Li J, Feng J, Wu L, Liu T, Yu Q, Xu S, Wang W, Lu X, Zhang Q, Chen K, Xia Y, Lu J, Zhou Y, Fan X, Xu L and Guo C. By inhibiting PFKFB3, aspirin overcomes sorafenib resistance in hepatocellular carcinoma: inhibiting PFKFB3, ASA overcomes sorafenib resistance. *Int J Cancer* 2017; 141: 2571-2584.
- [35] Dayton TL, Jacks T and Vander HM. PKM2, cancer metabolism, and the road ahead. *Embo Rep* 2016; 17: 1721-1730.
- [36] Zhang LF, Lou JT, Lu MH, Gao C, Zhao S, Li B, Liang S, Li Y, Li D and Liu MF. Suppression of miR-199a maturation by HuR is crucial for hypoxia-induced glycolytic switch in hepatocellular carcinoma. *EMBO J* 2015; 34: 2671-2685.
- [37] Xu Q, Dou C, Liu X, Yang L, Ni C, Wang J, Guo Y, Yang W, Tong X and Huang D. Oviductus ranae protein hydrolysate (ORPH) inhibits the growth, metastasis and glycolysis of HCC by targeting miR-491-5p/PKM2 axis. *Biomed Pharmacother* 2018; 107: 1692-1704.
- [38] Li Q, Pan X, Zhu D, Deng Z, Jiang R and Wang X. Circular RNA MAT2B promotes glycolysis and malignancy of hepatocellular carcinoma through the miR-338-3p/PKM2 axis under hypoxic stress. *Hepatology* 2019; 70: 1298-1316.
- [39] Xu Q, Zhang M, Tu J, Pang L, Cai W and Liu X. MicroRNA-122 affects cell aggressiveness and apoptosis by targeting PKM2 in human hepatocellular carcinoma. *Oncol Rep* 2015; 34: 2054-2064.
- [40] Song L, Zhang W, Chang Z, Pan Y, Zong H, Fan Q and Wang L. miR-4417 targets tripartite motif-containing 35 (TRIM35) and regulates pyruvate kinase muscle 2 (PKM2) phosphorylation to promote proliferation and suppress apoptosis in hepatocellular carcinoma cells. *Med Sci Monitor* 2017; 23: 1741-1750.
- [41] Zheng Y, Li L, Jia Y, Zhang B, Li J, Zhu Y, Li M, He J, Zeng T, Ban X, Yuan Y, Li Y and Guan X. LINC01554-mediated glucose metabolism reprogramming suppresses tumorigenicity in hepatocellular carcinoma via downregulating PKM2 expression and inhibiting Akt/mTOR signaling pathway. *Theranostics* 2019; 9: 796-810.
- [42] Zhang M, Zhang H, Hong H and Zhang Z. MiR-374b re-sensitizes hepatocellular carcinoma cells to sorafenib therapy by antagonizing PKM2-mediated glycolysis pathway. *Am J Cancer Res* 2019; 9: 765.
- [43] Sheng SL, Liu JJ, Dai YH, Sun XG, Xiong XP and Huang G. Knockdown of lactate dehydrogenase A suppresses tumor growth and metastasis of human hepatocellular carcinoma. *FEBS J* 2012; 279: 3898-3910.
- [44] Hua S, Liu C, Liu L and Wu D. miR-142-3p inhibits aerobic glycolysis and cell proliferation in hepatocellular carcinoma via targeting LDHA. *Biochem Bioph Res Co* 2018; 496: 947-954.
- [45] Fang Z, He L, Jia H, Huang Q, Chen D and Zhang Z. The miR-383-LDHA axis regulates cell proliferation, invasion and glycolysis in hepatocellular cancer. *Iran J Basic Med Sci* 2017; 20: 187-192.
- [46] Zhang H, Wang Y and Han Y. MicroRNA-34a inhibits liver cancer cell growth by reprogramming glucose metabolism. *Mol Med Rep* 2018; 17: 4483-4489.
- [47] Zhou Y, Huang Y, Hu K, Zhang Z, Yang J and Wang Z. HIF1A activates the transcription of lncRNA RAET1K to modulate hypoxia-induced glycolysis in hepatocellular carcinoma cells via miR-100-5p. *Cell Death Dis* 2020; 11: 176.
- [48] Liang Y, Zheng T, Song R, Wang J, Yin D, Wang L, Liu H, Tian L, Fang X, Meng X, Jiang H, Liu J and Liu L. Hypoxia-mediated sorafenib resistance can be overcome by EF24 through Von Hippel-Lindau tumor suppressor-dependent

NcRNAs orchestrates glucose metabolism in HCC

- HIF-1 α inhibition in hepatocellular carcinoma. *Hepatology* 2013; 57: 1847-1857.
- [49] Semenza GL. Targeting HIF-1 for cancer therapy. *Nat Rev Cancer* 2003; 3: 721-732.
- [50] Lu H, Forbes RA and Verma A. Hypoxia-inducible factor 1 activation by aerobic glycolysis implicates the Warburg effect in carcinogenesis. *J Biol Chem* 2002; 277: 23111-23115.
- [51] Jia Y, Zhao J, Li B, Gao K, Song Y, Liu M, Yang X, Xue Y, Wen A and Shi L. miR-592/WSB1/HIF-1 α axis inhibits glycolytic metabolism to decrease hepatocellular carcinoma growth. *Oncotarget* 2016; 7: 35257.
- [52] Li B, He L, Zuo D, He W, Wang Y, Zhang Y, Liu W and Yuan Y. Mutual regulation of mir-199a-5p and hif-1 α modulates the warburg effect in hepatocellular carcinoma. *J Cancer* 2017; 8: 940-949.
- [53] Chen Z, Zuo X, Zhang Y, Han G, Zhang L, Wu J and Wang X. MiR-3662 suppresses hepatocellular carcinoma growth through inhibition of HIF-1 α -mediated Warburg effect. *Cell Death Dis* 2018; 9: 549.
- [54] Takahashi K, Yan IK, Haga H and Patel T. Modulation of hypoxia-signaling pathways by extracellular linc-RoR. *J Cell Sci* 2014; 127: 1585-1594.
- [55] Han H, Li W, Shen H, Zhang J, Zhu Y and Li Y. microRNA-129-5p, a c-Myc negative target, affects hepatocellular carcinoma progression by blocking the Warburg effect. *J Mol Cell Biol* 2016; 8: 400-410.
- [56] Zhang Y, Zhang C, Zhao Q, Wei W, Dong Z, Shao L, Li J, Wu W, Zhang H, Huang H, Liu F and Jin S. The miR-873/NDP1 axis promotes hepatocellular carcinoma growth and metastasis through the AKT/mTOR-mediated Warburg effect. *Am J Cancer Res* 2019; 9: 927.
- [57] Yuan P, Yang T, Mu J, Zhao J, Yang Y, Yan Z, Hou Y, Chen C, Xing J, Zhang H and Li J. Circadian clock gene NPAS2 promotes reprogramming of glucose metabolism in hepatocellular carcinoma cells. *Cancer Lett* 2020; 469: 498-509.
- [58] Masri S, Kinouchi K and Sassone-Corsi P. Circadian clocks, epigenetics, and cancer. *Curr Opin Oncol* 2015; 27: 50-56.
- [59] Zhuang X, Chen Y, Wu Z, Xu Q, Chen M, Shao M, Cao X, Zhou Y, Xie M, Shi Y, Zeng Y and Bu H. Mitochondrial miR-181a-5p promotes glucose metabolism reprogramming in liver cancer by regulating the electron transport chain. *Carcinogenesis* 2019; 41: 972-983.
- [60] Fan S, Tian T, Chen W, Lv X, Lei X, Zhang H, Sun S, Cai L, Pan G, He L, Ou Z, Lin X, Wang X, Perez MF, Tu Z, Ferrone S, Tannous BA and Li J. Mitochondrial miRNA determines chemoresistance by reprogramming metabolism and regulating mitochondrial transcription. *Cancer Res* 2019; 79: 1069-1084.
- [61] Liu W, Kang L, Han J, Wang Y, Shen C, Yan Z, Tai Y and Zhao C. miR-342-3p suppresses hepatocellular carcinoma proliferation through inhibition of IGF-1R-mediated Warburg effect. *Onco Targets Ther* 2018; 11: 1643-1653.
- [62] Hu Y, Yang Z, Bao D, Ni J and Lou J. miR-455-5p suppresses hepatocellular carcinoma cell growth and invasion via IGF-1R/AKT/GLUT1 pathway by targeting IGF-1R. *Pathol Res Pract* 2019; 215: 152674.
- [63] Stincone A, Prigione A, Cramer T, Wamelink MM, Campbell K, Cheung E, Olin-Sandoval V, Grüning NM, Krüger A, Tauqeer Alam M, Keller MA, Breitenbach M, Brindle KM, Rabinowitz JD and Ralser M. The return of metabolism: biochemistry and physiology of the pentose phosphate pathway. *Biol Rev* 2015; 90: 927-963.
- [64] Kowalik MA, Guzzo G, Morandi A, Perra A, Menegon S, Masgras I, Trevisan E, Angioni MM, Fornari F, Quagliata L, Ledda-Columbano GM, Gramantieri L, Terracciano L, Giordano S, Chiarugi P, Rasola A and Columbano A. Metabolic reprogramming identifies the most aggressive lesions at early phases of hepatic carcinogenesis. *Oncotarget* 2016; 7: 32375.
- [65] Barajas JM, Reyes R, Guerrero MJ, Jacob ST, Motiwala T and Ghoshal K. The role of miR-122 in the dysregulation of glucose-6-phosphate dehydrogenase (G6PD) expression in hepatocellular cancer. *Sci Rep* 2018; 8: 9105.
- [66] Pilkis SJ and Granner DK. Molecular physiology of the regulation of hepatic gluconeogenesis and glycolysis. *Annu Rev Physiol* 1992; 54: 885-909.
- [67] Wang Z and Dong C. Gluconeogenesis in cancer: function and regulation of PEPCK, FB-Pase, and G6Pase. *Trends Cancer* 2019; 5: 30-45.
- [68] Vincent EE, Sergushichev A, Griss T, Gingras MC, Samborska B, Ntimbane T, Coelho PP, Blagih J, Raissi TC, Choinière L, Bridon G, Loginicheva E, Flynn BR, Thomas EC, Tavaré JM, Avizonis D, Pause A, Elder DJ, Artyomov MN and Jones RG. Mitochondrial phosphoenolpyruvate carboxykinase regulates metabolic adaptation and enables glucose-independent tumor growth. *Mol Cell* 2015; 60: 195-207.
- [69] Zhong L, Wang Y, Cheng Y, Wang W, Lu B, Zhu L and Ma Y. Circular RNA circC3P1 suppresses hepatocellular carcinoma growth and metastasis through miR-4641/PCK1 pathway. *Biochem Biophys Res Commun* 2018; 499: 1044-1049.
- [70] Shi H, Fang R, Li Y, Li L, Zhang W, Wang H, Chen F, Zhang S, Zhang X and Ye L. The oncoprotein HBXIP suppresses gluconeogenesis through modulating PCK1 to enhance the growth of hepatoma cells. *Cancer Lett* 2016; 382: 147-156.

NcRNAs orchestrates glucose metabolism in HCC

- [71] Hirata H, Sugimachi K, Komatsu H, Ueda M, Masuda T, Uchi R, Sakimura S, Nambara S, Saito T, Shinden Y, Iguchi T, Eguchi H, Ito S, Terashima K, Sakamoto K, Hirakawa M, Honda H and Mimori K. Decreased expression of fructose-1,6-bisphosphatase associates with glucose metabolism and tumor progression in hepatocellular carcinoma. *Cancer Res* 2016; 76: 3265-3276.
- [72] Zhang D, Li Z, Li T, Luo D, Feng X, Liu Y and Huang J. miR-517a promotes Warburg effect in HCC by directly targeting FBP1. *Onco Targets Ther* 2018; 11: 8025-8032.
- [73] Lee YM, Jun HS, Pan C, Lin SR, Wilson LH, Mansfield BC and Chou JY. Prevention of hepatocellular adenoma and correction of metabolic abnormalities in murine glycogen storage disease type Ia by gene therapy. *Hepatology* 2012; 56: 1719-1729.
- [74] Wang B, Hsu S, Frankel W, Ghoshal K and Jacob ST. Stat3-mediated activation of miR-23a suppresses gluconeogenesis in hepatocellular carcinoma by downregulating G6PC and PGC-1 α . *Hepatology* 2012; 56: 186-197.
- [75] Sharabi K, Lin H, Tavares CDJ, Dominy JE, Camporez JP, Perry RJ, Schilling R, Rines AK, Lee J, Hickey M, Bennion M, Palmer M, Nag PP, Bittker JA, Perez J, Jedrychowski MP, Ozcan U, Gygi SP, Kamenecka TM, Shulman GI, Schreiber SL, Griffin PR and Puigserver P. Selective chemical inhibition of PGC-1 α gluconeogenic activity ameliorates type 2 diabetes. *Cell* 2017; 169: 148-160.
- [76] Jeong H, Park S, Yang W and Lee W. The induction of miR-96 by mitochondrial dysfunction causes impaired glycogen synthesis through translational repression of IRS-1 in SK-Hep1 cells. *Biochem Biophys Res Commun* 2013; 434: 503-508.
- [77] Altman BJ, Stine ZE and Dang CV. From Krebs to clinic: glutamine metabolism to cancer therapy. *Nat Rev Cancer* 2016; 16: 619-634.
- [78] Sengupta D, Cassel T, Teng KY, Aljuhani M, Chowdhary VK, Hu P, Zhang X, Fan TW and Ghoshal K. Regulation of hepatic glutamine metabolism by miR-122. *Mol Metab* 2020; 34: 174-186.
- [79] Matsui M and Corey DR. Non-coding RNAs as drug targets. *Nat Rev Drug Discov* 2017; 16: 167-179.
- [80] Engreitz JM, Ollikainen N and Guttman M. Long non-coding RNAs: spatial amplifiers that control nuclear structure and gene expression. *Nat Rev Mol Cell Bio* 2016; 17: 756-770.
- [81] Hu H, Juvekar A, Lyssiotis CA, Lien EC, Albeck JG, Oh D, Varma G, Hung YP, Ullas S, Lauring J, Seth P, Lundquist MR, Tolan DR, Grant AK, Needleman DJ, Asara JM, Cantley LC and Wulf GM. Phosphoinositide 3-kinase regulates glycolysis through mobilization of aldolase from the actin cytoskeleton. *Cell* 2016; 164: 433-446.
- [82] Patitucci C, Couchy G, Bagattin A, Cañeque T, de Reyniès A, Scoazec J, Rodriguez R, Pontoglio M, Zucman-Rossi J, Pende M and Panasyuk G. Hepatocyte nuclear factor 1 α suppresses steatosis-associated liver cancer by inhibiting PPAR γ transcription. *J Clin Invest* 2017; 127: 1873-1888.
- [83] Zuo Q, He J, Zhang S, Wang H, Jin G, Jin H, Cheng Z, Tao X, Yu C, Li B, Yang C, Wang S, Lv Y, Zhao F, Yao M, Cong W, Wang C and Qin W. PGC1 α suppresses metastasis of HCC by inhibiting Warburg effect via PPAR γ -dependent WNT/ β -catenin/PDK1 axis. *Hepatology* 2020; [Epub ahead of print].
- [84] Li X, Zhao Q, Qi J, Wang W, Zhang D, Li Z and Qin C. lncRNA Ftx promotes aerobic glycolysis and tumor progression through the PPAR γ pathway in hepatocellular carcinoma. *Int J Oncol* 2018; 53: 551-566.
- [85] Guan Y, Huang Q, Ai Y, Chen Q, Zhao W, Wang X, Wu Q and Chen H. Nur77-activated lncRNA WFDC21P attenuates hepatocarcinogenesis via modulating glycolysis. *Oncogene* 2020; 39: 2408-2423.
- [86] Kolluri SK, Zhu X, Zhou X, Lin B, Chen Y, Sun K, Tian X, Town J, Cao X, Lin F, Zhai D, Kitada S, Luciano F, O'Donnell E, Cao Y, He F, Lin J, Reed JC, Satterthwait AC and Zhang X. A short Nur77-derived peptide converts Bcl-2 from a protector to a killer. *Cancer Cell* 2008; 14: 285-298.
- [87] Xiang S, Gu H, Jin L, Thorne RF, Zhang XD and Wu M. lncRNA IDH1-AS1 links the functions of c-Myc and HIF1 α via IDH1 to regulate the Warburg effect. *Proc Natl Acad Sci U S A* 2018; 115: E1465-E1474.
- [88] Malakar P, Stein I, Saragovi A, Winkler R, Stern-Ginossar N, Berger M, Pikarsky E and Karni R. Long noncoding RNA MALAT1 regulates cancer glucose metabolism by enhancing mTOR-mediated translation of TCF7L2. *Cancer Res* 2019; 79: 2480-2493.
- [89] Shao W, Wang D, Chiang Y, Ip W, Zhu L, Xu F, Columbus J, Belsham DD, Irwin DM, Zhang H, Wen X, Wang Q and Jin T. The Wnt signaling pathway effector TCF7L2 controls gut and brain proglucagon gene expression and glucose homeostasis. *Diabetes* 2013; 62: 789-800.
- [90] Grant SF, Thorleifsson G, Reynisdottir I, Benediktsson R, Manolescu A, Sainz J, Helgason A, Stefansson H, Emilsson V, Helgadottir A, Styrkarsdottir U, Magnusson KP, Walters GB, Palsdottir E, Jonsdottir T, Gudmundsdottir T, Gylfason A, Saemundsdottir J, Wilensky RL, Reilly MP, Rader DJ, Bagger Y, Christiansen C,

- Gudnason V, Sigurdsson G, Thorsteinsdottir U, Gulcher JR, Kong A and Stefansson K. Variant of transcription factor 7-like 2 (TCF7L2) gene confers risk of type 2 diabetes. *Nat Genet* 2006; 38: 320-323.
- [91] Vo JN, Cieslik M, Zhang Y, Shukla S, Xiao L, Zhang Y, Wu YM, Dhanasekaran SM, Engelke CG, Cao X, Robinson DR, Nesvizhskii AI and Chinnaiyan AM. The landscape of circular RNA in cancer. *Cell* 2019; 176: 869-881.
- [92] Memczak S, Jens M, Elefsinioti A, Torti F, Krueger J, Rybak A, Maier L, Mackowiak SD, Gregersen LH, Munschauer M, Loewer A, Ziebold U, Landthaler M, Kocks C, Le Noble F and Rajewsky N. Circular RNAs are a large class of animal RNAs with regulatory potency. *Nature* 2013; 495: 333-338.
- [93] Liang G, Ling Y, Mehrpour M, Saw PE, Liu Z, Tan W, Tian Z, Zhong W, Lin W, Luo Q, Lin Q, Li Q, Zhou Y, Hamai A, Codogno P, Li J, Song E and Gong C. Autophagy-associated circRNA circCDYL augments autophagy and promotes breast cancer progression. *Mol Cancer* 2020; 19: 65.
- [94] Han K, Wang F, Cao C, Ling H, Chen J, Chen R, Feng Z, Luo J, Jin X, Duan J, Li S, Ma N, Yun J, Guan X, Pan Z, Lan P, Xu R and Xie D. CircLONP2 enhances colorectal carcinoma invasion and metastasis through modulating the maturation and exosomal dissemination of microRNA-17. *Mol Cancer* 2020; 19: 60.
- [95] Jie M, Wu Y, Gao M, Li X, Liu C, Ouyang Q, Tang Q, Shan C, Lv Y, Zhang K, Dai Q, Chen Y, Zeng S, Li C, Wang L, He F, Hu C and Yang S. CircMRPS35 suppresses gastric cancer progression via recruiting KAT7 to govern histone modification. *Mol Cancer* 2020; 19: 56.
- [96] Yu J, Xu QG, Wang ZG, Yang Y, Zhang L, Ma JZ, Sun SH, Yang F and Zhou WP. Circular RNA cS-MARCA5 inhibits growth and metastasis in hepatocellular carcinoma. *J Hepatol* 2018; 68: 1214-1227.
- [97] Tang S, Tan G, Jiang X, Han P, Zhai B, Dong X, Qiao H, Jiang H, Sun X. An artificial lncRNA targeting multiple miRNAs overcomes sorafenib resistance in hepatocellular carcinoma cells. *Oncotarget* 2016; 7: 73257-73269.
- [98] Bonetta L. RNA-based therapeutics: ready for delivery? *Cell* 2009; 136: 581-584.
- [99] Wu K, Ho C, Chen Z, Chen L, Whang-Peng J, Lin T and Ho Y. The apple polyphenol phloretin inhibits breast cancer cell migration and proliferation via inhibition of signals by type 2 glucose transporter. *J Food Drug Anal* 2018; 26: 221-231.
- [100] Wu C, Ho Y, Tsai C, Wang Y, Tseng H, Wei P, Lee C, Liu R and Lin S. In vitro and in vivo study of phloretin-induced apoptosis in human liver cancer cells involving inhibition of type II glucose transporter. *Int J Cancer* 2009; 124: 2210-2219.
- [101] Wood TE, Dalili S, Simpson CD, Hurren R, Mao X, Saiz FS, Gronda M, Eberhard Y, Minden MD, Bilan PJ, Klip A, Batey RA and Schimmer AD. A novel inhibitor of glucose uptake sensitizes cells to FAS-induced cell death. *Mol Cancer Ther* 2008; 7: 3546-3555.
- [102] Chan DA, Sutphin PD, Nguyen P, Turcotte S, Lai EW, Banh A, Reynolds GE, Chi JT, Wu J, Solow-Cordero DE, Bonnet M, Flanagan JU, Bouley DM, Graves EE, Denny WA, Hay MP and Giaccia AJ. Targeting GLUT1 and the Warburg effect in renal cell carcinoma by chemical synthetic lethality. *Sci Transl Med* 2011; 3: 70r-94r.
- [103] Ojelabi OA, Lloyd KP, Simon AH, De Zutter JK and Carruthers A. WZB117 (2-Fluoro-6-(m-hydroxybenzoyloxy) Phenylm-Hydroxybenzoate) Inhibits GLUT1-mediated sugar transport by binding reversibly at the exofacial sugar binding site. *J Biol Chem* 2016; 291: 26762-26772.
- [104] Poordad F, Castro RE, Asatryan A, Aguilar H, Cacoub P, Dieterich D, Marinho RT, Carvalho A, Siddique A, Hu YB, Charafeddine M, Bondin M, Khan N, Cohen DE and Felizarta F. Long-term safety and efficacy results in hepatitis C virus genotype 1-infected patients receiving ombitasvir/paritaprevir/ritonavir + dasabuvir ± ribavirin in the TOPAZ-I and TOPAZ-II trials. *J Viral Hepatitis* 2020; 27: 497-504.
- [105] Siegel AB, Narayan R, Rodriguez R, Goyal A, Jacobson JS, Kelly K, Ladas E, Lunghofer PJ, Hansen RJ, Gustafson DL, Flaig TW, Tsai WY, Wu DP, Lee V and Greenlee H. A phase I dose-finding study of silybin phosphatidylcholine (Milk Thistle) in patients with advanced hepatocellular carcinoma. *Integr Cancer Ther* 2014; 13: 46-53.
- [106] Raez LE, Papadopoulos K, Ricart AD, Chiorean EG, DiPaola RS, Stein MN, Rocha Lima CM, Schlesselman JJ, Tolba K, Langmuir VK, Kroll S, Jung DT, Kurtoglu M, Rosenblatt J and Lampidis TJ. A phase I dose-escalation trial of 2-deoxy-d-glucose alone or combined with docetaxel in patients with advanced solid tumors. *Cancer Chemoth Pharm* 2013; 71: 523-530.
- [107] Mohanti BK, Rath GK, Anantha N, Kannan V, Das BS, Chandramouli BA, Banerjee AK, Das S, Jena A, Ravichandran R, Sahi UP, Kumar R, Kapoor N, Kalia VK, Dwarakanath BS and Jain V. Improving cancer radiotherapy with 2-deoxy-d-glucose: phase I/II clinical trials on human cerebral gliomas. *Int J Radiat Oncol Biol Phys* 1996; 35: 103-111.
- [108] Berruti A, Bitossi R, Gorzegno G, Bottini A, Alquati P, De Matteis A, Nuzzo F, Giardina G, Danese S, De Lena M, Lorusso V, Farris A, Sarobba MG, DeFabiani E, Bonazzi G, Castiglione F, Bumma C, Moro G, Bruzzi P and Dogliotti L.

- Time to progression in metastatic breast cancer patients treated with epirubicin is not improved by the addition of either cisplatin or lonidamine: final results of a phase III study with a factorial design. *J Clin Oncol* 2002; 20: 4150-4159.
- [109] Tao L, Wei L, Liu Y, Ding Y, Liu X, Zhang X, Wang X, Yao Y, Lu J, Wang Q and Hu R. Gen-27, a newly synthesized flavonoid, inhibits glycolysis and induces cell apoptosis via suppression of hexokinase II in human breast cancer cells. *Biochem Pharmacol* 2017; 125: 12-25.
- [110] Li W, Zheng M, Wu S, Gao S, Yang M, Li Z, Min Q, Sun W, Chen L, Xiang G and Li H. Benserazide, a dopadecarboxylase inhibitor, suppresses tumor growth by targeting hexokinase 2. *J Exp Clin Canc Res* 2017; 36: 58.
- [111] Dai W, Wang F, Lu J, Xia Y, He L, Chen K, Li J, Li S, Liu T, Zheng Y, Wang J, Lu W, Zhou Y, Yin Q, Abudumijiti H, Chen R, Zhang R, Zhou L, Zhou Z, Zhu R, Yang J, Wang C, Zhang H, Zhou Y, Xu L and Guo C. By reducing hexokinase 2, resveratrol induces apoptosis in HCC cells addicted to aerobic glycolysis and inhibits tumor growth in mice. *Oncotarget* 2015; 6: 13703-13717.
- [112] Xu D, Jin J, Yu H, Zhao Z, Ma D, Zhang C and Jiang H. Chrysin inhibited tumor glycolysis and induced apoptosis in hepatocellular carcinoma by targeting hexokinase-2. *J Exp Clin Canc Res* 2017; 36: 44.
- [113] Xuan Y, Hur H, Ham I, Yun J, Lee J, Shim W, Kim YB, Lee G, Han S and Cho YK. Dichloroacetate attenuates hypoxia-induced resistance to 5-fluorouracil in gastric cancer through the regulation of glucose metabolism. *Exp Cell Res* 2014; 321: 219-230.
- [114] Dunbar EM, Coats BS, Shroads AL, Langae T, Lew A, Forder JR, Shuster JJ, Wagner DA and Stacpoole PW. Phase 1 trial of dichloroacetate (DCA) in adults with recurrent malignant brain tumors. *Invest New Drugs* 2014; 32: 452-464.
- [115] Chu QS, Sangha R, Spratlin J, Vos LJ, Mackey JR, McEwan AJ, Venner P and Michelakis ED. A phase I open-labeled, single-arm, dose-escalation, study of dichloroacetate (DCA) in patients with advanced solid tumors. *Invest New Drugs* 2015; 33: 603-610.
- [116] Zhou M, Zhao Y, Ding Y, Liu H, Liu Z, Fodstad O, Riker AI, Kamarajugadda S, Lu J, Owen LB, Ledoux SP and Tan M. Warburg effect in chemosensitivity: targeting lactate dehydrogenase-A re-sensitizes taxol-resistant cancer cells to taxol. *Mol Cancer* 2010; 9: 33.
- [117] Le A, Cooper CR, Gouw AM, Dinavahi R, Maitra A, Deck LM, Royer RE, Vander Jagt DL, Semenza GL and Dang CV. Inhibition of lactate dehydrogenase A induces oxidative stress and inhibits tumor progression. *Proc Natl Acad Sci U S A* 2010; 107: 2037-2042.
- [118] Billiard J, Dennison JB, Briand J, Annan RS, Chai D, Colón M, Dodson CS, Gilbert SA, Greshock J, Jing J, Lu H, McSurdy-Freed JE, Orband-Miller LA, Mills GB, Quinn CJ, Schneck JL, Scott GF, Shaw AN, Waitt GM, Wooster RF and Duffy KJ. Quinoline 3-sulfonamides inhibit lactate dehydrogenase A and reverse aerobic glycolysis in cancer cells. *Cancer Metab* 2013; 1: 19.
- [119] Purkey HE, Robarge K, Chen J, Chen Z, Corson LB, Ding CZ, DiPasquale AG, Dragovich PS, Eigenbrot C, Evangelista M, Fauber BP, Gao Z, Ge H, Hitz A, Ho Q, Labadie SS, Lai KW, Liu W, Liu Y, Li C, Ma S, Malek S, O'Brien T, Pang J, Peterson D, Salphati L, Sideris S, Ultsch M, Wei B, Yen I, Yue Q, Zhang H and Zhou A. Cell active hydroxylactam inhibitors of human lactate dehydrogenase with oral bioavailability in mice. *ACS Med Chem Lett* 2016; 7: 896-901.
- [120] Kim E, Chung T, Han CW, Park SY, Park KH, Jang SB and Ha K. A novel lactate dehydrogenase inhibitor, 1-(phenylseleno)-4-(trifluoromethyl) benzene, suppresses tumor growth through apoptotic cell death. *Sci Rep* 2019; 9: 3969.
- [121] Chen J, Xie J, Jiang Z, Wang B, Wang Y and Hu X. Shikonin and its analogs inhibit cancer cell glycolysis by targeting tumor pyruvate kinase-M2. *Oncogene* 2011; 30: 4297-4306.
- [122] Zhou Y, Huang Z, Su J, Li J, Zhao S, Wu L, Zhang J, He Y, Zhang G, Tao J, Zhou J, Chen X and Peng C. Benserazide is a novel inhibitor targeting PKM2 for melanoma treatment. *Int J Cancer* 2020; 147: 139-151.
- [123] Clem B, Telang S, Clem A, Yalcin A, Meier J, Simmons A, Rasku MA, Arumugam S, Dean WL, Eaton J, Lane A, Trent JO and Chesney J. Small-molecule inhibition of 6-phosphofructo-2-kinase activity suppresses glycolytic flux and tumor growth. *Mol Cancer Ther* 2008; 7: 110-120.
- [124] Mondal S, Roy D, Sarkar Bhattacharya S, Jin L, Jung D, Zhang S, Kalogera E, Staub J, Wang Y, Xuyang W, Khurana A, Chien J, Telang S, Chesney J, Tapolsky G, Petras D and Shridhar V. Therapeutic targeting of PFKFB3 with a novel glycolytic inhibitor PFK158 promotes lipophagy and chemosensitivity in gynecologic cancers. *Int J Cancer* 2019; 144: 178-189.
- [125] Sun X, Sun G, Huang Y, Hao Y, Tang X, Zhang N, Zhao L, Zhong R and Peng Y. 3-Bromopyruvate regulates the status of glycolysis and BCNU sensitivity in human hepatocellular carcinoma cells. *Biochem Pharmacol* 2020; 177: 113988.
- [126] Kim ES. Enasidenib: first global approval. *Drugs* 2017; 77: 1705-1711.
- [127] Dhillon S. Ivosidenib: first global approval. *Drugs* 2018; 78: 1509-1516.
- [128] Nakagawa M, Nakatani F, Matsunaga H, Seki T, Endo M, Ogawara Y, Machida Y, Katsumoto

NcRNAs orchestrates glucose metabolism in HCC

- T, Yamagata K, Hattori A, Fujita S, Aikawa Y, Ishikawa T, Soga T, Kawai A, Chuman H, Yokoyama N, Fukushima S, Yahiro K, Kimura A, Shimada E, Hirose T, Fujiwara T, Setsu N, Matsumoto Y, Iwamoto Y, Nakashima Y and Kitabayashi I. Selective inhibition of mutant IDH1 by DS-1001b ameliorates aberrant histone modifications and impairs tumor activity in chondrosarcoma. *Oncogene* 2019; 38: 6835-6849.
- [129] Caravella JA, Lin J, Diebold RB, Campbell AM, Ericsson A, Gustafson G, Wang Z, Castro J, Clarke A, Gotur D, Josephine HR, Katz M, Kershaw M, Yao L, Toms AV, Barr KJ, Dinsmore CJ, Walker D, Ashwell S and Lu W. Structure-based design and identification of FT-2102 (Olutasidenib), a potent mutant-selective IDH1 inhibitor. *J Med Chem* 2020; 63: 1612-1623.
- [130] Mao MJ and Leonardi DE. Vascular-endothelial response to IDH1 mutant fibrosarcoma secretome and metabolite: implications on cancer microenvironment. *Am J Cancer Res* 2019; 9: 122-133.
- [131] Chaturvedi A, Gupta C, Gabdouliline R, Borchert NM, Goparaju R, Kaulfuss S, Görlich K, Schottmann R, Othman B, Welzenbach J, Panknin O, Wagner M, Geffers R, Ganser A, Thol F, Jeffers M, Haegebarth A and Heuser M. Synergistic activity of IDH1 inhibitor BAY1436032 with azacitidine in IDH1 mutant acute myeloid leukemia. *Haematologica* 2020; [Epub ahead of print].
- [132] Chaturvedi A, Goparaju R, Gupta C, Weder J, Klünemann T, Araujo Cruz MM, Kloos A, Görlich K, Schottmann R, Othman B, Struys EA, Bähre H, Grote-Koska D, Brand K, Ganser A, Preller M and Heuser M. In vivo efficacy of mutant IDH1 inhibitor HMS-101 and structural resolution of distinct binding site. *Leukemia* 2020; 34: 416-426.
- [133] Jia P, Wu Y, Du H, Yang L, Zhang Z, Ma T, Li S, Yuan S, Lu L and Zha X. I-8, a novel inhibitor of mutant IDH1, inhibits cancer progression in vitro and in vivo. *Eur J Pharm Sci* 2019; 140: 105072.