

Gut microbiota mediated molecular events and therapy in liver diseases

Xiaoqiang Qi, Ming Yang, Joseph Stenberg, Rahul Dey, Leslie Fogwe, Muhammad Shawkat Alam, Eric T Kimchi, Kevin F Staveley-O'Carroll, Guangfu Li

ORCID number: Xiaoqiang Qi 0000-0002-5631-4362; Ming Yang 0000-0002-4895-5864; Joseph Stenberg 0000-0003-4435-0240; Rahul Dey 0000-0003-3882-4251; Leslie Fogwe 0000-0001-9869-3375; Muhammad Shawkat Alam 0000-0002-1227-1305; Eric T Kimchi 0000-0002-5046-1142; Kevin F Staveley-O'Carroll 0000-0003-2207-0379; Guangfu Li 0000-0002-9817-568X.

Author contributions: Qi X, Yang M, and Stenberg J collected data, wrote the paper, and contributed equally; Dey R, Fogwe L, and Alam MS collected data and partly drafted the paper's first version; Kimchi ET, Staveley-O'Carroll KF, and Li G designed, revised, and finalized the manuscript, co-corresponding authors.

Supported by NIH Grant, No. R01CA208396; and VA Merit Award, No. I01 BX004065-1.

Conflict-of-interest statement: The authors declare no conflict of interest.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0)

Xiaoqiang Qi, Ming Yang, Joseph Stenberg, Rahul Dey, Leslie Fogwe, Muhammad Shawkat Alam, Eric T Kimchi, Kevin F Staveley-O'Carroll, Guangfu Li, Department of Surgery, University of Missouri, Columbia, MO 65212, United States

Xiaoqiang Qi, Ming Yang, Eric T Kimchi, Kevin F Staveley-O'Carroll, Guangfu Li, Ellis Fischel Cancer Center, University of Missouri, Columbia, MO 65212, United States

Xiaoqiang Qi, Ming Yang, Eric T Kimchi, Kevin F Staveley-O'Carroll, Guangfu Li, VA Hospital, Harry S Truman Memorial VA Hospital, Columbia, MO 65201, United States

Guangfu Li, Department of Molecular Microbiology and Immunology, University of Missouri, Columbia, MO 65212, United States

Corresponding author: Guangfu Li, DVM, PhD, Assistant Professor, Department of Surgery, University of Missouri, 230 Jesse Hall, Columbia, MO 65212, United States.

liguan@health.missouri.edu

Abstract

Gut microbiota is a community of microorganisms that reside in the gastrointestinal tract. An increasing number of studies has demonstrated that the gut-liver axis plays a critical role in liver homeostasis. Dysbiosis of gut microbiota can cause liver diseases, including nonalcoholic fatty liver disease and alcoholic liver disease. Preclinical and clinical investigations have substantiated that the metabolites and other molecules derived from gut microbiota and diet interaction function as mediators to cause liver fibrosis, cirrhosis, and final cancer. This effect has been demonstrated to be associated with dysregulation of intrahepatic immunity and liver metabolism. Targeting these findings have led to the development of novel preventive and therapeutic strategies. Here, we review the cellular and molecular mechanisms underlying gut microbiota-mediated impact on liver disease. We also summarize the advancement of gut microbiota-based therapeutic strategies in the control of liver diseases.

Key Words: Gut microbiota; Intrahepatic immunity; Metabolite; Fecal microbial transplantation; Probiotic; Antibiotic

©The Author(s) 2020. Published by Baishideng Publishing Group Inc. All rights reserved.

license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Invited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: United States

Peer-review report's scientific quality classification

Grade A (Excellent): 0
Grade B (Very good): 0
Grade C (Good): C, C
Grade D (Fair): 0
Grade E (Poor): 0

Received: October 20, 2020

Peer-review started: October 20, 2020

First decision: November 13, 2020

Revised: November 24, 2020

Accepted: December 6, 2020

Article in press: December 6, 2020

Published online: December 28, 2020

P-Reviewer: Yuan J

S-Editor: Huang P

L-Editor: A

P-Editor: Liu JH



Core Tip: Accumulating evidence shows that gut microbiota plays a critical role in liver pathophysiology and targeting gut microbiota is a potential treatment option for chronic liver disease. Herein, this review explores the cellular and molecular mechanisms of how gut microbiota contributes to liver diseases, including alcohol-induced and nonalcohol-induced liver fatty liver diseases, liver fibrosis, cirrhosis, and cancer. This review also summarizes the current gut microbiota-based therapeutic strategies and discusses future directions in promoting gut microbiota-based therapy for liver diseases.

Citation: Qi X, Yang M, Stenberg J, Dey R, Fogwe L, Alam MS, Kimchi ET, Staveley-O'Carroll KF, Li G. Gut microbiota mediated molecular events and therapy in liver diseases. *World J Gastroenterol* 2020; 26(48): 7603-7618

URL: <https://www.wjgnet.com/1007-9327/full/v26/i48/7603.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v26.i48.7603>

INTRODUCTION

The growing evidence of gut microbial roles in human diseases attracts researchers' attention in exploring gut microbiota-mediated therapy. The gut microbiota is defined as the entire community of microorganisms residing in the gastrointestinal tract, and it is dominated mainly by bacteria^[1]. In the earlier stages of investigation, people focused solely on the gut microbiota's function regarding modulation of human nutrition and metabolism, on which a healthy status relies. However, dysbiosis of the gut microbiome may cause inflammatory bowel disease and bacterial infection in addition to impairing human alimentation^[2]. Nowadays, growing evidence has emerged to demonstrate how gut microbiota profoundly and systemically influences human health and disease through various mechanisms, particularly influencing cancer. It has been discovered that gut microbiota and their associated metabolites are closely related to malignant tumor generation by developing chronic inflammation and immune surveillance dysregulation.

More interestingly, due to the close anatomical and physiological connection between the gut and liver, the role of gut microbiota and its associated metabolites were found in liver diseases from many studies (discussed in the following sections). By focusing our attention on the interactions between gut microbiota and liver diseases, we are pulling back the curtain on how gut microbiota and its associated metabolites influence liver health and disease. However, it seems as though these findings are but the tip of the iceberg. Many questions still remain. Some cause-specific and disease severity-specific microbiota in liver diseases have been reported, but the underlying mechanisms are still unclear or only show a piece of the whole picture. For example, there are distinct explanations for antibiotics cocktail (ABX) mediated suppression of hepatocellular carcinoma (HCC) tumor progression. Therefore, it is not surprising that many different independent studies targeted the same question to attain different answers, as gut-mediated effect on the liver is involved by a wide range of factors. This further confirms that interactions amongst the gut microbiota and liver work through multiple pathways. These interactions may have developed over a long period as a result of evolution. A systemic review of recent research findings in the gut-liver axis helps to dig the underlying mechanism of how the gut modulates liver function and disease treatment.

Here, we discuss the current findings in correlations between gut microbiota and categorized liver diseases. The perspectives of underlying mechanisms will be reviewed and summarized. Development in gut-microbiota-manipulation-based preventive and therapeutic strategies in liver diseases also will be considered. Given all the current progress, gaps in understanding the influence of gut microbiota on liver diseases are proposed, and the feasible directions are prospectively assumed.

DYSBIOSIS OF GUT MICROBIOTA FACILITATES LIVER DISEASES

Nonalcoholic fatty liver disease and alcoholic liver disease

Nonalcoholic fatty liver disease (NAFLD) and the advanced stage nonalcoholic steatohepatitis (NASH) are currently the most common types of liver disease in humans. Studies have indicated that alteration of gut microbiota is involved in the development of NAFLD and NASH^[3,4]. Endotoxemia induced by increased gut permeability has been observed in patients with NAFLD, suggesting that gut permeability-induced inflammatory pathways contribute to NAFLD pathogenesis^[5]. Small intestinal bacterial overgrowth (SIBO) is another often observed dysbiosis of gut microbiota occurring in NAFLD/NASH^[3,6]. Along with the altered intestinal microbiota profiles, SIBO was seen in most patients with cirrhosis^[7]. The presence of SIBOs may partially explain why the permeability of the gut increases. Gut permeability and SIBOs enhance the hepatic expression of Toll-like receptor 4 (TLR-4) and the production of interleukin 8 (IL-8). Therefore, determining the specific pathogenic bacteria species has been made a very appealing question. Clinical studies revealed that patients with NASH exhibited significantly lower concentrations of *Bacteroidetes* in their gut compared to healthy individuals^[4,8]. Meanwhile, another cohort study showed that high-alcohol-producing *Klebsiella pneumoniae* was associated with up to 60% of NAFLD patients^[9]. Selective depletion of *K. pneumoniae* before fecal microbiota transplant (FMT) from NASH patients into mice prevented NAFLD development in recipient mice. In contrast, the study on patients with alcohol-induced liver cirrhosis exhibits a decreased proportion of *Bacteroidaceae* family than healthy individuals^[10]. Also, the analysis of the gut microbiota profiles in patients with cirrhosis revealed an increase in pathogenic bacteria and a decrease in beneficial bacteria, such as a decreased *Bacteroidetes* along with overgrowth of the *Proteobacteria* and *Enterobacteriaceae* species^[11,12]. However, it remains unclear if this change drives disease or it comes from disease as a result. To answer this question, a well-designed, large-scale clinical and experimental investigation is needed. Moreover, in excessive alcohol intake induced gut microbiota dysbiosis, the intestinal innate immune responses, such as secreted antimicrobial molecules, may impact disease development. For example, the reduced bactericidal c-type lectins, Reg3b, and Reg3g in the small intestines were discovered after alcohol feeding, further causing SIBO and dysbiosis in mice^[13].

HCC

HCC, the final stage of chronic liver disease, has been evidenced to have gut microbiota involved in its initiation and progression. In DEN/CCl₄ induced HCC murine model, germ-free mice presented fewer and smaller tumors compared to its syngeneic parallel^[14], indicating that gut commensal flora is required for tumor progression and the activation of TLR-4 by lipopolysaccharide (LPS) promotes tumor generation. In HCC murine models, studies also demonstrated that the administration of ABX in tumor-bearing mice effectively slowed tumor progression^[15,16]. Even though the underlying mechanisms were distinct in those studies, which will be further discussed in the following sections, we can assume modulating gut microbiota is favorable to prevent liver cancer. However, there is still a lack of clinical data to support it. In addition, obesity-induced gut microbiota dysbiosis that produces DNA damaging bacterial metabolites has been indicated to facilitate HCC development in high-fat diet (HFD)/carcinogen 7,12-dimethylbenz[a]anthracene treated murine model. Furthermore, the administration of antibiotics was able to reduce the prevalence of tumors in the chemically treated mice^[17]. However, in those studies, the relevant specific bacteria species remain unclear. Clearly identifying either specific pathogenic or beneficial bacteria species may be the next mission.

MODULATION OF INTRAHEPATIC IMMUNITY, THE CENTRAL ROLE OF GUT MICROBIOTA IN LIVER DISEASE

Although etiologies vary in liver diseases, inflammation and liver fibrosis are the basis for most liver diseases. The intrahepatic immunity has been certified as an acting point, as gut microbiota can play different roles in liver inflammation and establishment of fibrosis through different liver infiltrating immune cells and resident hepatic cells^[18]. Furthermore, the immune checkpoints emerged as a central pivot to modulate disease progression with gut microbiota-derived metabolites as informatic messengers (Figure 1) and various TLRs as informatic receivers.




Figure 1 The development of liver diseases. Without effective treatment or preventive strategies, fatty liver disease, alcohol-induced liver disease, nonalcoholic fatty liver disease, and nonalcoholic steatohepatitis can result in liver fibrosis, cirrhosis, and hepatocellular carcinoma. Gut microbiota-derived molecules, including lipopolysaccharide, CpG, and secondary bile acids, initially activate liver resident cells to produce cytokines and chemokines. Quiescent hepatic stellate cells (HSCs) can be activated and transformed into myofibroblasts (MFBs), which is mediated by chemokines and cytokines released by liver-infiltrating macrophages, leukocytes, and other cell types, including fat cells and damaged hepatocytes. MFBs are the predominant source of collagen-producing cells and other extracellular matrix proteins (ECM). With effective treatment, such as fecal microbiota transplant, probiotics, and anti-fibrogenic factors, fibrosis is reversible. The treatments that induce apoptosis (or deactivation) of activated HSCs or MFBs and degrade the ECM proteins can reduce the stiffness of the liver, reverse liver fibrosis, and inhibit the progression of liver disease. FL: Fatty liver; ALD: Alcohol-induced liver disease; NAFLD: Nonalcoholic fatty liver disease; NASH: Nonalcoholic steatohepatitis; HCC: Hepatocellular carcinoma; ECM: Extracellular matrix proteins; HFD: High-fat diet; FMT: Fecal microbiota transplant; LPS: Lipopolysaccharide; SBAs: Secondary bile acids; HSC: Hepatic stellate cells.

Hepatic resident and infiltrating cells

Hepatic cells are comprised of parenchymal hepatocytes and the non-parenchymal cells (NPCs), including quiescent hepatic stellate cells (HSCs), Kupffer cells (KCs), liver sinusoidal endothelial cells (LSECs), and diverse lymphocytes (e.g., natural killer cells, NKT cells), which play crucial roles in the liver health and disease^[19,20]. Here, we discuss how gut microbiota modulates intrahepatic immune responses by different hepatic cell types.

Hepatocytes: Hepatocytes consist of two-thirds of the total liver cell population and are responsible for the primary metabolic functions and engage in the immune responses by interacting with NPCs^[21]. In different mouse models of liver fibrosis, either induced with alcohol or non-alcohol (e.g., CCl₄ or HFD), injured hepatocytes release reactive oxygen species and fibrogenic mediators, which recruit lymphocytes and promote the activation of HSCs^[22]. The Notch signaling pathway also plays a pivotal role in liver development, liver repair, and carcinogenesis^[23]. Notch activation in hepatocytes can induce and promote liver fibrosis in normal chow diet and NASH diet-fed mice^[24], respectively. Even though the mechanism by which the gut microbiota directly impacts the Notch signaling in hepatocytes is unclear, some studies have shown that dysbiosis of gut microbiota may increase the activation of the Notch signaling pathway, thus promoting tumor regression^[25]. In addition, the TLR signaling pathway of hepatocytes is directly involved in the gut-liver axis, which will be discussed in the following context.

HSCs: HSCs reside in the space of Disse (perisinusoidal space) between the hepatocytes and LSECs. Their main functions include the storage of vitamin A and lipid droplets^[26]. HSCs are the primary sources of liver myofibroblasts^[27] and play a key role in the initiation, progression, and regression of liver fibrosis^[28]. All other liver cells can directly or indirectly impact the activation, trans-differentiation of HSCs to

myofibroblasts, and their deactivation. There are multiple cellular and molecular signals involved in these reactions^[29]. The interaction of HSCs with other hepatic cells is bidirectional. For instance, upon activation, HSCs, in turn, can further activate macrophages by secreting macrophage colony-stimulating factor (M-CSF), MCP-1, IL-6, RANTES, and so on^[30].

KCs: Liver macrophages, including the liver resident KCs and circulating monocyte-derived macrophages, play essential roles in various liver diseases, including virus hepatitis, alcoholic liver disease (ALD), NAFLD, NASH, and HCC^[31,32]. The purpose of macrophages in NAFLD and NASH and the underlying mechanisms involved in liver inflammation and liver fibrosis have been well summarized recently^[33]. Gut-derived endotoxins (LPS), lipid metabolites, and hepatocyte damage-associated molecules are the main factors inducing macrophage activation in NAFLD^[33]. Furthermore, the activated macrophages secrete chemokines (*e.g.*, CCL2) and cytokines (*e.g.*, TGF- β 1), which leads to the activation of T cells and the transformation of HSCs into myofibroblasts.

LSECs: With the loss of fenestration, capillarized LSECs reduce the transfer of nutrients and other products from the blood to hepatic cells^[34,35], which occurs with liver fibrosis in animals and humans^[36]. The capillarized LSECs can secrete TGF- β 1 and contribute to the accumulation of extracellular matrix proteins, including fibronectin and laminin in the liver, promoting the activation of HSCs and contributing to the formation of liver fibrosis^[35]. Xie *et al*^[37] reported that the reversal of LSEC differentiation with the use of BYY 60-2770, an activator of soluble guanylate cyclase (sGC), prevented the progress of rat liver cirrhosis. LSECs are directly exposed to dietary and bacterial products from the gut through the portal circulation. It has been shown that there is a link between the fenestration of LSECs and diet-induced changes in the gut microbiome. The increase of fenestration of LSECs was associated with a high abundance of *Firmicutes* and a low amount of *Bacteroidetes*^[38].

NKT cells: *In vivo* animal studies showed that alteration of gut bacteria with ABX treatment induced a liver-selective anti-tumor effect with an increase of hepatic CXCR6⁺NKT cells and heightened IFN- γ production upon antigen stimulation^[15]. NKT cell accumulation was regulated by CXCL16 expression of LSECs, which is controlled by gut microbiome-mediated primary-to-secondary bile acid (SBAs) conversion^[15]. Similarly, Zhang *et al*^[39] reported a significant increase of primary bile acids (PBAs) in the large intestine of mice treated with antibiotics such as vancomycin and imipenem. This antibiotic treatment was also associated with the selective suppression of several bacterial species such as *Clostridium*, responsible for deconjugating PBAs into SBAs. Selective depletion of intestinal bacteria *Firmicutes* increases PBA levels, which further promotes hepatic NKT cell accumulation and NKT-mediated anti-tumor effect, indicating that modulating gut microbiota is an optional strategy for HCC treatment.

Immune checkpoints

The efficacy of anti-immune checkpoints (PD-1 or CTLA4) in cancer treatment was markedly reduced in germ-free mice or specific-pathogen-free mice treated with broad-spectrum antibiotics^[40]. For instance, the efficacy of anti-CTLA4 on melanoma immunotherapy in mice and patients was associated with *Bacteroides fragilis*, since gavage with *B. fragilis* or adoptive transfer of *B. fragilis*-specific T cells recovered the anti-tumor efficacy of CTLA4 blockade in antibiotic-treated or GM mice^[41]. Similarly, the abundance of *Akkermansia muciniphila* was associated with the effectiveness of anti-PD-1 immunotherapy^[42]. FMT from cancer patients who were not responding to anti-PD-1 treatment to tumor-bearing mice failed to ameliorate the anti-tumor effects of anti-PD-1. In contrast, oral supplementation of *A. muciniphila* restored the anti-tumor efficacy of anti-PD-1 in those mice^[42]. Meanwhile, Gopalakrishnan *et al*^[43] also reported the gut microbiota modulated therapeutic responses to anti-PD-1 immunotherapy in human melanoma patients. Iida *et al*^[44] concluded that antibiotic-induced disruption of gut microbiota impaired both CpG oligonucleotide immunotherapy and platinum chemotherapy in their subcutaneous breast cancer murine model. Investigation on anti-immune checkpoints on HCC has recently begun^[45,46]; however, the role of gut microbial effects on the blockage of immune checkpoints in liver diseases remains unclear. Notably, immune checkpoint inhibitors-associated hepatotoxicity should not be ignored^[47,48]. For example, clinical trials have shown that alanine aminotransferase elevations in grade 3-4 levels are observed in approximately 20% of patients who are treated with ipilimumab (CTLA-4 inhibitor) and nivolumab (PD-1/PD-L1 inhibitor)^[49].

Metabolites

Currently, the role of microbiota in liver diseases has been broadly investigated^[50]. Gut microbiota-associated metabolites such as fatty acids, amino acids, and carbohydrates are a group of contributory factors involved in the modulation of intrahepatic immune responses^[51]. Ma *et al*^[15] observed that manipulation of gut microbiota by treatment with an ABX can suppress liver tumor growth *via* accumulated NKT cells, which increased anti-tumor immunity in the liver. Modulating commensal gut microbiota composition with ABX treatment changed the bile acid metabolism and significantly increased the PBA production in the liver microenvironment. This rise in PBA production elevated CXCL16 expression on LSECs, which recruited more CXCR6 specific NKT cells to the liver. Conversely, SBAs have been shown to reverse this phenomenon and reduce CXCL16 expression on LSECs. As such, alteration of the ratio of PBAs to SBAs resulted in the growth of hepatic NKT cell accumulation in the ABX treated mice compared to control mice. In addition, the conversion of PBAs to SBAs can be significantly inhibited with the treatment of antibiotics. SBAs, such as deoxycholic acid, activated mTOR to promote the HCC development in NASH mice fed with a steatohepatitis-inducing high-fat diet (STHD-01)^[52]. Accumulation of free fatty acids in hepatocytes can also induce lipotoxicity, which causes hepatocellular injury and NAFLD progression^[53]. Similarly, excess carbohydrates from dietary intake can be *de novo* synthesized to lipids in the liver, causing lipotoxic liver injury^[54]. The characteristics of metabolites of gut bacteria with the association of liver diseases found in recent studies are summarized in **Table 1**. The mechanisms of how microbiota-derived metabolites might influence liver diseases or their potential roles as diagnostic markers have been summarized when we were drafting this review^[55-61].

TLRs

The TLR family is one of the best-characterized families of pattern recognition receptors^[62], which can activate the innate immune system by recognizing pathogen-associated molecular patterns and damage-associated molecular patterns^[63]. There are a total of 13 TLRs in mammals; only TLR1 to 10 exist in humans^[64]. TLRs, excluding TLR10, can bind with different bacteria-derived molecules to participate in the inflammatory and immune responses^[65]. The gut-liver axis defines the close anatomic connection, through the circulation of the portal vein and biliary tract, and the functional interaction of the gastrointestinal tract and the liver^[66]. Through the portal vein, bacteria-derived products, including LPS and CpG-containing DNA, might regularly activate TLR signaling pathway in the hepatocytes and other hepatic cells. Current studies have demonstrated that TLR signaling pathways play pivotal roles in liver inflammation, fibrosis, regeneration, and carcinogenesis^[67,68]. TLRs are broadly expressed in hepatic cell populations, including hepatocytes, LSECs, KCs, lymphocytes, DCs, biliary epithelial cells, and HSCs^[68], which link the gut-liver axis. In this review, we summarize the recent findings of the role of TLRs in liver diseases.

TLR1, TLR6, and TLR10: Currently, TLR10 is the only member of the human TLR family without a definite known ligand, function, and localization^[69]. Phylogenetic analysis shows that TLR10 is most related to TLR1 and TLR6, both of which mediate immune responses in cooperation with TLR2^[70]. The study of chimeric receptors of TLR10 indicated that it can sense triacylated-lipopeptides and other microbial-derived agonists, the ligands of TLR1^[70]. We will not describe the separate roles of these three TLRs at this point.

TLR2: Liver injury resulting from adenovirus or CCl₄ promoted more robust liver inflammation by releasing the TLR2/TLR4 ligands, including HSP60, gp96, HMGB1^[71]. The amplified non-autoimmune hepatitis (AIH)-inflammation could result in the initiation of AIH, a severe autoimmune liver disease, which can lead to fibrosis, cirrhosis, and HCC. Conversely, TLR2 deficiency promoted HCC development in C57BL/6 mice associated with the increase of Ly6C^{high}IL18R α ⁺ myeloid-derived suppressor cells and with impaired CD8⁺ T cells function^[72]. This result further confirmed the previous finding that TLR2 knockdown by siRNAs inhibited the proliferation of HCC cell line BLE-7402, which was associated with a reduction of IL-6 and IL-8 production^[73].

TLR3: TLR3 activation by agonist double-stranded RNA BM-06 or poly(I:C) induced apoptosis of HepG2.2.15 HCC cells^[74]. The analysis of human HCC tissue also indicated that the expression of TLR3 positively correlated with apoptosis of HCC cells and negatively correlated with the proliferation of HCC cells and angiogenesis^[75], suggesting that TLR3 expression may serve as a prognostic marker of HCC.

Table 1 Summary of metabolites associated with liver diseases

Liver disease	Sample sources	Metabolites	Gut microbiota	Functions in liver disease	Ref.
NAFLD	Serum	3-(4-hydroxyphenyl) lactate	Abundance in bacteria <i>Firmicutes</i> , <i>Bacteroidetes</i> , <i>Proteobacteria</i>	It is positively associated with liver fibrosis	[56]
NAFLD	Serum	Eight lipids (5 α -androstan-3 β monosulfate, pregnanediol-3-glucuronide, androsterone sulfate, epiandrosterone sulfate, palmitoleate, dehydroisoandrosterone sulfate, 5 α -androstan-3 β disulfate, glycocholate), one amino acid (taurine) and one carbohydrate (fucose)	Without special bacterial species	Glycocholate is positively associated with advanced liver fibrosis	[57]
FL	Stool	Tryptamine and I3A	Microbiota-dependent without exact bacterial species	They inhibited the pro-inflammatory cytokines in macrophages and hepatocytes. I3A attenuated inflammatory responses under lipid loading and reduced the expression of fatty acid synthase and sterol regulatory element-binding protein-1c	[58]
FL	Stool	Gly-MCA	Increases in the ratio of <i>Bacteroidetes</i> / <i>Firmicutes</i>	Gly-MCA is an intestinal FXR antagonist, which inhibits HFD-induced fatty liver	[59]
HCC	Stool	Secondary bile acids, such as deoxycholic acid	Increase of <i>Bacteroides</i> and <i>Clostridium</i> cluster XVIII. Low of <i>Streptococcus</i> , <i>Bifidobacterium</i> , and <i>Prevotella</i>	Bile acids derived from the increased microbiota promote the progression of HCC	[52]
HCC	Stool	Bile acids, such as primary bile acid CDCA	Increase in <i>Clostridium</i> species	Removing gram-positive bacteria by antibiotic treatment with vancomycin, which contains the bacteria mediating primary-to-secondary bile acid conversion, was sufficient to induce hepatic NKT cell accumulation and decrease liver tumor growth	[15]
Liver cirrhosis	Stool	Glutamic acid, fumaric acid, 4-aminobutyric acid, succinic acid, isoleucine, valine, lactic acid, mannitol, sorbitol, carbamide, 4-aminobutyric acid, 5-aminopyramine, glutamate, proline, hydroxyproline	High of <i>Enterobacteriaceae</i> and <i>Veillonella</i> . Low of <i>Bacteroidetes</i>	These metabolites are involved in the KEGG pathway in nitrogen metabolism alanine, aspartate, and glutamate metabolism, and valine, leucine, and isoleucine, pantothenate and CoA biosynthesis, glycolysis/ gluconeogenesis, fructose and mannose metabolism, arginine and proline metabolism	[60]
NAFL or NASH	Stool	1-pentanol and 2-butanone, and 4-methyl-2-pentanone	Abundance in <i>Oscillospira</i> , <i>Dorea</i> , and <i>Ruminococcus</i>	The results indicated that significantly lower levels of <i>Oscillospira</i> and higher levels of 1-pentanol and 2-butanone in NAFL patients compared to healthy ones. In NASH, lower levels of <i>Oscillospira</i> were associated with a higher abundance of <i>Dorea</i> and <i>Ruminococcus</i> and higher levels of 2-butanone and 4-methyl-2-pentanone compared to CTRLs	[4]
ALD	Stool, urine	<i>e.g.</i> , SCFAs butyrate and propionate	The main harmful bacterial species included altered <i>Bacteroides</i> phylum as well as <i>Bifidophila</i> , <i>Alistipes</i> , <i>Butyricimonas</i> , <i>Clostridium</i> , <i>Proteus</i> , and <i>Escherichia coli</i> . <i>Faecalibacterium</i>	Metabolites are affected by chronic ethanol feeding or consumption, including amino acids, steroids and their derivatives, fatty acids and conjugates	[61]

NAFLD: Nonalcoholic fatty liver disease; FL: Fatty liver; I3A: Indole-3-acetate; Gly-MCA: Glycine- β -muricholic acid; HFD: High-fat diet; HCC: Hepatocellular carcinoma; CDCA: Chenodeoxycholic acid; NAFL: Nonalcoholic fatty liver; NASH: Nonalcoholic steatohepatitis; ALD: Alcohol-induced liver disease; SCFAs: Short-chain fatty acids.

TLR4: TLR4 is one of the most well-known TLRs implicated in the liver inflammation-fibrosis-cancer axis. TLR4 Ligands, including LPS, can induce liver fibrosis by promoting the transformation of HSCs to myofibroblasts through the TLR4-MyD88-

NF- κ B signaling pathway^[76]. Dapito *et al*^[14] reported that TLR4 was required for HCC progression but not the initiation, which is relative not limited to the increase of hepatomitogen epiregulin. Suppressing the TLR4 signaling pathway in HSCs by Dectin-1, the major progenitors of myofibroblasts, inhibited hepatic fibrosis and hepatocarcinogenesis^[77]. Moreover, blockage of TLR4 also potentially inhibited the LPS induced inflammatory reaction and other immune responses^[78].

TLR5: TLR5 knockout can attenuate CCl₄-induced liver fibrosis in wild-type C57BL/6 mice and the activation of HSCs *via* inhibiting NF- κ B and MAPK signaling pathways^[79]. Through the same signaling pathway, TLR5 knockdown also can ameliorate hyperammonemia (HA)-induced liver injury in rats by inhibiting hepatocyte apoptosis, inflammation, and oxidative stress^[80].

TLR7: TLR7 activation by its natural ligand imiquimod can induce autophagy and release of insulin-like growth factor (IGF-1) and inhibit lipid accumulation in hepatocytes induced by unsaturated fatty acid (arachidonic acid: oleic acid = 1:1) *in vitro*^[81]. More so, *in vivo* experiments also showed that TLR7 knockout prevented NAFLD progression *via* inducing autophagy and the release of IGF-1 from the liver.

TLR8: Even though not many studies about the role of TLR8 are reported currently, its role in liver disease should not be ignored. Stimulation of liver intrasinusoidal cells with TLR agonist 1/2, 2, 2/6, 3, 4, 5, 7, 8, or 9 (respectively Pam3CSK4, HKLM, FSL-1, poly(I:C), LPS, flagellin, imiquimod, ssRNA40, or CpG ODN2216) indicated that only the TLR8 agonist ssRNA40 selectively can activate the innate immune cells to produce IFN- γ ^[82].

TLR9: Tak1 Δ Hep mice with hepatic deletion of transforming growth factor- β -activated kinase 1 (Tak1 Δ Hep) can develop a spontaneous liver injury, inflammation, fibrosis, and HCC, mimicking the progression of human HCC^[83]. Ablation of TLR9 or TLR4 suppressed the spontaneous process liver inflammation-fibrosis-cancer axis in Tak1 Δ Hep mice. The inhibition of HCC progression was associated with downstream signaling molecules MyD88 and TNFR.

MANIPULATION OF GUT MICROBIOTA, A PROMISING THERAPEUTIC STRATEGY IN LIVER DISEASE

We have previously highlighted the strong correlations between liver disease and the intestinal microbiota. Further, the modulation of gut microbiota and its products may prove to be a promising target for liver disease and HCC treatment. For example, the abundance of *Enterococcus faecalis* was associated with liver disease severity and the mortality of patients with alcoholic hepatitis due to cytolysin production^[84]. Treatment with *E. faecalis*-targeted bacteriophages can ameliorate alcohol-induced liver disease in humanized mice by reducing liver cytolysin. The intestinal microbiome contains seven-fold more genetic material than the human genome, providing an abundance of potential therapeutic targets. Here, we focus on current studies involving probiotics, FMT, and antibiotics in liver disease treatment.

Probiotics, a new use of an old remedy against liver diseases

Probiotics, prebiotics, and synbiotics have recently undergone investigation for their ability to influence the gut microbiota composition^[2]. Research has long established that probiotic yogurt can positively impact gut health^[85], and more so, the prebiotic insulin was shown to reduce hepatic lipogenesis and plasma triglyceride concentrations in human patients^[86]. Further, the use of probiotics to combat the altered gut microbiota profile in liver disease is currently being investigated in clinical trials^[87]. One recent randomized, double-blind, placebo-controlled study investigated the effects of probiotic treatment on intrahepatic fat fractions in 68 patients with NAFLD. They reported that patients exhibited decreased intrahepatic fat fractions after 12 wk of probiotic treatment (from 16.3% \pm 15.0% to 14.1% \pm 7.7%, $P = 0.032$), with an overall mean decrease of 2.61% when compared to baseline measurements^[88]. Although the argument that this statistical significance may not be clinically significant, the authors propose a mechanism by which specific probiotic agents may aid in correcting dysbiosis rather than directly decreasing hepatic fat. Moreover, this may lead others to investigate the use of probiotics for correcting dysbiosis in liver disease.

FMT, a star of tomorrow to control patient's gut microbiota in personalized therapy

FMT is also an old yet underrepresented area of investigation. This technique has gained considerable attention after its effectiveness in the last resort treatment of antibiotic-resistant infections such as *Clostridium difficile*^[89]. The advantage of FMT from healthy persons to patients is the resetting of gut microbiota composition, returning the flora to a "healthy" state^[87]. To investigate the effects of FMT in liver disease, one study used fecal material from patients with ALD to transplant into germ-free mice through oral gavage. The investigators reported increased intestinal permeability and bacterial translocation, resulting in severe liver inflammation in the microbiome repopulated mice^[90]. Similarly, a pilot study containing eight male patients with severe alcoholic hepatitis were treated with nasoduodenal FMT over the course of 7 d. Results revealed the resolution of ascites and hepatic encephalopathy (HE) within one week in most FMT treated patients, along with significantly improved 1-year survival rates (87.5% vs 33.3%; $P = 0.018$) compared to historical controls^[91]. Taken together, these results demonstrate the importance of the intestinal microbiome population in the development and treatment of liver diseases. While FMT provides distinct challenges for its commercial use in the clinical setting, its profound therapeutic effects can no longer be ignored when current first-line therapies lack adequate efficacy.

Antibiotics, perhaps another resolution in fighting liver diseases

Treatment with antibiotics is the most obvious approach for modulating the intestinal bacterial profile and its products. It has been well documented that microbiota depletion with ABX results in a significant reduction of tumor number, size, and fibrosis severity in several HCC murine models^[14,15,17,76]. While ABX treatment is unfit for human studies, antibiotics have undergone investigation to treat liver disease in many clinical trials. The overwhelming majority of current antibiotic studies utilize the antibiotic rifaximin. Further, this section will primarily report current findings related to rifaximin treatment and liver disease; however, ABX and norfloxacin treatments will also be briefly discussed.

Rifaximin: Rifaximin is a highly favorable antibiotic due to its safety profile and broad-spectrum effects against gram-positive and gram-negative aerobic and anaerobic bacteria. Its classification of a non-absorbable antibiotic makes it well suited for targeting the intestinal microbiome rather than treating systemic infections. Unique to most antibiotics, rifaximin works by binding to the beta-subunit of bacterial DNA-dependent RNA polymerase, consequently blocking bacterial transcription^[92]. Further, this results in endotoxin-lowering and anti-inflammatory effects rather than diminishing the gut flora composition^[93-95]. Multiple studies have correlated rifaximin treatment with reduced risk of death and cirrhotic complications in patients with HE^[96-98]. More so, a phase 3, open-label maintenance study revealed long-term treatment (≥ 24 mo) of rifaximin (550 mg, twice daily) showed no increased adverse events in HE patients compared to placebo controls, further confirming its safety of use^[98]. However, patients with cirrhosis lack average BA concentrations, resulting in decreased efficacy of rifaximin. To combat this issue, rifaximin soluble solid dispersion (SSD) was introduced. Unlike the previous generation, rifaximin SSD is water-soluble and no longer dependent on the patient's intestinal BA concentrations. Despite this distinct advantage, few clinical trials have implemented rifaximin SSD into their liver disease studies.

It is widely theorized that liver cancer progression is partly mediated by gut dysbiosis and other downstream effects. Further, we believe a longitudinal study should be performed to assess the preventative effects of rifaximin and rifaximin SSD on liver cancer development in a high-risk patient population. While it is highly unlikely that rifaximin would provide a therapeutic effect in patients who have already developed tumors, the benefits of long-term, preventative rifaximin treatment on patients with a high risk of developing HCC may provide a better response than the current treatment options.

ABX: Besides enhancing anti-tumor immunity in the liver, antibiotics may also weaken the tumor-promoting impact of specific intestinal bacteria. One study investigated how antibiotic alteration of the gut microbiota impacted liver carcinogenesis by depleting the intestinal flora of mice through regular treatment of ABX. Further, results indicated a significant reduction in tumor number, size, and liver-body weight ratio compared to the untreated control mice^[14]. More so, the authors observed that antibiotics mitigated tumorigenesis in a healthy liver but did not impact tumors that had already developed. These results suggest that the gut microbiota plays a role in preventing hepatic proliferation and fibrogenesis, leading to liver carcinogenesis. And

also, there was a decreased expression of NF- κ B regulated genes and increased apoptosis rates in gut-sterilized mice^[14], indicating that the gut flora and its products may promote liver inflammation and carcinogenesis *via* activating hepatic TLR-4, further exerting pro-survival signals on hepatocytes.

Norfloxacin: Norfloxacin antibiotics may also attenuate pro-inflammatory phenomena exerted by the immune system. Zapater *et al*^[99] observed that selective intestinal decontamination, facilitated by the antibiotic norfloxacin, in patients with cirrhosis reduced serum and ascitic concentrations of pro-inflammatory cytokines TNF- α , IFN- γ , and IL-12 by blunting the activation of serum NF- κ B. Furthermore, norfloxacin treated patients exhibited both decreased levels of neutrophilic oxidative burst and apoptotic events. The authors did not describe a precise mechanism by which the fluoroquinolone mitigated immunological processes, though they did propose that norfloxacin exerted a direct effect on cellular function.

CONCLUSION

Here we have described many reported mechanisms that the gut-liver axis plays in liver disease. Further, we aim to spread awareness of the existing knowledge gap that could be used to modulate the gut-liver axis to alter the disease state. The modulation of the intestinal microbiome has transitioned to clinical trials for its use in both preventative and therapeutic approaches in human liver disease. A summary of the current ongoing clinical trials can be found in [Table 2](#). It should be recognized that alteration of the gut microbiota is not a one size fits all solution due to its highly individual nature and convoluted intricacies. However, the bilateral relationship of the gut-liver axis cannot be ignored when its alteration possesses abundant therapeutic potential.

To aid in furthering the use of gut microbiota-based therapy against liver diseases, we will give brief suggestions for future directions in this field. Lacking from clinical data are well-designed, large-scale investigations of the gut flora profile for liver disease progression (*e.g.*, NAFLD, NASH, HCC, *etc.*). Understanding the underlying mechanisms of how the gut microbiota or its metabolic products alter liver immunity over time could help clarify liver disease development. Such profiles could provide a model to test preventative treatments or therapeutic approaches with antibiotics, probiotics, and FMT. Establishing which bacteria to eliminate and which to nurture would allow for tailored probiotics and specific antibiotics for chronic liver disease therapy. Continuing with this strategy comes the need for longitudinal studies to assess the preventative and therapeutic effects of antibiotics (*e.g.*, rifaximin and rifaximin SSD) on liver disease development. As stated previously, rifaximin SSD would be an ideal drug to study its preventative measures against high-risk liver cancer populations. Pre-, pro-, and synbiotic strategies are underwhelming in clinical studies due to a lack of high-quality study design. Our review highlighted how a decreased level of *Bacteroidetes* is seen in patients with NASH, cirrhosis, and other liver diseases. Finding commonalities in the abovementioned gut profiles (*e.g.*, decreased *Bacteroidetes*) will prompt the use of gut microbiome repopulation therapies in clinical trials. Lastly, the use of FMT treatment desperately needs a more substantial safety investigation. Conducting a proper prospective cohort study with a large study size and a longer observation period is crucial for its safe implementation in the clinical setting.

Collectively, modulating the population of the gut microbiome by biological products may be a new generation therapeutic strategy for liver disease. For now, we must wait to see the results of ongoing and scheduled clinical trials. As we wait in anticipation for the results of mentioned clinical trials, we hope that our current knowledge of the gut-liver-axis inspires new ideas for the next potential therapies in liver disease.

Table 2 Gut microbiota-based clinical trials

Study	Mechanism	Trial phase	Patients (n)	ClinicalTrials.gov identifier	Estimated study completion date
Nivolumab (Anti-PD1), Tadalafil and Oral Vancomycin in People with Refractory Primary Hepatocellular Carcinoma or Liver Dominant Metastatic Cancer from Colorectal or Pancreatic Cancers	Immunotherapy + Antibiotic	Phase II	27	NCT03785210	2022
Administration of Rifaximin to Improve Liver Regeneration and Outcome Following Major Liver Resection (ARROW)	Antibiotic	Phase II	96	NCT02555293	2020
Efficacy of the Combination of Simvastatin Plus Rifaximin in Patients with Decompensated Cirrhosis to Prevent ACLF Development (2018-001698-25)	Antibiotic	Phase III	240	NCT03780673	2021
Comparative Study of Rifaximin Versus Norfloxacin in the Secondary Prophylaxis of Spontaneous Bacterial Peritonitis (SBP)	Antibiotic	Phase III	100	NCT02120196	2023
Efficacy of Antibiotic Therapy in Severe Alcoholic Hepatitis Treated with Prednisolone (AntibioCor)	Antibiotic	Phase III	280	NCT02281929	2019
Rifaximin Reduces the Complications of Decompensated Cirrhosis: A Randomized Controlled Trial	Antibiotic	Phase IV	200	NCT02190357	2019
Steady-State Pharmacokinetics of Rifaximin 550 mg Tablets in Healthy and Hepatically Impaired Subjects	Antibiotic	Phase IV	18	NCT03818672	2019
Efficacy, Safety, And Pharmacokinetics of Rifaximin In Subjects with Severe Hepatic Impairment and Hepatic Encephalopathy	Antibiotic	Phase IV	100	NCT01846663	2021
Rifaximin Soluble Solid Dispersion (SSD) Tablets Plus Lactulose for the Treatment of Overt Hepatic Encephalopathy (OHE) (OHE)	Antibiotic	Phase II	325	NCT03515044	2019
Mastiha Treatment for Obese with NAFLD Diagnosis (MAST4HEALTH)	Prebiotic	Early Phase I	52	NCT03135873	2019
Study on the Optimal Strategy for Acute-on-chronic Liver Failure with Integrative Treatment	Prebiotic	N/A	510	NCT03577938	2020
Efficacy of Albumin Therapy with Standard Medical Treatment (SMT) as Compared to Standard Medical Treatment (SMT) in Improving Patient Survival and Immune Modulation in Patients with Acute on Chronic Liver Failure (ASIA Trial).	Prebiotic	N/A	200	NCT03754400	2020
Impact of a Specific Micronutrient-probiotic-supplement on Fatty Liver of Patients After Mini-Gastric Bypass Surgery (FMG-01)	Probiotic	Phase III	60	NCT03585413	2019
Probiotics in the Treatment of NAFLD	Probiotic	N/A	58	NCT02764047	2018
Probiotics in NASH Patients - PROBILIVER TRIAL (NASH)	Probiotic	N/A	46	NCT03467282	2021
Profermin®: Prevention of Progression in Alcoholic Liver Disease by Modulating Dysbiotic Microbiota (SYN-ALD)	Probiotic	N/A	40	NCT03863730	2031
Novel Therapies in Moderately Severe Acute Alcoholic Hepatitis (NTAH-Mod)	Probiotic	N/A	130	NCT01922895	2019
Probiotics in the Prevention of Hepatocellular Carcinoma in Cirrhosis	Probiotic	N/A	280	NCT03853928	2023
Dietary Modulation of Intestinal Microbiota as Trigger of Liver Health: Role of Bile Acids - "A Diet for Liver Health" (ADLH)	Synbiotic	N/A	84	NCT03897218	2020
Investigation of Synbiotic Treatment in NAFLD (INSYTE)	Synbiotic	N/A	100	NCT01680640	2019
Fecal Microbiota Transplantation (FMT) in Nonalcoholic Steatohepatitis (NASH). A Pilot Study	FMT	Phase I	5	NCT02469272	2018
Fecal Microbiota Transplantation for the Treatment of Non-Alcoholic Steatohepatitis (FMT-NASH)	FMT	Phase I	15	NCT03803540	2021
Fecal Microbial Transplant for Alcohol Misuse in Cirrhosis	FMT	Phase I	20	NCT03416751	2019
FMT in Cirrhosis and Hepatic Encephalopathy	FMT	Phase II	100	NCT03796598	2023
Fecal Microbiota Transplant as Treatment of Hepatic Encephalopathy	FMT	Phase II	30	NCT03420482	2021
Fecal Microbiota Transplantation (FMT) in the Management of	FMT	Phase II	10	NCT02255617	2019

Hepatic Encephalopathy (HE): A Pilot Study					
Fecal Transplant for Hepatic Encephalopathy	FMT	Phase II	30	NCT03439982	2021
Trial of Faecal Microbiota Transplantation in Cirrhosis (PROFIT)	FMT	Phase III	32	NCT02862249	2019
To Assess the Role of Fecal Microbiota Transplant in Acute Liver Failure	FMT	N/A	40	NCT03363022	2018

REFERENCES

- Ley RE, Peterson DA, Gordon JI. Ecological and evolutionary forces shaping microbial diversity in the human intestine. *Cell* 2006; **124**: 837-848 [PMID: 16497592 DOI: 10.1016/j.cell.2006.02.017]
- Azad MAK, Sarker M, Li T, Yin J. Probiotic Species in the Modulation of Gut Microbiota: An Overview. *Biomed Res Int* 2018; **2018**: 9478630 [PMID: 29854813 DOI: 10.1155/2018/9478630]
- Fialho A, Fialho A, Thota P, McCullough AJ, Shen B. Small Intestinal Bacterial Overgrowth Is Associated with Non-Alcoholic Fatty Liver Disease. *J Gastrointest Liver Dis* 2016; **25**: 159-165 [PMID: 27308646 DOI: 10.15403/jgld.2014.1121.252.iwg]
- Del Chierico F, Nobili V, Vernocchi P, Russo A, De Stefanis C, Gnani D, Furlanello C, Zandonà A, Paci P, Capuani G, Dallapiccola B, Miccheli A, Alisi A, Putignani L. Gut microbiota profiling of pediatric nonalcoholic fatty liver disease and obese patients unveiled by an integrated meta-omics-based approach. *Hepatology* 2017; **65**: 451-464 [PMID: 27028797 DOI: 10.1002/hep.28572]
- Chu H, Williams B, Schnabl B. Gut microbiota, fatty liver disease, and hepatocellular carcinoma. *Liver Res* 2018; **2**: 43-51 [PMID: 30416839 DOI: 10.1016/j.livres.2017.11.005]
- Miele L, Valenza V, La Torre G, Montalto M, Cammarota G, Ricci R, Mascianà R, Forgione A, Gabrieli ML, Perotti G, Vecchio FM, Rapaccini G, Gasbarrini G, Day CP, Grieco A. Increased intestinal permeability and tight junction alterations in nonalcoholic fatty liver disease. *Hepatology* 2009; **49**: 1877-1887 [PMID: 19291785 DOI: 10.1002/hep.22848]
- Schnabl B, Brenner DA. Interactions between the intestinal microbiome and liver diseases. *Gastroenterology* 2014; **146**: 1513-1524 [PMID: 24440671 DOI: 10.1053/j.gastro.2014.01.020]
- Mouzaki M, Comelli EM, Arendt BM, Bonengel J, Fung SK, Fischer SE, McGilvray ID, Allard JP. Intestinal microbiota in patients with nonalcoholic fatty liver disease. *Hepatology* 2013; **58**: 120-127 [PMID: 23401313 DOI: 10.1002/hep.26319]
- Yuan J, Chen C, Cui J, Lu J, Yan C, Wei X, Zhao X, Li N, Li S, Xue G, Cheng W, Li B, Li H, Lin W, Tian C, Zhao J, Han J, An D, Zhang Q, Wei H, Zheng M, Ma X, Li W, Chen X, Zhang Z, Zeng H, Ying S, Wu J, Yang R, Liu D. Fatty Liver Disease Caused by High-Alcohol-Producing *Klebsiella pneumoniae*. *Cell Metab* 2019; **30**: 675-688. e7 [PMID: 31543403 DOI: 10.1016/j.cmet.2019.08.018]
- Mutlu EA, Gillevet PM, Rangwala H, Sikaroodi M, Naqvi A, Engen PA, Kwasny M, Lau CK, Keshavarzian A. Colonic microbiome is altered in alcoholism. *Am J Physiol Gastrointest Liver Physiol* 2012; **302**: G966-G978 [PMID: 22241860 DOI: 10.1152/ajpgi.00380.2011]
- Giannelli V, Di Gregorio V, Iebba V, Giusto M, Schippa S, Merli M, Thalheimer U. Microbiota and the gut-liver axis: bacterial translocation, inflammation and infection in cirrhosis. *World J Gastroenterol* 2014; **20**: 16795-16810 [PMID: 25492994 DOI: 10.3748/wjg.v20.i45.16795]
- Oikonomou T, Papatheodoridis GV, Samarkos M, Goulis I, Cholongitas E. Clinical impact of microbiome in patients with decompensated cirrhosis. *World J Gastroenterol* 2018; **24**: 3813-3820 [PMID: 30228776 DOI: 10.3748/wjg.v24.i34.3813]
- Yan AW, Fouts DE, Brandl J, Stärkel P, Torralba M, Schott E, Tsukamoto H, Nelson KE, Brenner DA, Schnabl B. Enteric dysbiosis associated with a mouse model of alcoholic liver disease. *Hepatology* 2011; **53**: 96-105 [PMID: 21254165 DOI: 10.1002/hep.24018]
- Dapito DH, Mencin A, Gwak GY, Pradere JP, Jang MK, Mederacke I, Caviglia JM, Khiabanian H, Adeyemi A, Bataller R, Lefkowitz JH, Bower M, Friedman R, Sartor RB, Rabadan R, Schwabe RF. Promotion of hepatocellular carcinoma by the intestinal microbiota and TLR4. *Cancer Cell* 2012; **21**: 504-516 [PMID: 22516259 DOI: 10.1016/j.ccr.2012.02.007]
- Ma C, Han M, Heinrich B, Fu Q, Zhang Q, Sandhu M, Agdashian D, Terabe M, Berzofsky JA, Fako V, Ritz T, Longerich T, Theriot CM, McCulloch JA, Roy S, Yuan W, Thovarai V, Sen SK, Ruchirawat M, Korangy F, Wang XW, Trinchieri G, Greten TF. Gut microbiome-mediated bile acid metabolism regulates liver cancer via NKT cells. *Science* 2018; **360** [PMID: 29798856 DOI: 10.1126/science.aan5931]
- Sethi V, Kurtom S, Tarique M, Lavania S, Malchiodi Z, Hellmund L, Zhang L, Sharma U, Giri B, Garg B, Ferrantella A, Vickers SM, Banerjee S, Dawra R, Roy S, Ramakrishnan S, Saluja A, Dudeja V. Gut Microbiota Promotes Tumor Growth in Mice by Modulating Immune Response. *Gastroenterology* 2018; **155**: 33-37. e6 [PMID: 29630898 DOI: 10.1053/j.gastro.2018.04.001]
- Yoshimoto S, Loo TM, Atarashi K, Kanda H, Sato S, Oyadomari S, Iwakura Y, Oshima K, Morita H, Hattori M, Honda K, Ishikawa Y, Hara E, Ohtani N. Obesity-induced gut microbial metabolite promotes liver cancer through senescence secretome. *Nature* 2013; **499**: 97-101 [PMID: 23803760 DOI: 10.1038/nature12347]
- Koyama Y, Brenner DA. Liver inflammation and fibrosis. *J Clin Invest* 2017; **127**: 55-64 [PMID:

- 28045404 DOI: [10.1172/JCI88881](https://doi.org/10.1172/JCI88881)]
- 19 **Mohar I**, Brempelis KJ, Murray SA, Ebrahimkhani MR, Crispe IN. Isolation of Non-parenchymal Cells from the Mouse Liver. *Methods Mol Biol* 2015; **1325**: 3-17 [PMID: [26450375](https://pubmed.ncbi.nlm.nih.gov/26450375/) DOI: [10.1007/978-1-4939-2815-6_1](https://doi.org/10.1007/978-1-4939-2815-6_1)]
 - 20 **Seo W**, Jeong WI. Hepatic non-parenchymal cells: Master regulators of alcoholic liver disease? *World J Gastroenterol* 2016; **22**: 1348-1356 [PMID: [26819504](https://pubmed.ncbi.nlm.nih.gov/26819504/) DOI: [10.3748/wjg.v22.i4.1348](https://doi.org/10.3748/wjg.v22.i4.1348)]
 - 21 **Li F**, Tian Z. The liver works as a school to educate regulatory immune cells. *Cell Mol Immunol* 2013; **10**: 292-302 [PMID: [23604044](https://pubmed.ncbi.nlm.nih.gov/23604044/) DOI: [10.1038/cmi.2013.7](https://doi.org/10.1038/cmi.2013.7)]
 - 22 **Hayashi H**, Sakai T. Animal models for the study of liver fibrosis: new insights from knockout mouse models. *Am J Physiol Gastrointest Liver Physiol* 2011; **300**: G729-G738 [PMID: [21350186](https://pubmed.ncbi.nlm.nih.gov/21350186/) DOI: [10.1152/ajpgi.00013.2011](https://doi.org/10.1152/ajpgi.00013.2011)]
 - 23 **Geisler F**, Strazzabosco M. Emerging roles of Notch signaling in liver disease. *Hepatology* 2015; **61**: 382-392 [PMID: [24930574](https://pubmed.ncbi.nlm.nih.gov/24930574/) DOI: [10.1002/hep.27268](https://doi.org/10.1002/hep.27268)]
 - 24 **Zhu C**, Kim K, Wang X, Bartolome A, Salomao M, Dongiovanni P, Meroni M, Graham MJ, Yates KP, Diehl AM, Schwabe RF, Tabas I, Valenti L, Lavine JE, Pajvani UB. Hepatocyte Notch activation induces liver fibrosis in nonalcoholic steatohepatitis. *Sci Transl Med* 2018; **10** [PMID: [30463916](https://pubmed.ncbi.nlm.nih.gov/30463916/) DOI: [10.1126/scitranslmed.aat0344](https://doi.org/10.1126/scitranslmed.aat0344)]
 - 25 **Troll JV**, Hamilton MK, Abel ML, Ganz J, Bates JM, Stephens WZ, Melancon E, van der Vaart M, Meijer AH, Distel M, Eisen JS, Guillemin K. Microbiota promote secretory cell determination in the intestinal epithelium by modulating host Notch signaling. *Development* 2018; **145** [PMID: [29475973](https://pubmed.ncbi.nlm.nih.gov/29475973/) DOI: [10.1242/dev.155317](https://doi.org/10.1242/dev.155317)]
 - 26 **Senoo H**, Kojima N, Sato M. Vitamin A-storing cells (stellate cells). *Vitam Horm* 2007; **75**: 131-159 [PMID: [17368315](https://pubmed.ncbi.nlm.nih.gov/17368315/) DOI: [10.1016/s0083-6729\(06\)75006-3](https://doi.org/10.1016/s0083-6729(06)75006-3)]
 - 27 **Iwaisako K**, Brenner DA, Kisseleva T. What's new in liver fibrosis? *J Gastroenterol Hepatol* 2012; **27** Suppl 2: 65-68 [PMID: [22320919](https://pubmed.ncbi.nlm.nih.gov/22320919/) DOI: [10.1111/j.1440-1746.2011.07002.x](https://doi.org/10.1111/j.1440-1746.2011.07002.x)]
 - 28 **Zhang CY**, Yuan WG, He P, Lei JH, Wang CX. Liver fibrosis and hepatic stellate cells: Etiology, pathological hallmarks and therapeutic targets. *World J Gastroenterol* 2016; **22**: 10512-10522 [PMID: [28082803](https://pubmed.ncbi.nlm.nih.gov/28082803/) DOI: [10.3748/wjg.v22.i48.10512](https://doi.org/10.3748/wjg.v22.i48.10512)]
 - 29 **Hou W**, Syn WK. Role of Metabolism in Hepatic Stellate Cell Activation and Fibrogenesis. *Front Cell Dev Biol* 2018; **6**: 150 [PMID: [30483502](https://pubmed.ncbi.nlm.nih.gov/30483502/) DOI: [10.3389/fcell.2018.00150](https://doi.org/10.3389/fcell.2018.00150)]
 - 30 **Lee UE**, Friedman SL. Mechanisms of hepatic fibrogenesis. *Best Pract Res Clin Gastroenterol* 2011; **25**: 195-206 [PMID: [21497738](https://pubmed.ncbi.nlm.nih.gov/21497738/) DOI: [10.1016/j.bpg.2011.02.005](https://doi.org/10.1016/j.bpg.2011.02.005)]
 - 31 **Kolios G**, Valatas V, Kouroumalis E. Role of Kupffer cells in the pathogenesis of liver disease. *World J Gastroenterol* 2006; **12**: 7413-7420 [PMID: [17167827](https://pubmed.ncbi.nlm.nih.gov/17167827/) DOI: [10.3748/wjg.v12.i46.7413](https://doi.org/10.3748/wjg.v12.i46.7413)]
 - 32 **Tacke F**. Targeting hepatic macrophages to treat liver diseases. *J Hepatol* 2017; **66**: 1300-1312 [PMID: [28267621](https://pubmed.ncbi.nlm.nih.gov/28267621/) DOI: [10.1016/j.jhep.2017.02.026](https://doi.org/10.1016/j.jhep.2017.02.026)]
 - 33 **Kazankov K**, Jørgensen SMD, Thomsen KL, Møller HJ, Vilstrup H, George J, Schuppan D, Grønbaek H. The role of macrophages in nonalcoholic fatty liver disease and nonalcoholic steatohepatitis. *Nat Rev Gastroenterol Hepatol* 2019; **16**: 145-159 [PMID: [30482910](https://pubmed.ncbi.nlm.nih.gov/30482910/) DOI: [10.1038/s41575-018-0082-x](https://doi.org/10.1038/s41575-018-0082-x)]
 - 34 **Zapotoczny B**, Szafranska K, Kus E, Braet F, Wisse E, Chlopicki S, Szymonski M. Tracking Fenestrae Dynamics in Live Murine Liver Sinusoidal Endothelial Cells. *Hepatology* 2019; **69**: 876-888 [PMID: [30137644](https://pubmed.ncbi.nlm.nih.gov/30137644/) DOI: [10.1002/hep.30232](https://doi.org/10.1002/hep.30232)]
 - 35 **Hammoutene A**, Rautou PE. Role of liver sinusoidal endothelial cells in non-alcoholic fatty liver disease. *J Hepatol* 2019; **70**: 1278-1291 [PMID: [30797053](https://pubmed.ncbi.nlm.nih.gov/30797053/) DOI: [10.1016/j.jhep.2019.02.012](https://doi.org/10.1016/j.jhep.2019.02.012)]
 - 36 **DeLeve LD**. Liver sinusoidal endothelial cells in hepatic fibrosis. *Hepatology* 2015; **61**: 1740-1746 [PMID: [25131509](https://pubmed.ncbi.nlm.nih.gov/25131509/) DOI: [10.1002/hep.27376](https://doi.org/10.1002/hep.27376)]
 - 37 **Xie G**, Wang X, Wang L, Wang L, Atkinson RD, Kanel GC, Gaarde WA, Deleve LD. Role of differentiation of liver sinusoidal endothelial cells in progression and regression of hepatic fibrosis in rats. *Gastroenterology* 2012; **142**: 918-927. e6 [PMID: [22178212](https://pubmed.ncbi.nlm.nih.gov/22178212/) DOI: [10.1053/j.gastro.2011.12.017](https://doi.org/10.1053/j.gastro.2011.12.017)]
 - 38 **Cogger VC**, Mohamad M, Solon-Biet SM, Senior AM, Warren A, O'Reilly JN, Tung BT, Svistounov D, McMahon AC, Fraser R, Raubenheimer D, Holmes AJ, Simpson SJ, Le Couteur DG. Dietary macronutrients and the aging liver sinusoidal endothelial cell. *Am J Physiol Heart Circ Physiol* 2016; **310**: H1064-H1070 [PMID: [26921440](https://pubmed.ncbi.nlm.nih.gov/26921440/) DOI: [10.1152/ajpheart.00949.2015](https://doi.org/10.1152/ajpheart.00949.2015)]
 - 39 **Zhang Y**, Limaye PB, Renaud HJ, Klaassen CD. Effect of various antibiotics on modulation of intestinal microbiota and bile acid profile in mice. *Toxicol Appl Pharmacol* 2014; **277**: 138-145 [PMID: [24657338](https://pubmed.ncbi.nlm.nih.gov/24657338/) DOI: [10.1016/j.taap.2014.03.009](https://doi.org/10.1016/j.taap.2014.03.009)]
 - 40 **Gopalakrishnan V**, Helmink BA, Spencer CN, Reuben A, Wargo JA. The Influence of the Gut Microbiome on Cancer, Immunity, and Cancer Immunotherapy. *Cancer Cell* 2018; **33**: 570-580 [PMID: [29634945](https://pubmed.ncbi.nlm.nih.gov/29634945/) DOI: [10.1016/j.ccell.2018.03.015](https://doi.org/10.1016/j.ccell.2018.03.015)]
 - 41 **Vétizou M**, Pitt JM, Daillère R, Lepage P, Waldschmitt N, Flament C, Rusakiewicz S, Routy B, Roberti MP, Duong CP, Poirier-Colame V, Roux A, Becharaf S, Formenti S, Golden E, Cording S, Eberl G, Schlitzer A, Ginhoux F, Mani S, Yamazaki T, Jacquelinot N, Enot DP, Bérard M, Nigou J, Opolon P, Eggermont A, Woerther PL, Chachaty E, Chaput N, Robert C, Mateus C, Kroemer G, Raoult D, Boneca IG, Carbonnel F, Chamillard M, Zitvogel L. Anticancer immunotherapy by CTLA-4 blockade relies on the gut microbiota. *Science* 2015; **350**: 1079-1084 [PMID: [26541610](https://pubmed.ncbi.nlm.nih.gov/26541610/) DOI: [10.1126/science.aad1329](https://doi.org/10.1126/science.aad1329)]
 - 42 **Routy B**, Le Chatelier E, Derosa L, Duong CPM, Alou MT, Daillère R, Fluckiger A, Messaoudene

- M, Rauber C, Roberti MP, Fidelle M, Flament C, Poirier-Colame V, Opolon P, Klein C, Iribarren K, Mondragón L, Jacquelot N, Qu B, Ferrere G, Clémenson C, Mezquita L, Masip JR, Naltet C, Brosseau S, Kaderbhai C, Richard C, Rizvi H, Levenez F, Galleron N, Quinquis B, Pons N, Ryffel B, Minard-Colin V, Gonin P, Soria JC, Deutsch E, Loriot Y, Ghiringhelli F, Zalcman G, Goldwasser F, Escudier B, Hellmann MD, Eggermont A, Raoult D, Albiges L, Kroemer G, Zitvogel L. Gut microbiome influences efficacy of PD-1-based immunotherapy against epithelial tumors. *Science* 2018; **359**: 91-97 [PMID: 29097494 DOI: 10.1126/science.aan3706]
- 43 **Gopalakrishnan V**, Spencer CN, Nezi L, Reuben A, Andrews MC, Karpinets TV, Prieto PA, Vicente D, Hoffman K, Wei SC, Cogdill AP, Zhao L, Hudgens CW, Hutchinson DS, Manzo T, Petaccia de Macedo M, Cotechini T, Kumar T, Chen WS, Reddy SM, Szczepaniak Sloane R, Galloway-Pena J, Jiang H, Chen PL, Shpall EJ, Rezvani K, Alousi AM, Chemaly RF, Shelburne S, Vence LM, Okhuysen PC, Jensen VB, Swennes AG, McAllister F, Marcelo Riquelme Sanchez E, Zhang Y, Le Chatelier E, Zitvogel L, Pons N, Austin-Breneman JL, Haydu LE, Burton EM, Gardner JM, Sirmans E, Hu J, Lazar AJ, Tsujikawa T, Diab A, Tawbi H, Glitza IC, Hwu WJ, Patel SP, Woodman SE, Amaria RN, Davies MA, Gershenwald JE, Hwu P, Lee JE, Zhang J, Coussens LM, Cooper ZA, Futreal PA, Daniel CR, Ajami NJ, Petrosino JF, Tetzlaff MT, Sharma P, Allison JP, Jenq RR, Wargo JA. Gut microbiome modulates response to anti-PD-1 immunotherapy in melanoma patients. *Science* 2018; **359**: 97-103 [PMID: 29097493 DOI: 10.1126/science.aan4236]
- 44 **Iida N**, Dzutsev A, Stewart CA, Smith L, Bouladoux N, Weingarten RA, Molina DA, Salcedo R, Back T, Cramer S, Dai RM, Kiu H, Cardone M, Naik S, Patri AK, Wang E, Marincola FM, Frank KM, Belkaid Y, Trinchieri G, Goldszmid RS. Commensal bacteria control cancer response to therapy by modulating the tumor microenvironment. *Science* 2013; **342**: 967-970 [PMID: 24264989 DOI: 10.1126/science.1240527]
- 45 **El-Khoueiry AB**, Sangro B, Yau T, Crocenzi TS, Kudo M, Hsu C, Kim TY, Choo SP, Trojan J, Welling TH Rd, Meyer T, Kang YK, Yeo W, Chopra A, Anderson J, Dela Cruz C, Lang L, Neely J, Tang H, Dastani HB, Melero I. Nivolumab in patients with advanced hepatocellular carcinoma (CheckMate 040): an open-label, non-comparative, phase 1/2 dose escalation and expansion trial. *Lancet* 2017; **389**: 2492-2502 [PMID: 28434648 DOI: 10.1016/S0140-6736(17)31046-2]
- 46 **Kudo M**. Combination Cancer Immunotherapy with Molecular Targeted Agents/Anti-CTLA-4 Antibody for Hepatocellular Carcinoma. *Liver Cancer* 2019; **8**: 1-11 [PMID: 30815391 DOI: 10.1159/000496277]
- 47 **Reddy HG**, Schneider BJ, Tai AW. Immune Checkpoint Inhibitor-Associated Colitis and Hepatitis. *Clin Transl Gastroenterol* 2018; **9**: 180 [PMID: 30228268 DOI: 10.1038/s41424-018-0049-9]
- 48 **Jennings JJ**, Mandaliya R, Nakshabandi A, Lewis JH. Hepatotoxicity induced by immune checkpoint inhibitors: a comprehensive review including current and alternative management strategies. *Expert Opin Drug Metab Toxicol* 2019; **15**: 231-244 [PMID: 30677306 DOI: 10.1080/17425255.2019.1574744]
- 49 **Larkin J**, Hodi FS, Wolchok JD. Combined Nivolumab and Ipilimumab or Monotherapy in Untreated Melanoma. *N Engl J Med* 2015; **373**: 1270-1271 [PMID: 26398076 DOI: 10.1056/NEJMc1509660]
- 50 **Minemura M**, Shimizu Y. Gut microbiota and liver diseases. *World J Gastroenterol* 2015; **21**: 1691-1702 [PMID: 25684933 DOI: 10.3748/wjg.v21.i6.1691]
- 51 **Levy M**, Thaiss CA, Elinav E. Metabolites: messengers between the microbiota and the immune system. *Genes Dev* 2016; **30**: 1589-1597 [PMID: 27474437 DOI: 10.1101/gad.284091.116]
- 52 **Yamada S**, Takashina Y, Watanabe M, Nagamine R, Saito Y, Kamada N, Saito H. Bile acid metabolism regulated by the gut microbiota promotes non-alcoholic steatohepatitis-associated hepatocellular carcinoma in mice. *Oncotarget* 2018; **9**: 9925-9939 [PMID: 29515780 DOI: 10.18632/oncotarget.24066]
- 53 **Li M**, Xu C, Shi J, Ding J, Wan X, Chen D, Gao J, Li C, Zhang J, Lin Y, Tu Z, Kong X, Li Y, Yu C. Fatty acids promote fatty liver disease via the dysregulation of 3-mercaptopyruvate sulfurtransferase/hydrogen sulfide pathway. *Gut* 2018; **67**: 2169-2180 [PMID: 28877979 DOI: 10.1136/gutjnl-2017-313778]
- 54 **Neuschwander-Tetri BA**. Hepatic lipotoxicity and the pathogenesis of nonalcoholic steatohepatitis: the central role of nontriglyceride fatty acid metabolites. *Hepatology* 2010; **52**: 774-788 [PMID: 20683968 DOI: 10.1002/hep.23719]
- 55 **Chu H**, Duan Y, Yang L, Schnabl B. Small metabolites, possible big changes: a microbiota-centered view of non-alcoholic fatty liver disease. *Gut* 2019; **68**: 359-370 [PMID: 30171065 DOI: 10.1136/gutjnl-2018-316307]
- 56 **Caussy C**, Hsu C, Lo MT, Liu A, Bettencourt R, Ajmera VH, Bassirian S, Hooker J, Sy E, Richards L, Schork N, Schnabl B, Brenner DA, Sirlin CB, Chen CH, Loomba R; Genetics of NAFLD in Twins Consortium. Link between gut-microbiome derived metabolite and shared gene-effects with hepatic steatosis and fibrosis in NAFLD. *Hepatology* 2018; **68**: 918-932 [PMID: 29572891 DOI: 10.1002/hep.29892]
- 57 **Caussy C**, Ajmera VH, Puri P, Hsu CL, Bassirian S, Mgdshyan M, Singh S, Faulkner C, Valasek MA, Rizo E, Richards L, Brenner DA, Sirlin CB, Sanyal AJ, Loomba R. Serum metabolites detect the presence of advanced fibrosis in derivation and validation cohorts of patients with non-alcoholic fatty liver disease. *Gut* 2019; **68**: 1884-1892 [PMID: 30567742 DOI: 10.1136/gutjnl-2018-317584]
- 58 **Krishnan S**, Ding Y, Saedi N, Choi M, Sridharan GV, Sherr DH, Yarmush ML, Alaniz RC, Jayaraman A, Lee K. Gut Microbiota-Derived Tryptophan Metabolites Modulate Inflammatory

- Response in Hepatocytes and Macrophages. *Cell Rep* 2018; **23**: 1099-1111 [PMID: 29694888 DOI: 10.1016/j.celrep.2018.03.109]
- 59 **Zhang L**, Xie C, Nichols RG, Chan SH, Jiang C, Hao R, Smith PB, Cai J, Simons MN, Hatzakis E, Maranas CD, Gonzalez FJ, Patterson AD. Farnesoid X Receptor Signaling Shapes the Gut Microbiota and Controls Hepatic Lipid Metabolism. *mSystems* 2016; **1** [PMID: 27822554 DOI: 10.1128/mSystems.00070-16]
- 60 **Wei X**, Zhao J, Jia X, Zhao X, Li H, Lin W, Feng R, Yuan J. Abnormal Gut Microbiota Metabolism Specific for Liver Cirrhosis. *Front Microbiol* 2018; **9**: 3051 [PMID: 30619141 DOI: 10.3389/fmicb.2018.03051]
- 61 **Sarin SK**, Pande A, Schnabl B. Microbiome as a therapeutic target in alcohol-related liver disease. *J Hepatol* 2019; **70**: 260-272 [PMID: 30658727 DOI: 10.1016/j.jhep.2018.10.019]
- 62 **Takeuchi O**, Akira S. Pattern recognition receptors and inflammation. *Cell* 2010; **140**: 805-820 [PMID: 20303872 DOI: 10.1016/j.cell.2010.01.022]
- 63 **Hug H**, Mohajeri MH, La Fata G. Toll-Like Receptors: Regulators of the Immune Response in the Human Gut. *Nutrients* 2018; **10** [PMID: 29438282 DOI: 10.3390/nu10020203]
- 64 **Kiziltas S**. Toll-like receptors in pathophysiology of liver diseases. *World J Hepatol* 2016; **8**: 1354-1369 [PMID: 27917262 DOI: 10.4254/wjh.v8.i32.1354]
- 65 **Lee MS**, Kim YJ. Signaling pathways downstream of pattern-recognition receptors and their cross talk. *Annu Rev Biochem* 2007; **76**: 447-480 [PMID: 17328678 DOI: 10.1146/annurev.biochem.76.060605.122847]
- 66 **Tripathi A**, Debelius J, Brenner DA, Karin M, Loomba R, Schnabl B, Knight R. The gut-liver axis and the intersection with the microbiome. *Nat Rev Gastroenterol Hepatol* 2018; **15**: 397-411 [PMID: 29748586 DOI: 10.1038/s41575-018-0011-z]
- 67 **Seki E**, Park E, Fujimoto J. Toll-like receptor signaling in liver regeneration, fibrosis and carcinogenesis. *Hepatol Res* 2011; **41**: 597-610 [PMID: 21696522 DOI: 10.1111/j.1872-034X.2011.00822.x]
- 68 **Chen Y**, Sun R. Toll-like receptors in acute liver injury and regeneration. *Int Immunopharmacol* 2011; **11**: 1433-1441 [PMID: 21601014 DOI: 10.1016/j.intimp.2011.04.023]
- 69 **Kesar V**, Odin JA. Toll-like receptors and liver disease. *Liver Int* 2014; **34**: 184-196 [PMID: 24118797 DOI: 10.1111/liv.12315]
- 70 **Guan Y**, Ranoa DR, Jiang S, Mutha SK, Li X, Baudry J, Tapping RI. Human TLRs 10 and 1 share common mechanisms of innate immune sensing but not signaling. *J Immunol* 2010; **184**: 5094-5103 [PMID: 20348427 DOI: 10.4049/jimmunol.0901888]
- 71 **Chi G**, Feng XX, Ru YX, Xiong T, Gao Y, Wang H, Luo ZL, Mo R, Guo F, He YP, Zhang GM, Tian DA, Feng ZH. TLR2/4 ligand-amplified liver inflammation promotes initiation of autoimmune hepatitis due to sustained IL-6/IL-12/IL-4/IL-25 expression. *Mol Immunol* 2018; **99**: 171-181 [PMID: 29793131 DOI: 10.1016/j.molimm.2018.05.005]
- 72 **Li S**, Sun R, Chen Y, Wei H, Tian Z. TLR2 limits development of hepatocellular carcinoma by reducing IL18-mediated immunosuppression. *Cancer Res* 2015; **75**: 986-995 [PMID: 25600646 DOI: 10.1158/0008-5472.CAN-14-2371]
- 73 **Huang Y**, Cai B, Xu M, Qiu Z, Tao Y, Zhang Y, Wang J, Xu Y, Zhou Y, Yang J, Han X, Gao Q. Gene silencing of Toll-like receptor 2 inhibits proliferation of human liver cancer cells and secretion of inflammatory cytokines. *PLoS One* 2012; **7**: e38890 [PMID: 22815694 DOI: 10.1371/journal.pone.0038890]
- 74 **Chen L**, Xu YY, Zhou JM, Wu YY, E Q, Zhu YY. TLR3 dsRNA agonist inhibits growth and invasion of HepG2.2.15 HCC cells. *Oncol Rep* 2012; **28**: 200-206 [PMID: 22552584 DOI: 10.3892/or.2012.1791]
- 75 **Yuan MM**, Xu YY, Chen L, Li XY, Qin J, Shen Y. TLR3 expression correlates with apoptosis, proliferation and angiogenesis in hepatocellular carcinoma and predicts prognosis. *BMC Cancer* 2015; **15**: 245 [PMID: 25884709 DOI: 10.1186/s12885-015-1262-5]
- 76 **Seki E**, De Minicis S, Osterreicher CH, Kluwe J, Osawa Y, Brenner DA, Schwabe RF. TLR4 enhances TGF-beta signaling and hepatic fibrosis. *Nat Med* 2007; **13**: 1324-1332 [PMID: 17952090 DOI: 10.1038/nm1663]
- 77 **Seifert L**, Deutsch M, Allothman S, Alqunaibit D, Werba G, Pansari M, Pergamo M, Ochi A, Torres-Hernandez A, Levie E, Tippens D, Greco SH, Tiwari S, Ly NNG, Eisenthal A, van Heerden E, Avanzi A, Barilla R, Zambirinis CP, Rendon M, Daley D, Pachter HL, Hajdu C, Miller G. Dectin-1 Regulates Hepatic Fibrosis and Hepatocarcinogenesis by Suppressing TLR4 Signaling Pathways. *Cell Rep* 2015; **13**: 1909-1921 [PMID: 26655905 DOI: 10.1016/j.celrep.2015.10.058]
- 78 **Yang JM**, Han DW, Xie CM, Liang QC, Zhao YC, Ma XH. Endotoxins enhance hepatocarcinogenesis induced by oral intake of thioacetamide in rats. *World J Gastroenterol* 1998; **4**: 128-132 [PMID: 11819255 DOI: 10.3748/wjg.v4.i2.128]
- 79 **Shu M**, Huang DD, Hung ZA, Hu XR, Zhang S. Inhibition of MAPK and NF- κ B signaling pathways alleviate carbon tetrachloride (CCl₄)-induced liver fibrosis in Toll-like receptor 5 (TLR5) deficiency mice. *Biochem Biophys Res Commun* 2016; **471**: 233-239 [PMID: 26845355 DOI: 10.1016/j.bbrc.2016.01.119]
- 80 **Yan J**, Shen S, He Y, Li Z. TLR5 silencing reduced hyperammonaemia-induced liver injury by inhibiting oxidative stress and inflammation responses via inactivating NF- κ B and MAPK signals. *Chem Biol Interact* 2019; **299**: 102-110 [PMID: 30508503 DOI: 10.1016/j.cbi.2018.11.026]
- 81 **Kim S**, Park S, Kim B, Kwon J. Toll-like receptor 7 affects the pathogenesis of non-alcoholic fatty

- liver disease. *Sci Rep* 2016; **6**: 27849 [PMID: [27279075](#) DOI: [10.1038/srep27849](#)]
- 82 **Jo J**, Tan AT, Ussher JE, Sandalova E, Tang XZ, Tan-Garcia A, To N, Hong M, Chia A, Gill US, Kennedy PT, Tan KC, Lee KH, De Libero G, Gehring AJ, Willberg CB, Klenerman P, Bertoletti A. Toll-like receptor 8 agonist and bacteria trigger potent activation of innate immune cells in human liver. *PLoS Pathog* 2014; **10**: e1004210 [PMID: [24967632](#) DOI: [10.1371/journal.ppat.1004210](#)]
- 83 **Song IJ**, Yang YM, Inokuchi-Shimizu S, Roh YS, Yang L, Seki E. The contribution of toll-like receptor signaling to the development of liver fibrosis and cancer in hepatocyte-specific TAK1-deleted mice. *Int J Cancer* 2018; **142**: 81-91 [PMID: [28875549](#) DOI: [10.1002/ijc.31029](#)]
- 84 **Duan Y**, Llorente C, Lang S, Brandl K, Chu H, Jiang L, White RC, Clarke TH, Nguyen K, Torralba M, Shao Y, Liu J, Hernandez-Morales A, Lessor L, Rahman IR, Miyamoto Y, Ly M, Gao B, Sun W, Kiesel R, Huttmacher F, Lee S, Ventura-Cots M, Bosques-Padilla F, Verna EC, Abraldes JG, Brown RS Jr, Vargas V, Altamirano J, Caballería J, Shawcross DL, Ho SB, Louvet A, Lucey MR, Mathurin P, Garcia-Tsao G, Batailler R, Tu XM, Eckmann L, van der Donk WA, Young R, Lawley TD, Stärkel P, Pride D, Fouts DE, Schnabl B. Bacteriophage targeting of gut bacterium attenuates alcoholic liver disease. *Nature* 2019; **575**: 505-511 [PMID: [31723265](#) DOI: [10.1038/s41586-019-1742-x](#)]
- 85 **Adolfsson O**, Meydani SN, Russell RM. Yogurt and gut function. *Am J Clin Nutr* 2004; **80**: 245-256 [PMID: [15277142](#) DOI: [10.1093/ajcn/80.2.245](#)]
- 86 **Letexier D**, Diraison F, Beylot M. Addition of inulin to a moderately high-carbohydrate diet reduces hepatic lipogenesis and plasma triacylglycerol concentrations in humans. *Am J Clin Nutr* 2003; **77**: 559-564 [PMID: [12600843](#) DOI: [10.1093/ajcn/77.3.559](#)]
- 87 **Wiest R**, Albillos A, Trauner M, Bajaj JS, Jalan R. Targeting the gut-liver axis in liver disease. *J Hepatol* 2017; **67**: 1084-1103 [PMID: [28526488](#) DOI: [10.1016/j.jhep.2017.05.007](#)]
- 88 **Ahn SB**, Jun DW, Kang BK, Lim JH, Lim S, Chung MJ. Randomized, Double-blind, Placebo-controlled Study of a Multispecies Probiotic Mixture in Nonalcoholic Fatty Liver Disease. *Sci Rep* 2019; **9**: 5688 [PMID: [30952918](#) DOI: [10.1038/s41598-019-42059-3](#)]
- 89 **Rossen NG**, MacDonald JK, de Vries EM, D'Haens GR, de Vos WM, Zoetendal EG, Ponsioen CY. Fecal microbiota transplantation as novel therapy in gastroenterology: A systematic review. *World J Gastroenterol* 2015; **21**: 5359-5371 [PMID: [25954111](#) DOI: [10.3748/wjg.v21.i17.5359](#)]
- 90 **Llopis M**, Cassard AM, Wrzosek L, Bosch L, Bruneau A, Ferrere G, Puchois V, Martin JC, Lepage P, Le Roy T, Lefèvre L, Langelier B, Cailleux F, González-Castro AM, Rabot S, Gaudin F, Agostini H, Prévot S, Berrebi D, Ciocan D, Jousse C, Naveau S, Gérard P, Perlemuter G. Intestinal microbiota contributes to individual susceptibility to alcoholic liver disease. *Gut* 2016; **65**: 830-839 [PMID: [26642859](#) DOI: [10.1136/gutjnl-2015-310585](#)]
- 91 **Philips CA**, Pande A, Shasthry SM, Jamwal KD, Khillan V, Chandel SS, Kumar G, Sharma MK, Maiwall R, Jindal A, Choudhary A, Hussain MS, Sharma S, Sarin SK. Healthy Donor Fecal Microbiota Transplantation in Steroid-Ineligible Severe Alcoholic Hepatitis: A Pilot Study. *Clin Gastroenterol Hepatol* 2017; **15**: 600-602 [PMID: [27816755](#) DOI: [10.1016/j.cgh.2016.10.029](#)]
- 92 **Scarpignato C**, Pelosini I. Rifaximin, a poorly absorbed antibiotic: pharmacology and clinical potential. *Chemotherapy* 2005; **51** Suppl 1: 36-66 [PMID: [15855748](#) DOI: [10.1159/000081990](#)]
- 93 **Bajaj JS**, Heuman DM, Sanyal AJ, Hylemon PB, Sterling RK, Stravitz RT, Fuchs M, Ridlon JM, Daita K, Monteith P, Noble NA, White MB, Fisher A, Sikaroodi M, Rangwala H, Gillevet PM. Modulation of the metabiome by rifaximin in patients with cirrhosis and minimal hepatic encephalopathy. *PLoS One* 2013; **8**: e60042 [PMID: [23565181](#) DOI: [10.1371/journal.pone.0060042](#)]
- 94 **Ponziani FR**, Gerardi V, Pecere S, D'Aversa F, Lopetuso L, Zocco MA, Pompili M, Gasbarrini A. Effect of rifaximin on gut microbiota composition in advanced liver disease and its complications. *World J Gastroenterol* 2015; **21**: 12322-12333 [PMID: [26604640](#) DOI: [10.3748/wjg.v21.i43.12322](#)]
- 95 **Soldi S**, Vasileiadis S, Uggeri F, Campanale M, Morelli L, Fogli MV, Calanni F, Grimaldi M, Gasbarrini A. Modulation of the gut microbiota composition by rifaximin in non-constipated irritable bowel syndrome patients: a molecular approach. *Clin Exp Gastroenterol* 2015; **8**: 309-325 [PMID: [26673000](#) DOI: [10.2147/CEG.S89999](#)]
- 96 **Kang SH**, Lee YB, Lee JH, Nam JY, Chang Y, Cho H, Yoo JJ, Cho YY, Cho EJ, Yu SJ, Kim MY, Kim YJ, Baik SK, Yoon JH. Rifaximin treatment is associated with reduced risk of cirrhotic complications and prolonged overall survival in patients experiencing hepatic encephalopathy. *Aliment Pharmacol Ther* 2017; **46**: 845-855 [PMID: [28836723](#) DOI: [10.1111/apt.14275](#)]
- 97 **Flamm SL**, Mullen KD, Heimanson Z, Sanyal AJ. Rifaximin has the potential to prevent complications of cirrhosis. *Therap Adv Gastroenterol* 2018; **11**: 1756284818800307 [PMID: [30283499](#) DOI: [10.1177/1756284818800307](#)]
- 98 **Mullen KD**, Sanyal AJ, Bass NM, Poordad FF, Sheikh MY, Frederick RT, Bortey E, Forbes WP. Rifaximin is safe and well tolerated for long-term maintenance of remission from overt hepatic encephalopathy. *Clin Gastroenterol Hepatol* 2014; **12**: 1390-7. e2 [PMID: [24365449](#) DOI: [10.1016/j.cgh.2013.12.021](#)]
- 99 **Zapater P**, Caño R, Llanos L, Ruiz-Alcaraz AJ, Pascual S, Barquero C, Moreu R, Bellot P, Horga JF, Muñoz C, Pérez J, García-Peñarrubia P, Pérez-Mateo M, Such J, Francés R. Norfloxacin modulates the inflammatory response and directly affects neutrophils in patients with decompensated cirrhosis. *Gastroenterology* 2009; **137**: 1669-79. e1 [PMID: [19660462](#) DOI: [10.1053/j.gastro.2009.07.058](#)]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

