

Therapy-induced Deletion in 11q23 Leading to Fusion of *KMT2A* With *ARHGEF12* and Development of B Lineage Acute Lymphoblastic Leukemia in a Child Treated for Acute Myeloid Leukemia Caused by t(9;11)(p21;q23)/*KMT2A-MLLT3*

IOANNIS PANAGOPOULOS¹, KRISTIN ANDERSEN¹, MARTINE EILERT-OLSEN¹,
BERNWARD ZELLER², MONICA CHENG MUNTHE-KAAS², JOCHEN BUECHNER²,
LIV T.N. OSNES³, FRANCESCA MICCI¹ and SVERRE HEIM^{1,4}

¹Section for Cancer Cytogenetics, Institute for Cancer Genetics and Informatics,
The Norwegian Radium Hospital, Oslo University Hospital, Oslo, Norway;

²Department of Pediatric Hematology and Oncology, Oslo University Hospital, Rikshospitalet, Oslo, Norway;

³Department of Immunology, Oslo University Hospital, Rikshospitalet, Oslo, Norway;

⁴Institute of Clinical Medicine, Faculty of Medicine, University of Oslo, Oslo, Norway

Abstract. *Background/Aim:* Fusion of histone-lysine N-methyltransferase 2A gene (*KMT2A*) with the Rho guanine nucleotide exchange factor 12 gene (*ARHGEF12*), both located in 11q23, was reported in some leukemic patients. We report a *KMT2A-ARHGEF12* fusion occurring during treatment of a pediatric acute myeloid leukemia (AML) with topoisomerase II inhibitors leading to a secondary acute lymphoblastic leukemia (ALL). *Materials and Methods:* Multiple genetic analyses were performed on bone marrow cells of a girl initially diagnosed with AML. *Results:* At the time of diagnosis with AML, the t(9;11)(p21;q23)/*KMT2A-MLLT3* genetic abnormality was found. After chemotherapy resulting in AML clinical remission, a 2 Mb deletion in 11q23 was found generating a *KMT2A-ARHGEF12* fusion gene. When the patient later developed B lineage ALL, a t(14;19)(q32;q13), loss of one chromosome 9, and *KMT2A-ARHGEF12* were detected. *Conclusion:* The patient sequentially developed AML and ALL with three leukemia-specific genomic abnormalities in her bone marrow cells, two of which were *KMT2A*-rearrangements.

The histone-lysine N-methyltransferase 2A (*KMT2A*, also known as *MLL*) gene in 11q23 (1, 2) may fuse with more than 100 different partners in acute lymphoblastic leukemia (ALL), acute myeloid leukemia (AML), chronic myeloid leukemia, myelodysplastic syndromes, lymphomas, and solid tumors (3). Some of the resulting chimeras are common, such as the fusions with the AF4/FMR2 family member 1 (*AFF1*) and *MLLT3* super elongation complex subunit (*MLLT3*) genes generated by t(4;11)(q21;q23) in ALL (*KMT2A-AFF1*) and t(9;11)(p21;q23) in AML (*KMT2A-MLLT3*), while others have only been reported in few or single cases (4, 5). The prognostic impact of the frequent *KMT2A* fusions is well known (6, 7); however, knowledge about the clinical consequences of the infrequent chimeras is inadequate. For this reason, not just because of biological curiosity, the reporting of cases involving uncommon *KMT2A* fusions is important as recently exemplified by the description of rare *KMT2A-ELL* fusion transcripts in pediatric AML associated with myeloid sarcomas (8-11).

We herein report the first pediatric leukemia, and the sixth case overall, in which fusion of *KMT2A* with the Rho guanine nucleotide exchange factor 12 (*ARHGEF12*) gene was detected.

This article is freely accessible online.

Correspondence to: Ioannis Panagopoulos, Section for Cancer Cytogenetics, Institute for Cancer Genetics and Informatics, The Norwegian Radium Hospital, Oslo University Hospital, P.O. Box 4953 Nydalen, NO-0424 Oslo, Norway. Tel: +47 22934424, email: ioannis.panagopoulos@rr-research.no

Key Words: Pediatric leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, chemotherapy, fusion gene, *KMT2A*, *ARHGEF12*, *KMT2A-ARHGEF12*, *KMT2A-MLLT3*.

Materials and Methods

Ethics statement. The study was approved by the regional ethics committee (Regional komité for medisinsk forskningsetikk Sør-Øst, Norge, <http://helseforskning.etikkom.no>), and written informed consent was obtained from the patient's parents. The Ethics Committee's approval included a review of the consent procedure. All patient information has been anonymized.

Case report. The patient was a girl diagnosed with AML at an age of 9.5 years (Table I, Sample S0), after a period of three months

Table I. *G*-banding, fluorescence in situ hybridization (FISH) and molecular genetic results of the patient.

Sample	Days from AML-diagnosis	Karyotype	Nuclear <i>in situ</i> hybridization (nuc ish) based on <i>KMT2A</i> break-apart probe	Aberrant nuclei (%)	Additional experiments
S0	0	46,XX,t(9;11)(p21;q23)[9]/46,XX[2]	nuc ish (KMT2Ax2)(5'KMT2A sep 3'KMT2Ax1)[195/204]	96%	FISH with a <i>KMT2A-MLL3</i> dual fusion probe: Fusion in metaphase spreads and in 184 out of 190 (97%) nuclei Neither RT-PCR nor genomic PCR amplified <i>KMT2A-ARHGEF12</i> fusion fragments
S1	300 (9 months and 24 days)	Fail	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[160/200]	80%	Not done
S2	328 (10 months and 23 days)	46,XX	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[181/204]	89%	Not done
S3	389 (1 year and 23 days)	46,XX	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[184/200]	92%	RNA sequencing: Detection of a <i>KMT2A-ARHGEF12</i> fusion transcript RT-PCR/Sanger sequencing: Confirmation of the presence of the <i>KMT2A-ARHGEF12</i> fusion transcript
S4	466 (1 year, 3 months, and 8 days)	46,XX	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[199/219]	91%	Not done
S5	571 (1 year, 6 months, and 21 days)	46,XX	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[164/219]	89%	Not done
S6	718 (1 year, 11 months, and 17 days)	46,XX	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[158/200]	79%	FISH with a <i>KMT2A-MLL3</i> dual fusion: No fusion in 200 nuclei
S7	914 (2 years, 5 months, and 30 days)	46,XX	Not done		FISH with a <i>KMT2A-MLL3</i> dual fusion probe: No fusion in 200 nuclei
S8	1,198 (3 years, 3 months, and 10 days), at clinical diagnosis of B lineage ALL	45,XX,-9,t(14;19)(q32;q13)[8]/46,XX[3]	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[194/200]	97%	Genomic PCR/Sanger sequencing: Detection of the genomic breakpoint/junction of <i>KMT2A-ARHGEF12</i> fusion gene FISH with a <i>KMT2A-MLL3</i> dual fusion probe: No fusion in 200 nuclei FISH with <i>IGH</i> break-apart probe: Split of <i>IGH</i> probe in 195 out of 211 (92%) nuclei
S9	1,294 (3 years, 6 months, and 14 days)	Fail	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[17/200]	8.5%	FISH with <i>IGH</i> break-apart probe: No split of <i>IGH</i> in 200 nuclei
S10	1,539 (4 years, 2 months, and 17 days)	46,XX	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[164/203]	81%	Genomic PCR/Sanger sequencing: Detection of the genomic breakpoint/junction of <i>KMT2A-ARHGEF12</i> fusion gene aCGH detected a 2Mbp deletion between <i>KMT2A</i> and <i>ARHGEF12</i> genes
S11-A	1,686A (4 years, 7 months, and 11 days)	46,XX	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[87/212]	41%	Not done
S11-B	1,686B (4 years, 7 months, and 11 days)	Not done	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[85/110]	77%	Not done
S11-C	1,686C (4 years, 7 months, and 11 days)	Not done	nuc ish (5'KMT2Ax2,3'KMT2Ax1)(5'KMT2A con 3'KMT2Ax1)[115/335]	34%	Not done

Table I. *Continued*

Table I. *Continued*

Sample	Days from AML-diagnosis	Karyotype	Nuclear <i>in situ</i> hybridization (nuc ish) based on <i>KMT2A</i> break-apart probe	Aberrant nuclei (%)	Additional experiments
S12	1,945 (5 years, 3 months, 26 days), 3 months after stem cell transplantation	Not done	nuc ish (5'KMT2A,3'KMT2A)x 2(5'KMT2A con 3'KMT2Ax2)[200]	100%	Not done

Clinical diagnosis of acute myeloid leukemia (AML) at time point (sample) S0, diagnosis of B lineage acute lymphoblastic leukemia (ALL) at time point S8. Sample 11 (S11-A, S11-B, and S-11C) represents the last bone marrow before stem cell transplantation. Sample 12 represents bone marrow 3 months after stem cell transplantation. RT-PCR: reverse transcription-polymerase chain reaction. aCGH: array comparative genomic hybridization. A: Bone marrow. B: CD34+/CD117+ cells. C: Negative fraction.

with fatigue, pallor, headache, and dizziness. There was no history of hematologic diseases, leukemia or frequent cancers in the family. At admittance, she was in good shape. Physical examination was normal except for pallor. Spleen and liver size were normal. Her blood tests were abnormal with hemoglobin 5.8 g/dl, platelet count $95 \times 10^9/l$, white blood cell count $4.9 \times 10^9/l$, and neutrophils $0.1 \times 10^9/l$. Bone marrow investigation revealed findings indicative of acute myelomonocytic leukemia (AML M4), and cytogenetic analysis showed a t(9;11)(p21;q23) chromosome translocation (Table I, sample S0, see below).

The girl was treated according to the NOPHO-DBH 2012 protocol (12) where she was classified as a standard risk patient. She received five chemotherapy courses including the drugs liposomal daunorubicin, mitoxantrone, etoposide, cytarabine, and fludarabine. Following completion of therapy, she entered a prolonged phase with moderate pancytopenia. During this period which lasted two years and three months, repeated bone marrow examinations were performed (Table I, samples S1-S7). They revealed hematological remission and a normal karyotype but with a persisting *KMT2A* rearrangement, however one that was different from the original *KMT2A-MLL3* (Table I, samples S1-S7, see below). Flow cytometry showed a small persisting CD34+, CD38+ cell population with aberrant phenotype (reduced HLA-DR, CD33 weak/negative, CD11a weak, CD117 heterogenous), in a proportion decreasing from 1.5% to 0.05%, in addition to the phenotypically normal CD34+ CD38+ myeloid precursors, over eight months. The girl was always in good general health and lived a completely normal life. Following normalization of her peripheral blood values, no more bone marrow examinations were performed until she, two years and nine months after start of the last AML chemotherapy course (or three years and three months after the initial diagnosis of AML), again developed pancytopenia. She was now diagnosed with ALL (Table I, S8). The bone marrow contained 90% blasts of pre-B phenotype, whereas cytogenetic analysis revealed a t(14;19)(q32;q13) chromosome translocation together with loss of one chromosome 9 (Table I, sample S8, see below). Treatment according to the NOPHO-ALL 2008 protocol (13) was begun but, due to increased toxicity, had to be modified. Bone marrow controls during therapy confirmed a good and early treatment response with swift reduction of the lymphoblasts. However, the same small population of abnormal CD34+, CD38+ cells detected shortly after discontinuation of the primary AML

treatment and persisting ever since, was still detectable. It was therefore decided to give her one dose of Daratumomab (a CD38-antibody) 16 mg/kg with the aim of eradicating the abnormal clone before stem cell transplantation. After the one dose of Daratumomab CD38 was not detected on any CD34+ cells, using the standard monoclonal anti-CD38. However, using a multiepitope anti-CD38 the antigen could still be detected, indicating down-regulation of the epitope detected by the monoclonal antibody. Other markers demonstrated that the abnormal population was still present in the first controls after Daratumomab. The first bone marrow negative for CD34+ cells was confirmed 11 weeks after initiation of Daratumomab treatment, *i.e.*, more than four years after the clone was first detected. Six weeks after the first negative bone marrow, allogeneic stem cell transplantation was performed with a human leukocyte antigen (HLA) - matched, unrelated donor after standard ALL-conditioning regimen consisting of total body irradiation (TBI) and etoposide. The patient is presently in complete remission, three months after stem cell transplantation. The current bone marrow assessment did not detect abnormal CD34+CD38+ cells or rearrangement of the *KMT2A* gene (see below).

G-banding and fluorescence in situ hybridization (FISH) analyses. Bone marrow cells were short-term cultured, G-banded, and analyzed cytogenetically as previously described (14).

FISH analyses of bone marrow interphase nuclei and metaphase spreads were performed with the Cytocell *KMT2A* (*MLL*) and *IGH* break-apart probes, as well as with *KMT2A-MLL3* translocation dual fusion probes (Cytocell, Oxford Gene Technology, Begbroke, Oxfordshire, UK). Fluorescent signals were captured and analyzed using the CytoVision system (Leica Biosystems, Newcastle, UK).

DNA and RNA isolation and complementary DNA (cDNA) synthesis. Genomic DNA was extracted from the patient's bone marrow samples at diagnosis as well as 1,198 and 1,539 days after the diagnosis (Table I, samples S0, S8, and S10) using the Maxwell 16 Instrument System and the Maxwell 16 Cell DNA Purification Kit (Promega, Madison, WI, USA). Total RNA was extracted from the patient's bone marrow at diagnosis and after 389 days (Table I, samples S0, and S3) using the miRNeasy Mini Kit (Qiagen, Hilden, Germany) and the QiaCube automated purification system according to the manufacturer's instructions (Qiagen). The concentration and purity of DNA and RNA

Table II. Primers used for PCR amplification and Sanger sequencing analyses. The M13 forward and reverse primer sequences are in bold and italics.

Name	Sequence (5'->3')	Position	Reference sequence	Gene
M13For-MLL-4010F1	<i>TGTA AACGACGGCCAGT</i> TAAAGCAGCCTCCACCACCAGAAT	4010-4032	NM_005933.3	<i>KMT2A</i>
M13For-MLL-4115F1	<i>TGTA AACGACGGCCAGT</i> ACCACCTCCGGTCAATAAGCAGG	4115-4137	NM_005933.3	<i>KMT2A</i>
M13Rev-ARHGEF12-1515R1	<i>CAGGAAACAGCTATGACCC</i> AAAGTCGACGGGTCAAATTGTGA	1537-1515	NM_015313.2	<i>ARHGEF12</i>
M13Rev-ARHGEF12-1579R1	<i>CAGGAAACAGCTATGACCC</i> GGAAGATGCGACGAGTTTCTTTGG	1579-1602	NM_015313.2	<i>ARHGEF12</i>
MLL-4116-F1	CCACCTCCGGTCAATAAGCAGGAGAA	4116-4141	NM_005933.3	<i>KMT2A</i>
MLL-4202-F1	TCCAGCAGATGGAGTCCACAGGATCA	4202-4227	NM_005933.3	<i>KMT2A</i>
ARHGEF12-1437-R1	TTCAATGCTCTGGAAACAGCTGCACTG	1463-1437	NM_015313.2	<i>ARHGEF12</i>
ARHGEF12-1502-R1	GCAGGGTCAAATTGTGAAACTACATGGTGT	1531-1502	NM_015313.2	<i>ARHGEF12</i>

were measured with the QIAxpert microfluidic UV/VIS spectrophotometer (Qiagen). In addition, the Agilent 2100 bioanalyzer and the RNA Integrity Number (RIN) were used to assess the RNA quality (15). RIN of RNA was 6.6. cDNA was synthesized from one µg of total RNA using the iScript Advanced cDNA Synthesis Kit for RT-qPCR according to the manufacturer's instructions (Bio-Rad, Hercules, CA, USA). The quality of the cDNA synthesis was assessed by amplification of a cDNA fragment of the ABL protooncogene 1, non-receptor tyrosine kinase (*ABL1*) gene using the primer combination ABL1-91F1/ABL1-404R1 (16).

RNA sequencing. High-throughput paired-end RNA-sequencing was performed at the Genomics Core Facility, Norwegian Radium Hospital, Oslo University Hospital (<http://genomics.no/oslo/>). For library preparation from total RNA, the Illumina TruSeq RNA Access Library Prep kit was used according to Illumina's protocol (Illumina, San Diego, CA, USA). Sequencing was performed on a NextSeq 550 System (Illumina) and 76 million reads were generated. The FASTQC software was used for quality control of the raw sequence data (<https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>). The software FusionCatcher was used to find fusion transcripts (17, 18).

PCR analyses. The primers used for PCR amplification and Sanger sequencing are listed in Table II. For reverse transcription-polymerase chain reaction (RT-PCR) and cycle Sanger sequencing, the BigDye Direct Cycle Sequencing Kit was used (ThermoFisher Scientific, Waltham, MA, USA) according to the company's recommendations. As template, cDNA corresponding to 20 ng total RNA was used. The primer combinations were M13For-MLL-4010F1/M13Rev-ARHGEF12-1579R1 and M13For-MLL-4115F1/M13Rev-ARHGEF12-1515R1. For the detection of possible *KMT2A-ARHGEF12* chimeric cDNA fragments at initial diagnosis, the primer combination MLL-4116-F1/ARHGEF12-1502-R1 was used as described below.

Genomic PCR amplifications were performed in 25 µl reaction volume which contained 12.5 µl Premix Ex Taq™ DNA Polymerase Hot Start Version (Takara Bio Europe/SAS, Saint-Germain-en-Laye, France), 100 ng of genomic DNA, and 0.4 µM of each of the forward and reverse primers. The primer combinations were MLL-4116-F1/ARHGEF12-1502-R1 and MLL-4202-F1/ARHGEF12-1437-R1. The PCR cycling conditions were an initial denaturation step at 94°C for 30 sec followed by 35 cycles of 7 sec at 98°C and 2 min at 68°C, and a final extension for 5 min at 68°C. Three µl of

the PCR products were stained with GelRed (Biotium, Fremont, CA, USA), analyzed by electrophoresis through 1% agarose gel, and photographed. DNA gel electrophoresis was performed using lithium borate buffer (19). The remaining PCR products were purified using the MinElute PCR Purification Kit (Qiagen) and direct sequenced using the dideoxy procedure with the BigDye terminator v1.1 cycle sequencing kit following the company's recommendations (ThermoFisher Scientific).

Sequence analyses were performed on the Applied Biosystems SeqStudio Genetic Analyzer system (ThermoFisher Scientific). The basic local alignment search tool (BLAST) software (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) was used for computer analysis of sequence data (20). The BLAT alignment tool and the human genome browser at UCSC were also used to map the sequences on the Human GRCh37/hg19 assembly (21, 22).

Array comparative genomic hybridization (aCGH) analysis. aCGH was performed using the CytoSure array products (Oxford Gene Technology, Begbroke, Oxfordshire, UK) following the company's protocols. Thus, the CytoSure Genomic DNA Labelling Kit was used for the labelling of 1 µg of patient's and reference DNAs and the CytoSure Cancer +SNP array for hybridization. The patient's DNA was that isolated from a sample drawn 1,539 days after the initial diagnosis of AML (Table I, S10). The reference DNA was Promega's human genomic female DNA (Promega, Madison, WI, USA). The slides were scanned by an Agilent scanner using Agilent Feature Extraction Software (version 10.7.3.1). Data were analysed using the CytoSure Interpret analysis software (version 4.9.40). Annotations are based on human genome build 19.

Results

G-banding analyses. The data from G-banding and FISH analyses are summarized in Table I. The G-banding analysis of bone marrow cells at diagnosis yielded the karyotype 46,XX,t(9;11)(p21;q23)[9]/46,XX[2] (Figure 1A). Subsequently, bone marrow G-banding analyses always showed a normal 46,XX karyotype except for the sample obtained 1,198 days after diagnosis. At that point, the karyotype had become 45,XX,-9, t(14;19)(q32;q13)[8]/ 46,XX[3] and the patient had developed B lineage ALL (Table I, sample S8) (Figure 1B).




Figure 1. Cytogenetic analyses of the two pediatric leukemias. (A) Karyogram showing the $t(9;11)(p21;q23)$ found at diagnosis, when the patient had acute myeloid leukemia (AML). (B) Karyogram showing the loss of chromosome 9 and the $t(14;19)(q32;q13)$ found when the patient was diagnosed with acute lymphoblastic leukemia (ALL) 1,198 days after the initial AML. Breakpoint positions are indicated by arrows.

FISH analyses. Three FISH analyses were performed at initial diagnosis of AML. Interphase FISH with the *KMT2A* break-apart probe showed a normal (yellow) as well as split (separated red and green) signals of the probe in 195 out of 204 examined nuclei (96%) (Table I, sample S0) (Figure 2A). FISH analysis with a *KMT2A-MLLT3* translocation dual fusion probe on metaphase spreads showed the *KMT2A-MLLT3* and *MLLT3-KMT2A* fusion genes on chromosomes der(11) and der(9), respectively (Figure 2B). Interphase FISH with the same *KMT2A-MLLT3* translocation dual fusion probe showed a normal red signal corresponding to the *MLLT3* gene, a normal green signal corresponding to the *KMT2A* gene, and two yellow fusion signals corresponding to *KMT2A-MLLT3* and *MLLT3-KMT2A* in 184 out of 190 (97%) examined nuclei (Figure 2C). In addition, at another diagnostic laboratory, a *KMT2A-MLLT3* fusion transcript was detected (data not shown) which was in an agreement with the G-banding and interphase FISH results.

In the sample obtained 300 days after diagnosis, interphase FISH with a break-apart *KMT2A* probe for the first time showed loss of the distal part of the *KMT2A* probe, namely in 160 out of 200 nuclei (80%) (Table I, sample S1, Figure 2D). This pattern with deletion or loss of the distal part of *KMT2A* has since persisted throughout the entire follow-up period (Table I, samples S1-S11-A) until the patient was transplanted. Deletion of the distal part of the *KMT2A* probe was also found in 77% of the CD34+CD117+ cells and in 34% of the negative fraction of the sorted cells 1,686 days after the diagnosis (Table I, samples S11-B and S11-C).

In the sample obtained 90 days after allogeneic stem cell transplantation the *KMT2A* break-apart probe showed two normal (yellow) signals in 200 interphase nuclei (Table I, sample S12).

Interphase FISH with a *KMT2A-MLLT3* translocation dual fusion probe on samples obtained 718 days and 914 days after the initial diagnosis did not detect any *KMT2A-MLLT3* fusion in 200 examined nuclei (Table I, samples S6 and S7).

In the sample obtained 1,198 days after diagnosis, when a t(14;19)(q32;q13) was seen by karyotyping and the patient had developed ALL, the FISH results were as follows: No *KMT2A-MLLT3* fusion was seen in 200 nuclei, deletion of the distal part of the *KMT2A* probe was detected in 194 out of 200 nuclei (97%) and on metaphase spreads (Figure 2E), and splitting of the *IGH* probe was seen in 195 out of 211 (92%) nuclei as well as on metaphase spreads (Figure 2F and 2G) (Table I, sample S8). In the sample obtained 1,294 days after diagnosis, FISH analyses showed deletion of the distal part of the *KMT2A* probe in 17 out of 200 nuclei (8.5%) and no split of *IGH* in 200 nuclei (Table I, sample S9).

RNA-sequencing and molecular genetic confirmation of the fusion. After the G-banding/interphase FISH results were obtained from the sample drawn at day 1,539, we decided to perform retroactive RNA sequencing of the bone marrow

sample obtained 389 days after the diagnosis. At that point in time (389 days), the G-banding analysis was normal whereas FISH with the break apart *KMT2A* probe showed deletion of the distal part of the probe in 184 out of 200 (92%) examined nuclei (Table I, sample S3).

Using FusionCatcher on the raw sequencing data obtained from the Genomics Core Facility, a *KMT2A-ARHGEF12* fusion transcript was found in which exon 9 of *KMT2A* (nt 4241 in sequences with accession numbers NM_005933.3 and NM_001197104.1) was fused to exon 14 of *ARHGEF12* (nt 1428 in sequence with accession number NM_015313.2) (AATTCCAGCAGATGGAGTCCACAGGATCAGAGTGGA CTTTAAG*ATCAATGGACAGTGCAGCTGTTTCCAGAG CATTGAATTACTAA). No reciprocal *ARHGEF12-KMT2A* fusion transcript was found.

RT-PCR and cycle Sanger sequencing with the primers M13For-MLL-4010F1/M13Rev-ARHGEF12-1579R1 and M13For-MLL-4115F1/M13Rev-ARHGEF12-1515R1 confirmed the results obtained by RNA sequencing/Fusion Catcher analysis (Figure 3A).

Genomic PCR using as template DNA extracted from the patient's bone marrow cells sampled 1,198 days and 1,539 days after the initial diagnosis amplified a 1264 bp fragment with the primer combination MLL-4116-F1/ARHGEF12-1502-R1 and a 1133 bp fragment with the primer combination MLL-4202-F1/ARHGEF12-1437-R1 (Figure 3B and C). Direct sequencing showed that they were genomic *KMT2A-ARHGEF12* chimeric fragments in which a sequence from intron 9 of *KMT2A* was fused to a sequence of intron 13 from *ARHGEF12* (Figure 3D).

Neither RT-PCR nor genomic PCR detected *KMT2A-ARHGEF12* chimeric fragments when the template was RNA (cDNA) or genomic DNA obtained from the diagnostic (day 0) bone marrow (Figure 3C).

aCGH analysis. aCGH detected a deletion in the q arm of chromosome 11 (Figure 4). Based on the hg19 assembly, the deletion was from the probe at position Chr11:118355288-118355347 in *KMT2A* to the probe at position Chr11:120290981-120291040 in *ARHGEF12* (Figure 4). Thus, the aCGH data agreed with the results of FISH analyses, genomic PCR, and RT-PCR; they suggested that the *KMT2A-ARHGEF12* fusion gene was the result of a deletion within chromosome band 11q23.

Discussion

A fusion of *KMT2A* with the *ARHGEF12* gene was first reported in a 38-year-old male with a history of occupational exposure to herbicides who had developed AML (FAB-M4) and the abnormal karyotype 51,XY,+8,+19,+3mar (23). The patient received standard induction chemotherapy and achieved complete remission by morphological bone marrow




Figure 2. Fluorescence in situ hybridization (FISH) analyses of pediatric leukemia. (A) Interphase FISH at initial diagnosis of AML with the *KMT2A* break-apart probe showing a normal (yellow) and split (separated red and green) signals of the probe in 3 nuclei. (B) FISH analysis at initial diagnosis of AML with the *KMT2A-MLLT3* translocation dual fusion probe on metaphase spreads showing a normal green signal on chromosome 11, corresponding to *KMT2A*, a normal red signal on chromosome 9, corresponding to *MLLT3*, and two yellow fusion signals on der(11) and der(9) corresponding to the *KMT2A-MLLT3* and *MLLT3-KMT2A* fusion genes, respectively. (C) Interphase FISH at initial diagnosis of AML on two nuclei using the *KMT2A-MLLT3* translocation dual fusion probe showing a normal green signal, corresponding to *KMT2A*, a normal red signal, corresponding to *MLLT3*, and two yellow fusion signals corresponding to the *KMT2A-MLLT3* and *MLLT3-KMT2A* fusion genes. (D) Interphase FISH with the break-apart *KMT2A* probe on the sample obtained 300 days after diagnosis showing deletion of the distal part of the *KMT2A* probe (lack of red signal) in two nuclei and two normal (yellow) *KMT2A* signals in one nucleus. (E) FISH analysis with the break-apart *KMT2A* probe on metaphase spread from the sample obtained 1,198 days after diagnosis, when a $t(14;19)(q32;q13)$ was seen by karyotyping and the patient had developed ALL. The distal part (red signal) of the probe is absent in one of the two copies of chromosome 11. (F) Interphase FISH with the *IGH* break-apart probe on the sample obtained 1,198 days after diagnosis, when a $t(14;19)(q32;q13)$ was seen by karyotyping and the patient had developed ALL. A normal (yellow) and split (separated red and green) signals of the probe are shown in 2 nuclei. (G) FISH analysis with the *IGH* break-apart probe on a metaphase spread from the sample obtained 1,198 days after diagnosis, when a $t(14;19)(q32;q13)$ was seen by karyotyping and the patient had developed ALL. A normal (yellow) signal on chromosome 14 together with separate red, on der(14), and green, on der(19), probe signals are shown.




Figure 3. PCR analyses of the pediatric leukemias. (A) Partial sequence chromatogram showing the junction position of exon 9 of the *KMT2A* with exon 14 of *ARHGEF12* in the chimeric transcript. (B) Gel electrophoresis showing the amplified *KMT2A-ARHGEF12* genomic DNA fragment using two primer combinations and template DNA extracted from the patient's bone marrow cells 1,539 days after the initial diagnosis. Lane 1: Primer combination MLL-4116F1/*ARHGEF12*-1502R1. Lane 2: Primer combination MLL-4202-F1/*ARHGEF12*-1437-R1. (C) Gel electrophoresis showing the absence of amplified *KMT2A-ARHGEF12* cDNA (lane 1) and genomic DNA (lane 3) fragments at diagnosis when the leukemic cells had *t*(9;11)(p21;q23) and a *KMT2A-MLLT3* fusion gene, and the presence of *KMT2A-ARHGEF12* in the patient's bone marrow cells 1,198 days after the initial diagnosis, when the patient had developed ALL with *t*(14;19)(q32;q13) and rearrangement of the *IGH* locus (lane 4). For both cDNA and genomic DNA amplifications, the primer combination MLL-4116F1/*ARHGEF12*-1502R1 was used. Lane 2: Assessment of the quality of cDNA synthesis by amplification of a cDNA fragment of *ABL1* using the primer combination *ABL1*-91F1/*ABL1*-404R1. Lane 4: Amplification of a genomic *KMT2A-ARHGEF12* DNA fragment in the patient's bone marrow cells 1,198 days after the initial diagnosis. M, GeneRuler 1 Kb Plus DNA ladder (ThermoFisher Scientific). (D) Partial sequence chromatogram showing the junction position of intron 9 of the *KMT2A* with intron 13 of *ARHGEF12* in the chimeric amplified DNA fragment. The junction of positions 11:118355287-11:120311617 is based on the human genome GRCh37/hg19 assembly.

analysis, whereupon he received successful allogeneic bone marrow transplantation from an HLA identical sibling. He died 6 months later from interstitial pneumonia. There were no signs of relapse at autopsy (23).

The second case was a 77-year-old female with AML (FAB-M5a) and the karyotype 53,XX,+6,+8,+8,+9,+11,+13,+22 (24). The third patient with *KMT2A-ARHGEF12* was an adult female with therapy-related AML (t-AML) (4, 5). No




Figure 4. aCGH showing the deletion in the q arm of chromosome 11. Based on the hg19 assembly, the deletion started from the probe at position Chr11:118355288-118355347 in *KMT2A* and ended at position Chr11:120290981-120291040 in *ARHGEF12*. The deletion is approximately 2 Mbp.

information on the treatment and clinical outcome was provided for the second and third patients. The fourth patient with *KMT2A-ARHGEF12* fusion was a 69-year-old-male with B-cell ALL and a normal karyotype (25). This patient did not achieve complete remission after two courses of induction therapy and died from the disease (25). The fifth patient was a female with B-cell ALL and the bone marrow karyotype 46,XX,der(9)t(3;9)(p21;q34)/46,idem,der(9)t(3;9) (26). No further information was provided.

The present, sixth patient is the first reported pediatric case with a *KMT2A-ARHGEF12* fusion gene. The fusion was detected in the patient's bone marrow after completion of treatment for an AML with t(9;11)(p21;q23) and a *KMT2A-MLL3* fusion gene. The chemotherapy included, among other drugs, daunorubicin, mitoxantrone and etoposide which all are topoisomerase II inhibitors and associated with DNA double-strand breaks, the generation of *KMT2A* rearrangements, the

formation of *KMT2A*-fusion genes, and therapy-related acute leukemias (27-33). B-ALL with 11q23 abnormalities/*KMT2A* rearrangements developing after treatment of a primary malignancy with topoisomerase II inhibitors have been reported (34-40). The majority of cases had the t(4;11)(q21;q23) translocation resulting in formation of a *KMT2A-AFF1* fusion gene, but fusions of *KMT2A* with *MLL1* (19p13.3), *FOXO3A* (6q21), *MAML2* (11q21), *ACTN4* (19q13), *CEP164* (11q23.3), and *PRRC1* (5q23) were also found (5, 39). Therefore, we consider the chemotherapy to have caused the *KMT2A-ARHGEF12* fusion.

The first indication of a *KMT2A-ARHGEF12* fusion gene was seen in the patient's bone marrow cells at days 300 and 328 after the initial AML diagnosis when, in FISH experiments, deletion of the distal part of the *KMT2A* probe was detected in spite of a normal karyotype (Table I, samples S1 and S2). RNA sequencing performed later detected the

KMT2A-ARHGEF12 fusion gene, PCR amplifications verified it both at the transcriptional and genomic level (Figure 3), and aCGH showed that the cause of the fusion was indeed a submicroscopic deletion starting in intron 9 of *KMT2A* extending to intron 13 of *ARHGEF12* (Figure 4). These retrospective findings fit perfectly the early detection of a deleted distal part of the *KMT2A* probe, linking it to the emergence of the *KMT2A-ARHGEF12* fusion. Because the aberration was found in 77% of the CD34+CD117+ cells (Table I, sample S11-B), we concluded that the treatment-induced generation of *KMT2A-ARHGEF12* fusion probably took place in a multipotent progenitor cell. Indeed, *in vitro* experiments have shown that etoposide-induced rearrangements of *KMT2A* resulting in fusion genes can occur in mouse embryonic stem cells (27), human CD34+ hematopoietic stem cells from fetal liver (41), human CD34+ cells isolated from umbilical cord blood (28, 42), and in human embryonic stem cells (43). Furthermore, Libura *et al.* (42) showed that the etoposide-induced-*KMT2A* rearrangements were found at high frequency in human primitive hematopoietic stem cells with, *in vitro* and *in vivo*, long-term repopulating potential.

During a period of two years and three months (after completion of therapy of the initial AML), repeated bone marrow examinations were performed and in all of them, although there was no sign of malignancy, the genetic pattern was the same: normal karyotype and deletion of the distal part of the *KMT2A* probe in FISH experiments indicating, as we only later realized, the formation of *KMT2A-ARHGEF12* (Table I, samples S1-S7). Searching the relevant literature, we found two reports describing similar findings to ours (44, 45). In the first, a 5-year-old girl was diagnosed with AML carrying a variant of t(8;21) (44). The patient achieved clinical and hematologic remission which, 4 years later when the authors reported the case, was still unquestionable. Bone marrow examination 15 months after the initial diagnosis had revealed a clonal t(11;11)(q13;q23), however, and the translocation gave rise to a *KMT2A-ARHGEF17* and the reciprocal *ARHGEF17-KMT2A* fusion genes (44). Retrospective PCR analysis did not detect any *KMT2A-ARHGEF17* chimera in samples drawn at the time of diagnosis suggesting that the fusion was treatment-induced. The *KMT2A* aberration was found only in myeloid lineage cells and in numerous samples over a 30-month period without simultaneous phenotypic signs of a new malignancy. Because myeloid cells as mature as promyelocytes and even differentiated polymorphonuclear neutrophils carried the *KMT2A-ARHGEF17*, the authors concluded that the fusion had probably occurred in a myeloid stem cell but without blocking further differentiation (44).

The second case was a 3-year-old boy with B-ALL and a hyperdiploid karyotype: 58<2n>,XY,+X,+4,+6,+10,+15,+17,+18,+18,+20,+21,+21,+22 (45). During chemotherapy,

a t(3;11)(p21;q23) developed in bone marrow cells giving rise to a *KMT2A-SACMIL* fusion gene. The *KMT2A-SACMIL* fusion was detected in many samples over a 7-year period in which the patient was in hematological remission. A difference between the *KMT2A-ARHGEF17* and *KMT2A-SACMIL* fusion genes was that whereas the first resulted in a *KMT2A-ARHGEF17* fusion protein, the second encodes a truncated *KMT2A* protein (44, 45).

Our patient was, two years and nine months after the start of the last AML chemotherapy course (or 3 years and 3 months after the initial diagnosis of AML), diagnosed with B-ALL (Table I, sample S8). The bone marrow contained 90% blasts of the pre-B phenotype, cytogenetic analysis revealed a t(14;19)(q32;q13) translocation together with loss of one chromosome 9 (Figure 1B, Table I, sample S8, see below), and FISH analyses showed splitting of the *IGH* probe in 92% of the nuclei together with deletion of the distal part of the *KMT2A* probe in 97% of the nuclei. In addition, later molecular analysis demonstrated presence of the *KMT2A-ARHGEF12* fusion gene also at this time. Given the fact that so many bone marrow cells had the aberrations, most of them must have had both, leading us to conclude that the t(14;19)(q32;q13) translocation and loss of one chromosome 9 were secondary to the submicroscopic deletion in 11q23. Three months after treatment for B-ALL had been instituted, the (14;19)(q32;q13)/*IGH* aberration was no longer detectable while the *KMT2A* rearrangement was found in only 8.5% of the examined interphase nuclei. Evidently, the treatment given had considerable effect also on the cells carrying *KMT2A* rearrangement, but not enough to completely eradicate them. Summing up the extensive and somewhat complicating data – the high frequency of t(14;19)(q32;q13)/*IGH* rearrangement and *KMT2A* rearrangement in bone marrow cells when the patient developed B-ALL, and the absence of t(14;19)(q32;q13)/*IGH* rearrangement together with the low frequency of *KMT2A* rearrangement after treatment for this disease – strongly indicate that (most of) the cells carrying *KMT2A*-rearrangement were part of the leukemic clone which carried the t(14;19)(q32;q13)/*IGH* rearrangement when the patient had B-ALL.

Our data are also in agreement with the findings reported by Jeffries *et al.* (46). Examining 20 B-ALL patients, they showed that translocations involving the *IGH* locus coexisted with other primary chromosomal rearrangements either in the same or separate clones. In one of the studied patients, *KMT2A* rearrangement was the primary aberration whereas the *IGH*-translocation the secondary. Most cells had only *KMT2A* rearrangement but cells with only *IGH*-translocation and cells with both *KMT2A* aberration and *IGH*-translocation were also found (46).

The case we present has extensive similarities with the case reported by Zuna *et al.* (47). In brief, a 15-year-old girl with acute promyelocytic leukemia (APL, AML-M3) carrying

a *PML-RARA* fusion gene together with internal tandem duplication of *FLT3* (*FLT3/ITD*) was treated with drugs that included the topoisomerase II inhibitors idarubicin and etoposide. Thirty months after the diagnosis of APL and 13 months after completion of therapy, she was diagnosed with B-ALL. Cytogenetic analysis revealed a t(6;11)(q21;q23) translocation aberration with molecular analyses showing the presence of a *KMT2A-FOXO3A* fusion gene whereas both *PML-RARA* fusion gene and *FLT3/ITD* were absent (47).

Backtracking the B-ALL, just as we did in the present study, the authors found the *KMT2A-FOXO3A* fusion gene in up to 90% of bone marrow cells, including cells of the myeloid lineage, more than one year before the B-ALL diagnosis which suggested that the generation of the *KMT2A-FOXO3A* fusion occurred in a multipotent progenitor. During this preleukemic phase, no blasts were found in the bone marrow but a small population of CD19+ B-cells, comprising 0.2% of the bone marrow, was detected (47). In our case, similarly a small persisting aberrant CD34+, CD38+ cell population, decreasing in a proportion from 1.5% to 0.05% over eight months, was found whereas the deletion of the distal part of *KMT2A* was found in 79-92% of the cells (Table 1, samples S1-S7).

Zuna *et al.* (47) identified a 10 Mb gain on 19q13.32 as a potential “second hit” for the development of the B-ALL. The gain was present in B-ALL but was not found in the preleukemic specimen. By analogy, in our case t(14;19)(q32;q13) and loss of one chromosome 9 were secondary aberrations found at diagnosis of ALL but not in any previously examined samples (Table I, samples S1-S8).

Our patient also has similarities with the case presented by Jonveaux *et al* (48): A 35-year-old male had an AML (FAB M5b) with the chromosome aberration t(6;11)(q27;q23). FISH analysis showed that presence of a 400-600 kbp deletion downstream of the 11q23 breakpoint. Using Southern blot methodology, the breakpoint of *KMT2A* was found within intron 7 of the gene. The patient received chemotherapy which, among the drugs, included mitoxantrone and etoposide. Ten and a half months after complete remission, he was diagnosed with B-ALL. The cytogenetic analysis now revealed a t(4;11)(q21;q23) chromosome translocation, FISH showed absence of the deletion, and Southern blot detected a breakpoint in intron 9 of *KMT2A* (48). The authors considered etoposide as responsible for the development of the secondary B-ALL.

At the transcription level, the fusion of the present case was found to recombine exon 9 of *KMT2A* with exon 14 of *ARHGEF12*. This is at odds with what was reported by Kourlas *et al.* (23) who found fusion between exons 8 of *KMT2A* and 12 of *ARHGEF12*. No information on the exact *KMT2A-ARHGEF12* fusion point was given for the other three reported patients (4, 5, 24, 25). The *KMT2A-ARHGEF12* fusion of the present patient was brought about

by a 2Mbp deletion stretching from intron 9 of *KMT2A* to intron 13 of *ARHGEF12*. Deletions were also reported as the cause of *KMT2A-ARHGEF12* formation in the patients described by Kourlas (23) and Meyer (4, 5).

The exact genomic breakpoint position in *KMT2A* has been shown to correlate with clinical outcome in acute leukemias caused by fusions between *KMT2A* and some of the more common partner genes (5, 49, 50). Breakpoints in intron 10 (listed as intron 11 in the references (5, 49, 50)) interfere with the dimerization capacity of *KMT2A* plant homeodomains (PHD) 1-3, disabling binding to the BMI1 (B cell-specific Moloney murine leukemia virus integration site 1) repressor complex, and are associated with worse prognosis (5, 49-51). Breakpoints within *KMT2A* introns 8 and 9 (introns 9 and 10 in references (5, 49, 50), on the other hand, do not affect the three PHD finger domains and are associated with somewhat better clinical outcomes. The breakpoint in intron 9 of the present case could explain the relatively benign clinical behavior observed with long-term survival in spite of the late detection of a *KMT2A-ARHGEF12* fusion gene and the therapy-related leukemia it signals.

The *ARHGEF12* gene codes for the Rho guanine nucleotide exchange factor 12 which is a guanine nucleotide exchange factor that activates small GTPases of the Rho family (see RHOA, 165390) and catalyzes the exchange of GDP for GTP (52-54). *ARHGEF12* was shown to be involved in cell polarization, morphology, invasion, and cytokinetic abscission (55-61). *ARHGEF12* together with *ARHGEF2* (also known as GEF-H1) were found to be key molecules for cellular adaptation to force and regulate the mechanical response to force on integrins (62). *ARHGEF12* was found to regulate human megakaryocyte maturation, is critical for platelet function, regulates erythropoiesis, and is involved in erythroid regeneration after chemotherapy in ALL patients (63, 64). *ARHGEF12* was also reported as a candidate tumor suppressor gene in breast and colorectal cancers (65). Mutations in the coding region of the gene are registered in COSMIC, the catalogue of somatic mutations in cancer (cancer.sanger.ac.uk), in various disease types (66).

In conclusion, we herein report the complex and sequential genetic steps of a therapy-related pediatric B-ALL. The patient was initially diagnosed with AML carrying a t(9;11)(p21;q23) chromosome translocation and a *KMT2A-MLL3* fusion gene. The first secondary genetic step, resulting from chemotherapy that included treatment with topoisomerase II-inhibitors, was a submicroscopic 2 Mb deletion in 11q23 generating a *KMT2A-ARHGEF12* fusion gene in a multipotent progenitor cell. At that point, the patient entered a long “preleukemic” phase which lasted until yet another genetic aberration occurred, t(14;19)(q32;q13) accompanied by the loss of one chromosome 9. This resulted in the development of B-ALL.

Conflicts of Interest

The Authors declare that they have no potential conflicts of interest in regard to this study.

Authors' Contributions

IP designed and supervised the experiments, performed bioinformatics analysis, molecular genetic experiments, evaluated the data, and drafted the manuscript. KA performed cytogenetic, FISH and molecular experiments and evaluated the data. ME-O evaluated the cytogenetic and FISH data. BZ made clinical evaluations and treated the patient. MCM-K made clinical evaluations and treated the patient. JB made clinical evaluations and treated the patient. LTNO performed the flow cytometric evaluations. FM evaluated the cytogenetic and FISH data. SH evaluated the data and assisted with writing of the manuscript. All Authors read and approved the final manuscript.

Acknowledgements

This work was supported by Grants from Radiumhospitalets Legater.

References

- 1 Daser A and Rabbitts TH: The versatile mixed lineage leukaemia gene MLL and its many associations in leukaemogenesis. *Semin Cancer Biol* 15(3): 175-188, 2005. PMID: 15826832. DOI: 10.1016/j.semcancer.2005.01.007
- 2 Marschalek R: Mechanisms of leukemogenesis by MLL fusion proteins. *Br J Haematol* 152(2): 141-154, 2011. PMID: 21118195. DOI: 10.1111/j.1365-2141.2010.08459.x
- 3 Mitelman F, Johansson B and Mertens F: Mitelman Database of Chromosome Aberrations and Gene Fusions in Cancer, 2020. Available at: <https://mitelmandatabase.isb-cgc.org/>
- 4 Meyer C, Hofmann J, Burmeister T, Groger D, Park TS, Emerenciano M, Pombo de Oliveira M, Renneville A, Villares P, Macintyre E, Cave H, Clappier E, Mass-Malo K, Zuna J, Trka J, De Braekeleer E, De Braekeleer M, Oh SH, Tsaour G, Fechina L, van der Velden VH, van Dongen JJ, Delabesse E, Binato R, Silva ML, Kustanovich A, Aleinikova O, Harris MH, Lund-Aho T, Juvonen V, Heidenreich O, Vormoor J, Choi WW, Jarosova M, Kolenova A, Bueno C, Menendez P, Wehner S, Eckert C, Talmant P, Tondeur S, Lippert E, Launay E, Henry C, Ballerini P, Lapillone H, Callanan MB, Cayuela JM, Herbaux C, Cazzaniga G, Kakadiya PM, Bohlander S, Ahlmann M, Choi JR, Gameiro P, Lee DS, Krauter J, Cornillet-Lefebvre P, Te Kronnie G, Schafer BW, Kubetzko S, Alonso CN, zur Stadt U, Sutton R, Venn NC, Izraeli S, Trakhtenbrot L, Madsen HO, Archer P, Hancock J, Cerveira N, Teixeira MR, Lo Nigro L, Moricke A, Stanulla M, Schrappe M, Sedek L, Szczepanski T, Zwaan CM, Coenen EA, van den Heuvel-Eibrink MM, Strehl S, Dworzak M, Panzer-Grumayer R, Dingermann T, Klingebiel T and Marschalek R: The MLL recombinome of acute leukemias in 2013. *Leukemia* 27(11): 2165-2176, 2013. PMID: 23628958. DOI: 10.1038/leu.2013.135
- 5 Meyer C, Burmeister T, Groger D, Tsaour G, Fechina L, Renneville A, Sutton R, Venn NC, Emerenciano M, Pombo-de-Oliveira MS, Barbieri Blunck C, Almeida Lopes B, Zuna J, Trka J, Ballerini P, Lapillone H, De Braekeleer M, Cazzaniga G, Corral Abascal L, van der Velden VHJ, Delabesse E, Park TS, Oh SH, Silva MLM, Lund-Aho T, Juvonen V, Moore AS, Heidenreich O, Vormoor J, Zerkalenkova E, Olshanskaya Y, Bueno C, Menendez P, Teigler-Schlegel A, Zur Stadt U, Lentes J, Gohring G, Kustanovich A, Aleinikova O, Schafer BW, Kubetzko S, Madsen HO, Gruhn B, Duarte X, Gameiro P, Lippert E, Bidet A, Cayuela JM, Clappier E, Alonso CN, Zwaan CM, van den Heuvel-Eibrink MM, Izraeli S, Trakhtenbrot L, Archer P, Hancock J, Moricke A, Alten J, Schrappe M, Stanulla M, Strehl S, Attarbaschi A, Dworzak M, Haas OA, Panzer-Grumayer R, Sedek L, Szczepanski T, Caye A, Suarez L, Cave H and Marschalek R: The MLL recombinome of acute leukemias in 2017. *Leukemia* 32(2): 273-284, 2018. PMID: 28701730. DOI: 10.1038/leu.2017.213
- 6 Heim S and Mitelman F: *Cancer Cytogenetics: Chromosomal and Molecular Genetic Abberations of Tumor Cells*. Fourth Edition edn. Wiley-Blackwell, 2015.
- 7 Winters AC and Bernt KM: MLL-rearranged leukemias-an update on science and clinical approaches. *Front Pediatr* 5: 4, 2017. PMID: 28232907. DOI: 10.3389/fped.2017.00004
- 8 De Braekeleer E, Meyer C, Douet-Guilbert N, Morel F, Le Bris MJ, Marschalek R, Ferec C and De Braekeleer M: A complex 1;19;11 translocation involving the MLL gene in a patient with congenital acute monoblastic leukemia identified by molecular and cytogenetic techniques. *Ann Hematol* 88(8): 795-797, 2009. PMID: 19107484. DOI: 10.1007/s00277-008-0656-8
- 9 De Braekeleer E, Meyer C, Douet-Guilbert N, Basinko A, Le Bris MJ, Morel F, Berthou C, Marschalek R, Ferec C and De Braekeleer M: Identification of MLL partner genes in 27 patients with acute leukemia from a single cytogenetic laboratory. *Mol Oncol* 5(6): 555-563, 2011. PMID: 21900057. DOI: 10.1016/j.molonc.2011.08.003
- 10 De Braekeleer E, Douet-Guilbert N, Meyer C, Morel F, Marschalek R and De Braekeleer M: MLL-ELL fusion gene in two infants with acute monoblastic leukemia and myeloid sarcoma. *Leuk Lymphoma* 53(6): 1222-1224, 2012. PMID: 22149207. DOI: 10.3109/10428194.2011.648632
- 11 Panagopoulos I, Gorunova L, Kerndrup G, Spetalen S, Tierens A, Osnes LT, Andersen K, Muller LS, Hellebostad M, Zeller B and Heim S: Rare MLL-ELL fusion transcripts in childhood acute myeloid leukemia-association with young age and myeloid sarcomas? *Exp Hematol Oncol* 5: 8, 2015. PMID: 26949571. DOI: 10.1186/s40164-016-0037-2
- 12 Zwaan CM, Kolb EA, Reinhardt D, Abrahamsson J, Adachi S, Aplenc R, De Bont ES, De Moerloose B, Dworzak M, Gibson BE, Hasle H, Leverger G, Locatelli F, Ragu C, Ribeiro RC, Rizzari C, Rubnitz JE, Smith OP, Sung L, Tomizawa D, van den Heuvel-Eibrink MM, Creutzig U and Kaspers GJ: Collaborative efforts driving progress in pediatric acute myeloid leukemia. *J Clin Oncol* 33(27): 2949-2962, 2015. PMID: 26304895. DOI: 10.1200/JCO.2015.62.8289
- 13 Toft N, Birgens H, Abrahamsson J, Griskevicius L, Hallbook H, Heyman M, Klausen TW, Jonsson OG, Palk K, Pruunsild K, Quist-Paulsen P, Vaitkeviciene G, Vettenranta K, Asberg A, Frandsen TL, Marquart HV, Madsen HO, Noren-Nystrom U and Schmiegelow K: Results of NOPHO ALL2008 treatment for patients aged 1-45 years with acute lymphoblastic leukemia. *Leukemia* 32(3): 606-615, 2018. PMID: 28819280. DOI: 10.1038/leu.2017.265

- 14 Panagopoulos I, Gorunova L, Torkildsen S, Tjonnfjord GE, Micci F and Heim S: DEK-NUP214-fusion identified by RNA-sequencing of an acute myeloid leukemia with t(9;12)(q34;q15). *Cancer Genomics Proteomics* 14(6): 437-443, 2017. PMID: 29109093. DOI: 10.21873/cgp.20053
- 15 Schroeder A, Mueller O, Stocker S, Salowsky R, Leiber M, Gassmann M, Lightfoot S, Menzel W, Granzow M and Ragg T: The RIN: an RNA integrity number for assigning integrity values to RNA measurements. *BMC Mol Biol* 7: 3, 2006. PMID: 16448564. DOI: 10.1186/1471-2199-7-3
- 16 Torkildsen S, Brunetti M, Gorunova L, Spetalen S, Beiske K, Heim S and Panagopoulos I: Rearrangement of the chromatin organizer special AT-rich binding protein 1 gene, SATB1, resulting from a t(3;5)(p24;q14) chromosomal translocation in acute myeloid leukemia. *Anticancer Res* 37(2): 693-698, 2017. PMID: 28179318. DOI: 10.21873/anticancer.11365
- 17 Kangaspekka S, Hultsch S, Edgren H, Nicorici D, Murumagi A and Kallioniemi O: Reanalysis of RNA-sequencing data reveals several additional fusion genes with multiple isoforms. *PLoS One* 7(10): e48745, 2012. PMID: 23119097. DOI: 10.1371/journal.pone.0048745
- 18 Nicorici D, Satalan H, Edgren H, Kangaspekka S, Murumagi A, Kallioniemi O, Virtanen S and Kikku O: FusionCatcher – a tool for finding somatic fusion genes in paired-end RNA-sequencing data. *bioRxiv*, 2014. DOI: 10.1101/011650
- 19 Singhal H, Ren YR and Kern SE: Improved DNA electrophoresis in conditions favoring polyborates and lewis acid complexation. *PLoS One* 5(6): e11318, 2010. PMID: 20593002. DOI: 10.1371/journal.pone.0011318
- 20 Altschul SF, Gish W, Miller W, Myers EW and Lipman DJ: Basic local alignment search tool. *J Mol Biol* 215(3): 403-410, 1990. PMID: 2231712. DOI: 10.1016/S0022-2836(05)80360-2
- 21 Kent WJ: BLAT – the BLAST-like alignment tool. *Genome Res* 12(4): 656-664, 2002. PMID: 11932250. DOI: 10.1101/gr.229202
- 22 Kent WJ, Sugnet CW, Furey TS, Roskin KM, Pringle TH, Zahler AM and Haussler D: The human genome browser at UCSC. *Genome Res* 12(6): 996-1006, 2002. PMID: 12045153. DOI: 10.1101/gr.229102
- 23 Kourlas PJ, Strout MP, Becknell B, Veronese ML, Croce CM, Theil KS, Krahe R, Ruutu T, Knuutila S, Bloomfield CD and Caligiuri MA: Identification of a gene at 11q23 encoding a guanine nucleotide exchange factor: evidence for its fusion with MLL in acute myeloid leukemia. *Proc Natl Acad Sci USA* 97(5): 2145-2150, 2000. PMID: 10681437. DOI: 10.1073/pnas.040569197
- 24 Shih LY, Liang DC, Fu JF, Wu JH, Wang PN, Lin TL, Dunn P, Kuo MC, Tang TC, Lin TH and Lai CL: Characterization of fusion partner genes in 114 patients with de novo acute myeloid leukemia and MLL rearrangement. *Leukemia* 20(2): 218-223, 2006. PMID: 16341046. DOI: 10.1038/sj.leu.2404024
- 25 Liu YF, Wang BY, Zhang WN, Huang JY, Li BS, Zhang M, Jiang L, Li JF, Wang MJ, Dai YJ, Zhang ZG, Wang Q, Kong J, Chen B, Zhu YM, Weng XQ, Shen ZX, Li JM, Wang J, Yan XJ, Li Y, Liang YM, Liu L, Chen XQ, Zhang WG, Yan JS, Hu JD, Shen SH, Chen J, Gu LJ, Pei D, Li Y, Wu G, Zhou X, Ren RB, Cheng C, Yang JJ, Wang KK, Wang SY, Zhang J, Mi JQ, Pui CH, Tang JY, Chen Z and Chen SJ: Genomic profiling of adult and pediatric B-cell acute lymphoblastic leukemia. *EBioMedicine* 8: 173-183, 2016. PMID: 27428428. DOI: 10.1016/j.ebiom.2016.04.038
- 26 Sevov M, Bunikis I, Häggqvist S, Höglund M, Rosenquist R, Ameer A and Cavellier L: Targeted RNA sequencing assay efficiently identifies cryptic KMT2A (MLL)-fusions in acute leukemia patients. *Blood* 124(21): 2406-2406, 2014. DOI: 10.1182/blood.V124.21.2406.2406
- 27 Blanco JG, Edick MJ and Relling MV: Etoposide induces chimeric Mll gene fusions. *FASEB J* 18(1): 173-175, 2004. PMID: 14630694. DOI: 10.1096/fj.03-0638fje
- 28 Libura J, Slater DJ, Felix CA and Richardson C: Therapy-related acute myeloid leukemia-like MLL rearrangements are induced by etoposide in primary human CD34+ cells and remain stable after clonal expansion. *Blood* 105(5): 2124-2131, 2005. PMID: 15528316. DOI: 10.1182/blood-2004-07-2683
- 29 Felix CA, Kolaris CP and Osheroff N: Topoisomerase II and the etiology of chromosomal translocations. *DNA Repair* 5(9-10): 1093-1108, 2006. PMID: 16857431. DOI: 10.1016/j.dnarep.2006.05.031
- 30 Sung PA, Libura J and Richardson C: Etoposide and illegitimate DNA double-strand break repair in the generation of MLL translocations: new insights and new questions. *DNA Repair* 5(9-10): 1109-1118, 2006. PMID: 16809075. DOI: 10.1016/j.dnarep.2006.05.018
- 31 Cowell IG and Austin CA: Mechanism of generation of therapy related leukemia in response to anti-topoisomerase II agents. *Int J Environ Res Public Health* 9(6): 2075-2091, 2012. PMID: 22829791. DOI: 10.3390/ijerph9062075
- 32 Ezoe S: Secondary leukemia associated with the anti-cancer agent, etoposide, a topoisomerase II inhibitor. *Int J Environ Res Public Health* 9(7): 2444-2453, 2012. PMID: 22851953. DOI: 10.3390/ijerph9072444
- 33 Evison BJ, Sleebs BE, Watson KG, Phillips DR and Cutts SM: Mitoxantrone, more than just another topoisomerase II poison. *Med Res Rev* 36(2): 248-299, 2016. PMID: 26286294. DOI: 10.1002/med.21364
- 34 Pagano L, Pulsoni A, Tosti ME, Annino L, Mele A, Camera A, Martino B, Guglielmi C, Cerri R, Di Bona E, Invernizzi R, Castagnola C, Bassan R, Mele L, Todeschini G, Leone G and Mandelli F: Acute lymphoblastic leukaemia occurring as second malignancy: report of the GIMEMA archive of adult acute leukaemia. *Gruppo Italiano Malattie Ematologiche Maligne dell'Adulto. Br J Haematol* 106(4): 1037-1040, 1999. PMID: 10520009. DOI: 10.1046/j.1365-2141.1999.01636.x
- 35 Ishizawa S, Slovak ML, Popplewell L, Bedell V, Wrede JE, Carter NH, Snyder DS and Arber DA: High frequency of pro-B acute lymphoblastic leukemia in adults with secondary leukemia with 11q23 abnormalities. *Leukemia* 17(6): 1091-1095, 2003. PMID: 12764373. DOI: 10.1038/sj.leu.2402918
- 36 Shivakumar R, Tan W, Wilding GE, Wang ES and Wetzler M: Biologic features and treatment outcome of secondary acute lymphoblastic leukemia – a review of 101 cases. *Ann Oncol* 19(9): 1634-1638, 2008. PMID: 18467310. DOI: 10.1093/annonc/mdn182
- 37 Abdulwahab A, Sykes J, Kamel-Reid S, Chang H and Brandwein JM: Therapy-related acute lymphoblastic leukemia is more frequent than previously recognized and has a poor prognosis. *Cancer* 118(16): 3962-3967, 2012. PMID: 22180297. DOI: 10.1002/ncr.26735
- 38 Tang G, Zuo Z, Thomas DA, Lin P, Liu D, Hu Y, Kantarjian HM, Bueso-Ramos C, Medeiros LJ and Wang SA: Precursor B-acute lymphoblastic leukemia occurring in patients with a history of prior malignancies: is it therapy-related? *Haematologica* 97(6): 919-925, 2012. PMID: 22207681. DOI: 10.3324/haematol.2011.057752

- 39 Douet-Guilbert N, Eveillard JR, Meyer C, Ugo V, Le Bris MJ, Basinko A, Morel F, Marschalek R and De Braekeleer M: MLL partner genes in secondary acute lymphoblastic leukemia: report of a new partner PRRC1 and review of the literature. *Leuk Res* 38(11): 1316-1319, 2014. PMID: 25205603. DOI: 10.1016/j.leukres.2014.08.011
- 40 Aldoss I, Douer D and Pullarkat V: Therapy-related acute lymphoblastic leukemia: Where do we stand with regards to its definition and characterization? *Blood Rev* 37: 100584, 2019. PMID: 31235377. DOI: 10.1016/j.blre.2019.06.001
- 41 Moneypenny CG, Shao J, Song Y and Gallagher EP: MLL rearrangements are induced by low doses of etoposide in human fetal hematopoietic stem cells. *Carcinogenesis* 27(4): 874-881, 2006. PMID: 16377807. DOI: 10.1093/carcin/bgi322
- 42 Libura J, Ward M, Solecka J and Richardson C: Etoposide-initiated MLL rearrangements detected at high frequency in human primitive hematopoietic stem cells with *in vitro* and *in vivo* long-term repopulating potential. *Eur J Haematol* 81(3): 185-195, 2008. PMID: 18510699. DOI: 10.1111/j.1600-0609.2008.01103.x
- 43 Bueno C, Catalina P, Melen GJ, Montes R, Sanchez L, Ligerio G, Garcia-Perez JL and Menendez P: Etoposide induces MLL rearrangements and other chromosomal abnormalities in human embryonic stem cells. *Carcinogenesis* 30(9): 1628-1637, 2009. PMID: 19587093. DOI: 10.1093/carcin/bgp169
- 44 Teuffel O, Betts DR, Thali M, Eberle D, Meyer C, Schneider B, Marschalek R, Trakhtenbrot L, Amariglio N, Niggli FK and Schafer BW: Clonal expansion of a new MLL rearrangement in the absence of leukemia. *Blood* 105(10): 4151-4152, 2005. PMID: 15867425. DOI: 10.1182/blood-2005-01-0286
- 45 Mori T, Nishimura N, Hasegawa D, Kawasaki K, Kosaka Y, Uchide K, Yanai T, Hayakawa A, Takeshima Y, Nishio H and Matsuo M: Persistent detection of a novel MLL-SACM1L rearrangement in the absence of leukemia. *Leuk Res* 34(10): 1398-1401, 2010. PMID: 20553989. DOI: 10.1016/j.leukres.2010.05.001
- 46 Jeffries SJ, Jones L, Harrison CJ and Russell LJ: IGH@ translocations co-exist with other primary rearrangements in B-cell precursor acute lymphoblastic leukemia. *Haematologica* 99(8): 1334-1342, 2014. PMID: 24816234. DOI: 10.3324/haematol.2014.103820
- 47 Zuna J, Burjanivova T, Mejstrikova E, Zemanova Z, Muzikova K, Meyer C, Horsley SW, Kearney L, Colman S, Ptoszkova H, Marschalek R, Hrusak O, Sary J, Greaves M and Trka J: Covert preleukemia driven by MLL gene fusion. *Genes Chromosomes Cancer* 48(1): 98-107, 2009. PMID: 18932267. DOI: 10.1002/gcc.20622
- 48 Jonveaux P, Hillion J, Bernard O, Le Coniat M, Derre J, Flexor M, Larsen CJ and Berger R: Distinct MLL gene rearrangements associated with successive acute monocytic and lymphoblastic leukemias in the same patient. *Leukemia* 8(12): 2224-2227, 1994. PMID: 7808011.
- 49 Emerenciano M, Meyer C, Mansur MB, Marschalek R, Pombo-de-Oliveira MS and Brazilian Collaborative Study Group of Infant Acute Leukemia: The distribution of MLL breakpoints correlates with outcome in infant acute leukaemia. *Br J Haematol* 161(2): 224-236, 2013. PMID: 23432364. DOI: 10.1111/bjh.12250
- 50 Meyer C, Lopes BA, Caye-Eude A, Cave H, Arfeuille C, Cuccuini W, Sutton R, Venn NC, Oh SH, Tsaur G, Escherich G, Feuchtinger T, Kosasih HJ, Khaw SL, Ekert PG, Pombo-de-Oliveira MS, Bidet A, Djahanschiri B, Ebersberger I, Zaliova M, Zuna J, Zermanova Z, Juvonen V, Grumayer RP, Fazio G, Cazzaniga G, Larghero P, Emerenciano M and Marschalek R: Human MLL/KMT2A gene exhibits a second breakpoint cluster region for recurrent MLL-USP2 fusions. *Leukemia* 33(9): 2306-2340, 2019. PMID: 30899083. DOI: 10.1038/s41375-019-0451-7
- 51 Rössler T and Marschalek R: An alternative splice process renders the MLL protein either into a transcriptional activator or repressor. *Pharmazie* 68(7): 601-607, 2013. PMID: 23923644. DOI: 10.1055/s-0033-1343653
- 52 Fukuhara S, Chikumi H and Gutkind JS: Leukemia-associated Rho guanine nucleotide exchange factor (LARG) links heterotrimeric G proteins of the G(12) family to Rho. *FEBS Lett* 485(2-3): 183-188, 2000. PMID: 11094164. DOI: 10.1016/S0014-5793(00)02224-9
- 53 Booden MA, Siderovski DP and Der CJ: Leukemia-associated Rho guanine nucleotide exchange factor promotes G alpha q-coupled activation of RhoA. *Mol Cell Biol* 22(12): 4053-4061, 2002. PMID: 12024019. DOI: 10.1128/mcb.22.12.4053-4061.2002
- 54 Hikumi H, Fukuhara S and Gutkind JS: Regulation of G protein-linked guanine nucleotide exchange factors for Rho, PDZ-RhoGEF, and LARG by tyrosine phosphorylation: evidence of a role for focal adhesion kinase. *J Biol Chem* 277(14): 12463-12473, 2002. PMID: 11799111. DOI: 10.1074/jbc.M108504200
- 55 Swiercz JM, Kuner R, Behrens J and Offermanns S: Plexin-B1 directly interacts with PDZ-RhoGEF/LARG to regulate RhoA and growth cone morphology. *Neuron* 35(1): 51-63, 2002. PMID: 12123608. DOI: 10.1016/S0896-6273(02)00750-x
- 56 Dubash AD, Wennerberg K, Garcia-Mata R, Menold MM, Arthur WT and Burridge K: A novel role for Lsc/p115 RhoGEF and LARG in regulating RhoA activity downstream of adhesion to fibronectin. *J Cell Sci* 120(Pt 22): 3989-3998, 2007. PMID: 17971419. DOI: 10.1242/jcs.003806
- 57 Kitzing TM, Sahadevan AS, Brandt DT, Knieling H, Hannemann S, Fackler OT, Grosshans J and Grosse R: Positive feedback between Dial1, LARG, and RhoA regulates cell morphology and invasion. *Genes Dev* 21(12): 1478-1483, 2007. PMID: 17575049. DOI: 10.1101/gad.424807
- 58 Goulimari P, Knieling H, Engel U and Grosse R: LARG and mDial1 link Galpha12/13 to cell polarity and microtubule dynamics. *Mol Biol Cell* 19(1): 30-40, 2008. PMID: 17959834. DOI: 10.1091/mbc.e06-11-1045
- 59 Evelyn CR, Ferng T, Rojas RJ, Larsen MJ, Sondek J and Neubig RR: High-throughput screening for small-molecule inhibitors of LARG-stimulated RhoA nucleotide binding *via* a novel fluorescence polarization assay. *J Biomol Screen* 14(2): 161-172, 2009. PMID: 19196702. DOI: 10.1177/1087057108328761
- 60 Shi Y, Zhang J, Mullin M, Dong B, Alberts AS and Siminovich KA: The mDial1 formin is required for neutrophil polarization, migration, and activation of the LARG/RhoA/ROCK signaling axis during chemotaxis. *J Immunol* 182(6): 3837-3845, 2009. PMID: 19265163. DOI: 10.4049/jimmunol.0803838
- 61 Martz MK, Grabocka E, Beeharry N, Yen TJ and Wedegaertner PB: Leukemia-associated RhoGEF (LARG) is a novel RhoGEF in cytokinesis and required for the proper completion of abscission. *Mol Biol Cell* 24(18): 2785-2794, 2013. PMID: 23885121. DOI: 10.1091/mbc.E12-07-0533
- 62 Guilluy C, Swaminathan V, Garcia-Mata R, O'Brien ET, Superfine R and Burridge K: The Rho GEFs LARG and GEF-H1 regulate the mechanical response to force on integrins. *Nat Cell Biol* 13(6): 722-727, 2011. PMID: 21572419. DOI: 10.1038/ncb2254

- 63 Zou S, Teixeira AM, Yin M, Xiang Y, Xavier-Ferrucio J, Zhang PX, Hwa J, Min W and Krause DS: Leukaemia-associated Rho guanine nucleotide exchange factor (LARG) plays an agonist specific role in platelet function through RhoA activation. *Thromb Haemost* 116(3): 506-516, 2016. PMID: 27345948. DOI: 10.1160/TH15-11-0848
- 64 Xie Y, Gao L, Xu C, Chu L, Gao L, Wu R, Liu Y, Liu T, Sun XJ, Ren R, Tang J, Zheng Y, Zhou Y and Shen S: ARHGEF12 regulates erythropoiesis and is involved in erythroid regeneration after chemotherapy in acute lymphoblastic leukemia patients. *Haematologica* 105(4): 925-936, 2020. PMID: 31467124. DOI: 10.3324/haematol.2018.210286
- 65 Ong DC, Ho YM, Rudduck C, Chin K, Kuo WL, Lie DK, Chua CL, Tan PH, Eu KW, Seow-Choen F, Wong CY, Hong GS, Gray JW and Lee AS: LARG at chromosome 11q23 has functional characteristics of a tumor suppressor in human breast and colorectal cancer. *Oncogene* 28(47): 4189-4200, 2009. PMID: 19734946. DOI: 10.1038/onc.2009.266
- 66 Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, Boutselakis H, Cole CG, Creatore C, Dawson E, Fish P, Harsha B, Hathaway C, Jupe SC, Kok CY, Noble K, Ponting L, Ramshaw CC, Rye CE, Speedy HE, Stefancsik R, Thompson SL, Wang S, Ward S, Campbell PJ and Forbes SA: COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Res* 47(D1): D941-D947, 2019. PMID: 30371878. DOI: 10.1093/nar/gky1015

Received September 30, 2020

Revised October 25, 2020

Accepted October 27, 2020