

Initial Guidance on Use of Monoclonal Antibody Therapy for Treatment of COVID-19 in Children and Adolescents

Joshua Wolf^{1,2}, Mark J. Abzug³, Rachel L. Wattier⁴, Paul K. Sue⁵, Surabhi B. Vora⁶, Philip Zachariah⁷, Daniel E. Dulek⁸, Alpana Waghmare^{6,9}, Rosemary Olivero¹⁰, Kevin J. Downes¹¹, Scott H. James¹², Swetha G. Pinninti¹², April Yarbrough¹³, Margaret L. Aldrich¹⁴, Christine E. MacBrayne¹⁵, Vijaya L. Soma¹⁶, Steven P. Grapentine¹⁷, Carlos R. Oliveira¹⁸, Molly Hayes¹⁹, David W. Kimberlin²⁰, Sarah B. Jones²¹, Laura L. Bio²², Theodore H. Morton¹, Jane S. Hankins²³, Gabriella M. Marón-Alfaro¹, Kathryn Timberlake²⁴, Jennifer L. Young²⁵, Rachel C. Orscheln²⁶, Hayden T. Schwenk²⁷, David L. Goldman¹⁴, Helen E. Groves²⁸, W. Charles Huskins²⁹, Nipunie S. Rajapakse²⁹, Gabriella S. Lamb³⁰, Alison C. Tribble³¹, Elizabeth E. Lloyd³¹, Adam L. Hersh³², Emily A. Thorell³², Adam J. Ratner^{16,33}, Kathleen Chiotos^{11,34} and Mari M. Nakamura^{30,35}

1. Department of Infectious Diseases, St. Jude Children's Research Hospital, Memphis, Tennessee, USA; 2. Department of Pediatrics, University of Tennessee Health Science Center, Memphis, Tennessee, USA; 3. Division of Infectious Diseases, Department of Pediatrics, University of Colorado School of Medicine and Children's Hospital Colorado, Aurora, Colorado, USA; 4. Division of Infectious Diseases and Global Health, Department of Pediatrics, University of California–San Francisco, San Francisco, California, USA; 5. Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Texas Southwestern Medical Center, Dallas, TX, USA; 6. Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Washington, Seattle; Children's Hospital, Seattle, Washington, USA; 7. Division of Infectious Diseases, Department of Pediatrics, Columbia University, New York, New York, USA; 8. Division of Infectious Diseases, Department of Pediatrics, Vanderbilt University and Monroe Carell Jr. Children's Hospital, Nashville, Tennessee, USA; 9. Vaccine and Infectious Diseases Division, Fred Hutchinson Cancer Research Center, Seattle, Washington, USA; 10. Section of Infectious Diseases, Department of Pediatrics and Human Development, Helen DeVos

© The Author(s) 2021. Published by Oxford University Press on behalf of The Journal of the Pediatric Infectious Diseases Society. All rights reserved. For permissions, please e-mail: journals.permissions@oup.com.

Children's Hospital of Spectrum Health, Michigan State College of Human Medicine, Grand Rapids, Michigan, USA; 11. Division of Infectious Diseases, Department of Pediatrics, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA; 12. Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Alabama at Birmingham, Birmingham, AL, United States; 13. Department of Pharmacy, Children's of Alabama, Birmingham, Alabama, USA; 14. Division of Infectious Diseases, Department of Pediatrics, Children's Hospital at Montefiore, New York, New York, USA; 15. Department of Pharmacy, Children's Hospital Colorado, Aurora, Colorado, USA; 16. Department of Pediatrics, Division of Infectious Diseases, New York University Grossman School of Medicine and Hassenfeld Children's Hospital, New York, United States; 17. Department of Pharmacy, University of California–San Francisco Benioff Children's Hospital, San Francisco, California, USA; 18. Division of Infectious Diseases and Global Health, Department of Pediatrics, Yale University School of Medicine, New Haven, CT, United States; 19. Antimicrobial Stewardship Program, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA; 20. Division of Pediatric Infectious Diseases, Department of Pediatrics, University of Alabama at Birmingham, Birmingham, Alabama, USA; 21. Department of Pharmacy, Boston Children's Hospital, Boston, Massachusetts, USA; 22. Department of Pharmacy, Lucile Packard Children's Hospital Stanford, Palo Alto, California, USA; 23. Department of Hematology, St. Jude Children's Research Hospital, Memphis, Tennessee, USA; 24. Department of Pharmacy, The Hospital for Sick Children, Toronto, Ontario, Canada; 25. Department of Pharmacy, St. Louis Children's Hospital, St. Louis, Missouri, USA; 26. Division of Infectious Diseases, Department of Pediatrics, Washington University and St. Louis Children's Hospital, St. Louis, Missouri, USA; 27. Division of Infectious Diseases, Department of Pediatrics, Stanford University School of Medicine and Lucile Packard Children's Hospital Stanford, Stanford, California, USA; 28. Division of Infectious Diseases, Department of Pediatrics,; Hospital for Sick Children, Toronto, Ontario, Canada; 29. Division of Pediatric Infectious Diseases, Department of Pediatric and Adolescent Medicine, Mayo Clinic College of Medicine and Science, Rochester, Minnesota; 30. Division of Infectious Diseases, Department of Pediatrics, Boston Children's Hospital, Boston, Massachusetts, USA; 31. Department

of Pediatrics, Division of Infectious Diseases, University of Michigan and CS Mott Children's Hospital, Ann Arbor, Michigan, USA; 32. Division of Infectious Diseases, Department of Pediatrics, University of Utah and Primary Children's Hospital, Salt Lake City, Utah, USA; 33. Department of Microbiology, New York University Grossman School of Medicine, New York, New York, USA; 34. Division of Critical Care Medicine, Department of Anesthesia and Critical Care Medicine, Children's Hospital of Philadelphia, Philadelphia, Pennsylvania, USA; 35. Antimicrobial Stewardship Program, Boston Children's Hospital, Boston, Massachusetts, USA

Summary: Although monoclonal antibody therapies for COVID-19 have received emergency use authorization for treatment of adolescents in specified high risk categories, there is currently insufficient evidence for necessity, safety or efficacy to recommend routine use, even in those with specified conditions.

Conflicts of interest: JW reports support to his institution from Karius Inc., Merck Inc. and Astellas Inc. for participation in sponsored research unrelated to this work; KJD reports support to his institution from Merck Inc. for participation in sponsored research unrelated to this work; SHJ reports payment for work as a consultant to Bayer unrelated to this work; PKS reports support to his institution from Merck Inc., Allovir Inc. and Gilead Sciences Inc. for participation in sponsored research unrelated to this work; WCH reports payment for work as a member of an Advisory Board to ADMA Biologics and a member of an Adjudication Committee for Pfizer, both unrelated to this work; DWK reports support to his institution from Gilead Sciences Inc. for participation in sponsored research unrelated to this work; AW reports support from Kyorin Pharmaceutical as an Advisory Board Member, support to her institution from Ansun Biopharma and Allovir Inc. for participation in sponsored research, and research support from VB Tech and Amazon Inc. all unrelated to this work.; RLW reports employment of a family member by Lucence Diagnostics unrelated to this work; JSH reports payment for work as a consultant to Global Blood Therapeutics and MJH Lifesciences, and

research support from Global Blood Therapeutics, all unrelated to this work; AJR reports payment for work as a consultant to Pfizer Inc. unrelated to this work. No other conflicts of interest were reported.

Corresponding author:

Dr. Joshua Wolf MBBS, PhD, FRACP

Department of Infectious Diseases

St. Jude Children's Research Hospital

262 Danny Thomas Place, MS 320

Memphis, TN 38105, USA

Tel: 901 595 3300; Fax: 901 595 3099

Email: Joshua.Wolf@stjude.org

Co-Corresponding author:

Mari M. Nakamura, MD, MPH

Division of Infectious Diseases

Department of Pediatrics

Boston Children's Hospital,

300 Longwood Avenue, Mailstop BCH 3052

Boston, MA 02115

Tel: 617 355 1561

Email: mari.nakamura@childrens.harvard.edu

Abstract

Background. In November 2020, the US Food and Drug Administration (FDA) provided Emergency Use Authorizations (EUA) for two novel virus-neutralizing monoclonal antibody therapies, bamlanivimab, and REGN-COV2 (casirivimab plus imdevimab), for the treatment of mild to moderate COVID-19 in adolescents and adults in specified high-risk groups. This has challenged clinicians to determine the best approach to use of these products.

Methods. A panel of experts in pediatric infectious diseases, pediatric infectious diseases pharmacy, pediatric intensive care medicine, and pediatric hematology from 29 geographically diverse North American institutions was convened. Through a series of teleconferences and web-based surveys, a guidance statement was developed and refined based on review of the best available evidence and expert opinion.

Results. The course of COVID-19 in children and adolescents is typically mild and there is no high-quality evidence supporting any high risk groups. There is no evidence for safety and efficacy of monoclonal antibody therapy for treatment of COVID-19 in children or adolescents, limited evidence of modest benefit in adults, and evidence for potential harm associated with infusion reactions or anaphylaxis.

Conclusions. Based on evidence available as of December 20, 2020, the panel suggests against routine administration of monoclonal antibody therapy (bamlanivimab, or casirivimab and imdevimab), for treatment of COVID-19 in children or adolescents, including those designated by the FDA as at high risk of progression to hospitalization or severe disease. Clinicians and health systems choosing to use these agents on an individualized basis should consider risk factors supported by pediatric-specific evidence, and ensure implementation of a system for safe and timely administration that does not exacerbate existing healthcare disparities.

Keywords: COVID-19; Pediatric; Bamlanivimab; Casirivimab; Imdevimab

Accepted Manuscript

1. Introduction

The pandemic of Coronavirus disease 2019 (COVID-19) caused by severe acute respiratory syndrome coronavirus-2 (SARS-CoV-2) has been met with rapid development of novel therapeutics.[1] As new therapies are discovered and become available for use, careful evaluation of evidence is essential to provide guidance for safe and effective use.[2-5] The most recent additions to the COVID-19 armamentarium are virus-neutralizing monoclonal antibodies, which are human or humanized antibodies administered by intravenous infusion that bind to virus or to infected cells to treat SARS-CoV-2 infection.[6] The Food and Drug Administration (FDA) has issued Emergency Use Authorizations (EUA) for two such products, bamlanivimab (LY-CoV555) and REGN-COV2, a combination of casirivimab and imdevimab.[7, 8] Another monoclonal antibody, etesevimab (LY-CoV016), has not yet received authorization and will not be discussed further here.[9] In this guidance statement, we focus on the use of available SARS-CoV-2 neutralizing monoclonal antibodies in children and adolescents with COVID-19.

Bamlanivimab is a single monoclonal antibody that binds to the receptor binding domain (RBD) of the SARS-CoV-2 spike protein; it was authorized for use in the USA under an EUA on November 9, 2020.[8] Casirivimab and imdevimab (REGN-COV2) are monoclonal antibodies that bind to non-overlapping regions of the SARS-CoV-2 RBD, and were authorized for use in combination under an EUA on November 21, 2020.[7] Because of the rapid advancement of these products from discovery to clinical use, there is little published evidence about their use in humans, and no evidence for use in children or adolescents. We summarize available clinical data for each product below. In both cases, the EUAs allow use in pediatric patients ≥ 12 years of age and ≥ 40 kg with mild to moderate COVID-19, not requiring hospitalization or new/increased supplemental oxygen for COVID-19, who are deemed to be “at high risk for progressing to severe COVID-19 and/or hospitalization”.[7, 8] Neither agent is authorized for use in patients hospitalized or requiring supplemental oxygen

therapy for COVID-19 because of evidence that these products might cause harm in that setting.[7, 8] The criteria for those at high risk are defined identically in the EUA for each monoclonal antibody treatment and are also discussed below.

To develop guidance on the potential use of these agents for treatment of mild to moderate COVID-19 in high risk adolescents and young adults, as authorized by the current EUAs, we assembled a panel of experts in pediatric infectious diseases, pediatric infectious diseases pharmacy, pediatric intensive care medicine, and pediatric hematology from 29 geographically diverse North American institutions to evaluate the evidence for safety and efficacy of these agents in pediatric patients. This consensus statement has been reviewed and approved by all members, and endorsed by the Pediatric Infectious Diseases Society.

2. Guidance statement

Statement: Based on evidence available as of December 20, 2020, the panel suggests against routine administration of monoclonal antibody therapy (bamlanivimab, or casirivimab and imdevimab) for treatment of COVID-19 in children or adolescents, including those designated by the FDA as at high risk of progression to hospitalization or severe disease.

Remark: Clinicians and health systems choosing to use these agents on an individualized basis should consider risk factors supported by pediatric-specific evidence, and ensure implementation of a system for safe and timely administration that does not exacerbate existing healthcare disparities.

3. Rationale

This guidance statement is based primarily on the current lack of efficacy or safety data in pediatric patients, the generally lower risk of progression to severe disease in children and adolescents, and the apparently modest efficacy of these treatments in adults. Neither agent is authorized for use in patients hospitalized or requiring supplemental oxygen therapy for COVID-19 because of evidence that these products might cause harm in that setting.[7, 8] In this context, along with documented adverse events in adult studies, and plausibility for differential efficacy or safety in younger patients, the potential costs and risks of administration of these products might outweigh the benefits even in children or adolescents designated as being at higher risk of hospitalization or progression to severe disease. Moreover, while the FDA authorized use for adolescents only with specific comorbidities, there are neither sufficient data to support a high risk of severe illness in any pediatric population, nor comparative data to inform risk stratification across the identified groups. Similar reasoning may also apply to young adults, who are also at lower risk of severe disease than older adults, and for whom there are insufficient data to identify potentially high-risk groups and limited evidence for efficacy and safety of monoclonal antibody therapy.

The intent of this guidance is not to preclude the use of these agents in any pediatric patients, but to clarify that routine or standard use in patients meeting EUA criteria is not justified by currently available evidence. The term “suggest” is used to indicate that the panel concluded that the risks of routine use might outweigh the benefits, but that evidence is limited and guidance could change as more data become available. Individual clinicians or institutions may choose to administer these agents to children and adolescents who meet EUA criteria on a case-by-case basis, and should ensure appropriate infrastructure exists to support use of these agents, including a mechanism to rapidly test and treat eligible patients, ability to maintain strict infection control precautions in the

ambulatory setting, and an allocation system that is equitable and does not exacerbate healthcare disparities, particularly while resources are scarce.

4. Evidence Summary

Bamlanivimab

Available clinical evidence for bamlanivimab comprises published interim results from the Blocking Viral Attachment and Cell Entry with SARS-CoV-2 Neutralizing Antibodies (BLAZE-1) trial, a randomized, double-blind, placebo-controlled phase II study conducted at 41 US centers.[8, 10, 11] Participants were required to have positive SARS-CoV-2 viral testing and symptoms of mild or moderate COVID-19. This interim analysis compared 3 treatment groups receiving a single infusion of bamlanivimab at a dose of 700 mg, 2800 mg, or 7000 mg (101, 107, and 101 participants allocated, respectively) against a placebo group (143 participants). Participants evaluated in the interim analysis were all ≥ 18 years of age, with a median age of approximately 45. Almost 70% of subjects had some risk factor for severe disease, defined as age ≥ 65 years, body-mass index (BMI) ≥ 35 kg/m², or a comorbid condition.[10]

The primary outcome was change in log viral load from baseline to day 11 after the positive SARS-CoV-2 test. The study population, including the placebo group, had an overall mean decrease of -3.81 (6.36 at baseline, 2.56 at day 11). Compared with the placebo group, the 2800-mg dose group showed a statistically significant difference of -0.53 (95%CI -0.98 to -0.08; $p=0.02$). A smaller, non-significant difference was observed for participants who received the 700-mg dose specified in the EUA (-0.20, 95%CI -0.66 to 0.25, $p=0.38$), whereas the decrease experienced by the 7000-mg dose group was less than that of the placebo group and not statistically significant (0.09, 95% CI -0.37 to 0.55; $p=0.70$). The major clinical outcome was COVID-19 related hospitalization, emergency

department (ED) visit, or death. The rate of this outcome was 1.6% (5 of 309) among treated subjects versus 6.3% (9 of 143) among placebo recipients, with a number needed to treat (NNT) of 21. In a *post hoc* analysis of subjects who were ≥ 65 years of age and/or had a BMI ≥ 35 kg/m², the rates of hospitalization or ED visit were 4% (4 of 95) for those who received bamlanivimab and 15% (7 of 48) for those who received placebo, with an NNT of 10. No other results stratified by risk factors were reported.[10] There was no reported evaluation of effect on risk of severe or life-threatening illness. None of the bamlanivimab recipients experienced serious adverse events, and the most common adverse event in the bamlanivimab was nausea (3.9%). Infusion reactions were reported in 2.3% of bamlanivimab recipients and 1.4% of placebo recipients; most reactions were described as mild, but at least one case of anaphylaxis has been reported.[10, 11] To date, there are no reports of the safety, efficacy or use of bamlanivimab in children or adolescents.

Casirivimab and imdevimab (REGN-COV2)

Current clinical evidence for casirivimab and imdevimab is described in published interim results from a continual enrollment, multicenter, randomized, double-blind, placebo-controlled, phase 1-2 clinical trial.[7, 12-18] Data were presented for 275 participants, among whom 93 received placebo, 92 low-dose, and 90 high-dose casirivimab and imdevimab.[18] A larger number of participants (799) are reported in a press release, however, these data are not yet published.[13, 16]

Compared to placebo, there was a greater time weighted average reduction in viral load by day 7 for those treated with casirivimab and imdevimab (difference 0.56 log₁₀ copies/mL). *Post hoc* analysis demonstrated that participants who had a higher viral load and seronegative status at baseline appeared to have a larger reduction in viral load.[16, 18] Treatment with casirivimab and imdevimab reduced the absolute risk of need for a medically attended visit by 3% (3% vs 6%) overall with an NNT of 33, and by 9% (6% vs. 15%) in seronegative participants with an NNT of 11.[18] In data

presented in the FDA EUA, treatment with casirivimab and imdevimab reduced the absolute risk of COVID-19-related ER visits or hospitalization by 2% (2% vs. 4%; $p=0.078$) overall with an NNT of 50, and by 6% (3% vs. 9%; $p=0.049$) in participants considered to be at high risk with an NNT of 17.[13, 16]

Administration of combination casirivimab and imdevimab was reported to be safe, with few serious adverse events. However, adverse events in those who received the combination treatment included infusion reactions and anaphylaxis. [7, 12-18] To date, there are no reports of the efficacy, safety, or use of casirivimab or imdevimab in children or adolescents.

COVID-19 Pediatric High-Risk Groups

Under the EUAs, both monoclonal antibody products are authorized for use only in patients with mild to moderate COVID-19 who are “at high risk for progressing to severe COVID-19 and/or hospitalization”. [7, 8] The specified risk categories relevant to pediatric patients include: obesity (BMI ≥ 35 , or $\geq 85^{\text{th}}$ centile for age in adolescents); chronic kidney disease; diabetes; immunosuppressive disease or immunosuppressive treatment; sickle cell disease; congenital or acquired heart disease; neurodevelopmental disorder; medical-related technology dependence; and asthma, reactive airway disease, or other chronic respiratory disease that requires daily medication for control. [11, 16]

In general, children and adolescents are at relatively low risk for hospitalization or severe disease with COVID-19, with around 7% requiring hospitalization and 2% requiring admission to intensive care. [19] Because the overall rate of adverse infection outcomes is low, even those with a higher relative risk of severe disease may remain at low absolute risk. Therefore, data suggesting that a child or adolescent is at a relatively higher risk would not necessarily be sufficient to indicate use of

an untested intervention designed to prevent progression of disease, if the number needed to treat remains high.

We evaluated available evidence on criteria that might confer increased risk of severe disease in children and adolescents. Severe disease was not defined in the EUAs; in our analysis we considered 'severe COVID-19' as disease requiring supplemental oxygen or ICU admission.[3] We reviewed data from published literature where available, and registry data if sufficient published data were lacking. With respect to chronic respiratory conditions, we considered asthma separately from other conditions (e.g. cystic fibrosis or bronchiectasis). And, we interpreted "medical device dependence" to be a surrogate for 'medical complexity' as discussed in our earlier guidance, although this may not hold true for some patients, such as those with only gastrostomy tubes.[3]

There is limited observational evidence suggesting that some of the specified conditions, including obesity[19-30], profound immunocompromise or hypogammaglobulinemia[31-39], chronic cardiac disease[19, 21, 22, 34, 40-46], neurodevelopmental disorders or 'medical complexity'[19, 22, 34, 43, 45, 47, 48], and sickle cell disease[47, 49-58] increase the risk of hospitalization or severe COVID-19 in children and adolescents. Even for these conditions, it is difficult to determine the absolute risk, and it is unknown whether hospitalization or severe disease could be prevented by monoclonal antibody therapy. For some of the conditions, such as sickle cell disease or immunocompromise, the driving force for the apparent increased risk of hospitalization might be explained by protocolized treatment for fever or other symptoms rather than tendency to severe disease, and further investigation is needed.[59, 60] There is insufficient evidence to determine whether non-asthma lung disease[19, 21, 22, 34, 40-46] and diabetes mellitus[61-64] significantly increase risk of hospitalization or severe COVID-19. In contrast, available data suggest that some of the listed conditions do not independently affect the risk of hospitalization or progression to severe COVID-19. Although asthma was noted to be prevalent in hospitalized children with COVID-19 in two US multi-

state cohorts, studies that have specifically analyzed asthma as a risk factor for severe pediatric COVID-19, as distinct from other chronic respiratory conditions, have not shown an association with worse outcomes.[22, 41, 43, 65-69] Similarly, although prospective analysis of immunocompromised condition as a risk factor is sparse, studies to date suggest that children who are mild-moderately immunocompromised[3] are not at increased risk for severe COVID-19.[19, 32-36, 38, 39, 60] Lastly, a number of observational studies of children with chronic kidney disease who develop SARS-CoV-2 infection have not found it to be a risk factor for severe COVID-19.[34, 40, 44, 70-74]

COVID-19 and healthcare disparities

The current pandemic has highlighted longstanding disparities in healthcare, with a disproportionate negative effect on communities of color. Recent data have found that adults and children of racial and/or ethnic minority groups have, on average, a rate of SARS-CoV-2 infections that is more than four times higher than non-Hispanic whites.[75-77] Disparities in disease severity have also been well documented. Several studies have noted that, on average, three out of every four children hospitalized with COVID-19 and/or MIS-C come from racial or ethnic minority groups.[78, 79] Though consensus has not been established, it is likely these inequalities are driven by a combination of societal- and individual-level factors. Access to care and equitable distribution of therapeutic interventions is also likely to be an important driver of these disparities. Thus, to avoid further exacerbating this divide, attention should be given to the equitable distribution of these novel therapeutics when individualized decisions are being made on their use.

5. Conclusions and Research Priorities

Currently, there is insufficient evidence for utility, safety or efficacy to recommend routine use of monoclonal antibody therapy for children and adolescents with COVID-19, even those considered to be at higher risk of hospitalization or severe disease. At this time, neither bamlanivimab nor casirivimab plus imdevimab should be considered standard of care in any pediatric population, even in patients who meet high-risk criteria. There are no data supporting safety and efficacy in children or adolescents, and the evidence supporting use in the adult population (including young adults) is modest and/or unpublished, and has limited applicability to pediatrics or to many specified risk groups. More research is needed to identify pediatric patients at high absolute risk of severe COVID-19 and to determine the impact of monoclonal antibody therapies in this population. This guidance will be re-evaluated as more evidence becomes available.

References

1. Food and Drug Administration. Coronavirus Treatment Acceleration Program (CTAP). Available at: <https://www.fda.gov/drugs/coronavirus-covid-19-drugs/coronavirus-treatment-acceleration-program-ctap> Accessed November 30, 2020.
2. Bhimraj A, Morgan RL, Shumaker AH, et al. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19. Available at: <https://www.idsociety.org/practice-guideline/covid-19-guideline-treatment-and-management/> Accessed November 30, 2020 2020.
3. Chiotos K, Hayes M, Kimberlin DW, et al. Multicenter interim guidance on use of antivirals for children with COVID-19/SARS-CoV-2. Journal of the Pediatric Infectious Diseases Society. **2020**. PubMed PMID: 32918548. Pubmed Central PMCID: PMC7543452. Epub 2020/09/13. eng.
4. Chiotos K, Hayes M, Kimberlin DW, et al. Multicenter initial guidance on use of antivirals for children with COVID-19/SARS-CoV-2. Journal of the Pediatric Infectious Diseases Society. **2020**. PubMed PMID: 32318706. Pubmed Central PMCID: PMC7188128. Epub 2020/04/23. eng.
5. World Health Organization. Therapeutics and COVID-19: living guideline. Available at: <https://www.who.int/publications/i/item/therapeutics-and-covid-19-living-guideline> Accessed November 30, 2020.
6. DeFrancesco L. COVID-19 antibodies on trial. Nat Biotechnol. **2020**;38(11):1242-52. PubMed PMID: 33087898. eng.
7. Food and Drug Administration. Casirivimab and Imdevimab EUA Letter of Authorization. Available at: <https://www.fda.gov/media/143891/download> Accessed November 30, 2020.
8. Food and Drug Administration. Bamlanivimab EUA Letter of Authorization. Available at: <https://www.fda.gov/media/143602/download> Accessed November 30, 2020.
9. Lilly provides comprehensive update on progress of SARS-CoV-2 neutralizing antibody programs. Available at: <https://investor.lilly.com/news-releases/news-release-details/lilly-provides-comprehensive-update-progress-sars-cov-2> Accessed December 1, 2020.

10. Chen P, Nirula A, Heller B, et al. SARS-CoV-2 Neutralizing Antibody LY-CoV555 in Outpatients with Covid-19. *N Engl J Med*. **2020**.
11. Food and Drug Administration. Fact sheet for health care providers; emergency use authorization (EUA) of bamlanivimab. Available at: <https://www.fda.gov/media/143603/download> Accessed November 30, 2020.
12. Regeneron. Casirivimab and imdevimab. Available at: <https://www.regeneron.com/casirivimab-imdevimab> Accessed November 25, 2020.
13. Regeneron. Regeneron's COVID-19 outpatient trial prospectively demonstrates that REGN-COV2 antibody cocktail significantly reduced virus levels and need for further medical attention. Available at: <https://investor.regeneron.com/news-releases/news-release-details/regenerons-covid-19-outpatient-trial-prospectively-demonstrates/> Accessed November 25, 2020.
14. Regeneron. Regeneron's REGN-COV2 antibody cocktail reduced viral levels and improved symptoms in non-hospitalized covid-19 patients. Available at: <https://investor.regeneron.com/news-releases/news-release-details/regenerons-regn-cov2-antibody-cocktail-reduced-viral-levels-and/> Accessed November 25, 2020.
15. Regeneron. Regeneron's casirivimab and imdevimab antibody cocktail for COVID-19 is first combination therapy to receive FDA emergency use authorization. Available at: <https://investor.regeneron.com/news-releases/news-release-details/regenerons-regen-cov2-first-antibody-cocktail-covid-19-receive/> Accessed November 25, 2020.
16. Food and Drug Administration. Fact sheet for health care providers; emergency use authorization (EUA) of casirivimab and imdevimab. Available at: <https://www.fda.gov/media/143892/download> Accessed November 30, 2020.
17. Food and Drug Administration. Frequently Asked Questions on the Emergency Use Authorization of Casirivimab + Imdevimab. Available at: <https://www.fda.gov/media/143894/download> Accessed November 30, 2020.

18. Weinreich DM, Sivapalasingam S, Norton T, et al. REGN-COV2, a Neutralizing Antibody Cocktail, in Outpatients with Covid-19. *N Engl J Med*. **2020**. PubMed PMID: 33332778. Epub 2020/12/18. eng.
19. Bailey LC, Razzaghi H, Burrows EK, et al. Assessment of 135 794 Pediatric Patients Tested for Severe Acute Respiratory Syndrome Coronavirus 2 Across the United States. *JAMA pediatrics*. **2020**. PubMed PMID: 33226415. Epub 2020/11/24. eng.
20. Cai Q, Chen F, Wang T, et al. Obesity and COVID-19 Severity in a Designated Hospital in Shenzhen, China. *Diabetes Care*. **2020**;43(7):1392-8. PubMed PMID: 32409502. Epub 2020/05/16. eng.
21. Chao JY, Derespina KR, Herold BC, et al. Clinical Characteristics and Outcomes of Hospitalized and Critically Ill Children and Adolescents with Coronavirus Disease 2019 at a Tertiary Care Medical Center in New York City. *J Pediatr*. **2020**;223:14-9.e2. PubMed PMID: 32407719. Pubmed Central PMCID: PMC7212947. Epub 2020/05/15. eng.
22. DeBiasi RL, Song X, Delaney M, et al. Severe Coronavirus Disease-2019 in Children and Young Adults in the Washington, DC, Metropolitan Region. *The Journal of pediatrics*. **2020**;223:199-203.e1. PubMed PMID: 32405091. Epub 2020/05/13. eng.
23. Huang R, Zhu L, Xue L, et al. Clinical findings of patients with coronavirus disease 2019 in Jiangsu province, China: A retrospective, multi-center study. *PLoS Negl Trop Dis*. **2020**;14(5):e0008280. PubMed PMID: 32384078. Pubmed Central PMCID: PMC7239492. Epub 2020/05/10. eng.
24. Kalligeros M, Shehadeh F, Mylona EK, et al. Association of Obesity with Disease Severity Among Patients with Coronavirus Disease 2019. *Obesity (Silver Spring)*. **2020**;28(7):1200-4. PubMed PMID: 32352637. Epub 2020/06/12. eng.
25. Kass DA, Duggal P, Cingolani O. Obesity could shift severe COVID-19 disease to younger ages. *Lancet*. **2020**;395(10236):1544-5. PubMed PMID: 32380044. Pubmed Central PMCID: PMC7196905. Epub 2020/05/08. eng.

26. Leon-Abarca JA. Obesity and immunodeficiencies are the main pre-existing conditions associated with mild to moderate COVID-19 in children. *Pediatr Obes.* **2020**;15(12):e12713-e. PubMed PMID: 32789994. Epub 2020/08/12. eng.
27. Lighter J, Phillips M, Hochman S, et al. Obesity in Patients Younger Than 60 Years Is a Risk Factor for COVID-19 Hospital Admission. *Clin Infect Dis.* **2020**;71(15):896-7. PubMed PMID: 32271368. Pubmed Central PMCID: PMC7184372. Epub 2020/04/10. eng.
28. McMichael TM, Clark S, Pogojans S, et al. COVID-19 in a Long-Term Care Facility - King County, Washington, February 27-March 9, 2020. *MMWR Morb Mortal Wkly Rep.* **2020**;69(12):339-42. PubMed PMID: 32214083. Epub 2020/03/28. eng.
29. Ogden CL, Carroll MD, Fakhouri TH, et al. Prevalence of Obesity Among Youths by Household Income and Education Level of Head of Household - United States 2011-2014. *MMWR Morb Mortal Wkly Rep.* **2018**;67(6):186-9. PubMed PMID: 29447142. Pubmed Central PMCID: PMC5815488. Epub 2018/02/16. eng.
30. Zachariah P, Johnson CL, Halabi KC, et al. Epidemiology, Clinical Features, and Disease Severity in Patients With Coronavirus Disease 2019 (COVID-19) in a Children's Hospital in New York City, New York. *JAMA pediatrics.* **2020**:e202430. PubMed PMID: 32492092. Epub 2020/06/04. eng.
31. Global Registry of COVID-19 in Pediatric Cancer. Available at: <https://global.stjude.org/en-us/global-covid-19-observatory-and-resource-center-for-childhood-cancer/registry.html> Accessed November 30, 2020.
32. Goss MB, Galván NTN, Ruan W, et al. The pediatric solid organ transplant experience with COVID-19: An initial multi-center, multi-organ case series. *Pediatr Transplant.*n/a(n/a):e13868.
33. He W, Chen L, Chen L, et al. COVID-19 in persons with haematological cancers. *Leukemia.* **2020**;34(6):1637-45. PubMed PMID: 32332856. Pubmed Central PMCID: PMC7180672. Epub 2020/04/26. eng.

34. Shekerdemian LS, Mahmood NR, Wolfe KK, et al. Characteristics and Outcomes of Children With Coronavirus Disease 2019 (COVID-19) Infection Admitted to US and Canadian Pediatric Intensive Care Units. *JAMA pediatrics*. **2020**;174(9):868-73.
35. Verma S, Lumba R, Dapul HM, et al. Characteristics of Hospitalized Children With SARS-CoV-2 in the New York City Metropolitan Area. *Hospital Pediatrics*. **2020**:hpeds.2020-001917.
36. Vicent MG, Martinez AP, Trabazo del Castillo M, et al. COVID-19 in pediatric hematopoietic stem cell transplantation: The experience of Spanish Group of Transplant (GETMON/GETH). *Pediatr Blood Cancer*. **2020**;67(9):e28514.
37. Ho H-E, Mathew S, Peluso MJ, Cunningham-Rundles C. Clinical outcomes and features of COVID-19 in patients with primary immunodeficiencies in New York City. *J Allergy Clin Immunol Pract*. **2020**:S2213-198(20)31102-8. PubMed PMID: 33039649. eng.
38. Vijenthira A, Gong IY, Fox TA, et al. Outcomes of patients with hematologic malignancies and COVID-19: A systematic review and meta-analysis of 3377 patients. *Blood*. **2020**;136(25):2881-92. PubMed PMID: 33113551. Pubmed Central PMCID: PMC7746126. Epub 2020/10/29. eng.
39. Madhusoodhan PP, Pierro J, Musante J, et al. Characterization of COVID-19 disease in pediatric oncology patients: The New York-New Jersey regional experience. *Pediatr Blood Cancer*. **2020**:e28843. PubMed PMID: 33338306. Epub 2020/12/19. eng.
40. Parri N, Lenge M, Buonsenso D. Children with Covid-19 in Pediatric Emergency Departments in Italy. *N Engl J Med*. **2020**;383(2):187-90. PubMed PMID: 32356945. Pubmed Central PMCID: PMC7206930. Epub 2020/05/02. eng.
41. Kim L, Whitaker M, O'Halloran A, et al. Hospitalization Rates and Characteristics of Children Aged <18 Years Hospitalized with Laboratory-Confirmed COVID-19 - COVID-NET, 14 States, March 1- July 25, 2020. *MMWR Morb Mortal Wkly Rep*. **2020**;69(32):1081-8. PubMed PMID: 32790664. Pubmed Central PMCID: PMC7440125. Epub 2020/08/14. eng.
42. Göttinger F, Santiago-García B, Noguera-Julián A, et al. COVID-19 in children and adolescents in Europe: a multinational, multicentre cohort study. *The Lancet Child & adolescent health*.

2020;4(9):653-61. PubMed PMID: 32593339. Pubmed Central PMCID: PMC7316447. Epub 2020/07/01. eng.

43. Fernandes DM, Oliveira CR, Guerguis S, et al. SARS-CoV-2 Clinical Syndromes and Predictors of Disease Severity in Hospitalized Children and Youth. *The Journal of pediatrics*. **2020**:S0022-3476(20)31393-7. PubMed PMID: 33197493. eng.

44. Centers for Disease Control and Prevention. Coronavirus Disease 2019 in Children - United States, February 12-April 2, 2020. *MMWR Morb Mortal Wkly Rep*. **2020**;69(14):422-6. PubMed PMID: 32271728. Pubmed Central PMCID: PMC7147903. Epub 2020/04/10. eng.

45. Bellino S, Punzo O, Rota MC, et al. COVID-19 Disease Severity Risk Factors for Pediatric Patients in Italy. *Pediatrics*. **2020**;146(4):e2020009399.

46. Sanna G, Serrau G, Bassareo PP, Neroni P, Fanos V, Marcialis MA. Children's heart and COVID-19: Up-to-date evidence in the form of a systematic review. *Eur J Pediatr*. **2020**;179(7):1079-87. PubMed PMID: 32474800. Epub 2020/05/30. eng.

47. Oualha M, Bendavid M, Berteloot L, et al. Severe and fatal forms of COVID-19 in children. *Arch Pediatr*. **2020**;27(5):235-8. PubMed PMID: 32518045. Pubmed Central PMCID: PMC7269941. Epub 2020/06/11. eng.

48. Swann OV, Holden KA, Turtle L, et al. Clinical characteristics of children and young people admitted to hospital with covid-19 in United Kingdom: prospective multicentre observational cohort study. *BMJ*. **2020**;370:m3249.

49. Telfer P, de la Fuente J, Sohal M, et al. Real-time national survey of COVID-19 in hemoglobinopathy and rare inherited anemia patients. *Haematologica*. **2020**;105(11):2651-4.

50. Subarna C, Giselle P-P, Fester I, et al. COVID-19 in patients with sickle cell disease - a case series from a UK Tertiary Hospital. *Haematologica*. **2020**;105(11):2691-3.

51. Panepinto JA, Brandow A, Mucalo L, et al. Coronavirus Disease among Persons with Sickle Cell Disease, United States, March 20-May 21, 2020. *Emerg Infect Dis*. **2020**;26(10):2473-6. PubMed PMID: 32639228. Epub 2020/07/08. eng.

52. Odièvre MH, de Marcellus C, Ducou Le Pointe H, et al. Dramatic improvement after tocilizumab of severe COVID-19 in a child with sickle cell disease and acute chest syndrome. *Am J Hematol.* **2020**;95(8):E192-e4. PubMed PMID: 32358817. Pubmed Central PMCID: PMC7267654. Epub 2020/05/03. eng.
53. Morrone KA, Strumph K, Liszewski MJ, et al. Acute chest syndrome in the setting of SARS-COV-2 infections-A case series at an urban medical center in the Bronx. *Pediatr Blood Cancer.* **2020**:e28579. PubMed PMID: 32893954. Epub 2020/09/08. eng.
54. Jacob S, Dworkin A, Romanos-Sirakis E. A pediatric patient with sickle cell disease presenting with severe anemia and splenic sequestration in the setting of COVID-19. *Pediatr Blood Cancer.* **2020**;67(12):e28511. PubMed PMID: 32776396. eng.
55. Heilbronner C, Berteloot L, Tremolieres P, et al. Patients with sickle cell disease and suspected COVID-19 in a paediatric intensive care unit. *Br J Haematol.* **2020**;190(1):e21-e4. PubMed PMID: 32420608. Pubmed Central PMCID: PMC7276717. Epub 2020/05/19. eng.
56. Arlet J-B, de Luna G, Khimoud D, et al. Prognosis of patients with sickle cell disease and COVID-19: a French experience. *The Lancet Haematology.* **2020**;7(9):e632-e4. PubMed PMID: 32563282. Epub 2020/06/18. eng.
57. Appiah-Kubi A, Acharya S, Fein Levy C, et al. Varying presentations and favourable outcomes of COVID-19 infection in children and young adults with sickle cell disease: an additional case series with comparisons to published cases. *Br J Haematol.* **2020**. PubMed PMID: 32652551. eng.
58. Al-Hebshi A, Zolaly M, Alshengeti A, et al. A Saudi family with sickle cell disease presented with acute crises and COVID-19 infection. *Pediatr Blood Cancer.* **2020**;67(9):e28547. PubMed PMID: 32648991. eng.
59. Eisenbrown K, Ellison AM, Nimmer M, Badaki-Makun O, Brousseau DC. Practice Variation in Emergency Department Management of Children With Sickle Cell Disease Who Present With Fever. *Pediatr Emerg Care.* **2018**;34(8):574-7. PubMed PMID: 30020250. Epub 2018/07/19. eng.

60. Lehrnbecher T, Robinson P, Fisher B, et al. Guideline for the management of fever and neutropenia in children with cancer and hematopoietic stem-cell transplantation recipients: 2017 update. *J Clin Oncol.* **2017**;35(18):2082-94. PubMed PMID: 28459614.
61. Armann JP, Diffloth N, Simon A, et al. Hospital Admission in Children and Adolescents With COVID-19. *Dtsch Arztebl Int.* **2020**;117(21):373-4. PubMed PMID: 32519943. eng.
62. European Society for Paediatric Endocrinology. ESPE Patient Information on COVID-19 and Pediatric Endocrine Disease; Disease specific information and advice – Type 1 Diabetes. Available at: <https://www.eurospe.org/media/2314/espe-patient-information-on-covid-19-and-type-1-diabetes.pdf> Accessed November 19, 2020.
63. Kostopoulou E, Güemes M, Shah P. COVID-19 in Children and Adolescents with Endocrine Conditions. *Horm Metab Res.* **2020**;52(11):769-74. PubMed PMID: 32898896. Epub 2020/09/09. eng.
64. Rabbone I, Schiaffini R, Cherubini V, Maffei C, Scaramuzza A. Has COVID-19 Delayed the Diagnosis and Worsened the Presentation of Type 1 Diabetes in Children? *Diabetes Care.* **2020**;43(11):2870-2. PubMed PMID: 32778554. Epub 2020/08/12. eng.
65. Chao JY, Derespina KR, Herold BC, et al. Clinical Characteristics and Outcomes of Hospitalized and Critically Ill Children and Adolescents with Coronavirus Disease 2019 (COVID-19) at a Tertiary Care Medical Center in New York City. *J Pediatr.* **2020**. PubMed PMID: 32407719. Pubmed Central PMCID: PMC7212947. Epub 2020/05/15. eng.
66. Hurst JH, Heston SM, Chambers HN, et al. SARS-CoV-2 Infections Among Children in the Biospecimens from Respiratory Virus-Exposed Kids (BRAVE Kids) Study. *Clin Infect Dis.* **2020**. PubMed PMID: 33141180. Pubmed Central PMCID: PMC7665428. Epub 2020/11/04. eng.
67. Lovinsky-Desir S, Deshpande DR, De A, et al. Asthma among hospitalized patients with COVID-19 and related outcomes. *J Allergy Clin Immunol.* **2020**;146(5):1027-34.e4.
68. Zachariah P, Johnson CL, Halabi KC, et al. Epidemiology, Clinical Features, and Disease Severity in Patients With Coronavirus Disease 2019 (COVID-19) in a Children's Hospital in New York

City, New York. JAMA pediatrics. **2020**;174(10):e202430-e. PubMed PMID: 32492092. Epub 2020/10/05. eng.

69. Zhu Z, Hasegawa K, Ma B, Fujiogi M, Camargo CA, Jr., Liang L. Association of asthma and its genetic predisposition with the risk of severe COVID-19. J Allergy Clin Immunol. **2020**;146(2):327-9.e4. PubMed PMID: 32522462. Pubmed Central PMCID: PMC7423602. Epub 2020/06/12. eng.

70. Dong Y, Mo X, Hu Y, et al. Epidemiological Characteristics of 2143 Pediatric Patients With 2019 Coronavirus Disease in China. Pediatrics. **2020**:e20200702.

71. Lu X, Zhang L, Du H, et al. SARS-CoV-2 Infection in Children. N Engl J Med. **2020**;382(17):1663-5. PubMed PMID: 32187458.

72. Melgosa M, Madrid A, Álvarez O, et al. SARS-CoV-2 infection in Spanish children with chronic kidney pathologies. Pediatric nephrology (Berlin, Germany). **2020**;35(8):1521-4. PubMed PMID: 32435879. Epub 2020/05/20. eng.

73. Plumb L, Benoy-Deeney F, Casula A, et al. COVID-19 in children with chronic kidney disease: findings from the UK renal registry. Arch Dis Child. **2020**. PubMed PMID: 32709685. Epub 2020/07/28. eng.

74. Rawson A, Wilson AC, Schwaderer AL, et al. Coronavirus disease 2019 (COVID-19) in two pediatric patients with kidney disease on chronic immunosuppression: A case series. Hemodialysis international International Symposium on Home Hemodialysis. **2020**. PubMed PMID: 33015942. Pubmed Central PMCID: PMC7675317. Epub 2020/10/06. eng.

75. Centers for Disease Control and Prevention. Weekly Updates by Select Demographic and Geographic Characteristics: Provisional Death Counts for Coronavirus Disease 2019 (COVID-19). Available at: https://www.cdc.gov/nchs/nvss/vsrr/covid_weekly/index.htm Accessed Accessed December 15 2020.

76. Goyal MK, Simpson JN, Boyle MD, et al. Racial and/or Ethnic and Socioeconomic Disparities of SARS-CoV-2 Infection Among Children. Pediatrics. **2020**;146(4). PubMed PMID: 32759379. Epub 2020/08/08. eng.

77. Wadhera RK, Wadhera P, Gaba P, et al. Variation in COVID-19 Hospitalizations and Deaths Across New York City Boroughs. *JAMA*. **2020**;323(21):2192-5. PubMed PMID: 32347898. Pubmed Central PMCID: PMC7191469 from the National Heart, Lung, and Blood Institute (grant K23HL148525-1) and previously serving as a consultant for Regeneron. Dr Figueroa reported receiving research support from the National Center for Advancing Translational Sciences (grant KL2 TR002542) and the Commonwealth Fund. Dr Joynt Maddox reported receiving research support from the National Heart, Lung, and Blood Institute (grant R01HL143421) and the National Institute on Aging (grant R01AG060935); receiving grant support from the Commonwealth Fund; and having prior contract work with the Office of the Assistant Secretary for Planning and Evaluation. Dr Yeh reported receiving research support from the National Heart, Lung, and Blood Institute (grant R01HL136708); serving as a consultant to Biosense Webster; and serving as a consultant to and receiving grants from Abbott Vascular, AstraZeneca, Boston Scientific, and Medtronic. No other disclosures were reported. Epub 2020/04/30. eng.
78. Dufort EM, Koumans EH, Chow EJ, et al. Multisystem Inflammatory Syndrome in Children in New York State. *N Engl J Med*. **2020**;383(4):347-58. PubMed PMID: 32598830. Pubmed Central PMCID: PMC7346766. Epub 2020/07/01. eng.
79. Fernandes DM, Oliveira CR, Guerguis S, et al. Severe Acute Respiratory Syndrome Coronavirus 2 Clinical Syndromes and Predictors of Disease Severity in Hospitalized Children and Youth. *J Pediatr*. **2020**. PubMed PMID: 33197493. Pubmed Central PMCID: PMC7666535. Epub 2020/11/17. eng.