



Published in final edited form as:

*Pediatrics*. 2020 February ; 145(2): . doi:10.1542/peds.2019-1490.

## Subacute Neuropsychiatric Syndrome in Girls With *SHANK3* Mutations Responds to Immunomodulation

Alexandra L. Bey, MD, PhD<sup>a</sup>, Mark P. Gorman, MD<sup>b</sup>, William Gallentine, DO<sup>a,c</sup>, Teresa M. Kohlenberg, MD<sup>d</sup>, Jennifer Frankovich, MD, MS<sup>e</sup>, Yong-hui Jiang, MD, PhD<sup>f,g,h</sup>, Keith Van Haren, MD<sup>i</sup>

<sup>a</sup>Department of Psychiatry and Behavioral Sciences, Duke University, Durham, North Carolina

<sup>b</sup>Department of Neurology, Boston Children's Hospital and Harvard Medical School, Harvard University, Boston, Massachusetts

<sup>c</sup>Department of Neurology, Duke University, Durham, North Carolina

<sup>d</sup>Department of Psychiatry, Medical School, University of Massachusetts, Worcester, Massachusetts

<sup>e</sup>Division of Rheumatology, Department of Pediatrics, School of Medicine, Stanford University, Palo Alto, California

<sup>f</sup>Department of Pediatrics and Neurobiology, Duke University, Durham, North Carolina

<sup>g</sup>Department of Genomics and Genetics Graduate Program, Duke University, Durham, North Carolina

<sup>h</sup>Department of Duke Institute for Brain Sciences, Duke University, Durham, North Carolina

<sup>i</sup>Department of Neurology and Neurological Sciences, School of Medicine, Stanford University, Palo Alto, California

### Abstract

Phenotypic and biological characterization of rare monogenic disorders represents 1 of the most important avenues toward understanding the mechanisms of human disease. Among patients with SH3 and multiple ankyrin repeat domains 3 (*SHANK3*) mutations, a subset will manifest neurologic regression, psychosis, and mood disorders. However, which patients will be affected, when, and why are important unresolved questions. Authors of recent studies suggest neuronal *SHANK3* expression is modulated by both inflammatory and hormonal stimuli. In this case series, we describe 4 independent clinical observations of an immunotherapy responsive phenotype of peripubertal-onset neuropsychiatric regression in 4 girls with pathogenic *SHANK3* mutations.

---

Address correspondence to Keith Van Haren, MD, Pediatric Neuroimmunology Clinic, 730 Welch Rd, Palo Alto, CA 94304. kpv@stanford.edu.

Drs Jiang and Van Haren conceptualized the case series, provided clinical information, critically reviewed the manuscript, and contributed significant intellectual content; Dr Bey contributed clinical information, drafted the initial manuscript, and led in manuscript revision and submission; Drs Gorman, Gallentine, Kohlenberg, and Frankovich collected clinical data, provided important interpretations of the data, critically reviewed the manuscript, and contributed significant intellectual content; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

**POTENTIAL CONFLICT OF INTEREST:** The authors have indicated they have no potential conflicts of interest to disclose.

**FINANCIAL DISCLOSURE:** The authors have indicated they have no financial relationships relevant to this article to disclose.

Each child exhibited a history of stable, mild-to-moderate lifelong developmental disability until 12 to 14 years of age, at which time each manifested a similar, subacute-onset neurobehavioral syndrome. Symptoms included mutism, hallucinations, insomnia, inconsolable crying, obsessive-compulsive behaviors, loss of self-care, and urinary retention and/or incontinence. Symptoms were relatively refractory to antipsychotic medication but improved after immunomodulatory treatment. All 4 patients exhibited chronic relapsing courses during a period of treatment and follow-up ranging from 3 to 6 years. Two of the 4 girls recovered their premorbid level of functioning. We briefly review the scientific literature to offer a conceptual and molecular framework for understanding these clinical observations. Future clinical and translational investigations in this realm may offer insights into mechanisms and therapies bridging immune function and human behavior.

---

Characterization of monogenic variants represents an important avenue toward understanding human behavior and disease. Deletions or point mutations affecting SH3 and multiple ankyrin repeat domains 3 (*SHANK3*), which encodes a postsynaptic scaffolding protein, have been linked to autism spectrum disorder (ASD), Phelan-McDermid syndrome, and schizophrenia.<sup>1-3</sup> Peripubertal neurobehavioral regression, sometimes with mania and catatonia, has been reported in some patients with Phelan-McDermid syndrome, but the etiology is unknown.<sup>4-6</sup> Emerging literature reveals the ability of physiologic signals (eg, inflammatory stimuli, sex hormones) to regulate *SHANK3* expression.<sup>7-9</sup>

In this report, we describe 4 peripubertal girls with previously stable developmental disabilities who exhibited subacute disabling symptoms that included mutism, hallucinations, insomnia, inconsolable crying, diminished self-care, and urinary retention and/or incontinence. Each child was evaluated by independent teams for suspected autoimmune encephalopathy before detection of *SHANK3* point mutation. The families of these patients verbally consented to inclusion in this study.

## CASE 1

A 13-year-old girl had a history of mild-to-moderate developmental delay without ASD and autoimmune thyroiditis on levothyroxine. Her family history was notable for autoimmune disease in the father and brother. At 13 years, over several weeks, her symptoms progressed from irritability to audiovisual hallucinations, insomnia, compulsivity, aggression, catatonia, mutism, and urinary retention. Her symptoms were refractory to antipsychotics and benzodiazepines.

Serology and thyroid function studies prompted consideration of Hashimoto encephalopathy. At 14 years, 40 mg of oral prednisone taken daily for 2 weeks yielded no improvement, but 2 g/kg of intravenous immunoglobulin (IVIG) led to sustained improvement in language, restlessness, and agitation within 2 days. Symptoms recurred 3 weeks later prompting monthly IVIG, which demonstrated the same pattern of response. Attempts to increase treatment intervals led to symptom reemergence.

At 16 years, she continues on IVIG (1 g/kg) and methylprednisolone every 4 to 6 weeks. She has regained 2 of 6 independent activities of daily living (ADLs) from nadir and weaned off

daily psychotropic agents but remains substantially below her premorbid baseline. Whole-exome sequencing was pursued for unknown etiology of lifelong developmental delay and identified a pathogenic de novo frameshift mutation in exon 21 of *SHANK3*.

## CASE 2

A 14-year-old girl had a history of moderate intellectual disability without ASD. Her maternal grandmother had psychosis. At age 14 years, she developed 10-day episodes of mania with psychotic features coinciding with menstruation with poor response to antipsychotics. After her third event, she developed depression with catatonic features and developmental regression. Daily oral contraceptives to suppress menses, lamotrigine, and a switch from aripiprazole to olanzapine correlated with a gradual return to her neuropsychiatric baseline over 8 months. At age 15 years, she developed myoclonic jerks, athetoid, and choreiform movements. At age 16 years, she developed overnight onset of severe anxiety, compulsions, insomnia, urinary incontinence, facial tics, sobbing, and loss of self-care and academic skills. After 2 months of symptoms and an unrevealing workup, her symptoms prompted treatment of seronegative autoimmune encephalitis using 2 g/kg of IVIG, which was temporally associated with symptom resolution within 3 weeks. She was maintained on olanzapine, lamotrigine, N-acetyl cysteine, methylated B vitamins, and oral contraceptives. After 5 months, neuropsychiatric testing matched her premorbid baseline.

Two manic relapses occurred 6 months and 1 year later, both improving in conjunction with IVIG. The latter event was notable for a seizure, echolalia, and anxiety. Within 1 hour, 0.5 mg of lorazepam for suspected catatonia yielded dramatic improvement. Mania and catatonia relapsed 1 month later but resolved with daily lorazepam.

At age 18 years, she maintains all ADLs on a daily regimen of lorazepam, olanzapine, lamotrigine, oral contraceptive, N-acetyl cysteine, vitamin D, folate, omega-3 fatty acids, and monthly IVIG (1 g/kg). Whole-exome sequencing was pursued for unknown etiology of complex neuropsychiatric illness and revealed a pathogenic frameshift mutation in exon 21 of *SHANK3*; the inheritance of this mutation is unknown.

## CASE 3

A 12-year-old girl had a history of mild intellectual disability, ASD, tics, and attention-deficit/hyperactivity disorder. At age 12 years, 3 months after menarche, she developed abrupt onset of hyperactivity, paranoia, mutism, akathisia, audiovisual hallucinations, inconsolable screaming, and periodic staring spells. Her workup was notable for elevated antinuclear antibody and slight elevation of white blood cell count in cerebrospinal fluid. The symptoms were refractory to guanfacine and risperidone but improved with 30 mg/kg of intravenous methylprednisolone within days but reemerged over weeks. IVIG (2 g/kg) yielded significant improvement in language, social engagement, sleep, agitation, and concentration, but the effect consistently waned at approximately 3 weeks. Mycophenolate and rituximab were added. Over the next 2 years, she maintained stable premorbid language, cognition, and ADLs but relapsed within 6 months of weaning immunomodulation at 14 years, prompting reinitiation of a similar regimen. She maintained her premorbid baseline

for 3 years until at age 17 years, weaning of mycophenolate was associated with neuropsychiatric decline, prompting a 3-week ICU admission with reinitiation of IVIG, rituximab, mycophenolate, high-dose lorazepam, and ultimately electroconvulsive therapy for catatonia. Within a month, she reapproximated her baseline.

At 18 years, her current treatment includes monthly IVIG, mycophenolate, and rituximab for immunotherapy in addition to lorazepam, methylphenidate, trazodone, lithium, and clonidine. Whole-exome sequencing was pursued for unknown etiology of complex autoimmune encephalitis and lifelong developmental delay and identified a pathogenic de novo frameshift mutation in exon 21 of *SHANK3*.

## CASE 4

A 13-year-old girl who had a history of mild cognitive delay and learning disabilities without ASD developed severe acute anxiety within 4 days of starting minocycline for facial acne; the medication was discontinued. One month later, she developed overnight onset of anxiety, obsessiveness, insomnia, inability to chew or swallow, aphasia, agraphia, inconsolable screaming, impulsivity, and urinary incontinence. She was diagnosed with bipolar disorder and hospitalized 6 times in the next 6 months but was poorly responsive to antipsychotics, benzodiazepines, antidepressants, or anticonvulsants. Lithium conferred a moderate, mood-stabilizing effect. After 1 year of symptoms, an autoimmune evaluation was notable for elevated antinuclear and histone antibodies and low complement component 4, which led to treatment with IV methylprednisolone. Within 3 days, she manifested improved expressive language, transitioned to daily prednisone, and returned to her neuropsychiatric baseline for several months. However, symptoms recurred with steroid weaning. After transition to 2 g/kg of IVIG, mycophenolate, and rituximab, she maintained her premorbid baseline for the next 2 years despite weaning all psychiatric medications. However, when her immunosuppression was weaned off at 16 years, severe psychiatric symptoms returned. Despite reinitiating intravenous methylprednisolone, IVIG, mycophenolate, and rituximab, and a trial of plasmapheresis, her symptoms persisted. She was subsequently administered a 6-month course of cyclophosphamide and clozapine. Her symptoms began to improve only after her white blood cell counts began to fall.

At 20 years of age, she remains on rituximab and prednisone and has tolerated weaning of mycophenolate. She maintains a relatively stable baseline independent in all ADLs with intermittent “flares” of obsessive-compulsive/psychotic symptoms, often in the setting of a systemic infection; these symptoms often self-resolve after treating the infection, sometimes with the addition of steroids. Whole-exome sequencing was pursued because of unknown etiology of psychosis and revealed a pathogenic de novo frameshift mutation in exon 14 of *SHANK3*.

## DISCUSSION

We describe 4 peripubertal girls with stable, lifelong developmental disabilities with and without comorbid ASD caused by pathogenic mutations in *SHANK3* who developed a subacute stereotyped, profoundly disabling neuropsychiatric syndrome that prompted 4

independent clinical teams to initiate treatment of suspected autoimmune encephalopathy. Behavioral symptoms were chronic and/or relapsing but often improved with immunomodulatory therapies administered intermittently over several years (Table 1). Two patients recovered to their premorbid function.

Each patient demonstrated symptom improvement typically within days of IVIG followed in some by a waning effect within the following weeks. This pattern approximates a therapeutic rule for seronegative autoimmune encephalopathies.<sup>10</sup> Although the short-term treatment effect was reproducible within and across cases, the degree to which immunomodulatory therapy was responsible for long-term improvements in these patients can be debated. A precise comparison is confounded by the heterogeneity of each course including fluctuating symptoms, variation in immunomodulatory and psychotropic regimens, and incomplete return to baseline in 2 individuals. Although all patients were considered relatively refractory to antidepressant and antipsychotic medications, patients reported variable responses to lorazepam, lithium, and/or hormonal contraceptives.

Experimental data offer possible molecular and conceptual mechanisms supporting the notion of immune-mediated behavioral modulation via *SHANK3*. At the molecular level, *SHANK3* functions as a master scaffolding protein at the synapse, supporting several neurotransmitter receptors (Fig 1). Three cases (cases 1–3) had truncating mutations in exon 21 encoding the proline-rich domain, which mediates contacts with the metabotropic glutamate receptor, and 1 (case 4) had a truncating mutation in a well-characterized PDZ domain that interfaces with N-methyl-D-aspartate (NMDA) and  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptors.<sup>11</sup>

Exons 14 and 21 are subject to alternative splicing and intragenic promoters, such that mutations in these regions are predicted to result in isoform-specific disruption of *SHANK3* in the brain.<sup>11–13</sup> Autoantibodies against the synaptic proteins supported by *SHANK3* scaffolding have been implicated in the onset of classic autoimmune encephalitides.<sup>14,15</sup> Experimental models demonstrate neuronally mediated inflammatory signals capable of reducing *SHANK3* expression even in the absence of overt neuroinflammation.<sup>7,8</sup> Sex hormones modulate *SHANK3* expression in a similar fashion.<sup>9</sup>

The modulation of behavior via inflammatory signaling fits within the conceptual framework of “sickness behaviors” (Fig 1), which constitute a diverse suite of evolutionarily conserved responses (eg, irritability, mild encephalopathy, depressed mood, diminished self-care, social withdrawal) that occurs to varying degrees within the setting of systemic illness.<sup>16–18</sup> Links between sickness behaviors and psychiatric disease have been widely speculated.<sup>19</sup> The molecular mechanisms underlying sickness behaviors have been studied in animal models, which reveal a variety of molecular pathways linking systemic inflammatory stimuli with neuronal and behavioral modulation. Genetic variants conferring a gain or loss of function of key regulatory molecules within these pathways would be expected to yield a phenotypic change in behavior conferring resilience or vulnerability to inflammation. *SHANK3* perturbations, such as the putative loss of function mutations seen in these 4 cases, have been implicated in autism, schizophrenia, and bipolar disorder, all of which have been linked to immune dysregulation in genome-wide investigations, but the

genetic complexity warrants further investigation because mania or autism has been seen in patients with gain and loss of function mutations, and *SHANK3* has immense transcriptional complexity.<sup>20,21</sup> Thus, although experimental models<sup>7,8</sup> and a previous case<sup>22</sup> have implicated *SHANK3* in immune-behavior connections, the 4 cases described in this series may highlight the real-world clinical implications of this phenomenon.

Our cases, and others like them,<sup>5–7,22</sup> support targeted genetic sequencing of *SHANK3* in individuals with acute neuropsychiatric decline or catatonia in the context of preexisting intellectual disability, with or without comorbid ASD given the heterogeneous presentations of patients with *SHANK3* mutations. The experience of these 4 patients lends support to additional diagnostic evaluations, structured clinical trials, and further laboratory work to investigate new mechanisms and therapies for acute neuropsychiatric events in patients with *SHANK3* mutations. Immunomodulatory therapy carries risks and should only be pursued in accord with existing guidelines.<sup>10</sup> Insights gleaned from individuals with well-characterized molecular perturbations may clarify our understanding of the neurobiological basis of human behavior and highlight new therapeutic strategies.

## ACKNOWLEDGMENTS

We acknowledge Elise Brimble, MSc, MS, and Heather Van Mater, MD, MSc, for assistance reading and revising the article. We thank David Goldstein, Vandana Shashi, and Rebecca C. Spillmann for the whole-exome sequencing study of case 3 that was enrolled in protocol 32301: Genomic Study of Medical, Developmental, or Congenital Problems of Unknown Etiology. This study was approved by the Duke University Health System Institutional Review Board for Clinical Investigators (FWA 00009025), supported by the Duke University Health system, and partially funded by Union Chimique Belge Celltech. Dr Jiang is supported by National Institutes of Health grants MH117289, HD087795, and MH098114.

## ABBREVIATIONS


<b>ADL</b>	activity of daily living
<b>ASD</b>	autism spectrum disorder
<b>IVIG</b>	intravenous immunoglobulin
<b>NMDA</b>	N-methyl-D-aspartate
<b>SHANK3</b>	SH3 and multiple ankyrin repeat domains 3

## REFERENCES

1. Durand CM, Betancur C, Boeckers TM, et al. Mutations in the gene encoding the synaptic scaffolding protein SHANK3 are associated with autism spectrum disorders. *Nat Genet.* 2007;39(1):25–27 [PubMed: 17173049]
2. Gauthier J, Champagne N, Lafrenière RG, et al.; S2D Team. De novo mutations in the gene encoding the synaptic scaffolding protein SHANK3 in patients ascertained for schizophrenia. *Proc Natl Acad Sci USA.* 2010;107(17): 7863–7868 [PubMed: 20385823]
3. Leblond CS, Nava C, Polge A, et al. Meta-analysis of SHANK mutations in autism spectrum disorders: a gradient of severity in cognitive impairments. *PLoS Genet.* 2014;10(9):e1004580 [PubMed: 25188300]



4. Denayer A, Van Esch H, de Ravel T, et al. Neuropsychopathology in 7 patients with the 22q13 deletion syndrome: presence of bipolar disorder and progressive loss of skills. *Mol Syndromol*. 2012;3(1): 14–20 [PubMed: 22855650]
5. Serret S, Thümmler S, Dor E, Vesperini S, Santos A, Askenazy F. Lithium as a rescue therapy for regression and catatonia features in two SHANK3 patients with autism spectrum disorder: case reports. *BMC Psychiatry*. 2015;15:107 [PubMed: 25947967]
6. Zwanenburg RJ, Ruiter SA, van den Heuvel ER, Flapper BC, Van Ravenswaaij-Arts CM. Developmental phenotype in Phelan-McDermid (22q13.3 deletion) syndrome: a systematic and prospective study in 34 children. *J Neurodev Disord*. 2016;8:16 [PubMed: 27118998]
7. Grubucker AM. Environmental factors in autism. *Front Psychiatry*. 2013;3:118 [PubMed: 23346059]
8. Zhao Y, Lukiw WJ. Bacteroidetes neurotoxins and inflammatory neurodegeneration. *Mol Neurobiol*. 2018;55(12):9100–9107 [PubMed: 29637444]
9. Berkel S, Eltokhi A, Fröhlich H, et al. Sex hormones regulate SHANK expression. *Front Mol Neurosci*. 2018;11:337 [PubMed: 30319350]
10. Graus F, Titulaer MJ, Balu R, et al. A clinical approach to diagnosis of autoimmune encephalitis. *Lancet Neurol*. 2016;15(4):391–404 [PubMed: 26906964]
11. Wang X, Xu Q, Bey AL, Lee Y, Jiang YH. Transcriptional and functional complexity of SHANK3 provides a molecular framework to understand the phenotypic heterogeneity of SHANK3 causing autism and SHANK3 mutant mice. *Mol Autism*. 2014;5:30 [PubMed: 25071925]
12. Zhu L, Wang X, Li XL, et al. Epigenetic dysregulation of SHANK3 in brain tissues from individuals with autism spectrum disorders. *Hum Mol Genet*. 2014;23(6):1563–1578 [PubMed: 24186872]
13. De Rubeis S, Siper PM, Durkin A, et al. Delineation of the genetic and clinical spectrum of Phelan-McDermid syndrome caused by SHANK3 point mutations. *Mol Autism*. 2018;9:31 [PubMed: 29719671]
14. Armangue T, Petit-Pedrol M, Dalmau J. Autoimmune encephalitis in children. *J Child Neurol*. 2012;27(11): 1460–1469 [PubMed: 22935553]
15. Jiang YH, Ehlers MD. Modeling autism by SHANK gene mutations in mice. *Neuron*. 2013;78(1):8–27 [PubMed: 23583105]
16. Kluger MJ, Ringler DH, Anver MR. Fever and survival. *Science*. 1975;188(4184): 166–168 [PubMed: 1114347]
17. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nat Rev Neurosci*. 2008;9(1):46–56 [PubMed: 18073775]
18. Pecchi E, Dallaporta M, Jean A, Thirion S, Troadec JD. Prostaglandins and sickness behavior: old story, new insights. *Physiol Behav*. 2009;97(3–4): 279–292 [PubMed: 19275907]
19. Maes M, Berk M, Goehler L, et al. Depression and sickness behavior are Janus-faced responses to shared inflammatory pathways. *BMC Med*. 2012;10:66 [PubMed: 22747645]
20. Alexandrov PN, Zhao Y, Jaber V, Cong L, Lukiw WJ. Deficits in the proline-rich synapse-associated Shank3 protein in multiple neuropsychiatric disorders. *Front Neurol*. 2017;8:670 [PubMed: 29321759]
21. Gandal MJ, Haney JR, Parikshak NN, et al.; CommonMind Consortium; PsychENCODE Consortium; iPSYCH-BROAD Working Group. Shared molecular neuropathology across major psychiatric disorders parallels polygenic overlap. *Science*. 2018; 359(6376):693–697 [PubMed: 29439242]
22. Jungová P, umová A, Kramarová V, et al. Phelan-McDermid syndrome in adult patient with atypical bipolar psychosis repeatedly triggered by febrility. *Neurocase*. 2018;24(4):227–230 [PubMed: 30376408]



**FIGURE 1.**

Sickness behavior as a hypothetical model for conceptualizing *SHANK3*'s interface with the immune system. *SHANK3* functions as an important scaffolding protein for neurotransmitter receptors (see 1). A, In experimental models, systemic inflammatory stimuli can decrease *SHANK3* expression via nuclear factor  $\kappa$ -light-chain enhancer of activated B cells (NF- $\kappa$ B)-related signaling pathways (see 8). B, A variety of stereotyped behaviors occur as codified responses to nonhomeostatic physiologic states (eg, acute illness), each using unique molecular pathways to manifest adaptive, stimuli-specific behavioral responses. C, Alteration of proteins within these pathways, such as via genetic variants that confer a gain or loss of function, would be hypothesized to alter the associated behavioral response. The diverse behaviors that accompany states of acute infection, known as sickness behaviors, which include irritability, depressed mood, and social withdrawal, offer a template for conceptualizing the molecular and behavioral phenomena that may result from systemic inflammation and, conversely, be modified by its resolution (see 17). Considered together with our case histories, the combined observations offer a reasonable hypothesis implicating *SHANK3* as a candidate protein involved in 1 or more molecular pathways regulating clinically relevant sickness behaviors in humans. AMPAR,  $\alpha$ -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; mGluR, metabotropic glutamate receptor; NF- $\kappa$ B, nuclear factor kappa-light-chain-enhancer of activated B cells; NMDAR, N-methyl-D-aspartate receptor; NP<sub>0</sub>, common variant of a hypothetical neurobehavioral protein; NP<sub>1</sub>, gain of function variant of neurobehavioral protein; NP<sub>2</sub>, loss of function of neurobehavioral protein; NP, neurobehavioral protein.



TABLE 1

Clinical Case Summaries

	Case 1	Case 2	Case 3	Case 4
Baseline clinical features				
Age, sex	13-y-old girl	14-y-old girl	12-y-old girl	13-y-old girl
<i>SHANK3</i> mutation	c.3679dupG, p.A1227Gfs*69 (pathogenic de novo frameshift in exon 21)	c.3424_3425delCT, p.Leu1142Valfs*153 (pathogenic unknown inheritance frameshift in exon 21)	c.4116delC; p.Thr1373Glnfs*13 (pathogenic de novo frameshift in exon 21)	c.1864_1865delinsA, p.A622fs*XX (pathogenic de novo frameshift in exon 14)
Duration of follow-up	4 y	4 y	5 y	6 y
Neurocognitive baseline	Moderate intellectual disability (verbal IQ 78; performance IQ 53)	Moderate intellectual disability (IQ 40)	Moderate intellectual disability (IQ 55)	Mild cognitive and learning disabilities (IQ 78), delayed acquisition of speech, auditory processing disorder, and fine motor issues
Neuropsychiatric baseline and history	Happy, socially engaged child with no previous psychiatric illnesses	History of cyclic perimenstrual mania, catatonia, and depressive episodes beginning 2 y before intractable, subacute encephalopathy ADHD tics	ADHD tics	Happy, socially engaged, high-functioning child with no previous psychiatric illnesses or mood instability
Other medical history	History of autoimmune thyroiditis (elevated thyroid-stimulating hormone, elevated Abs against thyroglobulin and thyroid peroxidase)	Premature puberty at age 8 y treated with monthly Lupron injections from ages 8–12 y	Generalized tonic-clonic seizure (once) at 10 y	—
Family history	Father with multiple sclerosis and inflammatory bowel disease brother with lupus	Maternal grandmother with bipolar disorder	No relevant conditions	Maternal aunt with bipolar and maternal grandmother with mood disorder
Age at menarche	11 y 4 mo	12 y	11 y 10 mo	12 y
Subacute neuropsychiatric symptoms at presentation				
Age at subacute onset	13 y 10 mo	14 y 4 mo	12 y 1 mo	13 y 5 mo
Time from onset to symptomatic nadir	3-wk progressive loss of bowel and bladder control followed by overnight onset of psychosis	Overnight onset of severe obsessive-compulsive symptoms	Overnight onset of agitation, psychosis, and insomnia	Overnight onset of anxiety, severe sleep disruption, eating restriction, and eventual obsessive-compulsive symptoms
Audiovisual hallucinations	Yes	Yes	Yes	No
Inconsolable screaming and/or crying	Yes	Yes	Yes	Yes
Aphasia or mutism	Yes, complete aphasia	No	Yes, complete aphasia	Yes, near complete aphasia
Severe anxiety	Yes	Yes	Yes	Yes
Obsessive thinking	Yes	Yes	Yes	Yes

	Case 1	Case 2	Case 3	Case 4
Insomnia	Yes	Yes	Yes	Yes
Loss of self-care	Yes	Yes	Yes	Yes
Urinary retention or incontinence	Yes: profound urinary retention and incontinence	Yes: sporadic episodes of daytime urinary incontinence during illness	Yes: sporadic incontinence	Yes: profound polyuria followed by urinary incontinence
Hyperkinetic	Yes: akathisia movements	Yes: athetosis 2–3 mo before neuropsychiatric onset; tremor; facial tics	Yes: akathisia	Yes: chorea; tremor
Catatonia	Yes	Yes	Yes	No
Other	Aggression, bowel incontinence, pica	Exaggerated startle	Paranoia	Agraphia
Diagnosics studies				
Neurologic examination at first evaluation	Nonfocal	Nonfocal, choreiform movements of toes	Nonfocal	Choreiform hand movements and tremor
Neuroimaging	MRI of the brain with mild ventricular prominence	MRI of the brain unrevealing	MRI/MRA brain, CTA of the head unrevealing	MRI of the brain unrevealing
Elevated Ab titers	Antinuclear Ab + 1:80 antithyroxidase Ab (>830 U/mL; normal <4) antithyroglobulin Ab (208 IU/mL; normal <2)	Elevation in anti-calcium-dependent kinase II serology	Antinuclear Ab + 1:160	Antinuclear Ab + 1:320, positive histone Abs
Other abnormal study results	Complement C4 (low) persistent monocytosis C1q binding (elevated) thyroid-stimulating hormone (elevated) T4 alternately high and low at different intervals	Borderline (+) titer for <i>Bartonella</i>	EEG with diffuse slowing	Low complement C4 (subsequently normalized)
Cerebrospinal studies	White cells, red cells, glucose, protein, IgG index, and oligoclonal bands all normal	White cells, red cells, glucose, and protein all normal	9 white cells (67% neutrophils, 33% lymphocytes; normal range 0–5 white cells) protein, glucose, IgG index, oligoclonal bands, lactate, <i>Cryptococcus</i> antigen, India ink, Gram-stain, enterovirus PCR, human herpesvirus 6 PCR, mycoplasma PCR, tubeola IgG and IgM, herpes simplex 1 and 2 PCR, HIV PCR, NMDA receptor Ab, and paraneoplastic Ab panel were all normal	White cells, red cells, glucose, protein, IgG index, and oligoclonal bands all normal
Other normal or negative study results	Normal Ab serologies: Mayo autoimmune encephalopathy panel including NMDA receptor, antistreptolysin 0 B-cell and T-cell immunophenotyping panel, quantitative immunoglobulins Normal genetic and/or metabolic: single nucleotide polymorphism array, fragile X, urine organic acids, plasma amino acids, carnitine, ammonia, zinc, methylmalonic acid, acylcarnitine profile, urine glycosaminoglycans EEG unrevealing	NMDA receptor Ab EEG unrevealing Abdominal US: negative results for teratoma	Sedimentation rate, C-reactive protein, thyroid-stimulating hormone, antithyroglobulin Ab, serum immunoglobulins, C4, C3, paraneoplastic Ab panel, serum NMDA receptor Abs, angiotensin-converting enzyme, ANCA, <i>Bartonella</i> IgG and IgM, lysosomal and peroxisomal panel, urine oligosaccharide and mucopolysaccharide screen, and porphyrins Full-body PET: pelvic ultrasound	Normal Ab serologies: NMDA receptor; thyroxidase, thyroglobulin, double-stranded DNA, voltage-gated potassium channels, anti-Ro, anti-La, phospholipid, cardiolipin, and $\beta$ -2-glycoprotein I, neutrophils, paraneoplastic antigens Infectious studies: <i>Bartonella</i> species, <i>Ehrlichia chaffeensis</i> , <i>Babesia microti</i> , <i>Leishmania</i> , Lyme panel, West Nile virus, herpes simplex viruses 1 and 2, Epstein-Barr virus, syphilis, and enterovirus Genetic and/or metabolic: lactate,

	Case 1	Case 2	Case 3	Case 4
	<p>Pelvic ultrasound normal</p> <p>Abdominal MRI notable for mild splenic enlargement</p>			<p>pyruvate, ammonia, fatty acid profile, acylcarnitine profiles, amino acid profiles of urine, serum and CSF, mucopolysaccharide and oligo- and polysaccharide profiles, cytogenetic fluorescence in situ hybridization, fragile X, array comparative genomic hybridization, and mercury</p>
Therapeutic interventions				
Immunotherapy regimens and response	<p>Daily oral prednisone 2 times per wk: no improvement</p> <p>IVIG 2 g/kg monthly for 4 mo: consistent, temporally correlated improvement in neuropsychiatric symptoms with waning effect before next dose</p> <p>IVIG 1 g/kg and IV methylprednisolone 1 g coadministration monthly (ongoing): similar to above</p>	<p>IVIG 2 g/kg twice accompanied by azithromycin, correlated with complete resolution of obsessive-compulsive behaviors within 4 wk; improvement beyond premorbid baseline over 2 mo</p> <p>Second dose of IVIG during "manic relapse" correlated with resolution of mood symptoms and return to baseline</p>	<p>IV methylprednisolone 30 mg/kg per d for 3 d, then monthly for 3 mo</p> <p>IVIG 2 g/kg once followed by IVIG 1 g/kg monthly</p> <p>Mycophenolate 750 mg BID with improvement in many symptoms and multiple less severe relapses when weaned off above regimen</p>	<p>IV methylprednisolone 1000 mg daily for 3 d; language improved within 3 d</p> <p>IVIG: temporally correlated with symptom improvement</p> <p>Rituximab and mycophenolate: neuropsychiatric symptoms generally maintained at baseline while on rituximab and mycophenolate but relapsed when weaned</p> <p>Cyclophosphamide was used to treat refractory relapse.</p> <p>Rituximab and prednisone: near baseline but with mild psychotic exacerbations while on maintenance</p>
Other therapeutic modifications during immunotherapy course	<p>Weaned off risperidone and lorazepam during immunotherapy</p> <p>Addition of levonorgestrel-ethinyl after 1 y of immunotherapy</p> <p>Also received trazodone, vitamin D, propranolol, and melatonin for discrete periods</p>	<p>Lorazepam: coadministered with a course of IVIG with immediate, marked improvement</p> <p>Clonazepam: initiated as maintenance therapy after lorazepam course</p> <p>Olanzapine: variably stopped and started during treatment course</p> <p>Daily oral contraceptive: maintained through most of course</p> <p>Metformin (for olanzapine associated wt gain)</p> <p>Clonidine: variably stopped and started to treat agitation</p> <p>Lamotrigine: started after seizure</p>	<p>Sertraline: used for obsessive-compulsive features and anxiety</p> <p>Aripiprazole: started for irritability and/or repetitive behaviors</p> <p>Methylphenidate: for ADHD, continues to take</p> <p>Clonidine: used intermittently for sleep, ADHD symptoms</p> <p>Lorazepam: used for agitation and catatonia, currently weaning off this medication</p> <p>Lithium: started during most recent relapse to treat manic symptoms</p> <p>Metformin: for aripiprazole associated wt gain</p>	<p>Multiple trials of antipsychotics, antidepressants, sleep medications, and benzodiazepines</p> <p>Slowly weaned off all psychiatric medications after initiation of immunotherapy</p>
Basic ADL measures at baseline, nadir, and last follow-up				
At premorbid baseline	<p>Independent: 5 (transferring, dressing, eating, toileting, continence)</p> <p>Dependent: 1 (bathing)</p>	<p>Independent: 6 (transferring, bathing, dressing, eating, toileting, continence)</p> <p>Dependent: 0</p>	<p>Independent: 6 (transferring, bathing, dressing, eating, toileting, continence)</p> <p>Dependent: 0</p>	<p>Independent: 6 (transferring, bathing, dressing, eating, toileting, continence)</p> <p>Dependent: 0</p>
At symptomatic nadir	<p>Independent: 0</p> <p>Dependent: 6 (transferring, bathing, dressing, eating, toileting, continence)</p>	<p>Independent: 2 (transferring, toileting)</p> <p>Dependent: 4 (bathing, dressing, eating, continence)</p>	<p>Independent: 0</p> <p>Dependent: 6 (transferring, bathing, dressing, eating, toileting, continence)</p>	<p>Independent: 0</p> <p>Dependent: 6 (transferring, bathing, dressing, eating, toileting, continence)</p>

	Case 1	Case 2	Case 3	Case 4
Most recent	Independent: 2 (transferring, eating) Dependent: 4 (bathing, dressing, toileting, continence)	Independent: 6 (transferring, bathing, dressing, eating, toileting, continence) Dependent: 0	Independent: 6 (transferring, bathing, dressing, eating, toileting, continence) but impulsivity warrants supervision Dependent: 0	Independent: 6 (transferring, bathing, dressing, eating, toileting, continence) Dependent: 0

Ab, antibody; ADHD, attention-deficit/hyperactivity disorder; ANCA, antineutrophil cytoplasmic antibodies; BID, bis in die; C1q, complement component 1q; C3, complement component 3; C4, complement component 4; CSF, cerebrospinal fluid; CT, computed tomography; IgG, immunoglobulin G; IgM, immunoglobulin M; La, anti-La/SSB antibody; MRA, magnetic resonance angiography; PCR, polymerase chain reaction; PET, positron emission tomography; Ro.; anti-Ro/SSA antibody; T4, thyroxine T4; US, ultrasound; —, not applicable.