

Basic Study

Circular RNA AKT3 governs malignant behaviors of esophageal cancer cells by sponging miR-17-5p

Hong-Liang Zang, Fu-Jian Ji, Hai-Ying Ju, Xiao-Feng Tian

ORCID number: Hong-Liang Zang 0000-0002-9837-0805; Fu-Jian Ji 0000-0002-6041-6089; Hai-Ying Ju 0000-0001-8243-5224; Xiao-Feng Tian 0000-0002-6544-0395.

Author contributions: Zang HL contributed to the study design and reviewed the manuscript; Ji FJ and Tian XF analyzed the data and wrote the manuscript; Ju HY and Zang HL contributed to the data collection, data interpretation, and manuscript writing; all authors read and approved the final manuscript.

Institutional review board

statement: This research was reviewed and approved by the Ethics Committee of China-Japan Union Hospital of Jilin University and complied with the guidelines of Declaration of Helsinki.

Institutional animal care and use

committee statement: The procedures were in accordance with guidelines of Animal Use and Care Committee of China-Japan Union Hospital of Jilin University.

Conflict-of-interest statement: All the authors have no conflict of interest related to the manuscript.

Data sharing statement: No additional data are available.

Hong-Liang Zang, Xiao-Feng Tian, Department of Hepatobiliary and Pancreatic Surgery, China-Japan Union Hospital of Jilin University, Changchun 130033, Jilin Province, China

Fu-Jian Ji, Department of Colorectal Surgery, China-Japan Union Hospital of Jilin University, Changchun 130033, Jilin Province, China

Hai-Ying Ju, Department of Hematology, Jilin Province Blood Center, Changchun 130000, Jilin Province, China

Corresponding author: Xiao-Feng Tian, MD, Chief Doctor, Department of Hepatobiliary and Pancreatic Surgery, China-Japan Union Hospital of Jilin University, No. 126 Xiantai Street, Erdao District, Changchun 130033, Jilin Province, China. fengwykctut4210@163.com

Abstract**BACKGROUND**

Recent studies have demonstrated that circular RNA AKT3 (circAKT3) plays a crucial role in regulating the malignant phenotypes of tumor cells. However, the potential effects of circAKT3 on esophageal cancer have not been investigated.

AIM

To illuminate the role of circAKT3 in malignant behaviors of esophageal cancer cells and its underlying mechanism.

METHODS

Clinical samples were collected to detect the expression of circAKT3. The role of circAKT3 in proliferation, migration, invasion, and apoptosis of esophageal cancer cells was evaluated using Cell Counting Kit-8, wound healing assays, Transwell assays, and fluorescence analysis, respectively. The target of circAKT3 was screened and identified using an online database and luciferase reporter assay. A xenograft nude mouse model was established to investigate the role of circAKT3 *in vivo*.

RESULTS

In vitro assays showed that proliferative, migratory, and invasive capacities of esophageal cancer cells were significantly enhanced by circAKT3 overexpression. Furthermore, miR-17-5p was screened as the target of circAKT3, and miR-17-5p antagonized the effects of circAKT3 on esophageal cancer cells. Moreover, we identified RHOC and STAT3 as the direct target molecules of miR-17-5p, and

ARRIVE guidelines statement: The authors have read the ARRIVE guidelines, and the manuscript was prepared and revised according to the ARRIVE guidelines.

Open-Access: This article is an open-access article that was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution NonCommercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Manuscript source: Unsolicited manuscript

Specialty type: Gastroenterology and hepatology

Country/Territory of origin: China

Peer-review report's scientific quality classification

Grade A (Excellent): A
Grade B (Very good): 0
Grade C (Good): 0
Grade D (Fair): 0
Grade E (Poor): 0

Received: October 30, 2020

Peer-review started: October 30, 2020

First decision: November 30, 2020

Revised: December 4, 2020

Accepted: December 16, 2020

Article in press: December 16, 2020

Published online: January 21, 2021

P-Reviewer: Kravtsov V

S-Editor: Chen XF

L-Editor: Filipodia

P-Editor: Ma YJ



circAKT3 facilitated expression of *RHOC* and *STAT3* by inhibiting miR-17-5p. *In vivo* assays showed *circAKT3* knockdown inhibited growth of esophageal cancer.

CONCLUSION

CircAKT3 contributed to the malignant behaviors of esophageal cancer *in vitro* and *in vivo* by sponging miR-17-5p thus providing a potential target for treatment of esophageal cancer.

Key Words: Esophageal cancer; Circular RNA AKT3; miR-17-5p; Proliferation; Migration; Invasion

©The Author(s) 2021. Published by Baishideng Publishing Group Inc. All rights reserved.

Core Tip: Circular RNA AKT3 was overexpressed in esophageal cancer and enhanced proliferative, migratory, and invasive capacities of esophageal cancer cells by sponging miR-17-5p to exert protumor effects. Downregulation of circular RNA AKT3 inhibited xenograft tumor growth of esophageal cancer *in vivo*. This research provided a potential target for better understanding the underlying mechanism of esophageal cancer.

Citation: Zang HL, Ji FJ, Ju HY, Tian XF. Circular RNA AKT3 governs malignant behaviors of esophageal cancer cells by sponging miR-17-5p. *World J Gastroenterol* 2021; 27(3): 240-254

URL: <https://www.wjgnet.com/1007-9327/full/v27/i3/240.htm>

DOI: <https://dx.doi.org/10.3748/wjg.v27.i3.240>

INTRODUCTION

As one of the most malignant tumors, esophageal cancer ranks in the top ten in morbidity and mortality of the tumors worldwide^[1]. However, the exact mechanism is still in discussion. Currently, surgery is the predominant approach for treatment of esophageal cancer in combination with some other alternatives^[2,3]. Although great progress has been made in the diagnosis and treatment of esophageal cancer, the prognosis has not been improved satisfactorily with a 5-year survival rate of less than 30%^[4]. Thus, investigating the underlying mechanism is conducive to deepening the pathogenesis of esophageal cancer and developing targeted therapies.

Circular RNAs (circRNAs) belong to endogenous noncoding RNAs, which could be expressed in eukaryotic cells^[5]. Unlike other RNA molecules, circRNAs possess the structure of a closed loop and cannot be degraded by RNase R^[6]. Accumulating evidence has demonstrated that circRNAs participate in the modulation of protein expression by sponging microRNAs (miRNAs) or interacting with RNA binding proteins^[7,8]. In tumor tissues and cell lines, the aberrant expression of circRNAs is involved in the regulation of malignant phenotypes of tumor cells. Wang *et al*^[9] reported that circLMTK2 is overexpressed in gastric cancer, which could also be a predictor for a poor prognosis in patients with gastric cancer. In colorectal cancer, *circFBXW7* is significantly downregulated, and *circFBXW1* overexpression suppresses proliferative, migratory, and invasive abilities of colorectal cancer cells^[10]. Therefore, differential expression of circRNAs exerts the opposite effects on tumor cells. However, only a few studies have concentrated on the effects of circRNAs on esophageal cancer.

Circular RNA AKT3 (circAKT3) is a novel molecule, which is derived from the *AKT3* gene and identified in gastric cancer with cisplatin resistance^[11]. The *in vitro* and *in vivo* assays have revealed that circAKT3 is overexpressed in cisplatin resistant-gastric cancer and leads to the increased expression of *PIK3R1* to mediate resistance of gastric cancer cells to cisplatin^[11]. However, Xia *et al*^[12] demonstrated that glioblastoma expressed a low level of *circAKT3*, and *circAKT3* overexpression suppressed proliferation and promoted sensitivity to radiation of glioblastoma cells. The contradictory findings remind us of the potential role of circAKT3 in esophageal cancer.

MATERIALS AND METHODS

Collection of clinical samples

The clinical tissue samples were acquired from patients with esophageal cancer ($n = 82$) who underwent surgery at China-Japan Union Hospital of Jilin University. The relative expression of *circAKT3* less than the mean value was classified into the *circAKT3* low group ($n = 43$), and the relative expression of *circAKT3* more than the mean value was classified into the *circAKT3* high group ($n = 39$).

In vivo assay

The nude mice ($n = 12$) were purchased from Experimental Animal Center of China-Japan Union Hospital of Jilin University, housed in cages under the condition of the 12-h shift of light-dark cycles, and randomly divided into two groups. Negative control short hairpin RNA and short hairpin RNA against *circAKT3*, purchased from Integrated Biotech Solutions Co. (China), were transfected into TE-1 cells in the presence of puromycin (3 mg/mL; Thermo Fisher Scientific, United States) to establish the *circAKT3* knockdown cell lines. The mice injected with the negative control short hairpin RNA-transfected TE-1 cells were designated as the negative control short hairpin RNA group, and mice injected with the short hairpin RNA against *circAKT3*-transfected TE-1 cells were designated as the short hairpin RNA against *circAKT3* group. After feeding for 30 d, the mice were anesthetized with pentobarbital sodium (25 mg/kg; Sigma-Aldrich, United States) intraperitoneally and killed by decapitation. The xenografts were isolated carefully and weighed.

Cell culture

Human esophageal cancer cell lines (KYSE-150, TE-10, TE-1) and human embryonic kidney cell line (HEK-293T) were cultured in Dulbecco's Modified Eagle Medium containing 10% fetal bovine serum (Gibco, United States). Human normal esophageal epithelial cell line (HEEC) was cultured in Dulbecco's Modified Eagle Medium with high glucose containing 10% fetal bovine serum (Gibco, United States). All of the cells were cultured in a CO₂ incubator chamber (Thermo Fisher Scientific, United States).

Quantitative real time-polymerase chain reaction

TRIzol (Ambion, United States) was utilized to extract the total RNA. SuperScript III (Invitrogen, United States) was used to obtain cDNA. SYBR Premix Ex Taq II (Takara, Japan) was utilized to perform quantitative real time(qRT)-PCR. The primer sequences of *circAKT3* and U6 were adopted as previously reported^[11]. The primer sequences of miR-17-5p were obtained from recent research^[13]. The primer sequences of AKT3, RHOC, STAT3, and GAPDH were obtained from the NCBI online database (<https://www.ncbi.nlm.nih.gov/gene/>). *CircAKT3* (forward, 5'-TCCAAATAAACGCCT TGGTGG-3'; reverse, 5'-CCTCAGAGAACACCCGCTCT-3'); miR-17-5p (forward, 5'-CGGCGCAAAGTGCTTACAG-3'; reverse, 5'-GTGCAGGGTCCGAGGT-3'); U6 (forward, 5'-CTCGCTTCGGCAGCAC-3'; reverse, 5'-AACGCTTCACGA ATTGCGT-3'); AKT3 (forward, 5'-CGGCTTCTCACGGATCAC-3'; reverse, 5'-CGGGACACTTTCCTCCTCC-3'); RHOC (forward, 5'-AAGTTCCCTTTGCCCG TCTG-3'; reverse, 5'-ACAGTGGCAACTCAAGGGTC-3'); STAT3 (forward, 5'-CCAGGTACCGTGTGTCAAGC-3'; reverse, 5'-CAGACCTGACACCTGTGTTG-3'); GAPDH (forward, 5'-TGAAATGTGCACGCACCAAG-3'; reverse, 5'-GGGAAGCAGCATTTCAGGTCT-3'). All of the primers were synthesized by Sangon (China).

Cell transfection

The siRNAs against *circAKT3*, miR-17-5p mimics, miR-17-5p inhibitor, indicated negative control, and plasmids were synthesized and purchased from Sangon (China). Lipofectamine 3000 (Invitrogen, United States) was used to conduct the transfection. RNase R (Epicentre, United States) was used to digest linear RNAs (but not circRNAs) in line with the manufacturer's instruction.

Fluorescence analysis

The Hoechst Staining Kit (Beyotime, China) was used to evaluate apoptotic cells. In brief, TE-1 cells were seeded at a density of 2.0×10^5 cells per well. After cell transfection with indicated vectors, 4% paraformaldehyde was used to fix the cultured TE-1 cells, followed by using Hoechst 33258 to stain the cells. A fluorescence microscope (Olympus, Japan) was utilized to observe the cells.

Measurement of cell viability

The Cell Counting Kit-8 (CCK-8; Beyotime, China) was utilized to measure cell viability. After transfection, TE-1 cells were seeded at a density of 1×10^4 cells per well for the indicated time. CCK-8 solution was added into each well for 2 h, and the Microplate Reader (Bio-Rad, United States) was used to record the value at 450 nm.

Wound healing assay

The wound healing assay was used to measure the migratory ability. After transfection, TE-1 cells were seeded into a 6-well plate for 24 h. The cells were scratched by a 10 mL Finntip and cultured in serum-free medium. Images were captured at 0 h and 24 h and analyzed by ImageJ software.

Measurement of invasive ability

The Transwell assay was conducted to evaluate the invasive ability. TE-1 cells with indicated transfection were cultured in Millicell Standing Cell Culture 24 well (Millipore, United States). After incubation for 24 h, glutaraldehyde solution was used to fix the cells, and crystal violet was used to stain the cells.

Luciferase reporter assay

ENCORI database was used to acquire the predicted binding sites between circAKT3 and miR-17-5p. TargetScanHuman database was utilized to acquire the potential binding sites of miR-17-5p with RHOC or STAT3. The procedures of the luciferase reporter assay were conducted in line with the previous study^[11]. In our research, HEK-293T cells were utilized to detect the binding ability by cotransfecting the indicated vectors (Sangon, China) with Lipofectamine 3000 (Invitrogen, United States). The luciferase activity was evaluated by Bio-Glo™ Luciferase Assay System (Promega, United States).

Western blot

Western blot was carried out in line with the previous report^[14]. In brief, the total protein was lysed by RIPA lysis buffer (Beyotime, China), and the BCA Kit (Thermal Fisher Scientific, United States) was used to detect the precise concentration. The total protein was added into SDS-PAGE gel (Beyotime, China). Primary antibodies [STAT3 rabbit monoclonal antibody (Cell Signaling Technology, United States); RHOC rabbit monoclonal antibody (Cell Signaling Technology, United States); GAPDH mouse monoclonal antibody (Cell Signaling Technology, United States)] were used overnight at 4 °C. Secondary antibodies with horseradish peroxidase (Beyotime, China) were applied for 2 h, and the indicated proteins were monitored by ChemDoc™ XRS+ System (Bio-Rad, United States).

Statistical analysis

The data in our research were displayed as mean \pm standard deviation and analyzed by GraphPad Prism 8.0 Software (GraphPad, United States). The *t*-test, two-way ANOVA or Chi-squared test was used. Statistical significance was set as the *P* value less than 0.05.

RESULTS

CircAKT3 was overexpressed in esophageal cancer

First, we detected the expression of *circAKT3* in the esophageal cancer tissue samples and cell lines. The qRT-PCR results showed that *circAKT3* was significantly upregulated in esophageal cancer tissue samples (Figure 1A). The characteristic analysis of the patients with esophageal cancer indicated that the tumor size, lymphatic metastasis, and clinical tumor node metastasis staging were positively related to the upregulation of *circAKT3* in the tissue samples (Table 1). We also found that human esophageal cancer cell lines TE-1 cells, TE-10 cells, and KYSE-150 cells expressed a significantly higher level of *circAKT3* compared to human normal esophageal epithelial cell line HEEC cells (Figure 1B).

CircAKT3 promoted malignant behaviors of TE-1 cells

Next, siRNAs against *circAKT3* were transfected into TE-1 cells, and qRT-PCR analysis revealed a marked efficiency of *circAKT3* knockdown (Figure 1C). As the differential structure between *circAKT3* and AKT3 mRNA, *circAKT3* was resistant to RNase R,

Table 1 Characteristics of patients with esophageal cancer

Characteristic	CircAKT3 low	CircAKT3 high	P value
Age in yr			0.824
< 50	19	19	
≥ 50	24	20	
Tumor diameter in cm			0.026
< 3	27	14	
≥ 3	16	25	
Lymphatic metastasis			0.027
No	24	12	
Yes	19	27	
TNM staging			0.016
I-II	26	13	
III-IV	17	26	

CircAKT3: Circular RNA AKT3; TNM: Tumor node metastasis.


while AKT3 mRNA was sensitive to RNase R (Figure 1C and 1D), which confirmed the circular structure of circAKT3. Moreover, the *circAKT3* overexpression plasmids were transfected into TE-1 cells and could significantly increase the expression of *circAKT3* (Figure 1E). CCK-8 assay showed that *circAKT3* siRNAs inhibited proliferation of TE-1 cells, while *circAKT3* overexpression plasmids facilitated proliferation of TE-1 cells (Figure 1F). We also found that *circAKT3* knockdown suppressed migration and invasion of TE-1 cells, while *circAKT3* overexpression exerted the opposite role by conducting the wound healing assay and Transwell assay, respectively (Figure 1G and 1H). We also measured the effect of *circAKT3* on apoptosis. The fluorescence analysis indicated that *circAKT3* siRNAs increased the number of condensed nuclei in TE-1 cells; however, *circAKT3* overexpression had no effect on the apoptotic rate of TE-1 cells (Figure 1I). These findings suggested that *circAKT3* facilitated proliferative, migratory, and invasive capacities of esophageal cancer cells.




***miR-17-5p* acted as the downstream molecule of *circAKT3* to antagonize the effects of *circAKT3* on TE-1 cells**

Accumulating evidence has reported that circRNAs regulate cell functions by sponging miRNAs^[7]. The ENCORI database (<http://starbase.sysu.edu.cn/index.php>) was utilized to predict the potential target of *circAKT3*, and we identified *miR-17-5p* as the target of *circAKT3*. We carried out a luciferase reporter assay and found that TE-1 cells with wild-type *circAKT3* and *miR-17-5p* mimics cotransfection presented a remarkably decreased luciferase activity (Figure 2A). Next, we detected the expression of *miR-17-5p* and found a lower level of *miR-17-5p* in esophageal cancer tissue samples (Figure 2B). Moreover, *miR-17-5p* expression was inversely associated with the expression of *circAKT3* in clinical tissue samples (Figure 2C). Meanwhile, esophageal cancer cell lines showed decreased expression of *miR-17-5p* (Figure 2D). These results indicated that *miR-17-5p* was inhibited by *circAKT3*. We then explored the role of *miR-17-5p* functioning as a downstream molecule of *circAKT3* in TE-1 cells. CCK-8 assay showed that cell viability of TE-1 cells with cotransfection of *circAKT3* overexpression plasmids and *miR-17-5p* mimics was decreased compared to that with *circAKT3* overexpression plasmids transfection (Figure 2E). Moreover, *miR-17-5p* mimics suppressed migratory and invasive abilities of *circAKT3* overexpression plasmids-transfected TE-1 cells by the wound healing assay and Transwell assay, respectively (Figure 2F and 2G). In parallel, *miR-17-5p* mimics increased the apoptotic rate of *circAKT3* overexpression plasmids-transfected TE-1 cells (Figure 2H).

***RHOC* and *STAT3* were modulated by the *circAKT3/miR-17-5p* axis**

We then identified *RHOC* and *STAT3* as the potential targets of *miR-17-5p* by using TargetScanHuman database (http://www.targetscan.org/vert_72/). The luciferase








Figure 1 Circular RNA AKT3 promoted malignant phenotypes of TE-1 cells. A: The expression of *circular RNA AKT3* (*circAKT3*) in tissue samples; B: The expression of *circAKT3* human normal esophageal epithelial cell line (HEEC) and human esophageal cancer cell lines; C and D: TE-1 cells were transfected with control oligo and *circAKT3* siRNAs in the presence or absence of RNase R to detect the expression of *circAKT3* and *AKT3*; E: The expression of *circAKT3* in TE-1 cells with *circAKT3* overexpression (OE); F: Cell viability of TE-1 cells as detected by the Cell Counting Kit-8 assay was performed to detect cell viability of TE-1 cells with indicated transfection; G: Migratory capacity of TE-1 cells with *circAKT3* OE and knockdown; H: Invasive ability of TE-1 cells with *circAKT3* OE and knockdown; I: Apoptosis of TE-1 cells with *circAKT3* OE and knockdown by Hoechst staining. ^a $P < 0.05$; ^b $P < 0.01$; ^c $P < 0.001$. NC: Negative control; OD: Optical density.

reporter assay showed that TE-1 cells with cotransfection of wild-type RHOC and miR-17-5p mimics presented a lower luciferase activity (Figure 3A). In addition, the decreased luciferase activity was also shown in TE-1 cells with cotransfection of wild-type STAT3 and miR-17-5p mimics (Figure 3B). Next, we detected the mRNA level of RHOC and STAT3 in clinical samples and found that both RHOC and STAT3 were upregulated in esophageal cancer tissue samples (Figure 3C and 3D). Moreover, the expression of RHOC and STAT3 was increased in TE-1 cells, TE-10 cells, and KYSE-150 cells (Figure 3E and 3F). Western blot analysis showed that *circAKT3* promoted the expression of RHOC and STAT3 in TE-1 cells, and miR-17-5p led to the reduced expression of RHOC and STAT3, which was reversed by *circAKT3* overexpression plasmids cotransfection (Figure 3G). Conversely, *circAKT3* siRNA inhibited the expression of RHOC and STAT3 in TE-1 cells, and a miR-17-5p inhibitor promoted the expression of RHOC and STAT3, which was reverted by *circAKT3* siRNA cotransfection (Figure 3H). In consequence, *circAKT3* facilitated the expression of RHOC and STAT3 by sponging miR-17-5p in esophageal cancer cells.








***CircAKT3* knockdown suppressed growth of esophageal cancer in vivo**

CircAKT3 shRNA was transfected into TE-1 cells to establish the stable *circAKT3* knockdown cell line (Figure 4A). The *in vivo* assay showed that the size and weight of the xenograft tumor injected with *circAKT3* shRNA-transfected TE-1 cells were smaller and lighter (Figure 4B and 4C). The qRT-PCR analysis indicated that the xenograft tumor injected with *circAKT3* shRNA-transfected TE-1 cells expressed a lower level of *circAKT3*, RHOC, and STAT3 and a higher level of miR-17-5p (Figure 4D-G). In parallel, the protein levels of RHOC and STAT3 were decreased in the xenograft tumor injected with *circAKT3* shRNA-transfected TE-1 cells (Figure 4H).

DISCUSSION

Increased proliferative, migratory, and invasive abilities are hallmarks of tumor cells^[15]. Thus, exploring the exact mechanism of malignant phenotypes of tumor cells

A
 WT circAKT3 5'-gaAACUACUAGGUAAGGCACUUU-3'
 hsa-miR-17-5p 3'-gaUGGACGUGACAUU----CGUGAAA-5'
 Mut circAKT3 5'-gaGGAGCUUAGCGCGAGCUGUGCC-3'






Figure 2 miR-17-5p acted as the downstream molecule of *circular RNA AKT3* to antagonize the effects of *circular RNA AKT3* on TE-1 cells. A: Schematic of the miR-17-5p binding sites in wild-type (WT) *circular RNA AKT3* (*circAKT3*) and the mutant (Mut) *circAKT3* (up). Luciferase reporter assay was performed in HEK-293T cells to determine the binding ability between miR-17-5p and *circAKT3* (down); B: The expression of miR-17-5p in tissue samples; C: *CircAKT3* was negatively correlated with miR-17-5p in esophageal cancer tissue samples; D: miR-17-5p expression in indicated cell lines; E-H: Cell viability (E), migratory ability (F), invasive ability (G), and apoptosis (H) of *circAKT3* overexpression (OE)-transfected TE-1 cells in the presence or absence of miR-17-5p mimics. ^a $P < 0.05$; ^b $P < 0.01$; ^c $P < 0.001$. HEEC: Human normal esophageal epithelial cell line; NC: Negative control; OD: Optical density.

contributes to clarifying the pathogenesis of tumorigenesis. Herein, we reported the protumor role of *circAKT3* in esophageal cancer cells by measuring the proliferative, migratory, and invasive abilities. In addition, miR-17-5p acted as a sponge molecule to inhibit the effects of *circAKT3* on esophageal cancer cells. Furthermore, two protumor molecules, RHOC and STAT3, were the direct downstream targets of miR-17-5p, and *circAKT3* upregulated RHOC and STAT3 by inhibiting miR-17-5p in esophageal cancer cells. Finally, we found *circAKT3* knockdown inhibited growth of esophageal cancer *in vivo*.

CircRNAs function among the complex molecular network and are implicated in the expression of functional proteins^[5,7]. Mechanistically, the inhibitory effect of circRNAs on miRNAs is one of the predominant mechanisms involving regulation of proteins^[7]. Increasing evidence has focused on the effects of circRNAs on the modulation of malignant phenotypes of tumor cells; however, only a few studies explore the relationship between circRNAs and esophageal cancer. A recent study reported that *circZNF292* is upregulated in the esophageal cancer cell line Eca-109, and inhibition of *circZNF292* enhanced the expression of *miR-206* to limit the proliferative, migratory, and invasive ability of Eca-109 cells^[16]. In addition, Shi *et al*^[17] demonstrated that overexpressed *circPRKCI* promoted the expression of *AKT3* by inhibiting miR-3680-3p to facilitate proliferation and migration of esophageal cancer cells. In addition, *circSMAD7*^[18] and *circFOXO3*^[19] could sponge different miRNAs to suppress malignant phenotypes of esophageal cancer cells. Therefore, it can be inferred that circRNAs exert the opposite roles in development and progression of esophageal cancer.

Our research identified *circAKT3* as a motivator in development and progression of esophageal cancer cells. The gene symbol of *circAKT3* is *AKT3*, which is a key member of the PI3K/AKT signaling pathway^[11,20]. Previously, Huang *et al*^[11] reported that *circAKT3* overexpression in gastric cancer participated in cisplatin resistance by restraining miR-198. However, the contradiction about the effects of *circAKT3* on tumors is that *circAKT3* could suppress development of glioblastoma^[12] and




Figure 3 RHOC and STAT3 were modulated by the circular RNA AKT3/miR-17-5p axis. A: Schematic of the miR-17-5p binding sites in wild-type (WT) RHOC and the mutant (Mut) RHOC (up). Luciferase reporter assay was performed in HEK-293T cells to determine the binding ability between miR-17-5p and RHOC (down); B: Schematic of the miR-17-5p binding site in WT STAT3 and the Mut STAT3 (up). Luciferase reporter assay was performed in HEK-293T cells to determine the binding ability between miR-17-5p and STAT3 (down); C and D: The expression of RHOC (C) and STAT3 (D) in tissue samples; E and F: The expression of RHOC (E) and STAT3 (F) in indicated cell lines; G and H: Protein level of RHOC and STAT3 in TE-1 cells with indicated transfection. ^b*P* < 0.01; ^c*P* < 0.001. circAKT3: Circular RNA AKT3; HEEC: Human normal esophageal epithelial cell line; NC: Negative control; OE: Overexpression.

progression of clear cell renal cell carcinoma^[20]. These findings, combined with our results, indicate a tumor specificity of circAKT3. Moreover, we identified miR-17-5p as a downstream molecule of circAKT3 to suppress the effects of circAKT3 on esophageal cancer cells by evaluating proliferation, migration, and invasion of TE-1 cells and found that esophageal cancer tissue samples and cell lines showed a significantly lower level of miR-17-5p. It has been reported that the expression of *miR-17-5p* decreased esophageal cancer stem-like cells and mediated radiation resistance^[13], which further reinforced our findings.

Additionally, we screened two protumor molecules RHOC and STAT3 as the direct downstream targets of miR-17-5p. RHOC is one of the Ras homolog family members and functions as a protumor molecule in multiple tumors^[21,22]. Clinical data shows that RHOC is positively expressed in esophageal cancer tissue samples and could be regarded as a predictor for patients with a poor prognosis of esophageal cancer^[23,24]. The *in vitro* and *in vivo* experiments manifest that RHOC enhanced the proliferative and invasive capacity of esophageal cancer cells^[25]. Similar to RHOC, STAT3 is a pivotal molecule that accelerates development and progression of malignant tumors^[26,27]. In esophageal cancer cells, activation of STAT3 dramatically reduced apoptotic rate and contributed to increased proliferative ability^[28]. Moreover, STAT3 promoted the expression of programmed death ligand 1 to mediate immune evasion of esophageal cancer cells^[29].

CONCLUSION

Taken together, our research discovered that circAKT3 enhanced proliferative, migratory, and invasive capacities of esophageal cancer cells by sponging miR-17-5p to exert protumor effects, and downregulation of circAKT3 inhibited xenograft tumor growth of esophageal cancer *in vivo*, providing a potential target for better understanding the underlying mechanism of esophageal cancer.




Figure 4 Circular RNA AKT3 knockdown suppressed growth of esophageal cancer *in vivo*. A: The expression of circular RNA AKT3 (*circAKT3*) in TE-1 cells with *circAKT3* short hairpin (sh)RNA transfection; B and C: The size (B) and weight (C) of the xenograft tumor injected with *circAKT3* shRNA-transfected TE-1 cells; D-G: The expression of *circAKT3* (D), *miR-17-5p* (E), *RHOC* (F) and *STAT3* (G) in the xenograft tumor injected with *circAKT3* shRNA-transfected TE-1 cells by quantitative real time-polymerase chain reaction; H: The protein levels of *RHOC* and *STAT3* in the xenograft tumor injected with *circAKT3* shRNA-transfected TE-1 cells. * $P < 0.001$. sh-NC: Negative control shRNA; sh-*circAKT3*: Circular RNA AKT3 shRNA.

ARTICLE HIGHLIGHTS

Research background

Esophageal cancer is one of the most malignant tumors in the digestive system and exploring the underlying mechanism is a key issue that needs to be addressed. Circular RNA AKT3 (*circAKT3*) is a novel noncoding RNA participating in the development and progression of multiple tumors.

Research motivation

To identify biomarkers for the diagnosis and treatment of esophageal cancer.

Research objectives

To explore the mechanism of the role of *circAKT3* in the development of esophageal cancer.

Research methods

Cell Counting Kit-8, wound healing assay, Transwell assay, and fluorescence analysis

were used to evaluate the circAKT3 effect on proliferation, migration, invasion, and apoptosis of esophageal cancer cells.

Research results

In vitro assays results revealed that proliferative, migratory, and invasive capacities of esophageal cancer cells were significantly enhanced by circAKT3 overexpression. miR-17-5p was screened as the target of circAKT3, and miR-17-5p antagonized the effects of circAKT3 on esophageal cancer cells. Moreover, we identified RHOC and STAT3 as the direct target molecules of miR-17-5p, and circAKT3 facilitated expression of RHOC and STAT3 by inhibiting miR-17-5p. *In vivo* assays showed circAKT3 knockdown inhibited growth of esophageal cancer.

Research conclusions

CircAKT3 enhanced proliferative, migratory, and invasive capacities of esophageal cancer cells by sponging miR-17-5p to exert protumor effects, and downregulation of circAKT3 inhibited xenograft tumor growth of esophageal cancer *in vivo*, providing a potential target for better understanding the underlying mechanism of esophageal cancer.

Research perspectives

The role of circAKT3 in other tumors may be uncovered in the future, and its application in anticancer therapy will be promoted.

REFERENCES

- 1 Bray F, Ferlay J, Soerjomataram I, Siegel RL, Torre LA, Jemal A. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA Cancer J Clin* 2018; **68**: 394-424 [PMID: 30207593 DOI: 10.3322/caac.21492]
- 2 Lagergren J, Smyth E, Cunningham D, Lagergren P. Oesophageal cancer. *Lancet* 2017; **390**: 2383-2396 [PMID: 28648400 DOI: 10.1016/S0140-6736(17)31462-9]
- 3 van Rossum PSN, Mohammad NH, Vleggaar FP, van Hillegeersberg R. Treatment for unresectable or metastatic oesophageal cancer: current evidence and trends. *Nat Rev Gastroenterol Hepatol* 2018; **15**: 235-249 [PMID: 29235549 DOI: 10.1038/nrgastro.2017.162]
- 4 Ohashi S, Miyamoto S, Kikuchi O, Goto T, Amanuma Y, Muto M. Recent Advances From Basic and Clinical Studies of Esophageal Squamous Cell Carcinoma. *Gastroenterology* 2015; **149**: 1700-1715 [PMID: 26376349 DOI: 10.1053/j.gastro.2015.08.054]
- 5 Kristensen LS, Andersen MS, Stagsted LVW, Ebbesen KK, Hansen TB, Kjems J. The biogenesis, biology and characterization of circular RNAs. *Nat Rev Genet* 2019; **20**: 675-691 [PMID: 31395983 DOI: 10.1038/s41576-019-0158-7]
- 6 Xia L, Song M, Sun M, Wang F, Yang C. Circular RNAs as Biomarkers for Cancer. *Adv Exp Med Biol* 2018; **1087**: 171-187 [PMID: 30259366 DOI: 10.1007/978-981-13-1426-1_14]
- 7 Qu S, Yang X, Li X, Wang J, Gao Y, Shang R, Sun W, Dou K, Li H. Circular RNA: A new star of noncoding RNAs. *Cancer Lett* 2015; **365**: 141-148 [PMID: 26052092 DOI: 10.1016/j.canlet.2015.06.003]
- 8 Zang J, Lu D, Xu A. The interaction of circRNAs and RNA binding proteins: An important part of circRNA maintenance and function. *J Neurosci Res* 2020; **98**: 87-97 [PMID: 30575990 DOI: 10.1002/jnr.24356]
- 9 Wang S, Tang D, Wang W, Yang Y, Wu X, Wang L, Wang D. circLMTK2 acts as a sponge of miR-150-5p and promotes proliferation and metastasis in gastric cancer. *Mol Cancer* 2019; **18**: 162 [PMID: 31722712 DOI: 10.1186/s12943-019-1081-4]
- 10 Lu H, Yao B, Wen X, Jia B. FBXW7 circular RNA regulates proliferation, migration and invasion of colorectal carcinoma through NEK2, mTOR, and PTEN signaling pathways in vitro and in vivo. *BMC Cancer* 2019; **19**: 918 [PMID: 31519156 DOI: 10.1186/s12885-019-6028-z]
- 11 Huang X, Li Z, Zhang Q, Wang W, Li B, Wang L, Xu Z, Zeng A, Zhang X, Zhang X, He Z, Li Q, Sun G, Wang S, Li Q, Wang L, Zhang L, Xu H, Xu Z. Circular RNA AKT3 upregulates PIK3R1 to enhance cisplatin resistance in gastric cancer via miR-198 suppression. *Mol Cancer* 2019; **18**: 71 [PMID: 30927924 DOI: 10.1186/s12943-019-0969-3]
- 12 Xia X, Li X, Li F, Wu X, Zhang M, Zhou H, Huang N, Yang X, Xiao F, Liu D, Yang L, Zhang N. A novel tumor suppressor protein encoded by circular AKT3 RNA inhibits glioblastoma tumorigenicity by competing with active phosphoinositide-dependent Kinase-1. *Mol Cancer* 2019; **18**: 131 [PMID: 31470874 DOI: 10.1186/s12943-019-1056-5]
- 13 Lynam-Lennon N, Heavey S, Sommerville G, Bibby BA, Ffrench B, Quinn J, Gasch C, O'Leary JJ, Gallagher MF, Reynolds JV, Maher SG. MicroRNA-17 is downregulated in esophageal adenocarcinoma cancer stem-like cells and promotes a radioresistant phenotype. *Oncotarget* 2017; **8**: 11400-11413 [PMID: 28002789 DOI: 10.18632/oncotarget.13940]

- 14 **Zhan N**, Li B, Xu X, Xu J, Hu S. Inhibition of FASN expression enhances radiosensitivity in human non-small cell lung cancer. *Oncol Lett* 2018; **15**: 4578-4584 [PMID: 29541228 DOI: 10.3892/ol.2018.7896]
- 15 **Pastushenko I**, Blanpain C. EMT Transition States during Tumor Progression and Metastasis. *Trends Cell Biol* 2019; **29**: 212-226 [PMID: 30594349 DOI: 10.1016/j.tcb.2018.12.001]
- 16 **Liu Z**, Hu G, Zhao Y, Xiao Z, Yan M, Ren M. Silence of cZNF292 suppresses the growth, migration, and invasion of human esophageal cancer Eca-109 cells via upregulating miR-206. *J Cell Biochem* 2020; **121**: 2354-2362 [PMID: 31680303 DOI: 10.1002/jcb.29458]
- 17 **Shi N**, Shan B, Gu B, Song Y, Chu H, Qian L. Circular RNA circ-PRKCI functions as a competitive endogenous RNA to regulate AKT3 expression by sponging miR-3680-3p in esophageal squamous cell carcinoma. *J Cell Biochem* 2019; **120**: 10021-10030 [PMID: 30659640 DOI: 10.1002/jcb.28285]
- 18 **Zhang Y**, Wang Q, Zhu D, Rong J, Shi W, Cao X. Up-regulation of circ-SMAD7 inhibits tumor proliferation and migration in esophageal squamous cell carcinoma. *Biomed Pharmacother* 2019; **111**: 596-601 [PMID: 30611100 DOI: 10.1016/j.biopha.2018.12.116]
- 19 **Xing Y**, Zha WJ, Li XM, Li H, Gao F, Ye T, Du WQ, Liu YC. Circular RNA circ-Foxo3 inhibits esophageal squamous cell cancer progression via the miR-23a/PTEN axis. *J Cell Biochem* 2020; **121**: 2595-2605 [PMID: 31680314 DOI: 10.1002/jcb.29481]
- 20 **Xue D**, Wang H, Chen Y, Shen D, Lu J, Wang M, Zebibula A, Xu L, Wu H, Li G, Xia L. Circ-AKT3 inhibits clear cell renal cell carcinoma metastasis via altering miR-296-3p/E-cadherin signals. *Mol Cancer* 2019; **18**: 151 [PMID: 31672157 DOI: 10.1186/s12943-019-1072-5]
- 21 **Guan X**, Chen S, Zhao Y. The role of RhoC in malignant tumor invasion, metastasis and targeted therapy. *Histol Histopathol* 2018; **33**: 255-260 [PMID: 28664531 DOI: 10.14670/HH-11-915]
- 22 **Thomas P**, Pranatharthi A, Ross C, Srivastava S. RhoC: a fascinating journey from a cytoskeletal organizer to a Cancer stem cell therapeutic target. *J Exp Clin Cancer Res* 2019; **38**: 328 [PMID: 31340863 DOI: 10.1186/s13046-019-1327-4]
- 23 **Faried A**, Faried LS, Usman N, Kato H, Kuwano H. Clinical and prognostic significance of RhoA and RhoC gene expression in esophageal squamous cell carcinoma. *Ann Surg Oncol* 2007; **14**: 3593-3601 [PMID: 17896152 DOI: 10.1245/s10434-007-9562-x]
- 24 **Zhao ZH**, Tian Y, Yang JP, Zhou J, Chen KS. RhoC, vascular endothelial growth factor and microvascular density in esophageal squamous cell carcinoma. *World J Gastroenterol* 2015; **21**: 905-912 [PMID: 25624724 DOI: 10.3748/wjg.v21.i3.905]
- 25 **Faried A**, Faried LS, Kimura H, Nakajima M, Sohda M, Miyazaki T, Kato H, Usman N, Kuwano H. RhoA and RhoC proteins promote both cell proliferation and cell invasion of human oesophageal squamous cell carcinoma cell lines in vitro and in vivo. *Eur J Cancer* 2006; **42**: 1455-1465 [PMID: 16750623 DOI: 10.1016/j.ejca.2006.02.012]
- 26 **Huynh J**, Chand A, Gough D, Ernst M. Therapeutically exploiting STAT3 activity in cancer - using tissue repair as a road map. *Nat Rev Cancer* 2019; **19**: 82-96 [PMID: 30578415 DOI: 10.1038/s41568-018-0090-8]
- 27 **Thilakasiri PS**, Dmello RS, Nero TL, Parker MW, Ernst M, Chand AL. Repurposing of drugs as STAT3 inhibitors for cancer therapy. *Semin Cancer Biol* 2019 [PMID: 31711994 DOI: 10.1016/j.semcancer.2019.09.022]
- 28 **Liu Y**, Zhi Y, Song H, Zong M, Yi J, Mao G, Chen L, Huang G. S1PR1 promotes proliferation and inhibits apoptosis of esophageal squamous cell carcinoma through activating STAT3 pathway. *J Exp Clin Cancer Res* 2019; **38**: 369 [PMID: 31438989 DOI: 10.1186/s13046-019-1369-7]
- 29 **Zhang W**, Pang Q, Yan C, Wang Q, Yang J, Yu S, Liu X, Yuan Z, Wang P, Xiao Z. Induction of PD-L1 expression by epidermal growth factor receptor-mediated signaling in esophageal squamous cell carcinoma. *Onco Targets Ther* 2017; **10**: 763-771 [PMID: 28243112 DOI: 10.2147/OTT.S118982]



Published by **Baishideng Publishing Group Inc**
7041 Koll Center Parkway, Suite 160, Pleasanton, CA 94566, USA
Telephone: +1-925-3991568
E-mail: bpgoffice@wjgnet.com
Help Desk: <https://www.f6publishing.com/helpdesk>
<https://www.wjgnet.com>

