



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



# Polyphenols as promising biologically active substances for preventing SARS-CoV-2: A review with research evidence and underlying mechanisms

Taha Mehany<sup>a,1</sup>, Ibrahim Khalifa<sup>b,1</sup>, Hassan Barakat<sup>b,c,\*</sup>, Sami A. Althwab<sup>c</sup>,  
Yousef M. Alharbi<sup>d</sup>, Sobhy El-Sohaimy<sup>a,e</sup>

<sup>a</sup> Food Technology Department, Arid Lands Cultivation Research Institute, City of Scientific Research and Technological Applications, 21934, Alexandria, Egypt

<sup>b</sup> Food Technology Department, Faculty of Agriculture, Benha University, Moshohor, 13736, Egypt

<sup>c</sup> Department of Food Science and Human Nutrition, College of Agriculture and Veterinary Medicine, Qassim University, Buraydah, 51452, Saudi Arabia

<sup>d</sup> Department of Veterinary Medicine, College of Agriculture and Veterinary Medicine, Qassim University, Buraydah, 51452, Saudi Arabia

<sup>e</sup> Department of Technology and Organization of Public Catering, Institute of Sport, Tourism and Service, South Ural State University, 454080, Chelyabinsk, Russia

## ARTICLE INFO

### Keywords:

SARS-CoV2  
Polyphenols  
Functional foods  
Boosting immune functions

## ABSTRACT

Currently, antiviral drugs and/or vaccines are not yet available to treat or prevent severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). In this review, we narrated the available data, from credible publishers, regarding the possible role of polyphenols and natural extracts-containing polyphenols in the prevention of coronavirus disease 2019 (COVID-19), and their immune-boosting properties. It was revealed that polyphenols could be considered as promising biologically active substances for the prevention of COVID-19. The underlying potential mechanism behind this action is mostly due to the antiviral activities and the immune-regulation functions of polyphenols against COVID-19-infections. Antivirus polyphenolic-based medications can mitigate SARS-CoV-2-enzymes, which are vital for virus duplication and infection. It was also found that triterpenoid, anthraquinone, flavonoids, and tannins are possible keys to scheming antiviral therapies for inhibiting SARS-CoV-2-proteases. The identified pharmacophore structures of polyphenols could be utilized in the explanation of novel *anti*-COVID-19 designs. The advantage of using mixtures containing polyphenols is related to the high-safety profile without having major side-effects, but further randomized controlled trials are required in the upcoming studies.

## 1. Introduction

SARS-CoV-2 affects all peoples, especially those who had weak immune systems and/or weak immune-based responses. Coronaviruses (CoVs) belonged to the sub-family of *Ortho*-coronavirinae, under the Coronaviridae family and in order of Nidovirales, and this sub-family contains alpha-, beta-, gamma-, and delta-CoVs (Banerjee, Kulcsar, Misra, Frieman, & Mossman, 2019). The deadliness of CoVs has been identified by the outbreaks of severe observation of acute respiratory syndrome (SARS-CoV) in 2002/2003 and the Middle East Respiratory Syndrome (MERS-CoV) in 2012/2013. SARS-CoV and MERS-CoV are going to  $\beta$ -CoV (Zhang, Wu, Zhang, Deng, & Peng, 2020). Freshly, a new flu-like CoV (COVID-19) associated with SARS-CoV and MERS-CoV was

realized in the late of 2019 and spread worldwide with a pandemic (Cohen & Normile, 2020). The total number of cases globally was 91,322,850 and 1,952,979 confirmed deaths in WHO situation report COVID-19 in January 12, 2021 (worldmeters.info).

Hitherto, no vaccine and actual SARS-CoV-2 medications based on *anti*-SARS-CoV therapy by the scientific community were confirmed (Jo, Kim, Kim, Shin, & Kim, 2019), while scientists are trying to develop efficient vaccines or drugs for the SARS-CoV-2 virus. The general viral infection treatment involving dietary interferences and other immune enhancers which were utilized to promote immunity towards the viral poisons. Thus, they may also be utilized to contest SARS-CoV-2 by the nutritional status of each patient and should be assessed before applying the overall treatments (Zhang et al., 2020). Recently, numerous studies

**Abbreviations:** SARS-CoV-2, Severe acute respiratory syndrome coronavirus 2; COVID-19, Coronavirus disease 2019; CoVs, Coronaviruses; SARS-CoV, Severe observation of acute respiratory syndrome; MERS-CoV, Middle East Respiratory Syndrome.

\* Corresponding author. Food Technology Department, Faculty of Agriculture, Benha University, Moshohor, 13736, Egypt.

E-mail addresses: [hassan.barakat@fagr.bu.edu.eg](mailto:hassan.barakat@fagr.bu.edu.eg), [haa.mohamed@qu.edu.sa](mailto:haa.mohamed@qu.edu.sa) (H. Barakat).

<sup>1</sup> Authors are equally contributed.

<https://doi.org/10.1016/j.fbio.2021.100891>

Received 17 June 2020; Received in revised form 12 January 2021; Accepted 12 January 2021

Available online 20 January 2021

2212-4292/© 2021 Elsevier Ltd. All rights reserved.

have exposed that phytonutrients play a vital part in the stoppage of chronic syndromes, as most can be related to nourishments (Fraga, Croft, Kennedy, & Tomás-Barberán, 2019). Various countries have entered the notion of functional-based food not only essential for human life but also to boost the physiological functions and immunity responses (Del Bo et al., 2019; Quero, Marmol, Cerrada, & Rodríguez-Yoldi, 2020). The importance of the proper functioning of the body and the well-balanced immune system for the maintenance of human life has been remarkably evident over the last decades (Lopez-Varela, Gonzalez-Gross, & Marcos, 2002; Wichers, 2009).

In addition to phytonutrients mainly presented in fruit and vegetables, polyphenols have recently been targeted due to their role in the prevention of several diseases as described (Fraga et al., 2019). Interestingly, polyphenols have attracted much attention in recent years since their dietary consumption has been associated with the prevention of some chronic and degenerative diseases that are significant causes of death and incapacity in developing countries (Santos-Buelga, González-Paramás, Oludemi, Ayuda-Durán, & González-Manzano, 2019). In addition, due to the biological activities of polyphenols, particularly antimicrobial, antioxidant, anti-inflammatory, pro-apoptotic behavior, anti-proliferative, and hormonal control ability among polyphenols, countless plants have been considered and investigated because of the classification of new substances with nutraceutical possessions (Forni et al., 2019). Dietary polyphenols signify several bioactive substances broadly found in plant-based food which have the capabilities to prevent the beginning and development of several diseases, and to decrease and control their signs which become a central issue in food research (Del Bo et al., 2019; Quero et al., 2020).

The outbreak of the COVID-19 pandemic caused by SARS-CoV-2, along with the lack of targeted medicaments and vaccines, forced the scientific world to seek new antiviral formulations. This article reviews the feasibility of using polyphenols extracted from plants as a source of bioactive components (e.g., gallates and quercetin), effective in mitigating SARS-CoV-2 and boosting the immune system. There is potential for the utilization of plant-derived polyphenols as functional foods and pharmaceutical formulations. The method of employing such formulations is fast because the raw materials for their preparation, namely herbs and useable plants, are permitted for human consumption worldwide. Approximately 100 research papers were selected from Science Direct, Scopus, MDPI, PubMed, and Web of Science to review the recently published papers on the related topic with the aid of relative keywords.

### 1.1. Functional foods and their efficiency in boosting the immune response

In fact, malnutrition is the most common cause of immunodeficiency concerns worldwide (Schaible & Stefan, 2007). The immune system, which is joined into all physiological arrangements, defends the human body towards microorganisms' infections and other internal and external abuses (Castelo-Branco & Soveral, 2014; Maggini, Pierre, & Calder, 2018). And suitable nutrition at all phases of life is an important factor to guarantee a sufficient stock of energy, macro- and micro-nutrients essential for the growth, upkeep, and countenance of the immune response (Maggini, Wintergerst, Beveridge, & Hornig, 2007). Furthermore, undernutrition is leading to weakening immune function which can be due to consumption of fast food, insufficient intakes of energy and macronutrients, and/or shortages in the essential micro-nutrients (Alpert, 2017; Calder, 2013; Pereira, 2003). Fig S1. showed the sources of variation in the immune system and its relation with suboptimal nutrient status on an increased risk of infection, as previously described (Calder & Kew, 2007; Maggini et al., 2018), showing the vital role of the functional foods.

In this context, functional foods are used to strengthen other physiological functions including the immunity responses (Roberfroid, 2000). The characteristics of functional foods were tabulated in Table 1. Functional food components can be useful components that originate

**Table 1**

The Japanese "FOSHU" criteria for functional foods, (Iwatani & Yamamoto, 2019).

1. They can or should be consumed as a portion of a normal diet.
2. They are foods: not capsules, pills, or powder-based on natural food components.
3. They have a clear function on the human organism such as
3.1. Improving the immunity function.
3.2. Supporting recovery from specific diseases.
3.3. Preventing specific diseases.
3.4. Slowing down the aging process.
3.5. Controlling physical complaints.

logically in or complement food as functional elements such as vitamins, minerals, dietary fiber, fatty acids, amino acids, polyphenols, phyto-pigments, non-saponifiable fractions, and functional plant-based proteins (Guiné, Lima, & Barroca, 2009; London, 2010). Table 2 summarises the functional foods-based products which were quoted from the available data. These foods and/or ingredients boost the immune response by different mechanisms (Powell, Borchers, Yoshida, & Gershwin, 2000, pp. 21–31). Meanwhile, the deficiencies of one or more essential fatty and amino acids, vitamins, Se, Zn, Fe, Cu, and phytonutrients will affect almost all types of immunity (London, 2010; Maggini et al., 2018; Zhang et al., 2020). Regarding COVID-19 pandemic, these

**Table 2**

Overview of vital functional food components (non-polyphenols) in boosting up the immune response, virus targeted, and functions related, quoted from (London, 2010; Maggini et al., 2018; Zhang & Liu, 2020).

Food components	Health properties	Virus targeted and related functions
U-3 fatty acids Vitamin C	<ul style="list-style-type: none"> <li>Stimulation of the immune system.</li> <li>Stimulates production, function, and movement of leukocytes (e.g., lymphocytes, neutrophils, and phagocytes). Antimicrobial, natural killer cell activities, and chemotaxis. Regenerates other important antioxidants such as glutathione and vitamin E to their active state. Promotes collagen synthesis. Increases serum levels of complement proteins. Involved in apoptosis and clearance of spent neutrophils from sites of infection by macrophages.</li> </ul>	Influenza and human immunodeficiency virus. Avian coronavirus.
Vitamin D	<ul style="list-style-type: none"> <li>Stimulates immune cell proliferation and cytokine production and helps protect against infection caused by pathogens. Expressed in innate immune cells (e.g., monocytes, macrophages, dendritic cells).</li> </ul>	Bovine coronavirus.
Vitamin A	<ul style="list-style-type: none"> <li>Maintains the structural and functional integrity of mucosal cells in innate barriers (e.g., skin, and respiratory tract).</li> </ul>	Measles virus, human immunodeficiency virus, and avian coronavirus.
Vitamin B	<ul style="list-style-type: none"> <li>Boosting the immune response of the host.</li> </ul>	MERS-CoV.
Vitamin E	<ul style="list-style-type: none"> <li>Impairs humoral and cell-mediated aspects of adaptive immunity, i.e., B and T cell function.</li> </ul>	Coxsackievirus and bovine coronavirus.
Se	<ul style="list-style-type: none"> <li>Important for the antioxidant host defense system affecting leukocyte.</li> </ul>	Influenza virus and avian coronavirus.
Fe	<ul style="list-style-type: none"> <li>Stimulation of the immune system.</li> </ul>	Viral mutations.
Zn	<ul style="list-style-type: none"> <li>Stimulation of the immune system.</li> </ul>	Measles and SARS-CoV.

functional ingredients may mitigate SARS-CoV-2 through their immunoenhancing and antiviral activity. For example, Zhang et al. (2020) had claimed that vitamin A and/or its derivatives like retinoic acid could provide a promising alternative for CoV-therapy. Vitamin D has several immune-modulatory effects and might work as additional therapeutic against COVID-19 (Moriguchi and Muraga, 2000; Zhang et al., 2020). Zhang and Liu (2020) had stated that Zn may affect not only COVID-19-linked symptoms like inferior respiratory tract infection and diarrhea but also SARS-CoV-2 replication. Dosing the patients with Se could be an active choice for the cure of COVID-19 as previously stated by Zhang et al. (2020), for instance. These functional compounds are essential substances at the molecular and cellular levels that play critical functions.

## 2. The potential uses of plant extracts containing polyphenols against SARS-CoV-2

### 2.1. Overview of the health-benefits of polyphenols

Polyphenols constituents are categorized by the attendance of one or more hydroxy group on an aromatic circle. These components are categorized by their molecular mass, chemical construction, and intricacy to flavonoids (flavones, flavonols, flavanones, flavanonols, isoflavonoids, flavanols, anthocyanidins, and chalcones) and non-flavonoid ones (phenolic acids, stilbenes, curcuminoids, lignans, and tannins) (Quero et al., 2020). Fig S2. portrayed the plant polyphenolics classifications according to Phenol-Explorer (<http://www.phenol-explorer.eu>). Such polyphenols are valuable to human health and are capable to decrease the diseases perils (Rasouli, Farzaei, & Khodarahmi, 2017). Thus, polyphenols have recently attracted much attention to the scientific community due to their dietary intake which is related to the deterrence of some chronic and wasting diseases that establish the main reasons of death. The unique properties of their nature and position of the substituents have concerned for biological utilization, namely, antiviral, antibacterial, antioxidant, and anti-inflammatory activities. These polyphenols have been isolated from a great number of plants, such as fruits, vegetables, legumes, nuts, seeds, herbs, spices, tea, coffee, dark chocolate, and so on (Quero et al., 2020; Rasouli et al., 2017; Santos-Buelga et al., 2019; Silva & Pogačnik, 2020). Among the sources of polyphenols, citrus is a promising source due to its content from bioflavonoids including apigenin, acacetin, luteolin, diosmetin, chrysoeriol, and glycosides which play key roles in providing an extensive range of health useful effects. However, polyphenols' super-healthy properties depend on the amount ingested and their bioavailability (Silva & Pogačnik, 2020).

### 2.2. Polyphenols and immune system

Many factors cause immune disorders, including environmental stress and changes in the immune regulators that can lead to many human diseases. Bioactive components can play a vital role in immunity through intrusive with immune cell regulation and pro-inflammatory cytokine synthesis. Recently, a huge number of *in vitro* and *in vivo* studies demonstrated the immunomodulatory part of polyphenols. These studies have shown that people with a specific diet, especially polyphenols-rich ones, are at reduced risk for a variety of diseases (Rasouli et al., 2017; Yahfoufi, Alsadi, Jambi, & Matar, 2018). Additionally, polyphenols are developing therapeutic tools for controlling and protecting the body from several autoimmune disorders due to their immunomodulatory effects (Khalifa, Zhu, Mohammed, Dutta, & Li, 2020; Khan et al., 2019). The immune modulation effect of polyphenols is related to the impact on immune cells, pro-inflammatory gene expression, and modulate cytokines production (Yahfoufi et al., 2018). In addition to the antioxidant properties such as reactive oxygen species scavenging of polyphenols and their donations to the rule of inflammatory signing. Polyphenols also showed high-effects in increasing the

natural killers in fruits (Yahfoufi et al., 2018). For their effect on the balance between pro and anti-inflammatory cytokine secretion, polyphenols compounds such as catechins and quercetin have been added to various foods. Additionally, numerous flavonoids can inhibit the expression of various pro-inflammatory cytokines and chemokines (Comalada et al., 2006). Furthermore, the epigallocatechin gallate and curcumin can induce epigenetic changes in cells. Thus, polyphenols compounds can be efficiently used to control the immune response.

### 2.3. The antiviral activities of polyphenols

The literature review demonstrated that polyphenols have been investigated for their potential antiviral efficiency against SARS-CoV-2 in molecular modeling studies, cell-free polyphenol-protein interaction studies, and in cell-based virus infection studies (Chojnacka, Witek-Krowiak, Skrzypczak, Mikula, & Młynarz, 2020; Paraiso, Revel, & Stevens, 2020). Convincing data shows that polyphenols such as epigallocatechin gallate, resveratrol, and curcumin are prime candidates for preclinical and clinical trials. As a cautionary note, the efficacy of tested polyphenols against human viral disease *in silico* and *in vitro* screening methods are not validated. Potential regulation of COVID-19 intensity by polyphenols controlling ACE2 *in vivo* has been suggested (Horne & Vohl, 2020; Manoharan et al., 2020) but very few studies have been performed to examine the antiviral activity of polyphenols against SARS-CoV-2 *in vivo*. Pudilan Xiaoyan Oral Liquid (PDL), a traditional Chinese medicine containing four herbs and more than 180 ingredients, showed potent *anti*-SARSCoV-2 activity in infected hACE2 mice (Deng et al., 2020). It has also been stated that a nebulized formula of quercetin and *N*-acetylcysteine substantially relieved the respiratory symptoms of SARS-CoV-2 in a patient treated with hydroxychloroquine and antibiotics (Schettig et al., 2020). This reinforces the value of more clinical trials to determine the potential of polyphenol-based nutraceuticals as an adjuvant or key treatment for COVID-19. High-throughput screening methods will speed the *in vitro* development of lead candidates by reducing the number of polyphenol libraries. Screening of polyphenol-rich plant extracts is an alternative, commonly used method, but it has the drawback that extracts contain a multitude of natural products with inherent difficulties of not being able to reliably distinguish active concepts and the potential for pharmacological antagonism. These problems can be solved by integrating classical bioassay-guided fractionation with machine learning approaches that expose the identity of bioactive natural products in extracts without the need for homogeneity purification. In our view, the above hybrid method aims to accelerate research because many antiviral *in vitro* experiments can be conducted without the treatment of live viruses and because the identification of polyphenols (and other natural products) has been made simpler over the past few years thanks to developments in plant metabolism and ever-growing natural product databases such as phenol-explorer (Rothwell et al., 2016), knapsack (Shinbo et al., 2006), and the global natural product social networking database (Wang et al., 2016). (Palamara et al., 2005) (Ryu, Jeong, et al., 2010).

Interestingly, there are no known therapies for CoVs-infection and defensive vaccines are still being discovered even discovered still need extensive evaluation. Thus, the situation illustrates the need for appropriate antivirals and/or immune-enhancers to be produced for prophylaxis CoV-infection. Recently, the antiviral activity of numerous polyphenols from different sources was evaluated against several viruses (Khaerunnisa, Kurniawan, Awaluddin, Suhartati, & Soetjipto, 2020; Maryam et al., 2020). Table 3 shows the antiviral activities of polyphenols, their main sources, and mode of action (Khaerunnisa, 2020). In a related study, Sundararajan et al. (2010) evaluated three pomegranate extracts including pomegranate juice, concentrated liquid extract, and dried extract for their anti-influenza activity. He found that the anti-influenza activity of pomegranate polyphenols is substantially modified by changes in covering glycoproteins. Likewise, Vela et al. (2010) who utilized isoflavone genistein and general kinase inhibitor for

**Table 3**

The antiviral activities of polyphenols, some examples from the available studies.

Polyphenols	Main sources	Conclusion	References
Theaflavins	Black tea	Theaflavins inhibited the Sindbis Virus infection by 99% at a concentration of 14.6 mM. Act directly on HCV-viral particles and inhibit the ability to bind to the receptor surface.	Villagomez (2017) Chowdhury et al. (2018)
Epigallocatechin gallate, epicatechingallate and gallicocatechin-3-gallate	Green tea	EGCG effectively inhibits porcine reproductive and respiratory syndrome virus infection and replication in porcine alveolar macrophages. It prevents MARC-145 cells. EGCG directly interacts with the Dengue virus molecule causing virus deformation and thus preventing the virus from infecting further cells.	Ge et al. (2018) Raekiansyah, Buerano, Luz, and Morita (2018)
Different polyphenols	<i>Radix Sophorae</i> , <i>rhizome Acanthopanaxis</i> , <i>radix Sanguisorbae</i> and extract <i>Torilis fructus</i> <i>Houttuynia cordata</i> Thunb	Inhibits the effects on replication stages and/or their influence on cellular signal pathways of Mouse hepatitis virus.	Kim et al. (2010)
Quercetin and rutin		Inhibitory effects on the ATPase of mouse hepatitis and DENV type 2 virus.	Chiow et al. (2016)
(+)-Catechin	Teas and pome fruits	Inhibiting effect on transmissible gastroenteritis virus proliferation <i>in vitro</i> and associated with its anti-oxidation.	Liang et al. (2015)
Caffeic, chlorogenic, and gallic acids	<i>Sambucus Formosana Nakai</i>	Inhibition of the replication of HCoV NL63 in a cell-type independent manner.	Weng et al. (2019)
Anthocyanins	<i>Sambucus nigra</i>	Inhibition of avian infectious bronchitis virus replication	Chen et al. (2014)
Resveratrol	Skins of red fruits	Resveratrol inhibits the binding of Zika virus particles to cells and reduces circulating Zika virus particles, which underlines its potential of limiting disease severity during the viraemic phase. Resveratrol exhibited potent inhibitory effects against influenza in MDCK cells via the blockade of nuclear-cytoplasmic translocation of viral ribonucleoprotein complexes, the decrease in the expression of late viral proteins, and the inhibition of cellular protein kinase C activity and its dependent pathways.	Mohd, Zainal, Tan, and AbuBakar (2019) Palamara et al. (2005)
Resveratrol	Skins of red fruits	Resveratrol exhibited potent inhibitory effects against influenza in MDCK cells via the blockade of nuclear-cytoplasmic translocation of viral ribonucleoprotein complexes, the decrease in the expression of late viral proteins, and the inhibition of cellular protein kinase C activity and its dependent pathways.	Palamara et al. (2005)
Flavonoids	Elderberry and blueberry	Several blueberry varieties exert anti-influenza viral activities, especially viral adsorption, which positively correlated with the total polyphenol content in Madin-Darby canine kidney cells.	Sekizawa, Ikuta, Ohnishi-Kameyama, Nishiyama, and Suzutani (2019)

treating animals, and reported that general kinase inhibitor genistein may use as an *anti-arenavirus* component. It was also found that theaflavins, which are the polyphenols of black tea, had antiviral effects on vaccinia virus, influenza virus, poliovirus-1, herpes simplex virus, coxsackievirus, and human rotavirus. Morán-Santibañez et al. (2018) reported that polyphenol-rich extracts isolated from Mexican seaweed have *in vitro* significant virucidal effects on the Measles virus. Virucidal activity of these extracts is not only a prophylactic tactic before viral infection but can also be fruitful as a dealing after contagion and for avoiding virus spreading.

#### 2.4. The role of polyphenols against SARS-CoV, MERS-CoV, and SARS-CoV-2

Regarding SRARS-CoV, saikosaponins, naturally occurring triterpene glycosides derived from medicinal plants such as *Heteromorpha spp.*, *Bupleurum spp.*, and *Scrophularia scorodonia*, showed several biological activities, specifically antihepatoma, anti-hepatitis, antinephritis, anti-inflammation, immunomodulation, and antimicrobial effects. Thus, Cheng, Rodak, Wong, and Hayden (2006) recommended the use of saikosaponins as *in vitro* antiviral activity against CoVs-22E9. The extracts of *Cibotium barometz*, *Gentiana scabra*, *Dioscorea batatas*, *Cassia tora*, and *Taxillus chinensis* inhibited the SARS-CoV replication better than other 200 extracts (Wen et al., 2011). Likewise, the extracts of *Gentiana radix*, *Dioscoreae rhizoma*, *Cassiae Semen*, *Loranthi Ramus*, and *Rhizoma Cibotii* in the concentrations from 25 to 200  $\mu\text{g mL}^{-1}$  inhibited SARS-CoV. Bioflavonoids from *Torreya nucifera* inhibited SARS-CoV 3CLpro by 62% at 100  $\mu\text{g mL}^{-1}$  (Ryu, Jeong, et al., 2010), where he identified 8 diterpenoids and 4 bioflavonoids by fluorescence resonance energy transfer analysis. The activity of flavones (apigenin, luteolin, and quercetin) was also investigated with  $\text{IC}_{50}$ -values of 280.8, 20.2, and 23.8  $\mu\text{M}$ , separately, and the presence of the apigenin moiety at the C-30 position of flavones seems to be more effective. Ethyl acetate fraction of *Houttuynia cordata* Thunb. contribute to the superior antiviral efficacy *in vitro* and *in vivo* presenting the potential for the development of antiviral

agents against coronavirus and dengue infections (Chiow, Phoon, Putti, Tan, & Chow, 2016) with an  $\text{IC}_{50}$ -value of 0.98  $\text{mg mL}^{-1}$ . The  $\text{IC}_{50}$ -values for quercetin, epigallocatechin gallate, and galusatechin gallate against SARS-CoV were 73, 73, and 47  $\mu\text{M}$ , respectively, and the galooyl group should be present at 3-OH-position to be more efficient (Nguyen et al., 2012). Quercetin derivatives such as 7-O-arylmethylquercetin and quercetin-3- $\beta$ -galactoside could combat as *anti-SARS-CoV* agent (Chen et al., 2006; Park, Yoon, Kim, Lee, & Chong, 2012). Flavonoids found in *Galla chinensis* or *Veronica lin rifolia* extracts may interact with the surface of the spiky protein of the SARS-CoV, avoiding virus particles from powerful the cell (Wang & Liu, 2014). Phlorotannins isolated from brown algae display an inhibitory effect of SARS-CoV-3CLpro, regulating the virus replication. In particular, dieckol showed high inhibitory activity with an  $\text{IC}_{50}$  value of 2.7  $\mu\text{M}$  (Park et al., 2013). Tetra-O-galooil- $\beta$ -d-glucose and luteolin have been confirmed to be active against SARS-CoV (Lin et al., 2005). In particular, *Lycoris radiata* extract showed to be a good entrant against SARS-CoV as it showed the highest efficacy, mainly due to the lycorin component (Lin et al., 2005). In addition, Park et al. (2017) isolated 10 polyphenolics from *Broussonetia papyrifera* as CoVs-protease inhibitors against 3-chymotrypsin-like (3CLpro) and-papain-like-CoVs-cysteine-proteases (PLpro). It was also found that all polyphenols have more effectiveness against PLpro than 3CLpro; the results suggested polyphenols of *B. papyrifera* have *anti-CoVs* activity. Likewise, Ryu, Jeong, et al. (2010) examined the capacity for SARS-CoV inhibition of flavonoids from the *Torreya nucifera* leaves. The ethanolic extract of *T. nucifera* leaves-inhibited the SARS-CoV-3CLprotease with inhibition values of 62% at 100  $\text{g mL}^{-1}$ . Additionally, this study confirmed that amentoflavone is more effective than the respective flavones (luteolin and apigenin) and bioflavonoids derivatives on SARS-CoV-3CLpro. Thus, amentoflavone may be a good component for development as a natural therapeutic medicine against SARS-CoV infection. Jo et al. (2020) studied the antiviral activity of three flavonoids against SARS-CoV. Rhoifolin, herbacetin, and pectolinarin have been found to block SARS-CoV-3CLpro's enzyme activity. The interaction of these bioflavonoids was established using a tryptophan-based

fluorescence method, and the three flavonoids were recommended to be templates to strategy functionally developed inhibitors. Fig. 1A presented the binding pose of amentoflavone in SARS-CoV 3CLpro cited from the study of Ryu, Jeong, et al. (2010). As for MERS-CoV, flavonoids were considered as potential inhibitors of MERS-CoV (Jo et al., 2019), where each of herbacetin, isobavachalcone, quercetin 3- $\beta$ -D-glucoside, and helichristetine blocked the enzymatic activity of MERS-CoV-3CLpro. It was noted that flavonoid derivatives with hydrophobic or carbohydrate groups devoted to their core structures inhibit MERS-CoV. The computational and experimental study showed that chalcone and flavonol are desired supports to bind with the catalytic site of MERS-CoV-3CLpro. Therefore, this study suggested that flavonoids with these properties can be used as models to develop powerful MERS-CoV-3CLpro inhibitors. Regarding SARS-CoV-2, Negida and Bahbah (2020) reported that saikosaponins are considered as active substances for SARS-CoV-2 due to their immunomodulatory, anti-inflammatory, and antiviral activities. The water extract from *Isatis indigotica* root, containing a number of phenolic compounds, namely sinigrin ( $IC_{50}$ : 217  $\mu$ M) and hesperetin ( $IC_{50}$ : 8.3  $\mu$ M) showed anti-SARS-CoV-2-3CLpro. Through sequential matching, it has been proven that SARS-CoV-2 major protease has approximately a 96% similarity to SARS-CoV-2-Mpro (Xu et al., 2020). The inhibition of Mpro was ascribed

to floronas, flavonoids, and pseudo-peptides. The most hopeful inhibitors of SARS-CoV-2 were found to be oligomers of phloroglucin (1,3,5-trihydroxybenzene) derived from brown algae *Sargassum spinuligerum*. In turn, the most active inhibitors of SARS-CoV-2 are compounds from the florotannin group (8,8'-Bieckol, 6,6'-Bieckol, Dieckol) isolated from the brown algae *Ecklonia cava* (Gentile et al., 2020). Khalifa, Nawaz, et al. (2020) revealed that phacelianin, gentiodelphin, cyanodelphin, and tecophilin authentically interacted with the receptor binding site and catalytic dyad (Cys145 and His41) of SARS-CoV-2-3CLpro. Khalifa, Zhu, et al. (2020) also discovered that pedunculagin, tercatain, and castalin intensely interacted with the receptor binding site and catalytic dyad (Cys145 and His41) of SARS-CoV-2-3CLpro. These studies unleashed that anthocyanins and/or tannins with specific structure could be used as effective anti-COVID-19 natural components. Meanwhile, Fig. 1B portrayed the interaction between tannins with SARS-CoV-2-3CLpro (Khalifa, Nawaz, et al., 2020; Khalifa, Zhu, et al., 2020). We summarized the antiviral activities from the recent studies of natural polyphenols from several plants and their mode of action against CoVs (Table 4).




Fig. 1. The binding poses of amentoflavone in SARS-CoV 3CLpro. Ribbon plot of amentoflavone complexed to 3CLpro with hydrogen bonding (A). Fig B. represented the binding between tannins and 3CLpro of SARS-CoV-2, adopted from Khalifa, Nawaz, Sobhy, Althwab, and Barakat (2020) and Khalifa, Zhu, et al. (2020).

**Table 4**  
The role of polyphenols against SARS-CoV, MERS-CoV, and SARS-CoV-2.

Polyphenols	Representative	Form/Source	Virus	Mechanism	Reference	
Phenolic acids	Gallic acid	Tetra-O-galloyl- $\beta$ -D glucose from <i>Galla chinensis</i>	SARS-CoV	Avidly binds with surface spike protein of SARS-CoV.	Yi et al. (2004)	
	Hydrobenzoic acid	Desmethoxyreserpine	SARS-CoV-2	Inhibit replication of 3CLpro, and entry.	Kesel (2005)	
Flavonoids	Kaempferol	Kaempferol derivatives, Kaempferol	SARS-CoV	Inhibit 3a ion channel of CoVs.	Schwarz et al. (2014)	
			MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	Park et al. (2017)	
	Quercetin	Quercetin, Quercetin 3- $\beta$ -D-glucoside, isobavachalcone, and helichrysetin	MERS	Inhibit cleavage activity of MERS-3CLpro enzyme.	Jo et al. (2019)	
			MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	Park et al. (2017)	
			SARS-CoV	Inhibit the cellular entry of SARS-CoV.	Chen et al. (2008)	
	Myricetin	Myricetin	SARS-CoV	Inhibit the cellular entry of SARS-CoV.	Park et al. (2013)	
			SARS-CoV-2	PLpro and 3CLpro enzyme.	Park et al. (2013)	
	Herbacetin	Herbacetin	SARS-CoV	Inhibit nsP13 by affecting the ATPase activity.	Yu et al. (2012)	
			SARS-CoV	SARS-CoV helicase inhibitor.	Yu et al. (2012)	
	Papyriflavonol A	Kazinol A, B, F and J, and brousoflavan A	<i>Broussonetia papyrifera</i>	MERS-CoV, SARS-CoV	Inhibit cleavage activity of MERS-3CLpro enzyme.	Jo et al. (2019)
				SARS-CoV	block the enzymatic activity of SARS-CoV 3CLpro.	Jo et al. (2020)
	Flavones	Amentoflavone,	<i>Torreya nucifera</i> leaves	SARS-CoV 3CL pro inhibitor	Inhibit PLpro. Inhibit SARS-3CLpro activity.	Park et al. (2017)
		Kaempferol, quercetin, luteolin-7-glucoside, demethoxycurcumin, naringenin, apigenin-7-glucoside, oleuropein, curcumin, catechin, epicatechingallate, zingerol, gingerol, and allicin	Traditional herbs	Inhibitors of SARS-CoV-2-Mpro	block the enzymatic activity of SARS-CoV 3CLpro.	Khaerunnisa et al. (2020)
Baicalin		<i>Scutellaria baicalensis</i>	SARS-CoV	Inhibit Angiotensin-converting enzyme.	Chen and Nakamura (2004)	
Scutellarein		<i>Scutellaria lateriflora</i>	SARS-CoV	Inhibit nsP13 by affecting the ATPase activity.	Yu et al. (2012)	
Rhoifolin		<i>Rhus succedanea</i> .	SARS-CoV	Inhibit SARS-3CLpro activity.	(Gong et al., 2008; Nguyen et al., 2012)	
Luteolin		luteolin, from <i>Veronica linaria</i>	SARS-CoV	Avidly binds with surface spike protein of SARS-CoV.	Yi et al. (2004)	
Daidzein		Plant-derived phenolic compounds and Root extract of <i>Isatis indigotica</i>	SARS-CoV	Not active.	Lin et al. (2005)	
30-(3-methylbut-2-enyl)-30,4,7-trihydroxyflavone		<i>Broussonetia papyrifera</i>	MERS-CoV, SARS-CoV	Inhibition of cysteine proteases CoV	Park et al. (2017)	
neobavaisoflavone		<i>Psoralea corylifolia</i>	SARS-CoV	Inhibitory activity toward SARS-CoV PLpro.	Kim et al. (2014)	
Flavanones	Herbacetin, Rhoifolin pectolarin Tetra-O-galoyl- $\beta$ -d-glucose (TGG) luteoline	Plant-derived phenolic compounds and Root extract of <i>Isatis indigotica</i> ,	SARS-CoV	Inhibit the cleavage activity of the SARS-3CLpro enzyme.	Jo et al. (2020); Lin et al. (2005)	
			Pelargonidin	<i>Pimpinella anisum</i>	SARS-CoV-2	Binding affinities to 3C-like protease of SARS-CoV-2
Bavachinin	<i>Psoralea corylifolia</i>	SARS-CoV	Inhibitory activity toward SARS-CoV PLpro.	Kim et al. (2014)		
Anthocyanidins	10 polyacylated and monomeric anthocyanins	Bure components	SARS-CoV-2	Constructively network with catalytic dyad residues of 3CLpro of SARS-CoV-2.	Khalifa, Nawaz, et al. (2020)	
Flavanols	Epigallocatechin gallate	Green tea	SARS-CoV	Inhibit SARS-3CLpro activity.	(Gong et al., 2008; Nguyen et al., 2012)	

(continued on next page)

Table 4 (continued)

Polyphenols	Representative	Form/Source	Virus	Mechanism	Reference
	Gallocatechin gallate and epicatechingallate	Green tea	SARS-CoV	Inhibit SARS-3CLpro activity.	(Gong et al., 2008; Nguyen et al., 2012)
	gallocatechin-3-gallate	Green tea	SARS-CoV	Inhibit SARS-3CLpro activity.	Ghosh, Chakraborty, Biswas, and Chowdhuri (2020)
Chalcone	Isoliquiritigenin	<i>Glycyrrhiza glabra</i>	MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	Park et al. (2017)
	Broussonchalcone B, broussonchalcone A, and 4-hydroxyisolonchocarpin	<i>Broussonetia papyrifera</i>	MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	Park et al. (2017)
	isobavachalcone	<i>Psoralea corylifolia</i>	SARS-CoV	inhibitory activity toward SARS-CoV PLpro.	Kim et al. (2014)
	4'-O-methylbavachalcone	<i>Broussonetia papyrifera</i>	MERS-CoV, SARS-CoV	Inhibit PLpro. Inhibit SARS-3CLpro activity.	Park et al. (2017)
Tannins	19 hydrolysable tannins	Bure components	SARS-CoV-2	Efficacious and selective anti-COVID-19 therapeutic compounds.	Khalifa, Zhu, et al. (2020)

### 3. The underlying mechanism of the antiviral activity of polyphenols


After the outbreak of SARS-CoV-2, there is an abundance of literature on plant-based materials such as traditional Chinese herbs as a possible cure for symptoms related to SARS like fever control and faster chest clearing. Nonetheless, further clinical trials are needed in treating acute respiratory tract infections (Wu, Yang, Zeng, & Poole, 2008). It remains important to recognize new therapeutic and functional agents of infectious diseases using natural sources (Cragg & Newman, 2013). During the last decade, scientists have made significant efforts to classify several polyphenols with anti-SARS-CoV activity. However, because of SARS-CoV and SARS-CoV-2 homology, these earlier findings regard naturally occurring substances could capable of inhibiting SARS-CoV-2. A 3CLpro is essential for virus replication and is thus a potential drug goal for the production of therapeutic materials for SARS-CoV as well as SARS-CoV-2 (Yang, Islam, Wang, Li, & Chen, 2020). It could be explained that bioactive compounds extracted from herbs were effectively inhibited the enzymatic activity of SARS-3CLpro in a dose-dependent matter. Rhubarb extracts (an IC<sub>50</sub> of 13.76 ± 0.03 µg mL<sup>-1</sup>) water extract (Luo et al., 2009), *Houttuynia cordata* water extract (Lau et al., 2008), litchi seeds-flavonoid extracted (Gong et al., 2008), and extracted from *Isatis indigotica* root (Lin et al., 2005) were efficiently effective against CoVs. Further, subsequent herb-derived occurring complexes-includes herbacetin, -rhoifolin, pectolarin (Jo et al., 2020), sinigrin (IC<sub>50</sub>: 217 µM), indigo (IC<sub>50</sub>: 752 µM), -aloe-emodin (IC<sub>50</sub>: 366 µM), hesperetin- (IC<sub>50</sub>:8.3-µM) (Lin et al., 2005), quercetin- (IC<sub>50</sub>: 73 µM), epigallocatechin-gallate- (IC<sub>50</sub>: 73 µM), and gallocatechin-gallate (IC<sub>50</sub>: 47 µM) (Nguyen et al., 2012), inhibited SARS 3CLpro effects. In addition, flavonoids, including isobavachalcone, herbacetin, helichrysetin, and quercetin 3-β-D-glucoside, could inhibit the enzymatic action of MERS-CoV 3CLpro (Jo et al., 2019). Helicase protein is often thought to be a possible focus for the production of anti-HCoV agents. Yu et al. stated that myricetin and scutellarein had a strong inhibition of nsP13 (SARS-CoV-helicase-protein) *in vitro* by influencing the function of ATPase (Yu et al., 2012). Wu et al. conducted a massive-scale review of emerging medicines, natural products, and synthetic substances (>10000 substances) to classify successful anti-SARS-CoV agents via a cell-based technique of SARS-CoV and Vero-E6-cells. Ginsenoside Rb1 extracted from *Panax-ginseng*, aescin extracted from horse chestnut, reserpine in the eucalyptus, Rauwolfia genus extracts, and *Lonicera japonica* blocked SARS-CoV replication at non-toxic concentrations (Wu et al., 2004). Same as SARS-CoV and HCoV-NL63, SARS-CoV-2-uses the ACE2-host receptor for cellular entry (Letko, Marzi, & Munster, 2020). Thus, traditional Chinese medicine with ACE<sub>2</sub> goal capability keeps a pledge to avoid SARS-CoV-2 infection. In addition, disruption of the 3a

receptor by kaempferol-juglanin derivatives may avoid viral release from contaminated cells (Schwarz et al., 2014). Significantly, in compromised patients with SARS-CoV, MERS-CoV, and SARS-CoV-2 infections, the overpowering inflammatory responses have remarkably occurred. As a consequence, anti-inflammatory drugs are expected to reduce the incidence and mortality risk (Lu, 2020). In addition, a variety of anti-CoV agents have been reported from natural herbs, but the mechanisms of action have not yet been explained. Recently, the Chinese Health Commission has officially confirmed that natural medicinal herbs could be used in combination with traditional medicine for the treatment of COVID-19 patients (Yang et al., 2020). Currently, experimental research on the efficacy of natural and medicinal plants focuses on the therapeutic potential of polyphenols against SARS-CoV-2 (Yang et al., 2020). As the homogeneity of SARS-CoV is closely similar to SARS-CoV-2, many experiments demonstrating the anti-SARS-CoV-2 behavior of polyphenols or are planned to be released in the immediate future. Medicinal plants provide helpful and preventive treatments but are still desperately needed for the care of patients with SARS-CoV-2 infection, and experience with polyphenols is certainly worth researching. More analysis of polyphenols may contribute to the discovery of innovative anti-CoV compounds that could probably prove effective in the treatment of SARS-CoV-2 or other evolving lethal viral diseases as traditional therapeutic agents. Experimental searching with medicinal plant fields still has some risks and needs massive and further experiments. Polyphenols therapies traditionally used to treat the viral respiratory infection can include specific anti-SARS-CoV-2 substances. This research described many medicinal plants labeled as antiviral/pneumonia-effective which could specifically inhibit SARS-CoV-2 (Zhang et al., 2020). Letko et al. (2020) confirmed how dissimilar lineage B-viruses can recombine to increase passage into human cells and have agreed that human ACE<sub>2</sub> is the receptor for the recent SARS-CoV-2 development. However, polyphenols including flavonoids as antiviral agents against MERS, SARS-CoV, and SARS-CoV-2 are not hardly investigated and more practical experiments are required. Fig. 2 illustrated the role of polyphenols against MERS, SARS-CoV, and SARS-CoV-2.

### 4. Future needs

This review delivers a comprehensive range of evidence on the utilization of polyphenols extracted from medicinal plants for the reserve of CoVs. The cases obviously demonstrate that the designs with specific components of polyphenols and/or herb extracts that have been established for centuries in traditional medicine can be useful as the basis of anti-SARS-CoV components in the treating and/or deterrence of COVID-19. Natural polyphenols can act as anti-enzymes, antioxidants, and





**Block the enzymatic activity of PL and 3CLpro CoV**

**Fig. 2.** Underlying of the most important structural proteins of CoV are spike protein (S), membrane protein (M), envelop protein (E), and the nucleocapsid protein (N), the RNA genome of CoV is packed in the nucleocapsid protein and further covered with envelope. The hydroxyl group of flavonoids at 7-position looks important to bind at the binding site against 3-chymotrypsin-like protease (3CLpro), and papain-like protease (PLpro). Flavonoids as a vast reservoir of therapeutically active constituents as antiviral candidates against RNA viruses, adopted from Rasouli et al. (2017), Solnier et al. (2020), and Pillaiyar, Meenakshisundaram, and Manickam (2020).

through binding-blocking virus exterior protein receptors. Different plants can be utilized as a source of polyphenols with a wide range of biological activity. These components offer hope that polyphenol extracts accessible in this review may have antiviral activity, also against CoVs-family. The wide range of natural polyphenols presented in

different sources, which are undervalued in conformist medicine, may establish a nearly infinite source of remedies. The latest tendencies in medicine and biotechnology have allowed the usage of polyphenols to a greater level, than in the earlier decades, mainly as nutritional complements and nutraceuticals. Their toxicity altogether with ingesting

time is yet to be explored by aims of multidisciplinary tactics (proteomics, omics science-genomics, and metabolomics). New formulation technologies and processing might aid to improve the antiviral polyphenols solubility, delivery tactics, and therapeutic effects to adjust them as antiviral functional foods and remedies. The research achieved so far have added to the information of the chemical structure of antiviral polyphenols. Bioinformatic studies of the polyphenols-pharmacophores structures can chief to scheming new antiviral polyphenols drugs. The usage of plant extracts might deliver synergistic health useful effects i.e., anti-inflammation, antioxidant, antimutagenic, and antiviral among occurring polyphenol blends, which should be considered and must be intensely discovered. (Khalifa, Zhu, et al., 2020), (Khalifa, Zhu, et al., 2020), (Khalifa, Zhu, et al., 2020), (Pekmezci, 2011) (Ryu, Jeong, et al., 2010).

## 5. Conclusion

Numerous instances display that nature proposes the greatest solutions. Many plant species hold biologically active polyphenols, which in synergistic blends are active in battling many diseases. They already have been discovered by ancient medicine and/or folk. Recognizing individual components and revealing the underlying mechanisms is a challenge. This delays the product's standardization, namely since plant extracts hold a wide variety of different components, happening in fluctuating concentrations, with seemingly synergistic effects. Despite several unknowns, studies decidedly show that polyphenols inhibit the life cycle of CoVs. This review indicates the huge possibility of polyphenols in the expansion of functional foods and/or pharmaceutical preparations. Laboratory model trials are required to verify their efficiency in inhibiting CoVs-replication. The upcoming phase should be to conduct clinical trials on COVID-19-patients and remarking the possibility to decrease virus multiplication in the patient's organism and thus recover clinical signs. It appears that the addition of polyphenols into food and clinical practice should be fairly fast because those are mixtures of natural origin, attained from plants approved for use as herbs and food. It is worth stating that the advantage of using polyphenols is related to the high-safety profile without causing major side effects.

## CRedit authorship contribution statement

**Taha Mehany:** Conceptualization, Methodology, and, Writing - original draft. **Ibrahim Khalifa:** Conceptualization, Methodology, and, Writing - original draft. **Hassan Barakat:** Conceptualization, Methodology, and, Writing - original draft, Visualization, Writing - review & editing. **Sami A. Althwab:** Visualization, Writing - review & editing. **Yousef M. Alharbi:** Language editing, and proofing. **Sobhy El-Sohaimy:** Visualization, Writing - review & editing.

## Declaration of competing interest

No conflict of interest among authors.

## Acknowledgments

This work was self-funded.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.fbio.2021.100891>.

## References

Alpert, P. T. (2017). The role of vitamins and minerals on the immune system. *Home Health Care Management & Practice*, 29(3), 199–202.  
Banerjee, A., Kulcsar, K., Misra, V., Frieman, M., & Mossman, K. (2019). Bats and coronaviruses. *Viruses*, 11(1), 41.

Calder, P. C. (2013). Feeding the immune system. *Proceedings of the Nutrition Society*, 72(3), 299–309.  
Calder, P. C., & Kew, S. (2007). The immune system: A target for functional foods? *British Journal of Nutrition*, 88(S2), S165–S176.  
Castelo-Branco, C., & Soveral, I. (2014). The immune system and aging: A review. *Gynecological Endocrinology*, 30(1), 16–22.  
Cheng, Y. T., Rodak, D., Wong, C., & Hayden, C. (2006). Effects of micro-and nano-structures on the self-cleaning behaviour of lotus leaves. *Nanotechnology*, 17(5), 1359.  
Chen, L., Li, J., Luo, C., Liu, H., Xu, W., Chen, G., et al. (2006). Binding interaction of quercetin-3-β-galactoside and its synthetic derivatives with SARS-CoV 3CLpro: Structure-activity relationship studies reveal salient pharmacophore features. *Bioorganic & Medicinal Chemistry*, 14(24), 8295–8306.  
Chen, C. J., Michaelis, M., Hsu, H. K., Tsai, C. C., Yang, K. D., Wu, Y. C., et al. (2008). Toona sinensis roem tender leaf extract inhibits sars coronavirus replication. *Journal of Ethnopharmacology*, 120(1), 108–111.  
Chen, Z., & Nakamura, T. (2004). Statistical evidence for the usefulness of Chinese medicine in the treatment of sars. *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives*, 18(7), 592–594.  
Chen, C., Zuckerman, D. M., Brantley, S., Sharpe, M., Childress, K., Hoiczky, E., et al. (2014). *Sambucus nigra* extracts inhibit infectious bronchitis virus at an early point during replication. *BMC Veterinary Research*, 10(1), 1–12.  
Chio, K., Phoon, M., Putti, T., Tan, B. K., & Chow, V. T. (2016). Evaluation of antiviral activities of houttuynia cordata thub. Extract, quercetin, quercetin and cinanserin on murine coronavirus and dengue virus infection. *Asian Pacific journal of tropical medicine*, 9(1), 1–7.  
Chojnacka, K., Witek-Krowiak, A., Skrzypczak, D., Mikula, K., & Mlynarz, P. (2020). Phytochemicals containing biologically active polyphenols as an effective agent against covid-19-inducing coronavirus. *Journal of Functional Foods*, 73, 104146.  
Chowdhury, P., Sahuc, M. E., Rouillé, Y., Rivière, C., Bonneau, N., Vandeputte, A., ... Séron, K. (2018). Theaflavins, polyphenols of black tea, inhibit entry of hepatitis C virus in cell culture. *PLoS One*, 13(11), Article e0198226.  
Cohen, J., & Normile, D. (2020). New sars-like virus in China triggers alarm. In *Vol. 367. American association for the advancement of science. Science* (pp. 234–235), 6475.  
Comalada, M., Ballester, I., Bailón, E., Sierra, S., Xaus, J., Gálvez, J., et al. (2006). Inhibition of pro-inflammatory markers in primary bone marrow-derived mouse macrophages by naturally occurring flavonoids: Analysis of the structure-activity relationship. *Biochemical Pharmacology*, 72(8), 1010–1021.  
Cragg, G. M., & Newman, D. J. (2013). Natural products: A continuing source of novel drug leads. *Biochimica et Biophysica Acta (BBA) - General Subjects*, 1830(6), 3670–3695.  
Del Bo, C., Bernardi, S., Marino, M., Porrini, M., Tucci, M., Guglielmetti, S., et al. (2019). Systematic review on polyphenol intake and health outcomes: Is there sufficient evidence to define a health-promoting polyphenol-rich dietary pattern? *Nutrients*, 11(6), 1355.  
Deng, J., Tang, X., Wei, X., Yu, C., Zhang, X., Shi, L., et al. (2020). Analysis method for the operation health of all-fiber optical current transformer. In *2019 international conference on optical instruments and technology: Optical sensors and applications* (Vol. 11436, p. 114360T) (International Society for Optics and Photonics).  
Forni, C., Facchiano, F., Bartoli, M., Pieretti, S., Facchiano, A., D'Arcangelo, D., et al. (2019). Beneficial role of phytochemicals on oxidative stress and age-related diseases. *BioMed Research International*, 8748253, 2019.  
Fraga, C. G., Croft, K. D., Kennedy, D. O., & Tomás-Barberán, F. A. (2019). The effects of polyphenols and other bioactives on human health. *Food & Function*, 10(2), 514–528.  
Gentile, D., Patamia, V., Scala, A., Sciortino, M. T., Piperno, A., & Rescifina, A. (2020). Putative inhibitors of SARS-CoV-2 main protease from a library of marine natural products: A virtual screening and molecular modeling study. *Marine Drugs*, 18(4), 225.  
Ge, M., Xiao, Y., Chen, H., Luo, F., Du, G., & Zeng, F. (2018). Multiple antiviral approaches of (–)-epigallocatechin-3-gallate (EGCG) against porcine reproductive and respiratory syndrome virus infection in vitro. *Antiviral Research*, 158, 52–62.  
Ghosh, R., Chakraborty, A., Biswas, A., & Chowdhuri, S. (2020). Evaluation of green tea polyphenols as novel corona virus (SARS CoV-2) main protease (Mpro) inhibitors—an in silico docking and molecular dynamics simulation study. *Journal of Biomolecular Structure Dynamics*, 1–13.  
Gong, S., Su, X., Wu, H., Li, J., Qin, Y., Xu, Q., et al. (2008). A study on anti-SARS-CoV 3CL protein of flavonoids from litchi chinensis sonn core. *Chinese Pharmacological Bulletin*, 24(5), 699–700.  
Guiné, R., Lima, M., & Barroca, M. (2009). Role and health benefits of different functional food components. *Millennium*, 1–17.  
Hasan, A., Nahar, N., Jannat, K., Bondhon, T. A., Jahan, R., & Rahmatullah, M. (2020). In silico studies on phytochemicals of *pimpinella anisum* L.(apiaceae) as potential inhibitors of SARS-CoV-2 3C-like protease. *International Journal of Ayurveda*, 5(11), 14, 08.  
Horne, J. R., & Vohl, M.-C. (2020). Biological plausibility for interactions between dietary fat, resveratrol, ACE2, and SARS-CoV illness severity. *American Journal of Physiology-Endocrinology and Metabolism*, 318(5), E830–E833.  
Iwatani, S., & Yamamoto, N. (2019). Functional food products in Japan: A review. *Food Science and Human Wellness*, 8(2), 96–101.  
Jo, S., Kim, H., Kim, S., Shin, D. H., & Kim, M. S. (2019). Characteristics of flavonoids as potent MERS-CoV 3C-like protease inhibitors. *Chemical Biology & Drug Design*, 94(6), 2023–2030.  
Jo, S., Kim, S., Shin, D. H., & Kim, M. S. (2020). Inhibition of SARS-CoV 3CL protease by flavonoids. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 35(1), 145–151.

- Kesel, A. J. (2005). Synthesis of novel test compounds for antiviral chemotherapy of severe acute respiratory syndrome (SARS). *Current Medicinal Chemistry*, 12(18), 2095–2162.
- Khaerunnisa, S., Kurniawan, H., Awaluddin, R., Suhartati, S., & Soetjipto, S. (2020). Potential inhibitor of COVID-19 main protease (Mpro) from several medicinal plant compounds by molecular docking study. *Preprints*. doi10., 1–14, 20944/preprints202003.0226.v1.
- Khalifa, I., Nawaz, A., Sobhy, R., Althwab, S. A., & Barakat, H. (2020). Polyacylated anthocyanins constructively network with catalytic dyad residues of 3CLpro of 2019-NCoV than monomeric anthocyanins: A structural-relationship activity study with 10 anthocyanins using *in-silico* approaches. *Journal of Molecular Graphics and Modelling*, 100, 107690.
- Khalifa, I., Xia, D., Dutta, K., Peng, J., Jia, Y., & Li, C. (2020). Mulberry anthocyanins exert anti-AGEs effects by selectively trapping glyoxal and structural-dependently blocking the lysyl residues of  $\beta$ -lactoglobulins. *Bioorganic Chemistry*, 96, 103615.
- Khalifa, I., Zhu, W., Mohammed, H. H. H., Dutta, K., & Li, C. (2020). Tannins inhibit SARS-CoV-2 through binding with catalytic dyad residues of 3CLpro: An *in silico* approach with 19 structural different hydrolysable tannins. *Journal of Food Biochemistry*, 44(10), Article e13432.
- Khan, H., Sureda, A., Belwal, T., Çetinkaya, S., Sıntar, İ., Tejada, S., et al. (2019). Polyphenols in the treatment of autoimmune diseases. *Autoimmunity Reviews*, 18(7), 647–657.
- Kim, H. Y., Eo, E. Y., Park, H., Kim, Y. C., Park, S., Shin, H. J., et al. (2010). Medicinal herbal extracts of Sophora radix, Acanthopanax cortex, Sanguisorba radix and Torilis fructus inhibit coronavirus replication *in vitro*. *Antiviral Therapy*, 15(5), 697–709.
- Kim, D. W., Seo, K. H., Curtis-Long, M. J., Oh, K. Y., Oh, J. W., Cho, J. K., et al. (2014). Phenolic phytochemical displaying SARS-CoV papain-like protease inhibition from the seeds of psoralea corylifolia. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 29(1), 59–63.
- Lau, K. M., Lee, K. M., Koon, C. M., Cheung, C. S. F., Lau, C. P., Ho, H. M., et al. (2008). Immunomodulatory and anti-sars activities of houttuynia cordata. *Journal of Ethnopharmacology*, 118(1), 79–85.
- Letko, M., Marzi, A., & Munster, V. (2020). Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage b betacoronaviruses. *Nature Microbiology*, 5(4), 562–569.
- Liang, W., He, L., Ning, P., Lin, J., Li, H., Lin, Z., et al. (2015). (+)-Catechin inhibition of transmissible gastroenteritis coronavirus in swine testicular cells is involved its antioxidant. *Research in Veterinary Science*, 103, 28–33.
- Lin, C. W., Tsai, F. J., Tsai, C. H., Lai, C. C., Wan, L., Ho, T. Y., et al. (2005). Anti-sars coronavirus 3c-like protease effects of isatis indigotica root and plant-derived phenolic compounds. *Antiviral Research*, 68(1), 36–42.
- London, C. (2010). Functional foods that boost the immune system. *Functional Food Product Development*, 2, 295–321.
- Lopez-Varela, S., Gonzalez-Gross, M., & Marcos, A. (2002). Functional foods and the immune system: A review. *European Journal of Clinical Nutrition*, 56(3), S29–S33.
- Lu, H. (2020). Drug treatment options for the 2019-new coronavirus (2019-ncov). *BioScience Trends*, 14(1), 69–71.
- Luo, W., Su, X., Gong, S., Qin, Y., Liu, W., Li, J., et al. (2009). Anti-sars coronavirus 3c-like protease effects of rheum palmatum L. Extracts. *Bioscience Trends*, 3(4), 124–126.
- Maggini, S., Pierre, A., & Calder, P. C. (2018). Immune function and micronutrient requirements change over the life course. *Nutrients*, 10(10), 1531.
- Maggini, S., Wintergerst, E. S., Beveridge, S., & Hornig, D. H. (2007). Selected vitamins and trace elements support immune function by strengthening epithelial barriers and cellular and humoral immune responses. *British Journal of Nutrition*, 98(Suppl 1), S29–S35.
- Manoharan, Y., Haridas, V., Vasanthakumar, K., Muthu, S., Thavoorullah, F. F., & Shetty, P. (2020). Curcumin: A wonder drug as a preventive measure for covid19 management. *Indian Journal of Clinical Biochemistry*, 35(3), 373–375.
- Maryam, M., Te, K. K., Wong, F. C., Chai, T. T., Gary, K., & Gan, S. C. (2020). Antiviral activity of traditional Chinese medicinal plants dryopteris crassirhizoma and morus alba against dengue virus. *Journal of Integrative Agriculture*, 19(4), 1085–1096.
- Mohd, A., Zainal, N., Tan, K. K., & AbuBakar, S. (2019). Resveratrol affects Zika virus replication *in vitro*. *Scientific Reports*, 9(1), 1–11.
- Morán-Santibañez, K., Peña-Hernández, M. A., Cruz-Suárez, L. E., Ricque-Marie, D., Skouta, R., Vasquez, A. H., et al. (2018). Virucidal and synergistic activity of polyphenol-rich extracts of seaweeds against measles virus. *Viruses*, 10(9), 465.
- Negida, A., & Bahbah, E. I. (2020). Comments on the hydroxychloroquine/azithromycin study and lessons for future clinical trials of covid-19 treatments. *Preprint*. [https://www.researchgate.net/profile/Ahmed\\_Negida/publication/340102911\\_Comments\\_on\\_the\\_HydroxychloroquineAzithromycin\\_Study\\_and\\_Lessons\\_for\\_Future\\_Clinical\\_Trials\\_of\\_COVID-19\\_Treatments/links/5e78e56f4585158bd50068d3/Comments-on-the-Hydroxychloroquine-Azithromycin-Study-and-Lessons-for-Future-Clinical-Trials-of-COVID-19-Treatments.pdf](https://www.researchgate.net/profile/Ahmed_Negida/publication/340102911_Comments_on_the_HydroxychloroquineAzithromycin_Study_and_Lessons_for_Future_Clinical_Trials_of_COVID-19_Treatments/links/5e78e56f4585158bd50068d3/Comments-on-the-Hydroxychloroquine-Azithromycin-Study-and-Lessons-for-Future-Clinical-Trials-of-COVID-19-Treatments.pdf).
- Nguyen, T. T. H., Woo, H.-J., Kang, H.-K., Kim, Y.-M., Kim, D.-W., Ahn, S.-A., et al. (2012). Flavonoid-mediated inhibition of sars coronavirus 3c-like protease expressed in pichia pastoris. *Biotechnology Letters*, 34(5), 831–838.
- Palamara, A. T., Nencioni, L., Aquilano, K., De Chiara, G., Hernandez, L., Cozzolino, F., ... Garaci, E. (2005). Inhibition of Influenza A virus replication by resveratrol. *The Journal of Infectious Diseases*, 191(10), 1719–1729.
- Pandey, P., Rane, J. S., Chatterjee, A., Kumar, A., Khan, R., Prakash, A., et al. (2020). Targeting SARS-CoV-2 spike protein of covid-19 with naturally occurring phytochemicals: An *in silico* study for drug development. *Journal of Biomolecular Structure Dynamics*, 1–11.
- Paraiso, I. L., Revel, J. S., & Stevens, J. F. (2020). Potential use of polyphenols in the battle against covid-19. *Current Opinion in Food Science*, 32, 149–155.
- Park, J.-Y., Kim, J. H., Kwon, J. M., Kwon, H.-J., Jeong, H. J., Kim, Y. M., et al. (2013). Dieckol, a SARS-CoV 3CLpro inhibitor, isolated from the edible brown algae ecklonia cava. *Bioorganic & Medicinal Chemistry*, 21(13), 3730–3737.
- Park, H. R., Yoon, H., Kim, M. K., Lee, S. D., & Chong, Y. (2012). Synthesis and antiviral evaluation of 7-o-arylmethylquercetin derivatives against sars-associated coronavirus (scv) and hepatitis c virus (hcv). *Archives of Pharmacol Research*, 35(1), 77–85.
- Park, J.-Y., Yuk, H. J., Ryu, H. W., Lim, S. H., Kim, K. S., Park, K. H., et al. (2017). Evaluation of polyphenols from broussonetia papyrifera as coronavirus protease inhibitors. *Journal of Enzyme Inhibition and Medicinal Chemistry*, 32(1), 504–512.
- Pekmezci, D. (2011). Vitamin E and immunity. In *Vitamins & hormones* (Vol. 86, pp. 179–215). New York, USA: Academic Press.
- Pereira, P. C. M. (2003). Interaction between infection, nutrition and immunity in tropical medicine. *Journal of Venomous Animals and Toxins Including Tropical Diseases*, 9, 163–173.
- Pillaiyar, T., Meenakshisundaram, S., & Manickam, M. (2020). Recent discovery and development of inhibitors targeting coronaviruses. *Drug Discovery Today*, 25(4), 668–688.
- Powell, J., Borchers, A. T., Yoshida, S., & Gershwin, M. E. (2000). Evaluation of the immune system in the nutritionally at-risk host. *Nutrition and immunology*. Totowa, NJ: Humana Press.
- Quero, J., Mármol, I., Cerrada, E., & Rodríguez-Yoldi, M. J. (2020). Insight into the potential application of polyphenol-rich dietary intervention in degenerative disease management. *Food & Function*, 11(4), 2805–2825.
- Raekiansyah, M., Buerano, C. C., Luz, M. A. D., & Morita, K. (2018). Inhibitory effect of the green tea molecule EGCG against dengue virus infection. *Archives of Virology*, 163(6), 1649–1655.
- Rasouli, H., Farzaei, M. H., & Khodarahmi, R. (2017). Polyphenols and their benefits: A review. *International Journal of Food Properties*, 20(sup2), 1700–1741.
- Roberfroid, M. B. (2000). Concepts and strategy of functional food science: The European perspective. *American Journal of Clinical Nutrition*, 71(6), 1660S–1664S.
- Rothwell, J. A., Urpi-Sarda, M., Boto-Ordoñez, M., Llorach, R., Farran-Codina, A., Barupal, D. K., et al. (2016). Systematic analysis of the polyphenol metabolome using the phenol-explorer database. *Molecular Nutrition & Food Research*, 60(1), 203–211.
- Ryu, Y. B., Jeong, H. J., Kim, J. H., Kim, Y. M., Park, J.-Y., Kim, D., et al. (2010). Biflavonoids from torreyia nucifera displaying SARS-CoV 3CLpro inhibition. *Bioorganic & Medicinal Chemistry*, 18(22), 7940–7947.
- Ryu, Y. B., Park, S.-J., Kim, Y. M., Lee, J.-Y., Seo, W. D., Chang, J. S., et al. (2010). SARS-CoV 3CLpro inhibitory effects of quinine-methide triterpenes from tripterygium regelii. *Bioorganic & Medicinal Chemistry Letters*, 20(6), 1873–1876.
- Santos-Buelga, C., González-Paramás, A. M., Oludemi, T., Ayuda-Durán, B., & González-Manzano, S. (2019). Plant phenolics as functional food ingredients. *Advances in Food & Nutrition Research*, 90, 183–257.
- Schaible, U. E., & Stefan, H. (2007). Malnutrition and infection: Complex mechanisms and global impacts. *PLoS Medicine*, 4(5), e115.
- Schettig, R., Sears, T., Klein, M., Tan-Lim, R., Matthias, R., Jr., Aussemes, C., et al. (2020). Covid-19 patient with multifocal pneumonia and respiratory difficulty resolved quickly: Possible antiviral and anti-inflammatory benefits of quercinex (nebulized quercetin-nac) as adjuvant. *Advances in Infectious Diseases*, 10(3), 45–55.
- Schwarz, S., Sauter, D., Wang, K., Zhang, R., Sun, B., Karioti, A., et al. (2014). Kaempferol derivatives as antiviral drugs against the 3a channel protein of coronavirus. *Planta Medica*, 80, 177–182, 02/03.
- Sekizawa, H., Ikuta, K., Ohnishi-Kameyama, M., Nishiyama, K., & Suzutani, T. (2019). Identification of the components in a vaccinium oldhamii extract showing inhibitory activity against influenza virus adsorption. *Food*, 8(5), 172.
- Shinbo, Y., Nakamura, Y., Altaf-Ul-Amin, M., Asahi, H., Kurokawa, K., Arita, M., ... Kanaya, S. (2006). KNAPsAcK: A comprehensive species-metabolite relationship database. In *Plant metabolomics* (pp. 165–181). Berlin, Heidelberg: Springer.
- Silva, R. F. M., & Pogačnik, L. (2020). Polyphenols from food and natural products: Neuroprotection and safety. *Antioxidants*, 9(1), 61.
- Solnier, J., & Fladerer, J. P. (2020). Flavonoids: A complementary approach to conventional therapy of COVID-19? *Phytochemistry Reviews*, 1–23.
- Sundararajan, A., Ganapathy, R., Huan, L., Dunlap, J. R., Webby, R. J., Kotwal, G. J., et al. (2010). Influenza virus variation in susceptibility to inactivation by pomegranate polyphenols is determined by envelope glycoproteins. *Antiviral Research*, 88(1), 1–9.
- Vela, E. M., Knostman, K. A., Mott, J. M., Warren, R. L., Garver, J. N., Vela, L. J., et al. (2010). Genistein, a general kinase inhibitor, as a potential antiviral for arenaviral hemorrhagic fever as described in the pirital virus-syrian golden hamster model. *Antiviral Research*, 87(3), 318–328.
- Villagomez, J. (2017). *Vitro antiviral activity of black tea polyphenols on sindbis virus in vero cells*. New Jersey, USA: Montclair State University. MSc Thesis.
- Wang, M., Carver, J. J., Phelan, V. V., Sanchez, L. M., Garg, N., Peng, Y., ... Porto, C. (2016). Sharing and community curation of mass spectrometry data with global natural products social molecular networking. *Nature Biotechnology*, 34(8), 828–837.
- Wang, X., & Liu, Z. (2014). Prevention and treatment of viral respiratory infections by traditional Chinese herbs. *Chinese Medical Journal*, 127(7), 1344–1350.
- Weng, J. R., Lin, C. S., Lai, H. C., Lin, Y. P., Wang, C. Y., Tsai, Y. C., et al. (2019). Antiviral activity of sambucus formosananakai ethanol extract and related phenolic acid constituents against human coronavirus NL63. *Virus Research*, 273, 197767.
- Wen, C.-C., Shyur, L.-F., Jan, J.-T., Liang, P.-H., Kuo, C.-J., Arulselvan, P., et al. (2011). Traditional Chinese medicine herbal extracts of cibotium barometz, gentiana scabra, dioscorea batatas, cassia tora, and taxillus chinensis inhibit SARS-CoV replication. *Journal of traditional and complementary medicine*, 1(1), 41–50.

- Wichers, H. (2009). Immunomodulation by food: Promising concept for mitigating allergic disease? *Analytical and Bioanalytical Chemistry*, 395(1), 37–45.
- Wu, C.-Y., Jan, J.-T., Ma, S.-H., Kuo, C.-J., Juan, H.-F., Cheng, Y.-S. E., et al. (2004). Small molecules targeting severe acute respiratory syndrome human coronavirus. *Proceedings of the National Academy of Sciences*, 101(27), 10012–10017.
- Wu, T., Yang, X., Zeng, X., & Poole, P. (2008). Traditional Chinese medicine in the treatment of acute respiratory tract infections. *Respiratory Medicine*, 102(8), 1093–1098.
- Xu, Z., Peng, C., Shi, Y., Zhu, Z., Mu, K., Wang, X., et al. (2020). Nelfinavir was predicted to be a potential inhibitor of 2019-ncov main protease by an integrative approach combining homology modelling, molecular docking and binding free energy calculation. *BioRxiv*, 2020.01.27.921627 <https://www.biorxiv.org/content/10.1101/2020.01.27.921627v1>.
- Yahfoufi, N., Alsadi, N., Jambi, M., & Matar, C. (2018). The immunomodulatory and anti-inflammatory role of polyphenols. *Nutrients*, 10(11), 1618.
- Yang, Y., Islam, M. S., Wang, J., Li, Y., & Chen, X. (2020). Traditional Chinese medicine in the treatment of patients infected with 2019-new coronavirus (SARS-CoV-2): A review and perspective. *International Journal of Biological Sciences*, 16(10), 1708–1717.
- Yi, L., Li, Z., Yuan, K., Qu, X., Chen, J., Wang, G., et al. (2004). Small molecules blocking the entry of severe acute respiratory syndrome coronavirus into host cells. *Journal of Virology*, 78(20), 11334–11339.
- Yu, M.-S., Lee, J., Lee, J. M., Kim, Y., Chin, Y.-W., Jee, J.-G., et al. (2012). Identification of myricetin and scutellarein as novel chemical inhibitors of the sars coronavirus helicase, nsp13. *Bioorganic & Medicinal Chemistry Letters*, 22(12), 4049–4054.
- Zhang, L., & Liu, Y. (2020). Potential interventions for novel coronavirus in China: A systematic review. *Journal of Medical Virology*, 92(5), 479–490.
- Zhang, D. H., Wu, K. L., Zhang, X., Deng, S. Q., & Peng, B. (2020). *In silico* screening of Chinese herbal medicines with the potential to directly inhibit 2019 novel coronavirus. *Journal of Integrative Medicine*, 18(2), 152–158.