



Corrigendum: Unmet Psychosocial Needs of Patients Newly Diagnosed with Ulcerative Colitis: Results from the Nationwide Prospective Cohort Study in Korea

Jung Rock Moon¹, Chang Kyun Lee¹, Sung Noh Hong², Jong Pil Im³, Byong Duk Ye⁴, Jae Myung Cha⁵, Sung-Ae Jung⁶, Kang-Moon Lee⁷, Dong Il Park⁸, Yoon Tae Jeen⁹, Young Sook Park¹⁰, Jae Hee Cheon¹¹, Hyesung Kim¹², BoJeong Seo¹², Youngdoe Kim¹², and Hyo Jong Kim¹, the MOSAIK study group of the Korean Association for the Study of Intestinal Diseases (KASID)

¹Center for Crohn's and Colitis, Department of Gastroenterology, Kyung Hee University College of Medicine, ²Department of Medicine, Sungkyunkwan University School of Medicine, ³Department of Internal Medicine, Seoul National University College of Medicine, ⁴Department of Gastroenterology, University of Ulsan College of Medicine, ⁵Department of Internal Medicine, Kyung Hee University Hospital at Gangdong, ⁶Department of Internal Medicine, Ewha Womans University College of Medicine, ⁷Department of Internal Medicine, College of Medicine, The Catholic University of Korea, ⁸Department of Internal Medicine, Kangbuk Samsung Hospital, Sungkyunkwan University School of Medicine, ⁹Department of Gastroenterology and Hepatology, Korea University College of Medicine, ¹⁰Department of Internal Medicine, Eulji University School of Medicine, ¹¹Department of Internal Medicine, Yonsei University College of Medicine, and ¹²Medical Affairs, Janssen Korea, Seoul, Korea

<https://doi.org/10.5009/gnl19107>

Gut Liver 2020;14(4):459-467

In the version of this article initially published, Table 2 contains an error in the data of p-value. On page 462, the table should be corrected as follows (marked in bold):

From:

Table 2. Mean Values of Each Patient-Reported Outcome Measure

PROs	Category	No.	Overall (n=355)	Disease severity		p-value
				Moderate (n=330)*	Severe (n=25) [†]	
HADS	Depression	349	6.6±4.5	6.5±4.4	8.9±5.2	0.610
	Anxiety	348	6.2±4.4	6.2±4.4	6.6±4.5	0.638
WPAI:UC	Absenteeism [‡]	149	29.1±36.9	27.3±35.8	54.1±43.5	0.102
	Presenteeism [§]	143	37.8±31.4	36.8±30.3	62.2±39.6	0.037
	Overall work impairment	130	45.5±32.7	44.1±31.9	70.7±37.8	0.048
	Social activity impairment [¶]	344	46.4±31.2	45.2±31.0	61.2±31.0	0.014
IBDQ	Bowel system	344	41.7±13.0	42.0±13.1	36.1±10.9	0.027
	Emotional health	347	51.7±14.9	52.0±14.9	46.8±13.6	0.097
	Systemic symptoms	347	20.5±6.8	20.7±6.8	17.0±6.0	0.009
	Social function	309	23.2±7.7	22.8±7.9	18.0±6.8	0.003
	Total score	305	137.3±38.8	137.5±39.0	117.9±33.2	0.002

Data are presented as the mean±SD.

PRO, patient-reported outcome; HADS, Hospital Anxiety and Depression Scale; WPAI, Work Productivity and Activity Impairment; UC, ulcerative colitis; IBDQ, Inflammatory Bowel Disease Questionnaire.

*Total Mayo score 6–10 points; [†]Total Mayo score ≥11; [‡]Absenteeism: percentage of work time missed=(Q2/[Q2+Q4])×100; [§]Presenteeism: percentage of impairment while working=(Q5/10)×100; ^{||}Percentage of overall work impairment=(Q2/(Q2+Q4)+[1-(Q2/(Q2+Q4))]×(Q5/10))×100; [¶]Percentage of social activity impairment=(Q6/10)×100; Q, questionnaire item.

Copyright © Gut and Liver.



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

To:

Table 2. Mean Values of Each Patient-Reported Outcome Measure

PROs	Category	No.	Overall (n=355)	Disease severity		p-value
				Moderate (n=330)*	Severe (n=25) [†]	
HADS	Depression	349	6.6±4.5	6.5±4.4	8.9±5.2	0.024
	Anxiety	348	6.2±4.4	6.2±4.4	6.6±4.5	0.638
WPAI:UC	Absenteeism [‡]	149	29.1±36.9	27.3±35.8	54.1±43.5	0.102
	Presenteeism [§]	143	37.8±31.4	36.8±30.3	62.2±39.6	0.037
	Overall work impairment	130	45.5±32.7	44.1±31.9	70.7±37.8	0.048
	Social activity impairment [¶]	344	46.4±31.2	45.2±31.0	61.2±31.0	0.014
IBDQ	Bowel system	344	41.7±13.0	42.0±13.1	36.1±10.9	0.027
	Emotional health	347	51.7±14.9	52.0±14.9	46.8±13.6	0.097
	Systemic symptoms	347	20.5±6.8	20.7±6.8	17.0±6.0	0.009
	Social function	309	23.2±7.7	22.8±7.9	18.0±6.8	0.003
	Total score	305	137.3±38.8	137.5±39.0	117.9±33.2	0.002

Data are presented as the mean±SD.

PRO, patient-reported outcome; HADS, Hospital Anxiety and Depression Scale; WPAI, Work Productivity and Activity Impairment; UC, ulcerative colitis; IBDQ, Inflammatory Bowel Disease Questionnaire.

*Total Mayo score 6–10 points; [†]Total Mayo score ≥11; [‡]Absenteeism: percentage of work time missed=(Q2/[Q2+Q4])×100; [§]Presenteeism: percentage of impairment while working=(Q5/10)×100; ^{||}Percentage of overall work impairment=(Q2/[Q2+Q4])+[1-(Q2/[Q2+Q4])×(Q5/10)]×100; [¶]Percentage of social activity impairment=(Q6/10)×100; Q, questionnaire item.