



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to the Editor

Cardiovascular complications in COVID-19: A systematic review and meta-analysis

To the Editor

The coronavirus disease 2019 (COVID-19) pandemic poses the most significant modern-day public health challenge since the Spanish flu of 1918, causing substantial morbidity and mortality worldwide.¹ Coronavirus disease 2019 predominantly affects the respiratory system, causing severe pneumonia and respiratory distress syndrome. There is also involvement of multiple organs, and the cardiovascular system has been implicated. In a recent study to investigate characteristics and prognostic factors in 339 elderly patients with COVID-19, Wang and colleagues observed a high proportion of severe and critical cases as well as high fatality rates.² The cardiovascular complications recorded were acute cardiac injury (21%), arrhythmia (10.4%) and cardiac insufficiency (17.4%). However, not all cardiovascular manifestations of COVID-19 are clearly defined by published studies and it is also not clear if these conditions are directly caused by COVID-19 or are just unspecific complications.³ There is a need to understand the interplay between COVID-19 and its cardiovascular manifestations to assist in the optimum management of patients. In this context, we conducted a systematic meta-analysis to attempt to address the following questions: (i) what are the cardiovascular complications associated with COVID-19?; (ii) what is the incidence of these complications?; and (iii) are patients with pre-existing cardiovascular morbidities more susceptible to these cardiovascular complications?

The protocol for this review was registered in the PROSPERO International prospective register of systematic reviews (CRD42020184851). The review was conducted in accordance with PRISMA and MOOSE guidelines^{4, 5} (Supplementary Materials 1-2). We searched MEDLINE, Embase, and The Cochrane library from 2019 to 27 May 2020 for published studies reporting on cardiovascular outcomes in patients with COVID-19. Details of the search strategy are reported in Supplementary Material 3. The prevalence of comorbidities (pre-existing hypertension and cardiovascular disease, CVD) and incidence of cardiovascular complications across studies with their 95% confidence intervals (CIs) were pooled using Freeman-Tukey variance stabilising double arcsine transformation and random-effects models. STATA release MP 16 (StataCorp LP, College Station, TX, USA) was used for all statistical analyses.

Seventeen retrospective cohort studies comprising of 5,815 patients with COVID-19 were included (Table 1; Supplementary Materials 4-5). Eleven studies were based in China, four in the USA, one in South Korea and one in the Netherlands. The average age at baseline ranged from 47 to 71 years.

Across 15 studies, the pooled prevalence of pre-existing hypertension (95% CI) in COVID-19 patients was 29.3% (25.5-33.4; I^2

=87%; 95% CI 79, 91%; p for heterogeneity <0.01) (Supplementary Material 6). The prevalence (95% CI) of pre-existing CVD across 16 studies was 14.6% (11.0-18.4; I^2 =91%; 95% CI 87, 94%; p for heterogeneity <0.01) (Supplementary Material 7).

Over hospital stays ranging from 2 to 28 days, the pooled incidence was 17.6% (14.2-21.2; I^2 =32%; 95% CI 0, 76%; p for heterogeneity=0.20) for heart failure (HF) ($n=4$ studies); 16.3% (11.8-21.3; I^2 =87%; 95% CI 79, 92%; p for heterogeneity <0.01) for myocardial injury ($n=11$ studies); 9.3% (5.1-14.6; I^2 =78%; 95% CI 52, 90%; p for heterogeneity <0.01) for cardiac arrhythmia ($n=6$ studies); 6.2% (1.8-12.3) for acute coronary syndrome (ACS) ($n=2$ studies); 5.7% (2.7-9.6) for cardiac arrest ($n=2$ studies); and 5.6% (3.4-8.3) for disseminated intravascular dissemination (DIC) ($n=2$ studies) (Fig. 1A). Subgroup analyses suggested that the incidence of myocardial injury was higher in older age groups and groups with a higher prevalence of pre-existing hypertension; however, the incidence of myocardial injury was similar in groups with high or low prevalence of pre-existing CVD (Fig. 1B). Over hospital stays ranging from 2 to 28 days following admission, mortality rate ranged from 0.7% to 52.4%, with a pooled rate of 15.3% (10.7-20.5).

The current data based on up-to-date evidence suggests that the most common cardiovascular complications of COVID-19 are HF, myocardial injury and cardiac arrhythmias. Though the mechanisms for cardiovascular manifestations of COVID-19 are still yet to be elucidated, the following multiple pathways have been proposed: (i) direct cardiotoxicity; (ii) systemic inflammation; (iii) myocardial demand-supply mismatch; (iv) plaque rupture and coronary thrombosis; (v) adverse effects of therapies during hospitalisation; (vi) sepsis leading to DIC; (vii) increased systemic thrombogenesis; and (viii) electrolyte imbalances.^{6, 7} Myocardial injury is reported to mainly result from direct viral involvement of cardiomyocytes and the effects of systemic inflammation.⁶ Though venous thromboembolism incidence was based on a single report, patients with COVID-19 are at increased risk of hypercoagulable states due to prolonged immobilisation, systemic inflammation and risk for DIC.⁷

In addition to pre-existing comorbidities including CVD being associated with worse outcomes in patients with COVID-19,^{8, 9} cardiovascular complications such as myocardial injury have also been shown to be associated with increased risk of severe COVID-19 and fatal outcomes.¹⁰ Myocardial injury is commonly defined as substantial elevation of high-sensitivity cardiac troponin levels and it has been reported that elevated troponin levels are associated with greater risk of severe disease and mortality.¹⁰ Monitoring of markers of cardiac damage such as troponin, N-terminal pro B-type natriuretic peptide and creatine kinase during hospitalisation for COVID-19 could help in the identification of patients with possible cardiac manifestations, to enable early and more aggressive intervention.

Table 1
Characteristics of included studies

Author, year of publication	Source of data	Country	Dates of data collection	Mean/median age (years)	Male %	Hospitalization (days)	Number of patients	CVD complications	NOS
Guo, 2020	Seventh Hospital of Wuhan City	China	Jan - Feb 2020	58.5	48.7	16.3	187	Myocardial injury; ventricular arrhythmia	5
Wang, 2020	Zhongnan Hospital of Wuhan University	China	Jan, 2020	56.0	54.3	7.0	138	Myocardial injury; cardiac arrhythmia	4
Huang, 2020	Jin Yintan Hospital, Wuhan	China	Dec - Jan 2020	49.0	73.0	7.0	41	Myocardial injury	4
Zhou, 2020	Jinyintan Hospital & Wuhan Pulmonary Hospital	China	Dec - Jan 2020	56.0	62.0	11.0	191	Myocardial injury; HF	5
Shi, 2020	Renmin Hospital of Wuhan University	China	Jan - Feb 2020	64.0	49.3	NR	416	Myocardial injury	6
Arentz, 2020	Evergreen Hospital in Kirkland, Washington	USA	Feb - March 2020	70.0	52.0	5.2	21	Cardiomyopathy	4
Chen, 2020	Tongji Hospital in Wuhan	China	Jan - Feb 2020	62.0	62.0	13.0	274	Myocardial injury; HF; DIC	4
Du, 2020	Hannan Hospital and Wuhan Union Hospital	China	Jan - Feb 2020	65.8	72.9	10.1	85	Cardiac arrest; ACS; arrhythmia; DIC	4
Wang, 2020b	Renmin Hospital of Wuhan University	China	Jan - Feb 2020	71.0	49.0	28.0	339	Myocardial injury, arrhythmia HF	4
Cao, 2020	Zhongnan Hospital of Wuhan University	China	Jan - Feb 2020	54.0	52.0	11.0	102	Myocardial injury; arrhythmia; cardiac arrest	4
Klok, 2020	Dutch Univesity Hospitals	Netherlands	March - April 2020	64.0	76.0	7.0	184	PE; VTE; stroke	4
Aggarwal, 2020	UnityPoint Clinic	USA	March - April 2020	67.0	75.0	2.0	16	ACS; cardiac arrhythmia; HF	4
Wang, 2020c	Zhongnan Hospital of Wuhan University and Xishui People's Hospital	China	Up to Feb, 2020	51.0	53.3	11.0	107	Myocardial injury	5
Hong, 2020	Yeungnam University Medical Center	South Korea	Up to March, 2020	55.4	38.8	7.7	98	Myocardial injury	4
Wan, 2020	Northeast Chongqing	China	Jan - Feb 2020	47.0	53.3	5.0	135	Myocardial injury	4
Price-Haywood, 2020	Ochsner Health in Louisiana	Asia	March - April, 2020	55.5	45.7	7.0	1,030	Cardiomyopathy/HF	6
Price-Haywood, 2020	Ochsner Health in Louisiana	Asia	March - April, 2020	53.6	37.7	6.0	2,451	Cardiomyopathy/HF	6

ACS, acute coronary syndrome; DIC, disseminated intravascular coagulation; HF, heart failure; NOS, Newcastle Ottawa Scale; NR, not reported; PE, pulmonary embolism; VTE, venous thromboembolism




Fig. 1. (A) Incidence of cardiovascular complications in COVID-19 patients; (B) Incidence of myocardial injury in COVID-19 patients, by clinically relevant characteristics CI, confidence interval (bars); CVD, cardiovascular disease; DIC, disseminated intravascular coagulation; PE, pulmonary embolism; VTE, venous thromboembolism; *, p-value for meta-regression

The inherent limitations of this review included the low sample sizes and methodological design of some of the studies, which was expected given the urgency to report and gain a better understanding of COVID-19; limited number of studies available, hence some of the findings were based on single reports; assays for cardiac injury and their time of assessment during hospitalisation may vary between studies, hence estimates may be biased; and the possibility of patient overlap given that the majority of studies were conducted from China and reports of duplicate publication of study participants in articles.⁹

Aggregate analysis of the literature suggests that the most frequent cardiovascular complications among patients hospitalised with COVID-19 are HF, myocardial injury, cardiac arrhythmias and ACS. Early identification and monitoring of cardiac complications could help in the prediction of more favourable outcomes. The causes of these cardiovascular manifestations warrant further investigation as more data becomes available.

Declaration of Competing Interest

None.

Acknowledgements

SKK acknowledges support from the NIHR Biomedical Research Centre at University Hospitals Bristol NHS Foundation Trust and the University of Bristol. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health and Social Care. These sources had no role in design and conduct of the study; collection, management, analysis, and interpretation of the data; and preparation, review, or approval of the manuscript.

Supplementary materials

Supplementary material associated with this article can be found, in the online version, at [doi:10.1016/j.jinf.2020.05.068](https://doi.org/10.1016/j.jinf.2020.05.068).

References

- Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020; **395**(10229):1054–62. doi:10.1016/S0140-6736(20)30566-3.
- Wang L, He W, Yu X, et al. Coronavirus disease 2019 in elderly patients: Characteristics and prognostic factors based on 4-week follow-up. *J Infect* 2020. doi:10.1016/j.jinf.2020.03.019.

- Ferrari R, Di Pasquale G, Rapezzi C. 2019 CORONAVIRUS: What are the implications for cardiology? *Eur J Prev Cardiol*. 2020; **27**(8):793–6. doi:10.1177/2047487320918102.
- Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of Observational Studies in Epidemiology. *JAMA: The Journal of the American Medical Association* 2000; **283**(15):2008–12. doi:10.1001/jama.283.15.2008.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *PLoS Med* 2009; **6**(7):e1000097. doi:10.1371/journal.pmed.1000097.
- Bansal M. Cardiovascular disease and COVID-19. *Diabetes & metabolic syndrome* 2020; **14**(3):247–50. doi:10.1016/j.dsx.2020.03.013.
- Kang Y, Chen T, Mui D, et al. Cardiovascular manifestations and treatment considerations in covid-19. *Heart* 2020. doi:10.1136/heartjnl-2020-317056.
- Weiss P, Murdoch DR. Clinical course and mortality risk of severe COVID-19. *Lancet* 2020; **395**(10229):1014–15. doi:10.1016/S0140-6736(20)30633-4.
- Kunutsor SK, Laukkanen JA. Markers of liver injury and clinical outcomes in COVID-19 patients: A systematic review and meta-analysis. *J Infect* 2020. doi:10.1016/j.jinf.2020.05.045.
- Guo T, Fan Y, Chen M, et al. Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiol* 2020. doi:10.1001/jamacardio.2020.1017.

Setor K. Kunutsor*

National Institute for Health Research Bristol Biomedical Research Centre, University Hospitals Bristol NHS Foundation Trust and University of Bristol, Bristol, UK
Musculoskeletal Research Unit, Translational Health Sciences, Bristol Medical School, University of Bristol, Learning & Research Building (Level 1), Southmead Hospital, Bristol, UK

Jari A. Laukkanen

Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland
Institute of Clinical Medicine, Department of Medicine, University of Eastern Finland, Kuopio, Finland
Central Finland Health Care District Hospital District, Department of Medicine, Jyväskylä, Finland District, Jyväskylä, Finland

*Corresponding author. Translational Health Sciences, Bristol Medical School, University of Bristol, Learning & Research Building (Level 1), Southmead Hospital, Bristol, BS10 5NB, UK; Phone: +44-7539589186; Fax: +44-1174147924. E-mail address: skk31@cantab.net (S.K. Kunutsor)