




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Review Article

The global path forward – Healthy Living for Pandemic Event Protection (HL – PIVOT)☆



Ross Arena<sup>a,b,\*</sup>, Carl J. Lavie<sup>b,c</sup>, On behalf of the HL-PIVOT Network

<sup>a</sup> Department of Physical Therapy, College of Applied Science, University of Illinois, Chicago, IL, USA

<sup>b</sup> Healthy Living for Pandemic Event Protection (HL – PIVOT) Network, Chicago, IL, USA

<sup>c</sup> Department of Cardiovascular Diseases, John Ochsner Heart and Vascular Institute, Ochsner Clinical School-University of Queensland School of Medicine, New Orleans, LA, USA

Contents

Appendix 1. Founding members of the HL-PIVOT Network . . . . .	100
References . . . . .	

The Merriam-Webster dictionary defines a pandemic as “an outbreak of a disease that occurs over a wide geographic area and affects an exceptionally high proportion of the population”.<sup>1</sup> The world has again become acutely aware of this term because of a disease (COVID-19) caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) that has dramatically changed our daily behaviors and habits. As of May 30th, 2020, there have been 5,775,043 confirmed cases and 361,220 confirmed deaths in 216 countries.<sup>2</sup> There will undoubtedly be lasting effects as a result of the COVID-19 pandemic, although how these lasting effects will impact daily life for the foreseeable future, as we begin to control viral spread and recover, remains uncertain at this time. One hope is that some of the lasting effects resulting from COVID-19 will be positive; we must learn from challenging times and try to make the world a better, healthier, and safer place for all inhabitants.

While certainly different from the current health crisis induced by COVID-19, we have been living with several other significant health conditions that continue to have catastrophic effects on a global scale. In fact, obesity,<sup>3</sup> physical inactivity,<sup>4</sup> diabetes<sup>5</sup> and other noncommunicable diseases (NCDs)<sup>6</sup> (e.g., cardiovascular disease, certain forms of cancer, and respiratory disease) have all been characterized as pandemics. **Table 1** lists key metrics for each of these conditions according to the World Health Organization (WHO).<sup>7–13</sup> Clearly, the data listed in **Table 1** make a strong case for each condition being characterized as a pandemic.

*Abbreviations and acronyms:* SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; NCDs, noncommunicable diseases; WHO, World Health Organization; HL, healthy living; CRF, cardiorespiratory fitness; HL-PIVOT, Healthy Living for Pandemic Event Protection.

☆ There is no conflict of interest of any of the listed authors.

\* Address reprint requests to Ross Arena, PhD, PT, FAHA, Professor, Founder, Department of Physical Therapy, College of Applied Health Sciences, University of Illinois Chicago, 1919 W. Taylor Street (MC 898), Chicago, IL 60612.

E-mail address: [raarena@uic.edu](mailto:raarena@uic.edu) (R. Arena).

These health conditions are also interrelated; for example, physical inactivity increases obesity risk which then increases risk for diabetes and other NCDs.<sup>14,15</sup> What is worse is that, while we have long recognized aggressively addressing the conditions in **Table 1** is of paramount importance, we are making little impact in reversing trends; excess body weight, for example (i.e., obesity), continues to increase on a global scale.<sup>16</sup>

We have come to the point, based on overwhelming scientific evidence spanning several decades, that the importance and value of healthy living (HL) (i.e., physical activity, good nutrition, appropriate body weight and not smoking) in preventing (primary prevention) and treating (secondary prevention) the conditions listed in **Table 1** is beyond dispute.<sup>17–22</sup> In fact, HL is the primary medicine to treat the obesity, physical inactivity/sedentarism, diabetes and NCD pandemics.<sup>17,23</sup> Moreover, and critically important to the current perspective, any movement away from unhealthy behaviors towards HL has significant health benefits. Moving from no physical activity to two 30-minute walks per week or replacing processed foods with two servings of fruits and vegetables per day, while not meeting ideal goals, significantly improves health outcomes.<sup>17–19,22</sup> For too long we have created a false and, for a large proportion of the population, unattainable *all or nothing* view of HL behaviors. If approached from the perspective that some HL behaviors are better than none and more is even better, we have a renewed opportunity to make significant progress in improving global trends in unhealthy lifestyles patterns and the resultant poor health trajectory.

As data from COVID-19 continues to evolve rapidly, one pattern is becoming clear - the presence of preexisting medical conditions, such as obesity, diabetes, hypertension and cardiovascular and respiratory diseases, significantly increase the risk of poor outcomes due to the viral infection.<sup>24–27</sup> Again, many of these preexisting medical conditions have also been described as pandemics and are listed in **Table 1**. Publications regarding the troubling interrelationships between COVID-19

**Table 1**  
Major health conditions described as pandemics.

Condition	Key Global Statistics From the World Health Organization
Overweight and obesity	<ul style="list-style-type: none"> <li>Global obesity has approximately tripled since 1975</li> <li>39% (1.9 billion) of women and men who are 18 or older were classified as overweight in 2016</li> <li>The global prevalence of obesity (BMI <math>\geq</math> 30 kg/m<sup>2</sup>) was 13% (650 million) in adults and 18% (340 million) in children and adolescents (5–19 years) in 2016</li> </ul>
Physical inactivity	<ul style="list-style-type: none"> <li>An estimated 40 million children under the age of 5 years are overweight or obese</li> <li>In 2016, 1 in 4 adults (1.4 billion people) and 3 in 4 children currently do not meet current physical activity recommendations</li> <li>Physical inactivity in some countries are as high as 70%, attributed to changes in transportation, technology, and urbanization</li> <li>The global cost of physical inactivity is estimated to be INT\$ 54 billion in 2013 with an additional INT\$ 14 billion attributed to lost productivity.</li> </ul>
Diabetes	<ul style="list-style-type: none"> <li>Global prevalence has increased from 108 million (4.7% adults 18 years or older) in 1980 to 422 million (8.5% of adults 18 years or older) in 2014</li> <li>Approximately 1.6 million deaths were directly attributed to diabetes in 2016</li> </ul>
Noncommunicable diseases	<ul style="list-style-type: none"> <li>Noncommunicable diseases kill 41 million people annually - 71% of all global deaths</li> <li>15 million people die from a noncommunicable disease each year who are between 30 and 69 years</li> <li>Cardiovascular disease accounts for the most annual deaths - 17.9 million people - 31% of all deaths worldwide</li> </ul>

and pre-existing medical conditions are already emerging. For instance, Dietz and Santos<sup>25</sup> discussed the increased hospitalization and mortality risk with H<sub>1</sub>N<sub>1</sub> influenza infection in patients who are obese or severely obese, citing the impact significant excess body mass has on pulmonary function as one of the reasons for this association. Given the impact of COVID-19 on the pulmonary system as a primary manifestation and reason for hospitalization, concerns over the increased health risks for individuals who are obese is certainly justified. Moreover, individuals who are obese Fig. 1 present with other detrimental alterations relevant to the COVID-19 pandemic including: 1) A baseline increase in systemic inflammation that, when a viral infection is present, blunts macrophage activation and pro-inflammatory cytokine production with macrophage stimulation, contributing to compromised immune function; 2) increased viral shedding time (i.e., contagious for a longer period than normal weight individuals); 3) the potential for more virulent viral strains induced by the obese microenvironment; and 4) a positive correlation between body mass index and viral spread through exhaled breath.<sup>24</sup> These concerns are exponentially magnified given the global prevalence of excess body mass and obesity (Table 1). From the perspective of multimorbidity, Muniyappa and Gubbi<sup>28</sup> discussed COVID-19 in the context of patients with diabetes, drawing attention to the hypertension-obesity-diabetes triad that is common and the severe health risks (i.e., hospital admission, acute respiratory distress syndrome, mechanical ventilation, or death) with SARS-CoV-2 infection when an individual expresses this phenotype. Moreover, while the independent contribution of increased plasma glucose on increased risk following SARS-CoV-2 infection is unclear at this point, it has been shown to be a predictor of morbidity and mortality in those infected with SARS.<sup>29</sup> In recently reported findings from Wuhan China, Chen et al<sup>27</sup> found that older patients hospitalized with SARS-CoV-2 infection had a higher prevalence of comorbidities, greater symptom severity, and a higher likelihood of multi-organ involvement and mortality compared to younger patients. It has recently been estimated that  $\approx$ 33% of the global population are affected by multimorbidity, particularly in those individuals who are 65 years or older.<sup>30</sup> In a recent CDC Monthly and Mortality Weekly Report,<sup>26</sup> the percentage of patients with SARS-CoV-2 infection and at least one pre-existing medical condition that required intensive care unit (ICU) admission (358 of 457, 78%) or hospitalization without ICU admission (732 of 1037, 71%) was significantly higher compared to those who were not hospitalized (1388 of 5143, 27%). Diabetes, as well as cardiovascular and chronic respiratory diseases, was the most reported pre-existing medical conditions. In >5000 patients admitted with SARS-CoV-2 infection to 12 New York (New York City, Long Island, and Westchester County) hospitals, Richardson et al<sup>31</sup> reported >90% had a pre-existing medical condition while 88% had two or more; the most common pre-existing conditions were hypertension ( $\approx$ 57%), obesity ( $\approx$ 42%), and diabetes ( $\approx$ 34%). These trends are disturbing in the context of the COVID-19 pandemic given the prevalence of the pre-existing health conditions listed in

Table 1, indicating a large percentage of the global population present with a baseline health status that substantially increases the risk of hospitalization and poor health outcomes with SARS-CoV-2 infection.

Conversely, there is strong evidence base that indicates high cardiorespiratory fitness (CRF) as well as regular physical activity and a healthy diet improve immune function, affording protection against viral infections.<sup>32–35</sup> The COVID-19 pandemic, which has precipitated widespread shelter in place and social distancing policies, has also caused great concern regarding physical activity and exercise patterns.<sup>36–39</sup> The world was already largely physically inactive and sedentary pre COVID-19 and trends were not improving.<sup>36</sup> A reasonable question post-COVID-19 becomes will these trends in physical inactivity and sedentarism further accelerate, leading to a greater prevalence of the obesity, diabetes and NCD pandemics in the years to come? Initial evidence is emerging that indicates the answer to the first part of this question is yes. Zhang et al<sup>40</sup> recently reported a significant reduction in physical activity and a significant increase in screen time during the COVID-19 pandemic in >2400 children and adolescents in Shanghai, China. In >3000 adults in the United States, Meyer et al<sup>41</sup> found that, in people who were previously active, physical activity was reduced by  $\approx$ 33% during the COVID-19 pandemic as well as a substantial increase in sedentary time regardless of pre-COVID-19 activity patterns. Decreased physical activity and increased sedentary time were both associated with worsening mental health. Interestingly, Irfan Ahmed<sup>42</sup> recently hypothesized that: 1) a measure of CRF, when available, may be valuable in risk stratifying individuals in the context of future viral pandemics; and 2) exercise training may serve a role in “preconditioning” individuals prior to viral infection. Both hypotheses should be investigated thoroughly and, if supported, would further bolster the importance of HL.

Like COVID-19, if we view the conditions listed in Table 1 as pandemics and we simultaneously recognize HL has profound benefits in preventing, mitigating and treating these pandemics, there becomes an opportunity to view a new global path forward for HL medicine. In this context, we would like to introduce the *HealthyLiving forPandemic Event Protection* (HL-PIVOT) network. The Merriam-Webster noun and verb definitions for *pivot* are “a person, thing, or factor having a major or central role, function, or effect” and “to turn on or as if on a pivot”, respectively.<sup>27</sup> We feel these definitions support our choosing of the PIVOT acronym for the proposed new initiative: 1) the COVID-19 pandemic is playing the major role in impacting daily life patterns and how we perceive population health and; 2) this impact warrants turning or pivoting in a different direction with the goal of meaningfully increasing HL behaviors on a global level. The overarching goal of this network is to *promote human resilience and quality of life by increasing healthy living behaviors*. Table 2 lists areas of focus in knowledge discovery, education, policy, and implementation. We acknowledge that such a broad focus will require the network to consist of members from differing areas of expertise and interests. In this context, the American Heart Association,

**Table 2**  
Major areas of focus for the HL-PIVOT Network.

Knowledge discovery	<ul style="list-style-type: none"> <li>• Conduct healthy living research that furthers our understanding of the field and has real world applicability</li> <li>• Whenever possible, collaborate on research projects within the network to increase broad applicability and impact</li> <li>• Disseminate findings through peer-reviewed publications and conference presentations</li> </ul>
Education	<ul style="list-style-type: none"> <li>• Advocate for increased research funding for healthy living and collaboratively apply for funding within the network</li> <li>• K-12 and general college/university education: Promote and support initiatives that enhance students' understanding of the importance of healthy living and creates a culture of health and wellness</li> <li>• Health Professional Education: Promote and support the inclusion of healthy living medicine training in all health professional education programs</li> </ul>
Policy Implementation	<ul style="list-style-type: none"> <li>• Propose, support, and promote policies that increase the likelihood of healthy living behaviors on global, national, state, and local levels</li> <li>• Support and promote initiatives and messaging that creates a culture of health and wellness where individuals work, live, and go to school</li> <li>• Support and promote initiatives that increase the use of healthy living medicine in all aspects of health care delivery, moving from a reactionary care model to a proactive, preventive care model</li> </ul>

Abbreviations: HL-PIVOT, Healthy Living for Pandemic Event Protection.


European Society of Cardiology, European Association for Cardiovascular Prevention and Rehabilitation, and American College of Preventive Medicine jointly published a policy statement in 2015 entitled “Healthy Lifestyle Interventions to Combat Noncommunicable Disease: A Novel Nonhierarchical Connectivity Model for Key Stakeholders”.<sup>20</sup> This policy statement took an *all hands on deck* approach to increase HL initiatives and behaviors. Fig 1 illustrates the central figure from this publication, identifying key stakeholders from multiple sectors who can work collaboratively to promote HL from a population to family/individual level. This policy statement, encouraging collaboration across multiple sectors in innovative ways, serves as a foundational framework for the HL-PIVOT network; any activity that promotes HL falls within the scope of the network. Admittedly, the novelty of HL-PIVOT is not to establish the importance that HL behaviors have on health trajectory; this premise is already beyond dispute. Rather, our hope is that HL-PIVOT becomes one of the important and lasting positive initiatives to emerge from the COVID-19 pandemic. Future goals regarding increasing HL behaviors and reducing the prevalence of unfavorable health conditions listed in Table 1 have been previously established; on our current trajectory we will not meet these goals.<sup>28</sup> A change in optics and a new approach for collaboration are primary driving forces for inception of the HL-PIVOT network. Healthy living has previously been described as a polypill that can come in many “shapes and sizes”<sup>18</sup> and still have tremendous benefits. Moreover, the HL polypill is an essential medicine for all the pandemics described in this commentary; no other medicine in the world can claim to have the same impact in treating multiple health conditions at the same time. Given the universal nature of HL medicine, working collaboratively, across professions in an unprecedented way, is warranted and perhaps the only way a meaningful impact on global health can be made at this point. Initially, founding HL-PIVOT network members will be encouraged to bring ongoing HL projects/initiatives under the HL-PIVOT umbrella. Subsequently, we intend to expand the HL-PIVOT network and continue to promote/facilitate collaboration and develop projects/initiatives in all areas listed in Table 2. It is our hope that a unified voice, promoting HL on a global scale, will greatly move the field forward and translate to meaningful improvements in HL behaviors and health outcomes.

In conclusion, the COVID-19 pandemic has caused a dramatic shift in how individuals live on a global scale. This change has dramatic implications for lifestyle behaviors (e.g., physical activity and nutrition), of which we will not understand the true lasting impact for some time. The COVID-19 pandemic also has the potential for valuable, long lasting lessons to be learned. Perhaps there is an opportunity to realize that we have been living in a world of *invisible pandemics* for some time now, specifically the obesity, physical inactivity, diabetes and NCD pandemics. Simultaneously, we are beginning to appreciate the relationship between the COVID-19 pandemic and pandemics that have been existing for years; those with obesity, diabetes or one or more NCD diagnoses who contract SARS-CoV-2 have a significantly higher risk for adverse health events. In this context, increasing HL behaviors across the lifespan, thereby building human resilience, can protect individuals

from NCDs as well as a worsening health trajectory because of a viral infection. Now more than ever, HL is of paramount importance. Our hope is that the HL-PIVOT initiative brings this fact to the forefront and facilitates a new global path forward.

### Appendix 1. Founding members of the HL-PIVOT Network

- Antonio Abbate, MD, PhD**, Virginia Commonwealth University, Richmond, Virginia, USA  
**Gopala Krishna Alaparathi, PhD, PT**, University of Sharjah, United Arab Emirates  
**Leslie D. Austford, MN, MBA, CMPE**, TotalCardiology Research Network, and TotalCardiology™, Calgary, Alberta, Canada  
**Robert S. Axtell, PhD**, Southern Connecticut State University, New Haven, Connecticut, USA  
**Abraham Samuel Babu, PhD, PT**, Manipal College of Health Professions, Manipal Academy of Higher Education, Manipal, India  
**Dr. Simon L. Bacon, PhD, FTOS, FCCS, FABMR**, Concordia University and CIUSSS-NIM, Montreal, Quebec, Canada  
**James R. Bagley, PhD**, San Francisco State University, San Francisco, California, USA  
**Kathy Berra, MSN, NP-BC, FAANP, FPCNA, FAHA, FAAN**, The LifeCare Company, Redwood City, California and Stanford University, Menlo Park, California, USA  
**Jarett D. Berry, MD**, UT Southwestern Medical Center, Dallas, TX, USA  
**Samantha Bond, MS**, University of Illinois at Chicago, Chicago, Illinois, USA  
**Sandra A Billinger, PhD**, University of Kansas Medical Center, Kansas City, Kansas, USA  
**Audrey Borghi-Silva, PhD, PT**, Federal University of Sao Carlos, Sao Paulo, Brazil  
**Harry Brar**, POWERbreathe International Ltd, Warwickshire, UK  
**Patrice Brassard, PhD**, Université Laval, Québec, Canada  
**Peter H. Brubaker PhD**, Wake Forest University, Winston Salem, North Carolina, USA  
**Paolo Brugnoli**, CosMed, Inc., Rome, Italy  
**Marco Brugnoli**, CosMed, Inc., Rome, Italy  
**Tavis S. Campbell, PhD**, University of Calgary, Calgary, Alberta, Canada  
**Salvatore Carbone, PhD, MS**, Virginia Commonwealth University, Richmond, Virginia, USA  
**Kalyana Chakravarthy B, PhD, PT**, University of Sharjah, United Arab Emirates  
**Dr. Cabrera-Lazarini, MBA, Monterrey Institute of Technology, Monterrey, Mexico**  
**Paul J. Chase, PhD**, Ohio University, Athens, Ohio, USA  
**Sundeep Chaudhry, MD, MET-TEST**, Roswell, Georgia, USA



**Fig 1.** Conceptual model for a comprehensive approach to healthy lifestyle promotion, education and interventions.

With permission: Arena, R., et al. Healthy lifestyle interventions to combat noncommunicable disease—a novel nonhierarchical connectivity model for key stakeholders: a policy statement from the American Heart Association, European Society of Cardiology, European Association for Cardiovascular Prevention and Rehabilitation, and American College of Preventive Medicine. *Mayo Clin Proc*. 2015. 90(8): p. 1082–103.

**Jeffrey W Christle, PhD, CEP**, Stanford University, Stanford, CA, USA

**Andrew J Stewart Coats**, Monash University, Melbourne, Australia and University of Warwick, Coventry, United Kingdom

**Naama W. Constantini, MD, DFM, FACSM**, Heidi Rothberg Sport Medicine Center, Shaare Zedek Medical Center, Jerusalem, Israel

**Jeff S. Coombes, PhD**, University of Queensland, Brisbane, Queensland, Australia

**Andréa Lúcia Gonçalves da Silva, PhD, PT**, University of Santa Cruz do Sul, Santa Cruz do Sul, Brazil

**Luiz Carlos Soares de Carvalho Junior, PhD, PT**, Federal University of Recôncavo of Bahia, Cruz das Almas, Bahia, Brazil

**Mildred Vanessa López Cabrera, PhD**, Monterrey Institute of Technology, Monterrey, Mexico

**Victor Zuniga Dourado, PhD, PT**, Federal University of São Paulo, Santos, São Paulo, Brazil

**Daniela Bassi Dibai, PhD, PT**, Ceuma University, São Luís, Brazil

**Snehil Dixit, PhD, PT**, King Khalid University, Abha, Saudi Arabia

**Victor Zuniga Dourado, PhD, PT**, Federal University of São Paulo, Santos, SP, Brazil

**Brandon Dykstra, PhD**, Taylor University, Upland, Indiana, USA

**Patrick Dunn, PhD**, American Heart Association and Walden University, Dallas, Texas, USA

**Diane Eastabrook**, Freelance Journalist, Chicago, Illinois, USA

**Barry A. Franklin, PhD**, Beaumont Health, Royal Oak, Michigan, USA

**Sherry L. Grace, PhD**, York University & University Health Network, Toronto, Canada and International Council of Cardiovascular Prevention and Rehabilitation, Markham, Ontario, Canada

**Gregory J. Grosicki, PhD**, Georgia Southern University (Armstrong Campus), Savannah, Georgia, USA

**Solange Guizilini, PhD, PT**, Federal University of Sao Paulo, Sao Paulo, Brazil.

**Grenita Hall, PhD**, University of Illinois at Chicago, Chicago, Illinois, USA

**Matthew P. Harber, PhD**, Ball State University, Muncie, Indiana, USA

**Erik Hayes, PhD**, Taylor University, Upland, Indiana, USA

**Samuel. A. E. Headley, PhD, FACSM**, Springfield College, Springfield, Massachusetts, USA

**John P. Heybach, PhD**, Altus Academy, Chicago, Illinois, USA

**Andrew P. Hills, PhD**, University of Tasmania, Tasmania, Australia  
**Anne E Holland PhD, PT**, Monash University and Alfred Health, Melbourne, Australia  
**Panniyammakal Jeemon, Ph.D, MPH, FESC**, Sree Chitra Tirunal Institute for Medical Sciences and Technology, Trivandrum, Kerala, India  
**Richard A Josephson, MS, MD**, Case Western Reserve University and University Hospitals of Cleveland, Cleveland, Ohio, USA  
**Leonard A. Kaminsky, PhD**, Ball State University, Muncie, Indiana, USA  
**Bradley Kendall, PhD, CSCS**, Taylor University, Upland, Indiana, USA  
**Aditya Khetan, MD**, McMaster University, Hamilton, Ontario, Canada  
**Peter Kokkinos, PhD**, Rutgers University, New Brunswick, NJ, and Veterans Affairs Medical Center Washington, DC, USA  
**Deepika R. Laddu, PhD**, University of Illinois at Chicago, Chicago, Illinois, USA  
**Jari Laukkanen, MD, PhD**, Institute of Clinical Medicine, University of Eastern Finland, Kuopio, Finland, **Dr. Lavoie, Ph.D., FCPA, FABMR, University of Quebec at Montreal, Montreal, Quebec, Canada**  
**Gregory D. Lewis, MD**, Massachusetts General Hospital, Boston, Massachusetts, USA  
**Andrew T. Lovering, PhD**, University of Oregon, Eugene, Oregon, USA  
**Sara Maldonado-Martín, PhD**, Department of Physical Education and Sports, University of the Basque Country (UPV/EHU), Basque Country, Spain  
**Susan Marzolini, RKin, PhD**, KITE Research Institute, Toronto Rehabilitation and University Health Network, Cardiovascular Prevention and Rehabilitation Program, Toronto, Canada  
**Norman Morris, PhD, PT**, Griffith University and Metro North Hospital and Health Service, The Prince Charles Hospital, Queensland, Australia  
**Jonathan Myers, PhD**, VA Palo Alto Health Care System and Stanford University, Palo Alto, California, USA  
**Jagat Narula, MD, PhD**, Mount Sinai Heart, New York, New York, USA  
**Javaid Nauman, PhD**, United Arab Emirates University, Al-Ain, UAE, Norwegian University of Science and Technology, Trondheim, Norway  
**Josef Niebauer, MD, PhD, MBA**, Institute of Sports Medicine, Prevention and Rehabilitation, Paracelsus Medical University, Salzburg, Austria  
**James H. O'Keefe, MD**, University of Missouri, Kansas City, Kansas, USA  
**Silvia Lizett Olivares Olivars, PhD**, Monterrey Institute of Technology, Monterrey, Mexico  
**Francisco B. Ortega, PhD**, University of Granada, Granada, Spain and Karolinska Institute, Stockholm, Sweden  
**Cemal Ozemek, PhD**, University of Illinois at Chicago, Chicago, Illinois, USA  
**Ambarish Pandey, MD, MSCS**, University of Texas Southwestern Medical Center, Dallas, Texas, USA  
**James E. Peterman, PhD**, Ball State University, Muncie, Indiana, USA  
**Shane A. Phillips, PhD, PT**, University of Illinois at Chicago, Chicago, Illinois, USA  
**Dejana Popovic, MD, PhD**, Clinic for Cardiology, Clinical Center of Serbia, University of Belgrade, Belgrade, Serbia

**Axel Pressler, MD, PhD, FESC**, Private Center of Sports Cardiology & Prevention, Munich, Germany  
**Nicolaas P. Pronk, PhD, MA**, HealthPartners Institute, Bloomington, Minnesota, and Harvard TH Chan School of Public Health, Boston, Massachusetts, USA  
**Chathuranga Ranasinghe MBBS, D.Sp.Med, PhD**, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka  
**Mollie Rose, BS**, University of Illinois at Chicago, Chicago, Illinois, USA  
**Robert Ross, PhD, RKin**, Queen's University, Kingston, Ontario, Canada  
**Michel Silva Reis, PhD, PT**, Federal University of Rio de Janeiro, Rio de Janeiro, Brazil  
**Fabrizio Ricci, MD, PhD, FEACVI**, G.d'Annunzio University, Chieti, Italy  
**Daniel S. Rubin, M.D., M.S.**, University of Chicago, Chicago, Illinois, USA  
**Ahmad Sabbahi, PT, PhD**, University of Illinois at Chicago, Chicago, Illinois, USA and Cairo University, Giza, Egypt  
**Patrick Savage, MS**, University of Vermont Medical Center, Cardiac Rehabilitation Program, South Burlington, Vermont, USA  
**Richard Severin, PT, DPT, CCS**, University of Illinois at Chicago, Chicago, Illinois and Baylor University, Waco, Texas, USA  
**Dorthe Stensvold, PhD**, Norwegian University of Science and Technology, Trondheim, Norway  
**Lindsey Strieter, MS**, University of Illinois at Chicago, Chicago, Illinois, USA  
**Xing Guo Sun, MD**, Beijing Fuwai Hospital National Center for Cardiovascular Diseases, Beijing, China  
**Renata Trimer, PhD, PT**, University of Santa Cruz do Sul, Santa Cruz do Sul, Brazil  
**Jean-Luc Vachieri, MD**, Cliniques Universitaires de Bruxelles-Hôpital, Brussels, Belgium  
**Laurie P. Whitset, PhD**, Ligonier, Pennsylvania, USA  
**Mark A. Williams, PhD**, Creighton University, Omaha, Nebraska, USA  
**Ulrik Wisløff, PhD**, Norwegian University of Science and Technology, Trondheim, Norway, and School of Human Movement and Nutrition Science, University of Queensland, Queensland, Australia.

## References

- Merriam-Webster. Definition of pandemic. <https://www.merriam-webster.com/dictionary/pandemic> 2020. Accessed April 17, 2020.
- World Health Organization. Coronavirus disease (COVID-19) pandemic. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019> 2020. Accessed May 30, 2020. [2020].
- Meldrum DR, Morris MA, Gambone JC. Obesity pandemic: causes, consequences, and solutions-but do we have the will? *Fertil Steril* 2017;107(4):833-839.
- Kohl 3rd HW, Craig CL, Lambert EV, et al. The pandemic of physical inactivity: global action for public health. *Lancet* 2012;380(9838):294-305.
- The Lancet. The diabetes pandemic. *Lancet* 2011;378(9786):99.
- Allen L. Are we facing a noncommunicable disease pandemic? *J Epidemiol Glob Health* 2017;7(1):5-9.
- World Health Organization. Physical inactivity: a global public health problem. [https://www.who.int/dietphysicalactivity/factsheet\\_inactivity/en/](https://www.who.int/dietphysicalactivity/factsheet_inactivity/en/) 2020. Accessed March 30, 2020. [2020].
- World Health Organization. Noncommunicable diseases. <https://www.who.int/news-room/fact-sheets/detail/noncommunicable-diseases> 2020. Accessed April 17, 2020.
- World Health Organization. Insufficient physical activity. [https://www.who.int/gho/ncd/risk\\_factors/physical\\_activity\\_text/en/](https://www.who.int/gho/ncd/risk_factors/physical_activity_text/en/) 2020. Accessed April 17, 2020.
- World Health Organization. Global health observatory data: overweight and obesity. [https://www.who.int/gho/ncd/risk\\_factors/overweight\\_obesity/adults/en/](https://www.who.int/gho/ncd/risk_factors/overweight_obesity/adults/en/) 2020. [4/17/2020].
- World Health Organization. Diabetes. <https://www.who.int/news-room/fact-sheets/detail/diabetes> 2020. Accessed April 17, 2020.
- World Health Organization. Cardiovascular diseases. [https://www.who.int/health-topics/cardiovascular-diseases/#tab=tab\\_1](https://www.who.int/health-topics/cardiovascular-diseases/#tab=tab_1) 2020. Accessed April 17, 2020.

13. World Health Organization. Overweight and obesity. <https://www.who.int/en/news-room/fact-sheets/detail/obesity-and-overweight> 2020. Accessed April 18, 2020.
14. Vidal-Puig A, Enerback S. "Obesity-associated metabolic complications, the second wave of the tsunami" 14th Key Symposium. *J Intern Med* 2018;284(5):447–449.
15. Karakurt M, Acar B, Ozeke O, et al. From the obesity tsunami to the diabetes avalanche: primordial prevention of the diabetes-related cardiovascular epidemic by diabeto-cardiologists. *Angiology* 2019;70(4):371–373.
16. The Lancet Public Health. Tackling obesity seriously: the time has come. *Lancet Public Health* 2018;3(4), e153.
17. Sagner M, McNeil A, Puska P, et al. The P4 Health Spectrum - a predictive, preventive, personalized and participatory continuum for promoting healthspan. *Prog Cardiovasc Dis* 2017;59(5):506–521.
18. Arena R, Lavie CJ, Guazzi M. Prescribing a healthy lifestyle poly pill with high therapeutic efficacy in many shapes and sizes. *Am J Lifestyle Med* 2017;11(6):476–478.
19. Arena R, Lavie CJ. Preventing bad and expensive things from happening by taking the healthy living poly pill: everyone needs this medicine. *Mayo Clin Proc* 2017. [S0025-6196(17)30121-0].
20. Arena R, Guazzi M, Lianov L, et al. Healthy lifestyle interventions to combat noncommunicable disease—a novel nonhierarchical connectivity model for key stakeholders: a policy statement from the American Heart Association, European Society of Cardiology, European Association for Cardiovascular Prevention and Rehabilitation, and American College of Preventive Medicine. *Mayo Clin Proc* 2015;90(8):1082–1103.
21. Gaye B, Canonico M, Perier MC, et al. Ideal cardiovascular health, mortality, and vascular events in elderly subjects: the three-city study. *J Am Coll Cardiol* 2017;69(25):3015–3026.
22. Aneni EC, Crippa A, Osondu CU, et al. Estimates of mortality benefit from ideal cardiovascular health metrics: a dose response meta-analysis. *Journal of the American Heart Association* 2017;6(12).
23. Fani Marvasti F, Stafford RS. From sick care to health care—reengineering prevention into the U.S. system. *N Engl J Med* 2012;367(10):889–891.
24. Luzi L, Radaelli MG. Influenza and obesity: its odd relationship and the lessons for COVID-19 pandemic. *Acta Diabetol* 2020;57(6):759–764. <https://doi.org/10.1007/s00592-020-01522-8>. Epub 2020 Apr 5.
25. Dietz W, Santos-Burgoa C. Obesity and its implications for COVID-19 mortality. *Obesity (Silver Spring, Md)* 2020;28(6):1005. <https://doi.org/10.1002/oby.22818>. Epub 2020 Apr 18.
26. Chow N, Fleming-Dutra K, Gierke R, et al. Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019 – United States, February 12–March 28, 2020. *MMWR Morbidity and Mortality Weekly Report* 2020;69(13).
27. Chen T, Dai Z, Mo P, et al. Clinical characteristics and outcomes of older patients with coronavirus disease 2019 (COVID-19) in Wuhan, China (2019): a single-centered, retrospective study. *The Journals of Gerontology Series A, Biological Sciences and Medical Sciences* 2020, glaa089. <https://doi.org/10.1093/gerona/glaa089>. Online ahead of print.
28. Muniyappa R, Gubbi S. COVID-19 pandemic, corona viruses, and diabetes mellitus. *Am J Physiol Endocrinol Metab* 2020;318(5):E736–E741. <https://doi.org/10.1152/ajpendo.00124.2020>. Published online 2020 Mar 31.
29. Yang JK, Feng Y, Yuan MY, et al. Plasma glucose levels and diabetes are independent predictors for mortality and morbidity in patients with SARS. *Diabet Med* 2006;23(6):623–628.
30. Nguyen H, Manolova G, Daskalopoulou C, Vitoratou S, Prince M, Prina AM. Prevalence of multimorbidity in community settings: a systematic review and meta-analysis of observational studies. *J Comorb* 2019;9. [2235042X19870934-12235042X19870934].
31. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting characteristics, comorbidities, and outcomes among 5700 patients hospitalized with COVID-19 in the New York City area. *Jama* 2020;323(20):2052–2059. <https://doi.org/10.1001/jama.2020.6775>. Online ahead of print.
32. Laddu DR, Lavie CJ, Phillips SA, Arena R. Physical activity for immunity protection: inoculating populations with healthy living medicine in preparation for the next pandemic. *Progress in Cardiovascular Diseases* Apr 9 2020. <https://doi.org/10.1016/j.pcad.2020.04.006>. [S0033-0620(20)30078-5, online ahead of print].
33. De Rosa V, Galgani M, Santopalo M, Colamatteo A, Laccetti R, Matarese G. Nutritional control of immunity: balancing the metabolic requirements with an appropriate immune function. *Semin Immunol* 2015;27(5):300–309.
34. Butler MJ, Barrientos RM. The impact of nutrition on COVID-19 susceptibility and long-term consequences. *Brain, Behavior, and Immunity* 2020. <https://doi.org/10.1016/j.bbi.2020.04.040>. Online ahead of print.
35. Zbinden-Foncea H, Francaux M, Deldicque L, Hawley JA. Does high cardiorespiratory fitness confer some protection against pro-inflammatory responses after infection by SARS-CoV-2? *Obesity*. [n/a(n/a)]. doi: 10.1002/oby.22849. Online ahead of print.
36. Hall G, Laddu DR, Phillips SA, Lavie CJ, Arena R. A tale of two pandemics: how will COVID-19 and global trends in physical inactivity and sedentary behavior affect one another? *Progress in Cardiovascular Diseases* Apr 8 2020. <https://doi.org/10.1016/j.pcad.2020.04.005>. [S0033-0620(20)30077-3, online ahead of print].
37. Sim YJ, Yu S, Yoon KJ, Loiacono CM, Kohut ML. Chronic exercise reduces illness severity, decreases viral load, and results in greater anti-inflammatory effects than acute exercise during influenza infection. *J Infect Dis* 2009;200(9):1434–1442.
38. Wong CM, Lai HK, Ou CQ, et al. Is exercise protective against influenza-associated mortality? *PLoS One* 2008;3(5), e2108.
39. Nieman DC, Wentz LM. The compelling link between physical activity and the body's defense system. *J Sport Health Sci* 2019;8(3):201–217.
40. Xiang M, Zhang Z, Kuwahara K. Impact of COVID-19 pandemic on children and adolescents' lifestyle behavior larger than expected. *Progress in Cardiovascular Diseases* 2020. <https://doi.org/10.1016/j.pcad.2020.04.013>. [S0033-0620(2020)30096-30097].
41. Meyer J, McDowell C, Lansing J, et al. Changes in physical activity and sedentary behaviour due to the COVID-19 outbreak and associations with mental health in 3,052 US adults. *Cambridge Open Engage* 2020. <https://doi.org/10.33774/coe-2020-h0b8g>.
42. Ahmed I. COVID-19 - does exercise prescription and maximal oxygen uptake (VO2 max) have a role in risk-stratifying patients? *Clinical Medicine (London, England)* 2020. <https://doi.org/10.7861/clinmed.2020-0111>.