



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Policy Statement

Policy statements and clinical policies are the official policies of the American College of Emergency Physicians and, as such, are not subject to the same peer review process as articles appearing in the journal. Policy statements and clinical policies of ACEP do not necessarily reflect the policies and beliefs of *Annals of Emergency Medicine* and its editors.

Care of Patients With Behavioral Health Emergencies and Suspected or Confirmed COVID-19

Approved April 2020

Originally approved April 2020

0196-0644/\$-see front matter

Copyright © 2020 by the American College of Emergency Physicians.

Care of Patients With Behavioral Health Emergencies and Suspected or Confirmed COVID-19



[Ann Emerg Med. 2020;76:e69-e70.]

A joint policy statement of the American Association for Emergency Psychiatry, American College of Emergency Physicians (ACEP), American Psychiatric Association, Coalition on Psychiatric Emergencies, Crisis Residential Association, and the Emergency Nurses Association

As with environmental disasters and other crisis events, a pandemic may exceed people's usual coping skills and capacity, which, in turn, may lead to problems with anxiety, depression, and increased use of substances, as well as exacerbation of underlying psychiatric disorders. Factors including, but not limited to, social and physical isolation, uncertainty, fear, evolving facts, changes in how individuals access outpatient care, and public health recommendations contribute to this stress. This affects people with and without preexisting psychiatric illnesses and can contribute to a number of challenges for our already taxed emergency and crisis health care system.

The most severely ill people with psychiatric illness have high rates of baseline medical comorbidity, have reduced access to primary care medical resources, and may lack resources to participate in telehealth services. As a result, this group may have elevated vulnerability to coronavirus disease 2019 (COVID-19) and have limitations in accessing services other than in emergency and crisis settings.¹

The following are guidelines for care of the behavioral health patient with suspected or confirmed COVID-19:

1. Encourage preparedness by supporting education and training on the treatment of psychiatric disorders and best practices for the care of the behavioral health patient.
2. Ensure that staff have access to appropriate, adequate personal protective equipment.
3. Encourage the use of existing, available behavioral health crisis services to mitigate unnecessary visits to the emergency department for psychiatric emergencies or for diverting people from psychiatric hospitals whenever possible.
4. Support medical screening through telehealth/telephonic and clinical preadmission screenings and assessments by qualified, licensed professionals. Additionally, we advocate expanded use of telehealth, including prescribing of controlled substances for opioid use disorder through telemedicine, for patient and provider safety in line with infectious disease recommendations (ie, social distancing). Encourage novel use of telehealth in high-risk environments for diversion and mitigation of unnecessary ED visits.
5. Recognize that patients who present with psychiatric complaints may also have co-occurring medical disorders that should have proper medical evaluation. Use preexisting, evidence-based recommendations and screening algorithms to perform appropriate and directed medical evaluations. Encourage providers to identify alternate methods and modalities to make those assessments in the current COVID-19 environment.
6. Understand that people will present in acute psychiatric crisis who are at risk of, have symptoms consistent with, or have tested positive for COVID-

- 19, who will not meet medical admission criteria but will meet criteria for further psychiatric care. Mental health and substance use care, based on the needs of the individual, must remain available.
7. Discourage the use of restraints while keeping people in the least restrictive setting possible that corresponds to their condition or presenting symptoms.
 8. Ensure that medical personnel are evaluating for signs of domestic violence in children, partners and spouses, the elderly, those with intellectual and developmental disabilities, and other vulnerable populations because implementation of social distancing and home-based self-quarantine could increase those risks.
 9. Encourage staff to formulate aftercare services that are based on existing resources and partnerships in the community.
 10. Provide individuals at risk of suicide with local and national resources of people to talk to if they are suicidal (local crisis call center number, National Suicide Prevention Lifeline, Trans LifeLine, The Trevor Project, and Crisis Text Line).
 11. Encourage the creation and use of psychiatric advanced directives by patients, wherever local jurisdictions permit, that will help provide treatment guidance for providers by patients before their symptoms worsen to the point of impairment in psychiatric medical decisionmaking.
 12. Encourage and promote self-care among individuals providing care to patients and their families. Acknowledge that health care workers will be committed to assisting all shortages/vacancies during these times of crisis, and that it is just as important to maintain one's individual health and wellness for the overall stability of the patients and the care delivery system. In addition to using their own internal coping skills and resources, staff should be made aware of all other local, state, and regional options for care.
 13. Ensure that there is adequate funding, governmental, nongovernmental, and private, to support all activities noted above and ensure that all insurance agencies, public and private, provide appropriate and reasonable reimbursement for the care and treatment of patients with behavioral emergencies.

Approved April 2020
Originally approved April 2020

REFERENCE

1. Osborn DPJ. The poor physical health of people with mental illness. *West J Med.* 2001;175:329-332.

<https://doi.org/10.1016/j.annemergmed.2020.06.015>