



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Antiviral activity of sulfated polysaccharides from marine algae and its application in combating COVID-19: Mini review

Nidhi Hans, Anushree Malik, Satyanarayan Naik^{*}

Centre for Rural Development and Technology, Indian Institute of Technology Delhi, New Delhi, India

ARTICLE INFO

Keywords:

Sulfated galactans
Virucidal activity
Respiratory syndrome
Immune response
Structural classification
Drug development

ABSTRACT

Marine-derived sulfated polysaccharides possess various antiviral activities against a broad range of enveloped and non-enveloped viruses. It has become the potential source of antiviral drugs for pharmaceutical development. In this review, we will discuss the different types of sulfated polysaccharides and their structural classification. Some of the major sulfated polysaccharides with potent antiviral activity, including carrageenan, agar, ulvan, fucoidan, and alginates, are considered in this review. The mechanism of these sulfated polysaccharides in inhibiting the different stages of the viral infection process inside the host cell is also demonstrated. It involves blocking the initial entry of the virus or inhibiting its transcription and translation by modulating the immune response of the host cell. In addition, we explore the potential of sulfated polysaccharides as antiviral agents in preventing recent Corona Virus Disease-2019 (COVID-19).

1. Introduction

In recent years, there is a sudden outburst of some viral diseases that have caused severe harm to human health (Wang et al., 2020). In the discovery and development of drugs, natural products play a vital role during the last decades. Various natural products, mostly from plant sources (root, bark, flowers, or essential oils) and algae-derived compounds are considered effective alternative against multiple diseases (Dutta et al., 2019; Kushwaha et al., 2020; Singh et al., 2019). These natural products contain structurally diverse active substances with a wide range of biological activities (Goswami et al., 2020; Kushwaha et al., 2015). The use of algae to prevent or treat numerous diseases has been exploited for many years and is still used in healthcare in many countries (Kushwaha et al., 2017; Xian et al., 2020). Seaweeds have been recognized as rich and valuable sources of bioactive compounds because of their various biological activities (Goswami et al., 2019a; Khalid et al., 2018). Marine sulfated polysaccharides are considered as a potential source of biologically active compounds for drug development (Bind et al., 2019; Hans et al., 2019). These compounds have been reported to have varieties of pharmacological activities such as antitumor, antiviral, antioxidant, antimicrobial, anticoagulant, and immune-inflammatory effects (Bind et al., 2018; Fedorov et al., 2013; Goswami et al., 2019b).

Sulfated polysaccharides (SPs) are natural complex polymers found

majorly in the cell walls of marine algae. Some of the essential SPs include carrageenan and agar from red macroalgae, ulvan from green macroalgae, and fucoidan and laminarian from brown macroalgae (Wijesekara et al., 2011). The sulfated polysaccharides of seaweeds have been shown to exhibit antiviral activity against a broad spectrum of viruses. It has been reported to inhibit antiviral activity against Herpes Simplex virus (HSV) (Gomaa and Elshoubaky, 2016), Human immunodeficiency virus type-1 (HIV-1) (Besednova et al., 2019), chikungunya virus (Cirne-Santos et al., 2019), and many other enveloped and non-enveloped viruses. Also, the antiviral activity of SPs against the currently on-going pandemic coronavirus disease 2019 (COVID-19) is reported (Chen et al., 2020). COVID-19 is a severe acute respiratory syndrome that may cause illness in animals or humans. This virus primarily spreads between people during close contact, often via small droplets from the nose or mouth, which are expelled by coughing, sneezing, or talking. According to worldometer, as of 20th November 2020, over 57,019,580 identified cases of COVID 19 worldwide in 218 countries and territories (Worldometer, 2020). There is no specific medicine to prevent or treat this disease. Marine sulfated polysaccharide exerts virucidal effect by intervening in different stages of viral infection. Thus, it raises the possibility for the advancement of antiviral agents in therapeutics (Wang et al., 2012). These natural nontoxic and effective antiviral drugs are currently investigated to analyze their potential role in the prevention of contagious diseases in humans.

^{*} Corresponding author at: Centre for Rural Development and Technology, Indian Institute of Technology, Delhi, New Delhi 110016, India.
E-mail address: snn@rdat.iitd.ac.in (S. Naik).

In this mini-review, the main focus is on the neoteric development in the field of antiviral activity of sulfated polysaccharides isolated from the marine biomasses, their molecular significances, and their mechanism of action in preventing and regulating immune responses of the host cell. Also, the role of sulfated polysaccharides in preventing COVID-

19 is briefly discussed. In the last decade, the research and developments on the structural or molecular characteristics and biological potency of sulfated polysaccharides have been done to a more considerable extent. From this pile of research findings, a glimpse of information is tried to be illustrated in this review paper.




Fig. 1. (a-k): Chemical structure of Marine derived-sulfated polysaccharides.




Fig. 1. (continued).

2. Worldwide production of marine sulfated polysaccharides

The global production of marine macroalgae is approximately 33 million tonnes in 2017, out of which 32 million tonnes were harvested from the culture sector (Ferdouse et al., 2018). The leading macroalgae producing countries are China, Norway, Indonesia, France, Ireland, and India (FAO, 2019). The universal seaweed market size was valued at \$4097.93 million in 2017 and is estimated to reach \$9075.65 million by 2024, which will lead to an increase in annual growth rate to 12.0% in 2024 (Allied market research, 2018). In 2012, the highest production of some of the common red seaweeds was 6.1 million tonnes of *Eucheuma* species (worth 1.2 billion USD), 0.8 million tonnes of *Gracilaria* species (worth 34 million USD), 2.1 million tonnes of *Kappaphycus* species (371 million USD), and 0.7 million tonnes of *Porphyra* species (worth 13 billion USD) (Bjerrregaard et al., 2016). Some of the brown seaweed production is 5.7 million tonnes of *Saccarina* species worth 330 million USD and 2.1 million tonnes of *Undaria pinnatifida* worth 0.9 billion USD. Green seaweed production is 2 million tonnes of *Ulva* species, 1.2 million tonnes of *Caulerpa* species (Ferdouse et al., 2018).

Red macroalgae are commercially more important than green and brown macroalgae as it covers three-fourth of the global value. There is an increase in demand for red seaweeds in the manufacturing of hydrocolloids such as agar and carrageenan. These hydrocolloids have various applications in food, pharmaceutical, and biotechnological industries. Worldwide production of agar in 2014 was about 10,600 tonnes, with a wholesale value of some \$191 million. World carrageenan

production exceeded 60,000 tonnes in 2014, with a value of over US \$626 million. In 2014 production of alginates was 30,000 tonnes with a value of about US\$339 million (Rhein-Knudsen et al., 2015). Many other macroalgae derived products like biomass, protein or organic chemicals have growing needs globally. Large-scale seaweed production offers research and development to expand the range of algae-derived products to upgrade the industrial market.

3. Marine-derived sulfated polysaccharides

3.1. Carrageenan and agar from red macroalgae

Sulfated galactans are the main polysaccharide component of red algae, which have a linear backbone of alternating 3-linked β -D-galactopyranose and 4-linked α -galactopyranose. There exist two major instances of the sulfated galactans isolated from the red algae, where agarans possess the 4-linked α -galactose moiety with levorotatory (L-) configuration and the other one termed as carrageenans consist of the similar linkage with dextrorotatory (D-) configuration (Al-Alawi et al., 2011). Thus carrageenans consist of linear chains of alternating β -D-galactopyranose units (G-units) with a linkage between the 1 and 3 positions of the monomeric units and α -galactopyranose units (D-units) with a linkage between 1 and 4 positions or 3, 6- α -galactopyranose (Anhydrous) units (AnGal units) (Ferreira et al., 2012). The most relevant carrageenans which are commercially produced are kappa (κ) [Fig. 1a], iota (ι) [Fig. 1b] and lambda (λ) [Fig. 1c], which differ

according to the number and position of sulfate ester groups (S) and the occurrence of 3, 6-anhydro-D-galactopyranose units. κ -carrageenan has one (25–30%), ι -carrageenan has two (28–30%), and λ -carrageenan has three (32–39%) of sulfate ester groups (Muthukumar et al., 2020). The composition of carrageenan varies among species such as *Kappaphycus alvarezii* is a significant source of κ -carrageenan (Rudke et al., 2020), *Euचेuma denticulatum* consists of ι -carrageenan (Jönsson et al., 2020) and *Gigartina skottsbergii* and *Chondrus crispus* comprises of λ -carrageenan (Zhu et al., 2018). Biological precursor carrageenan named μ (μ) and ν (ν) is subjected to hot alkaline treatment, which leads to the cyclization of 3,6 anhydro rings. This pretreatment converts μ and ν into commercial κ - and ι -carrageenan, respectively (Ortiz-Tena et al., 2017). During carrageenan preparation, galactose and sulfate can be either substituted by xylose, glucose, uronic acid, methyl ethers, or pyruvate groups (Yu et al., 2010).

Agar from red macroalgae is composed of two polysaccharides, agarose [Fig. 1d] and agaropectin [Fig. 1e]. It is mainly found in *Gelidium* and *Gracilaria* species. It is built of repeating alternating chains of 1 \rightarrow 3- β -D-galactopyranose unit and 1 \rightarrow 4- α -L-galactopyranose or 3 \rightarrow 6- α -L-galactopyranose (Anhydrous) unit (Usov, 2011). The β -D-galactopyranose unit can be substituted either by sulfate esters, methoxy groups, or pyruvic acid acetals groups (Lee et al., 2017a). For example, structural analysis of sulfated agar in *Polysiphonia nigrescens* is mainly replaced with sulfate on C6, with minor proportions of methyl ether and β -D-xylose (Bouhlal et al., 2011). Similarly, the sulfate group is present at the C2 position of agar isolated from *Acanthophora spicifera* with minor substitutions of pyruvylated, sulfated, and sulfated/pyruvylated disaccharide alditols (Gonçalves et al., 2002).

3.2. Ulvan from green macroalgae

Ulvan is one of the most complex sulfated polysaccharides representing about 9–36% of the algae dry weight and found in *Ulva*, *Gayralia*, and *Monostroma* species. Ulvan mainly consists of L-rhamnose constituting 5.0–92.2 M%, D-glucuronic acid comprising 2.6–52.0 M%, D-xylose constituting 0.0–38.0 M%, L-iduronic acid, which constitutes 0.6–15.3 M%, and sulfate in different proportions (Kim, 2015). The ulvan backbone mostly consists of α - and β -(1 \rightarrow 4)-linked sugars with characteristic repeating disaccharide units. Aldobiuronic acids, also known as ulvanobiuronic acid (types A and B) [Fig. 1f, g], and aldobiases, known as ulvanobiases (type U) are two major disaccharide repeating units [Fig. 1h, i] (Kidgell et al., 2019). One of the most common disaccharide units, Ulvanobiuronic acid type A_{3S} consists of β -D-glucuronic acid (1,4)-linked to α -L-rhamnose 3-sulfate, while in type B_{3S} α -L-iduronic acid is (1 \rightarrow 4)-linked to α -L-rhamnose 3-sulfate. Ulvanobiose U_{3S} consists of β -D-xylose 2-sulfate (1 \rightarrow 4)-linked to α -L-rhamnose 3-sulfate and type U_{2S,3S} consist of β -D-xylose (1 \rightarrow 4)-linked to α -L-rhamnose 3-sulfate (Figueira et al., 2020). Some of the marine algae synthesized these sulfated polysaccharides. Such as *Monostroma nitidum* composed of rhamnan sulfate polysaccharides (Lee et al., 2010). *Gayralia oxysperma* is consisting of heterorhamnan sulfate polysaccharides units (Li et al., 2012). *Enteromorpha compressa* composed of sulfated heteroglycuronan units, which contains rhamnose with terminal linkages between 1 and 4 position or 1 to (2,4) positions, xylopyranose unit with the linkage between 1 and 4 positions, and (1,4)-glucuronic acid units linked terminally (Ray, 2006).

3.3. Fucoidan from brown macroalgae

Fucoidan is a heterogeneous sulfated polysaccharide that constitutes 25–30% of the dry algal weight. It consists of a backbone of α -(1 \rightarrow 3)-L-fucopyranose residues [Fig. 1j] or alternating α -(1 \rightarrow 3) and α -(1 \rightarrow 4)-linked L-fucopyranosyls [Fig. 1k]. These residues are either substituted with sulfate, acetate, or glycosyl units such as glucuronic acid. It also contains minimal quantities of monosaccharides like D-xylose, D-galactose, D-mannose, and uronic acids (Ale et al., 2011). Fucoidan from *Fucus*

evanescens and *Ascophyllum nodosum* contain alternative units of α -(1 \rightarrow 3)-linked and α -(1 \rightarrow 4)-linked L-fucopyranose. In *Fucus evanescens*, the sulfate group is substituted on C-2 and C-4 fucopyranose residues, and *Ascophyllum nodosum* consists of sulfate at the C-2 position of the 3-linked fucose and C-2- and C-3 positions of the 4-linked fucose (Yuguchi et al., 2016).

4. The antiviral mechanism of sulfated polysaccharide

Marine sulfated polysaccharides exhibit unique structures that exert antiviral effects. It obstructs different phases of the viral life cycle by directly inactivating virions before infection or by inhibiting its replication inside the host cell. Thus, seaweeds rich in polysaccharides largely contribute to the discovery and development of antiviral drugs. The primary stages in the life cycle of the virus are attachment, penetration, uncoating, biosynthesis, viral assembly, and release, which differ with species [Fig. 2].

4.1. Inhibition of virus attachment

The initial contact occurs by the ionic interaction between positively charged external glycoproteins that are present on the enveloped virus surface and the negatively charged constituents of the host cell surface. A high density of negative charge is present on the cell surface due to the existence of sulfate residues that interact with the positively charged domain of viral glycoprotein that disrupts the initial virus-cell contact (Sepúlveda-Crespo et al., 2017). Carrageenan might prevent virus infection into the host cell by directly inhibiting its binding to the cell surface.

4.2. Inhibition of virus penetration

After binding of the virus onto the host cell, the subsequent virus invasion process is followed by irreversible adsorption via electrostatic interaction between the host cell and virus receptors. To cease this penetration operation, some of the sulfated marine polysaccharides interact with virus receptors, which block its interaction with the host cell surface or directly interact with virions to prevent virus infection. Several studies have shown that negative charges residing over the sulfate group of carrageenan interact with the virus by covering up the positive charge of the viral receptors (Wang et al., 2012).

4.3. Inhibit interiorization and uncoating of the virus

The virus is penetrated inside the host cell by invagination of the outer membrane to form a vacuole. Then it is transported through the intracellular fluid, i.e., cytoplasm, and delivered to endosomes and other intracellular organelles. After endocytosis, the virus interacts with the cell membrane or creates an intracellular compartment enclosing virus, thus changing the structure of the viral capsid. When a virus interacts with receptor protein around the endosome, specific signals are produced, which uncoats and discharges the virions (Mercer et al., 2010). The certain sulfated marine polysaccharides are reported to interfere with the virus internalization by interacting with virus membrane proteins. They bind with carbohydrate groups linked to the polypeptide chains of the virus to inhibit its penetration. Also, sulfated polysaccharides bind at the allosteric site of the viral capsid, which prevents the uncoating of the virus inside the host cell.

4.4. Inhibition of virus transcription and translation process

After internalization and uncoating, the virus is replicated inside the host cell. The replication of the virus involves the synthesis of viral messenger RNA, viral protein synthesis, and then viral genome replication mediated by regulatory protein expression. Many marine polysaccharides can inhibit the virus transcription and replication process




Fig. 2. Stages in life cycle of virus (Black) - (i) virus attachment, (ii) virus penetration and uncoating (iii) biosynthesis (iv) viral assembly (v) viral release. Mechanism of antiviral actions of Sulfated polysaccharides (SPs) (Red)- (i) inhibit virus attachment (ii) inhibit virus penetration (iii) inhibit virus internalization and uncoating (iv) Inhibit virus transcription and replication process. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

after entering into the host cells by interfering with the replication enzymes such as reverse transcriptase or by preventing the formation of proteins from messenger RNA in the host cell (Queiroz et al., 2008).

5. Antiviral activity of marine-derived sulfated polysaccharides

Studies conducted over 50 years ago showed the effects of polysaccharides from marine algae in the inhibition of influenza B and mumps virus. These findings have introduced algae-derived polysaccharides as a potent source of antiviral agents. The scientific research on antiviral compounds and various biological activities of marine sulfated polysaccharides towards several deadly viruses has eventually increased in the past decade. Since then, a vast number of studies have been published on the antiviral property of various sulfated polysaccharides. They prevent a wide range of contagion diseases such as human immunodeficiency virus (HIV), hepatitis-C virus (HCV), dengue virus (DENV), herpesvirus (HSV-1 and HSV-2), human papillomavirus (HPV), and many respiratory tract viruses. Antiviral activity of some of the sulfated polysaccharides and their mode of action are discussed below [Table 1].

5.1. The antiviral potential of sulfated polysaccharides carrageenan from red macroalgae

The presence of 3,6-anhydrogalactopyranose and allocation of sulfate groups on carrageenan show different inhibitory effects on different viruses (Jiao et al., 2011). Buck et al. (2006) reported that iota-carrageenan is a potent infection inhibitor for a specific sexually transmitted human papillomavirus (HPV) by preventing the binding of HPV virions to cells. It has shown a thousand-fold more susceptibility, with a 50% inhibitory concentration value (IC_{50}) at low concentration. They have also reported that contraceptives lubricated with carrageenans could effectively block HPV infection transmission through sexual contact. Besides, carrageenans incorporated within infant formulas have

shown restricted vertical transmission of HPV from mother to baby.

It is reported that λ -carrageenan and hybrid μ/λ carrageenan isolated from *Gigartina skottsbergii* can inhibit the activity of type-1 and type-2 herpes simplex virus (HSV). It interferes between cell surface heparin sulfate and HSV glycoprotein interlinkage at an early stage of infection. Carrageenan and its oligosaccharide derivatives bind with the glycoprotein present on the virus surface. This linkage leads to the denaturation and inactivation of HSV glycoprotein. Thus it inhibits the viral adsorption and replication inside the host cell (Carlucci et al., 2002).

Linear chains of galactopyranosyl residues of λ - and ν -carrageenan are found to be potent inhibitors of dengue virus type-2 (DENV-2) and type-3 (DENV-3) in monkey kidney cells and human hepatic cells. Results have shown that the dengue virus can be inhibited by carrageenan either at the early stage of virus adsorption or after the encapsulation of the capsid of virus-containing nucleic acid into the cytoplasm (Talarico and Damonte, 2007). Yamada and his co-workers demonstrated the anti-HIV activity of carrageenan. Carrageenan was modified by depolymerization and sulfation to increase its activity. Depolymerized and sulfated κ and ν carrageenan show the highest anti-HIV activity (Yamada et al., 2000). Also, low molecular weight κ -carrageenan obtained by microwave-aided acid hydrolysis was acetylated and sulfated. The obtained derivative of carrageenan displayed the highest activity against the influenza virus (H1N1) in mice (Tang et al., 2013). Wang and his co-workers have reported that carrageenan could inhibit mRNA and protein expression of the influenza A virus. Their result shows that after the virus is internalized inside the host cell, carrageenan blocks the viral replication cycle. It inhibited the formation of nucleic acids and proteins, which cease the release of the virus outside the host cell (Wang et al., 2011).

Therefore, the antiviral activity of carrageenan has distinct inhibitory actions on different viruses because of their molecular mass and level of sulfation, which comprises of sulfate content and the position of sulfation.

Table 1
Antiviral activity of some sulfated polysaccharides from marine algae.

Organism	Compounds	Activity	References
Red macroalgae <i>Acanthophora spicifera</i>	Agaran	^a HSV-1, HSV-2, inhibit initial attachment of virus to the cells	Bedoux et al., 2017
<i>Meristiella gelidium</i>	Iota/kappa/nu-hybrid carrageenan	HSV-2, ^b DENV-2, no cytotoxicity on Vero cells	De Sf-Tischer et al., 2006
<i>Acanthophora spicifera</i>	Carrageenan	HSV-1, ^c RVFV, inhibit virus replication	Gomaa and Elshoubaky, 2016
Sigma–Aldrich	Kappa-carrageenan	Enterovirus 71 prevent viral replication before and after adsorption	Chiu et al., 2012b
Purchased from FMC Biopolymers	Iota-carrageenan	^d HRV, prevent binding and replication of virions	Grassauer et al., 2008
<i>Gymnogongrus griffithsiae</i> , <i>Cryptonemia crenulata</i>	Sulfated galactans	HSV-1, HSV-2, inhibit virus adsorption	Talarico et al., 2004
<i>Gracilaria corticata</i>	Sulfated galactan	HSV-1, HSV-2, inhibition of virus attachment	Mazumder et al., 2002
Purchased from FMC Biopolymers	Iota-carrageenan	Pandemic ^e H1N1/2009 reduced viral replication	Leibbrandt et al., 2010
<i>Gymnogongrus torulosus</i>	DL-galactan	HSV-2, DENV-2, inhibit virus adsorption	Pujol et al., 2002
<i>Sebdenia polydactyla</i>	Sulfated xylomannans	HSV-1	Ghosh et al., 2009
<i>Lithothamnion muelleri</i>	Sulfated xylogalactans	HSV-1, HSV-2, inhibit viral adsorption and penetration into the cells	Malagoli et al., 2014
<i>Scinaia hatei</i>	Sulfated xylomannan	HSV-1, HSV-2, inhibit virus binding	Mandal et al., 2008
<i>Sphaerococcus coronopifolius</i> , <i>Boergeseniella thuyoides</i>	Water-soluble sulfated polysaccharides	HSV-1, ^f HIV-1, inhibit adsorption and replication of the virus	Bouhhal et al., 2011
Green macroalgae <i>Enteromorpha compressa</i>	Ulvan	HSV, inhibit adsorption and replication of the virus	Lopes et al., 2017
<i>Ulva intestinalis</i>	Ulvan	Measles virus, reduction of syncytia formation and low cytotoxicity	Morán-Santibañez et al., 2016
<i>Ulva armoricana</i>	Enzyme assisted-Ulvan	HSV-1, no cytotoxicity on Vero cells	Hardouin et al., 2016
<i>Monostroma latissimum</i>	Rhamnan sulfate	HSV-1, HIV-1, ^g HCMV, virus adsorption and replication	Wang et al., 2018
Brown macroalgae <i>Adenocystis utricularis</i>	Fucoidan (galactofucan)	HSV-1, HSV-2, no cytotoxicity	Ponce et al., 2003
<i>Dictyota dichotoma</i>	Fucoidan (galactofucan)	HSV-1, reduction in plaque formation	Rabanal et al., 2014
<i>Cladosiphon okamuranus</i>	Fucoidan (glucuronic acid, sulfated fucose)	DENV-2, direct binding	Hidari et al., 2008

Table 1 (continued)

Organism	Compounds	Activity	References
<i>Adenocystis utricularis</i>	Fucoidan	HIV-1, block entry of the virus	Trincherio et al., 2009
<i>Cystoseira indica</i>	Sulfated fucans	HSV-1, HSV-2, inhibit virus adsorption	Mandal et al., 2007
<i>Sargassum mcclurei</i>	Fucoidan	HIV-1, block entry of the virus	Thuy et al., 2015
<i>Caulerpa brachypus</i>	Xylan-fucoidan	HSV-1, inhibit attachment, penetration, and later stage replication	Lee et al., 2004
<i>Fucus vesiculosus</i>	Fucoidan	^h BVDV, inhibit binding of the virus	Güven et al., 2020
<i>Laminaria japonica</i>	Fucoidan	ⁱ H5N1, inhibit entry of the virus	Makarenkova et al., 2010
<i>Undaria pinnatifida</i>	Galactofucan	HSV-1, HSV-2, HCMV, inhibit virus entry, and host-virus binding	Hemmingson et al., 2006
<i>Sargassum trichophyllum</i>	Fucoidan	HSV-2, inhibit virus adsorption, penetration and replication	Lee et al., 2011

^a Herpes simplex virus.

^b Dengue virus.

^c Rift valley fever virus.

^d Human rhinovirus.

^e Influenza A virus.

^f Human immunodeficiency virus.

^g Human cytomegalovirus.

^h Bovine viral diarrhea virus.

ⁱ Avian influenza virus.

5.2. The antiviral potential of sulfated polysaccharide ulvan from green macroalgae

Chiu and his colleagues have tested the antiviral effects of ulvan from *Ulva lactuca* on the Japanese encephalitis virus (JEV). It is a mosquito-borne flavivirus that causes acute encephalitis in humans. They reported that ulvan could inhibit JEV infection in Vero cells by blocking virus adsorption and making the virus unable to enter cells (Chiu et al., 2012a).

Aguilar-Briseño and his teammates have isolated Ulvan from *Ulva clathrata* and fucoidan from *Cladosiphon okamuranus*. They evaluated the antiviral activity and mode of action of isolated Ulvan, fucoidan, and their mixture on Newcastle disease virus (NDV). Their antiviral activity was assessed by syncytia reduction assay. They stated that syncytia formation could only be inhibited if ulvan and its mixture were added before protein cleavage as once protein is cleaved via trypsin digestion; ulvan will lose the ability to inhibit syncytia formation. Their result shows that ulvan was able to inhibit 67% of syncytia formation, fucoidan inhibits 47%, and their mixture inhibits 59% of syncytia formation (Aguilar-Briseño et al., 2015).

5.3. The antiviral potential of sulfated polysaccharides fucoidan from brown macroalgae

Queiroz and his associates have examined the potential of fucoidan to inhibit HIV reverse transcriptase (RT) enzyme before integrating the virus into the host cell. RT enzyme is encoded by a retrovirus that catalyzes the viral RNA (ribonucleic acid) genome into DNA (deoxy-ribonucleic acid). They have also studied the role of sulfate and carboxyl groups present on the fucoidan to inhibit HIV. They have assessed that fucoidan has a prominent effect on reverse transcriptase in vitro, even at low concentration values. Their results have also shown that fucoidan polysaccharide, modified by carboxy-reduction and desulfation, has

reduced the inhibitory activity for HIV RT four times compared to unmodified ones. Therefore, they have suggested that chemical structures of molecules such as degree of sulfation, disaccharides linkages, polymeric backbone, and the degree of polymer chemical reactions could play an important function in affecting antiviral properties of fucoidan polysaccharides (Queiroz et al., 2008).

Dinesh et al. (2016) reported the anti-HIV-1 property of fucoidan extracted from *Sargassum swartzii*. They observed that it exhibit inhibitory activity against HIV-1 p24 antigen ($95.6 \pm 1.1\%$) and reverse transcriptase ($78.9 \pm 1.43\%$) at a concentration of 25 $\mu\text{g/ml}$. Yuguchi et al. (2016) reported that fucoidan obtained from certain brown seaweeds displayed similar anti-HIV activity by blocking the early steps of HIV entry into target cells. They have suggested that sulfate content and its position in the fucoidan backbone is not associated with antiviral activity. Hayashi and his colleagues have reported that fucoidan from *Undaria pinnatifida* can prevent the replication of influenza A virus inside the host cell by blocking the transcription process. They have also said that fucoidan helps in stimulating both innate and adaptive immune functions by regulating B- lymphocytes (bone marrow cells) and T- (thymus cells) lymphocytes in virus-infected mice (Hayashi et al., 2013).

6. Modulation of host antiviral immune responses with seaweed polysaccharides

The viral infection can induce the antiviral immune responses of the host cell. Type I interferon system (IFN), host natural killer (NK) cells, and phagocytic cells contribute significantly against viral infection. IFN interacts with receptors on the cell surface that generates the antiviral proteins and enhances the activity of NK cells, T lymphocytes, and macrophages (Sozzani et al., 2010). Several seaweed polysaccharides help indirectly in the inhibition of viral infection and speed up the process of virus removal by activating the NK cells and stimulating antiviral immune factors.

The λ -carrageenan polysaccharide has been shown to induce the synthesis of interferon, which contributes to the biological effects on the immune response. Microwave degradation of λ -carrageenan from *Chondrus ocellatus* has been reported to inhibit the growth of tumors by improving the activity of interferons and enhancing lymphocyte multiplication (Liu et al., 2019). Fucoidan has various immune-modulating effects, such as stimulating the production of NK cells, development of dendritic cells, and the function of cytotoxic lymphocytes. Moreover, it enhances the Th1-type immune response by producing antibodies against determinants of specific antigen and generates memory T cells against the particular virus (Zhang et al., 2015). Carrageenan from *Kappaphycus striatum* modulates the immune response by enhancing the activity of macrophages and NK cells and increases the expression of interleukin-2 (IL-2) and tumor necrosis factor-alpha (TNF- α) (Yuan et al., 2006). The sulfated polysaccharides from *Enteromorpha prolifera* can stimulate macrophages cells. It can also induce the production of various cytokines such as interferon-gamma (IFN- γ) and interleukin-2 (IL-2), thus upregulating the Th-1 response (Kim et al., 2011). Polysaccharides from brown and green macroalgae are also reported to improve the antiviral immune response of the host cell. Alginate polysaccharide 911 can block the absorption of the virus and inhibit its replication by modulating the immune response of T and B lymphocytes (Xianliang et al., 2000).

In conclusion, marine sulfated polysaccharides can impede the virus adsorption and its replication and, thus, activate the immune response of the host directly by accelerating the process of virus clearance.

7. Potential antiviral application of marine polysaccharide in combating COVID-19

In late December 2019, an outbreak of novel coronavirus diseases (COVID-19) in Wuhan, China has spread quickly nationwide. According to the World Health Organization (WHO), it is a pandemic disease

caused by a unique beta-coronavirus, one of the four genera of coronavirus (alpha, beta, delta, and gamma). At present, it is formally known as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) (Gorbalenya et al., 2020). It is transmitted through the air (cough or sneeze), physical contact, contaminated objects, or mass gathering. Infected people experience a dry cough, high fever, sore throat, and face difficulty in breathing.

The COVID-19 is a novel enveloped beta-coronavirus which mainly caused infections in the pulmonary and digestive tract (Zhu et al., 2020). The outer membrane of the virus is made up of spike proteins that protrude out of the cell surface. The inside virus contains capsid protein along with its genome and single-stranded RNA. It is reported that glycoproteins on the surface of coronavirus bind to the receptors of host cells. This attachment leads to the entrance of the virus inside the cell. Spike proteins on the virus surface bind with blood vessel cells of the heart and kidney and with epithelial cells of lungs and intestine via angiotensin-converting enzyme (Xu et al., 2020).

There are currently no vaccine or effective antiviral treatments for COVID-19 in general, although hundreds of clinical trials are underway. Recently, marine sulfated polysaccharides have gain attention because of their antiviral effects [Table 2]. It has been proved that the antiviral properties of carrageenan have shown promising inhibitory effects on many types of viruses. It can effectively disrupt the interaction between virus and host cell receptors, thus block the internalization of virus particles. Koenighofer and co-workers have developed carrageenan-based nasal spray, which was effective in a patient with a common cold infected by a human coronavirus OC43 (beta) and human coronavirus 229E (alpha). Its application has shown a 2.5-fold reduction in relapses of symptoms and increased viral clearance compared to placebo-treated patients during clinical trials (Koenighofer et al., 2014). To treat throat infection caused by a human coronavirus OC43, Morokutti-Kurz and his colleagues have prepared lozenges comprising 10 mg iota-carrageenan as active pharmaceutical constituents. Their results show that glycoproteins on the coronavirus surface were

Table 2
Some of the marine sulfated polysaccharides reported to combat COVID-19.

Compound	Efficacy	Effect/mechanism	Reference
Iota-carrageenan based nasal spray	Concentration as low as 6 $\mu\text{g/ml}$	Inhibit SARS-CoV-2 in vitro	Bansal et al., 2020
Carrageenan based nasal spray	–	2.5-Fold reduction in relapses of symptoms	Koenighofer et al., 2014
Fucoidan and carrageenan	At 3.90-500 $\mu\text{g/ml}$ concentration	Prevent SARS-CoV-2 entry into the cell by binding to the S-glycoprotein	Song et al., 2020
Iota-carrageenan based lozenges	At a concentration of 10 mg	Denaturation of coronavirus glycoprotein	Morokutti-Kurz et al., 2017
Fucoidan Kappa carrageenan Iota-carrageenan	Concentration of 0.01–10% by weight	Prevent or treat respiratory tract infections caused by the virus	Grassauer and Prieschl-Grassauer, 2019
Fucoidan	At approx. 83 nM concentration	SP bind to spike protein of SARS-CoV-2 in vitro, preventing its binding to the host cell	Kwon et al., 2020
Iota-carrageenan based nasal spray	At the concentration of 0.12%	Relieve nasal congestion in the upper respiratory tract	Graf et al., 2018
Iota-carrageenan	Total 1 mg dose daily	2.40 fold increase in recovery rate from coronavirus infection	Hemilä and Chalker, 2020
Lamda-carrageenan	At 0.3–1.4 $\mu\text{g/ml}$ concentration	Target viral attachment to cell surface receptors	Jang et al., 2020

denatured during the residence time of lozenge inside the mouth. Carrageenan based lozenges were responsible for the morphological changes of glycoproteins due to the low pH maintained inside the mouth. Denaturation of glycoproteins leads to inhibit the virucidal actions of coronavirus (Morokutti-Kurz et al., 2017). Graf and his partners have formed a nasal spray formulation comprising 0.05% of xylometazoline hydrochloride and 0.12% of iota-carrageenan. This formulation has been reported to relieve nasal congestion in the upper respiratory tract and simultaneously provides antiviral protection of respiratory mucosa (Graf et al., 2018).

Grassauer and Prieschl-Grassauer have investigated the effect of carrageenan on the inhibition of coronavirus mediated cell death in cat kidney cells. Iota-carrageenan shows significant inhibition at low levels (4 µg/ml), but kappa- and lambda-carrageenan are not sufficient. They reported that pretreatment of the host cell with the highest experimental carrageenan concentration of 400 µg/ml shows only 35% inhibition. This result indicates that pretreatment is not significant enough to protect cells from coronavirus infection. Therefore, they suggested that antiviral active agent carrageenan should be present at the time of infection, i.e., when virus and host cell interaction is about to take place. Thus, carrageenan facilitates protection against coronavirus. It also suggests that carrageenan can be coated or impregnated to a solid surface of hygiene or sanitary item such as gloves, tissue paper, cotton swab, and facial masks (Grassauer and Prieschl-Grassauer, 2019).

8. Benefits of marine SPs over other natural compounds

Several potential vaccines are being studied for COVID 19, but no specific effective treatment is developed. Although many natural compounds other than sulfated polysaccharides from marine algae also shows promising results on COVID 19 patients. Various herbal traditional medicines, plant extracts containing flavonoids and phenolic compounds, and essential oils from medicinal plants are reported to have antiviral bioactivities for COVID 19. Shree et al. (2020) have reported that phytochemicals from medicinal plants such as Ashwagandha (*Withania somnifera*), Giloy (*Tinospora cordifolia*), and Tulsi (*Ocimum sanctum*) showed potential inhibitor against SARS-CoV-2 main protease. Studies show that essential oil from Eucalyptus and Neem (*Azadirachta indica*) helps in the symptomatic treatment of COVID 19 (Roy and Bhattacharyya, 2020). Wahedi et al. (2020) have reported that phenolic compounds from plants such as Stilbene can disrupt the affinity binding of COVID spike protein on the human ACE2 receptor. Jo et al. (2020) has reported that the antiviral activity of some of the flavonoids can efficiently block the enzymatic activity of coronavirus 3C-like protease. Other plant secondary metabolites such as alkaloids, terpenoids, and polyphenols were also reported to prevent the binding of the virus to the host cell (Jahan and Onay, 2020).

Although potential anti-SARS-CoV-2 activity is exhibited by both algae-based and plant-based compounds, they both have their own merits and demerits. Marine macroalgae have been explored by many researchers as an excellent opportunity to become an inexhaustible source of biologically active compounds for the discovery of novel and useful therapeutic drugs. Algae-based and plant-based compounds are safe, biocompatible, and biodegradable, but the production cost of algae-based SPs is less than plant-based natural compounds due to their abundance in the ocean (Ruocco et al., 2016). Marine SPs are water-soluble and can easily be extracted using an aqueous extraction method, unlike plant-based compounds extracted using harmful organic compounds. Physicochemical and mechanical properties of SPs can easily be modified, which increases its use in pharmaceutical industries (Lee et al., 2017b). No clear health risk has been identified in the literature; however, initiatives are required to study the chemical composition, biological potency, bioavailability, toxicity, and associated mechanisms of sulfated polysaccharides in pharmaceutical sectors.

9. Conclusion

The sulfated polysaccharides are considered as a promising antiviral drug in the future. The diverse structure of SPs has an essential role in boosting the host antiviral response by interfering with virus attachment, adsorption, and its replication process. There are several antiviral studies of marine polysaccharides have been performed in vitro or on specific animal models. Across 40 compounds are commercially available in the market, and many more are being implemented at the pre-clinical or clinical stages of human trials. More initiatives are required to study the chemical composition, biological potency, and associated mechanisms of SPs in pharmaceutical sectors.

CRediT authorship contribution statement

Nidhi Hans: Conceptualization, Methodology, Validation, Investigation, Writing an original draft, Visualization. **Anushree Malik:** Conceptualization, Resources, Writing- review & editing, Supervision, Project administration, funding acquisition. **Satyanarayan Naik:** Conceptualization, Resources, Writing- review & editing, Supervision, Project administration, funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgment

The authors sincerely acknowledge the Centre for Rural Development and Technology, Indian Institute of Technology Delhi, New Delhi, India, for financial and technical support to carry out this research work.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

References

- Aguilar-Briseño, J.A., Cruz-Suarez, L.E., Sassi, J.F., Ricque-Marie, D., Zapata-Benavides, P., Mendoza-Gamboa, E., Rodríguez-Padilla, C., Trejo-Avila, L.M., 2015. Sulphated polysaccharides from *Ulva clathrata* and *Cladophora okamuranus* seaweeds both inhibit viral attachment/entry and cell-cell fusion, in NDV infection. *Mar. Drugs* 13, 697–712. <https://doi.org/10.3390/md13020697>.
- Al-Alawi, A.A., Al-Marhubi, I.M., Al-Belushi, B.S.M., Soussi, B., 2011. Characterization of carrageenan extracted from *Hypnea bryoides* in Oman. *Mar. Biotechnol.* 13, 893–899. <https://doi.org/10.1007/s10126-010-9350-7>.
- Ale, M.T., Mikkelsen, J.D., Meyer, A.S., 2011. Review of structure-function relations and extraction methods for fucose-containing sulfated polysaccharides from brown seaweeds. *Mar. Drugs* 9, 2106–2130.
- Allied market research, 2018. Seaweed Market by Product (Red, Brown, and Green) and Application (Human Food, Hydrocolloids, Fertilizers, Animal Feed Additives, and Others) - Global Opportunity Analysis and Industry Forecast. (Accessed 14 April 2020).
- Bansal, S., Jonsson, C.B., Taylor, S.L., Figueroa, J.M., Dugour, A.V., Palacios, C., Vega, J. C., 2020. Iota-carrageenan and xylitol inhibit SARS-CoV-2 in cell culture. *BioRxiv*. <https://doi.org/10.1101/2020.08.19.225854>.
- Bedoux, G., Caamal-Fuentes, E., Boulhou, R., Marty, C., Bourgougnon, N., Freile-Pelegrin, Y., Robledo, D., 2017. Antiviral and cytotoxic activities of polysaccharides extracted from four tropical seaweed species. *Nat. Prod. Commun.* 12 (6) <https://doi.org/10.1177/1934578X1701200602>, 1934578X1701200602.
- Besednova, N.N., Zvyagintseva, T.N., Kuznetsova, T.A., Makarenkova, I.D., Smolina, T.P., Fedyanina, L.N., Kryzhanovsky, S.P., Zaporozhets, T.S., 2019. Marine algae metabolites as promising therapeutics for the prevention and treatment of HIV/AIDS. *Metabolites*. 9 (5), 87. <https://doi.org/10.3390/metabo9050087>.
- Bind, A., Goswami, L., Prakash, V., 2018. Comparative analysis of floating and submerged macrophytes for heavy metal (copper, chromium, arsenic and lead) removal: sorbent preparation, characterization, regeneration and cost estimation. *Geol. Ecol. Landsc.* 2 (2), 61–72. <https://doi.org/10.1080/24749508.2018.1452460>.
- Bind, A., Kushwaha, A., Devi, G., Goswami, S., Sen, B., Prakash, V., 2019. Biosorption valorization of floating and submerged macrophytes for heavy-metal removal in a multi-component system. *Appl Water Sci* 9 (4), 95. <https://doi.org/10.1007/s13201-019-0976-y>.

- Bjerregaard, R., Valderrama, D., Radulovich, R., Diana, J., Capron, M., Mckinnie, C.A., Cedric, M., Hopkins, K., Yarish, C., Goudey, C., Forster, J., 2016. Seaweed Aquaculture for Food Security, Income Generation and Environmental Health in Tropical Developing Countries. World Bank Group, Washington, DC.
- Bouhhal, R., Haslin, C., Chermann, J.C., Collic-Jouault, S., Sinquin, C., Simon, G., Cerantola, S., Riadi, H., Bourgougnon, N., 2011. Antiviral activities of sulfated polysaccharides isolated from *Sphaerococcus coronopifolius* (Rhodophyta, Gigartinales) and *Boergeseniella thuyoides* (Rhodophyta, Ceramiales). *Mar. Drugs* 9 (7), 1187–1209. <https://doi.org/10.3390/md9071187>.
- Buck, C.B., Thompson, C.D., Roberts, J.N., Müller, M., Lowy, D.R., Schiller, J.T., 2006. Carrageenan is a potent inhibitor of papillomavirus infection. *PLoS Pathog.* 2 (7), e69 <https://doi.org/10.1371/journal.ppat.0020069>.
- Carlucci, M.J., Scolaro, L.A., Damonte, E.B., 2002. Herpes simplex virus type 1 variants arising after selection with an antiviral carrageenan: lack of correlation between drug susceptibility and syn phenotype. *J. Med. Virol.* 68, 92–98. <https://doi.org/10.1002/jmv.10174>.
- Chen, X., Han, W., Wang, G., Zhao, X., 2020. Application prospect of polysaccharides in the development of anti-novel coronavirus drugs and vaccines. *Int. J. Biol. Macromol.* <https://doi.org/10.1016/j.ijbiomac.2020.07.106>.
- Chiu, Y.H., Chan, Y.L., Li, T.L., Wu, C.J., 2012a. Inhibition of Japanese encephalitis virus infection by the sulfated polysaccharide extracts from *Ulva lactuca*. *Mar. Biotechnol.* 14, 468–478. <https://doi.org/10.1007/s10126-011-9428-x>.
- Chiu, Y.H., Chan, Y.L., Tsai, L.W., Li, T.L., Wu, C.J., 2012b. Prevention of human enterovirus 71 infection by kappa carrageenan. *Antivir. Res.* 95, 128–134. <https://doi.org/10.1016/j.antiviral.2012.05.009>.
- Cirne-Santos, G.C., Barros, C.D.S., Nogueira, C.C.R., Azevedo, R.C., Yamamoto, K.A., Meira, G.L.S., Vasconcelos, Z.F.M.D., Ratcliffe, N.A., Teixeira, V.L., Schmidt-Chanasit, J., Ferreira, D.F., Paixão, I.C.N.D.P., 2019. Inhibition by marine algae of chikungunya virus isolated from patients in a recent disease outbreak in Rio de Janeiro. *Front. Microbiol.* 10, 2426. <https://doi.org/10.3389/fmicb.2019.02426>.
- De SFTischer, P.C., Talarico, L.B., Nosedá, M.D., Guimarães, S.M.P.B., Damonte, E.B., Duarte, M.E.R., 2006. Chemical structure and antiviral activity of carrageenans from *Meristiella gelidium* against herpes simplex and dengue virus. *Carbohydr. Polym.* 63, 459–465. <https://doi.org/10.1016/j.carbpol.2005.09.020>.
- Dinesh, S., Menon, T., Hanna, L.E., Suresh, V., Sathuvan, M., Manikannan, M., 2016. In vitro anti-HIV-1 activity of fucoidan from *Sargassum swartzii*. *Int. J. Biol. Macromol.* 82, 83–88. <https://doi.org/10.1016/j.ijbiomac.2015.09.078>.
- Dutta, M., Kushwaha, A., Kalita, S., Devi, G., Bhuyan, M., 2019. Assessment of bioaccumulation and detoxification of cadmium in soil-plant-insect food chain. *Bioresour. Technol. Rep.* 7, 100242. https://doi.org/10.1007/978-981-15-6564-9_11.
- FAO., 2019. FAO yearbook. Fishery and Aquaculture Statistics 2017/FAO annuaire.
- Fedorov, S.N., Ermakova, S.P., Zvyagintseva, T.N., Stonik, V.A., 2013. Anticancer and cancer preventive properties of marine polysaccharides: some results and prospects. *Mar. Drugs* 11, 4876–4901. <https://doi.org/10.3390/md11124876>.
- Ferdouse, F., Holdt, S. L., Smith, R., Murdia, P., & Yang, Z., 2018. The global status of seaweed production, trade and utilization. *Globefish Research Programme*. 124, I.
- Ferreira, L.G., Nosedá, M.D., Gonçalves, A.G., Ducatti, D.R., Fujii, M.T., Duarte, M.E., 2012. Chemical structure of the complex pyruvylated and sulfated agaran from the red seaweed *Palisada flagellifera* (Ceramiales, Rhodophyta). *Carbohydr. Res.* 347, 83–94. <https://doi.org/10.1016/j.carres.2011.10.007>.
- Figueira, T.A., da Silva, A.J.R., Enrich-Prast, A., Yoneshigue-Valentin, Y., de Oliveira, V. P., 2020. Structural characterization of ulvan polysaccharide from cultivated and collected *Ulva fasciata* (Chlorophyta). *Adv. Biosci. Biotechnol.* 11 (5), 206–216. <https://doi.org/10.4236/abb.2020.115016>.
- Ghosh, T., Pujol, C.A., Damonte, E.B., Sinha, S., Ray, B., 2009. Sulfated xylomannans from the red seaweed *Sebdenia polydactyla*: structural features, chemical modification and antiviral activity. *Antivir. Chem. Chemother.* 19, 235–242. <https://doi.org/10.1177/095632020901900603>.
- Gomaa, H.H., Elshoubaky, G.A., 2016. Antiviral activity of sulfated polysaccharides carrageenan from some marine seaweeds. *Int. J. Curr. Pharm. Rev. Res.* 7, 34–42.
- Gonçalves, A.G., Ducatti, D.R., Duarte, M.E.R., Nosedá, M.D., 2002. Sulfated and pyruvylated disaccharide alditols obtained from a red seaweed galactan: ESIMS and NMR approaches. *Carbohydr. Res.* 337, 2443–2453. [https://doi.org/10.1016/S0008-6215\(02\)00318-X](https://doi.org/10.1016/S0008-6215(02)00318-X).
- Gorbalyeva, A.E., Baker, S.C., Baric, R., Groot, R.J.D., Drosten, C., Gulyaeva, A.A., Haagmans, B.L., Lauber, C., Leontovich, A.M., Neuman, B.W., Penzar, D., Perlman, S., Poon, L., Samborskiy, D., Sidorov, I.A., Solá, G.I., Ziebuhr, J., 2020. Severe acute respiratory syndrome-related coronavirus: the species and its viruses—a statement of the Coronavirus Study Group. *BioRxiv*. <https://doi.org/10.1101/2020.02.07.937862>.
- Goswami, L., Kumar, R.V., Manikandan, N.A., Pakshirajan, K., Pugazhenthii, G., 2019a. Anthracene biodegradation by Oleaginous *Rhodococcus opacus* for biodiesel production and its characterization. *Polycycl. Aromat. Compd.* <https://doi.org/10.1080/10406638.2017.1302971>.
- Goswami, L., Kumar, R.V., Pakshirajan, K., Pugazhenthii, G., 2019b. A novel integrated biodegradation—microfiltration system for sustainable wastewater treatment and energy recovery. *J. Hazard. Mater.* 365, 707–715. <https://doi.org/10.1016/j.jhazmat.2018.11.029>.
- Goswami, L., Pakshirajan, K., Pugazhenthii, G., 2020. Biological treatment of biomass gasification wastewater using hydrocarbonoclastic bacterium *Rhodococcus opacus* in an up-flow packed bed bioreactor with a novel waste-derived nano-biochar based bio-support material. *J. Clean. Prod.* 256, 120253. <https://doi.org/10.1016/j.jclepro.2020.120253>.
- Graf, C., Bernkop-Schnürch, A., Egyed, A., Koller, C., Prieschl-Grassauer, E., Morokutti-Kurz, M., 2018. Development of a nasal spray containing xylometazoline hydrochloride and iota-carrageenan for the symptomatic relief of nasal congestion caused by rhinitis and sinusitis. *Int. J. Gen. Med.* 11, 275. <https://doi.org/10.2147/IJGM.S167123>.
- Grassauer, A., & Prieschl-Grassauer, E., 2019. U.S. Patent No. 10,342,820. Washington, DC: U.S. Patent and Trademark Office.
- Grassauer, A., Weinmuellner, R., Meier, C., Prieschl-Grassauer, E., Unger, H., 2008. Iota-Carrageenan is a potent inhibitor of rhinovirus infection. *Virol. J.* 5, 107. <https://doi.org/10.1186/1743-422X-5-107>.
- Güven, K.C., Coban, B., Özdemiş, O., 2020. Pharmacology of marine macroalgae. *Encycl. Mar. Biotechnol.* 1, 585–615. <https://doi.org/10.1002/9781119143802.ch20>.
- Hans, N., Malik, A., Naik, S.N., 2019. Platform molecules from algae by using supercritical CO₂ and subcritical water extraction. In: Ravishanker, G.A., Ambati, R. R. (Eds.), *Handbook of Algal Technologies and Phytochemicals: Volume I Food, Health and Nutraceutical Applications*. CRC Press. <https://doi.org/10.1201/9780429054242>.
- Hardouin, K., Bedoux, G., Burlot, A.S., Donnay-Moreno, C., Bergé, J.P., Nyvall-Collén, P., Bourgougnon, N., 2016. Enzyme-assisted extraction (EAE) for the production of antiviral and antioxidant extracts from the green seaweed *Ulva armoricana* (Ulvales, Ulvophyceae). *Algal Res.* 16, 233–239. <https://doi.org/10.1016/j.algal.2016.03.013>.
- Hayashi, K., Lee, J.B., Nakano, T., Hayashi, T., 2013. Anti-influenza A virus characteristics of a fucoidan from sporophyll of *Undaria pinnatifida* in mice with normal and compromised immunity. *Microbes Infect.* 15, 302–309. <https://doi.org/10.1016/j.micinf.2012.12.004>.
- Hemilä, H., Chalker, E., 2020. Carrageenan Nasal Spray May Double the Rate of Recovery From Coronavirus and Influenza Virus Infections: Re-analysis of Randomized Trial Data. <https://doi.org/10.21203/rs.3.rs-108775/v1>.
- Hemmingson, J.A., Falshaw, R., Furneaux, R.H., Thompson, K., 2006. Structure and antiviral activity of the galactofucan sulfates extracted from *Undariapinnatifida* (Phaeophyta). *J. Appl. Phycol.* 18, 185. <https://doi.org/10.1007/s10811-006-9096-9>.
- Hidari, K.I., Takahashi, N., Arihara, M., Nagaoka, M., Morita, K., Suzuki, T., 2008. Structure and anti-dengue virus activity of sulfated polysaccharide from a marine alga. *Biochem. Biophys. Res. Commun.* 376, 91–95. <https://doi.org/10.1016/j.bbrc.2008.08.100>.
- Jahan, I., Onay, A., 2020. Potentials of plant-based substance to inhibit and probable cure for the COVID-19. *Turk. J. Biol.* 44 (3), 228. <https://doi.org/10.3906/biy-2005-114>.
- Jang, Y.J., Shin, H., Lee, M.K., Kwon, O.S., Shin, J.S., Kim, Y., Kim, M., 2020. Antiviral activity of lambda-carrageenan against influenza viruses in mice and severe acute respiratory syndrome coronavirus 2 in vitro. *BioRxiv*. <https://doi.org/10.1101/2020.08.23.255364>.
- Jiao, G., Yu, G., Zhang, J., Ewart, H.S., 2011. Chemical structures and bioactivities of sulfated polysaccharides from marine algae. *Mar. Drugs* 9, 196–223. <https://doi.org/10.3390/md9020196>.
- Jo, S., Kim, S., Shin, D.H., Kim, M.S., 2020. Inhibition of SARS-CoV 3CL protease by flavonoids. *J. Enzyme Inhib. Med. Chem.* 35 (1), 145–151. <https://doi.org/10.1080/14756366.2019.1690480>.
- Jönsson, M., Allahgholi, L., Sardari, R.R., Hreggviðsson, G.O., Nordberg Karlsson, E., 2020. Extraction and modification of macroalgal polysaccharides for current and next-generation applications. *Molecules*. 25 (4), 930. <https://doi.org/10.3390/molecules25040930>.
- Khalid, S., Abbas, M., Saeed, F., Bader-Ul-Ain, H., Suleria, H.A.R., 2018. Therapeutic potential of seaweed bioactive compounds. *Seaweed Biomater.* 1, 7–26. <https://doi.org/10.5772/intechopen.74060>.
- Kidgell, J.T., Magnusson, M., de Nys, R., Glasson, C.R., 2019. Ulvan: a systematic review of extraction, composition and function. *Algal Res.* 39, 101422. <https://doi.org/10.1016/j.algal.2019.101422>.
- Kim, J.K., Cho, M.L., Karnjanapratum, S., Shin, I.S., You, S.G., 2011. In vitro and in vivo immunomodulatory activity of sulfated polysaccharides from *Enteromorpha prolifera*. *Int. J. Biol. Macromol.* 49, 1051–1058. <https://doi.org/10.1016/j.ijbiomac.2011.08.032>.
- Kim, S.K. (Ed.), 2015. *Springer Handbook of Marine Biotechnology*. Springer. <https://doi.org/10.1007/978-3-642-53971-8>.
- Koenighofer, M., Lion, T., Bodenteich, A., Prieschl-Grassauer, E., Grassauer, A., Unger, H., Mueller, C.A., Fazekas, T., 2014. Carrageenan nasal spray in virus confirmed common cold: individual patient data analysis of two randomized controlled trials. *Multidiscip. Respir. Med.* 9, 57. <https://doi.org/10.1186/2049-6958-9-57>.
- Kushwaha, A., Rani, R., Kumar, S., Gautam, A., 2015. Heavy metal detoxification and tolerance mechanisms in plants: implications for phytoremediation. *Environ. Rev.* 24 (1), 39–51. <https://doi.org/10.1139/er-2015-0010>.
- Kushwaha, A., Rani, R., Kumar, S., 2017. Mechanism of soil-metal-microbe interactions and their implication on microbial bioremediation and phytoremediation. In: Kumar, P., Gujjar, B.R., Govil, J.N. (Eds.), *Environmental Science and Engineering, 8*. Studium Press LLC, U.S.A., p. 462.
- Kushwaha, A., Rani, R., Patra, J.K., 2020. Adsorption kinetics and molecular interactions of lead [Pb (II)] with natural clay and humic acid. *Int. J. Environ. Sci. Technol.* 17 (3), 1325–1336. <https://doi.org/10.1007/s13762-019-02411-6>.
- Kwon, P.S., Oh, H., Kwon, S.J., Jin, W., Zhang, F., Fraser, K., Hong, J.J., Linhardt, R.J., Dordick, J.S., 2020. Sulfated polysaccharides effectively inhibit SARS-CoV-2 in vitro. *Cell Discov.* 6 (1), 1–4. <https://doi.org/10.1038/s41421-020-00192-8>.
- Lee, J.B., Hayashi, K., Maeda, M., Hayashi, T., 2004. Antihyperthermic activities of sulfated polysaccharides from green algae. *Planta Med.* 70, 813–817. <https://doi.org/10.1055/s-2004-827228>.

- Lee, J.B., Koizumi, S., Hayashi, K., Hayashi, T., 2010. Structure of rhamnan sulfate from the green alga *Monostroma nitidum* and its anti-herpetic effect. *Carbohydr. Polym.* 81, 572–577. <https://doi.org/10.1016/j.carbpol.2010.03.014>.
- Lee, J.B., Takeshita, A., Hayashi, K., Hayashi, T., 2011. Structures and antiviral activities of polysaccharides from *Sargassum trichophyllum*. *Carbohydr. Polym.* 86, 995–999. <https://doi.org/10.1016/j.carbpol.2011.05.059>.
- Lee, W.K., Lim, Y.Y., Leow, A.T.C., Namasivayam, P., Abdullah, J.O., Ho, C.L., 2017a. Biosynthesis of agar in red seaweeds: a review. *Carbohydr. Polym.* 164, 23–30. <https://doi.org/10.1016/j.carbpol.2017.01.078>.
- Lee, Y.E., Kim, H., Seo, C., Park, T., Lee, K.B., Yoo, S.Y., Hong, S.C., Kim, J.T., Lee, J., 2017b. Marine polysaccharides: therapeutic efficacy and biomedical applications. *Arch. Pharm. Res.* 40 (9), 1006–1020. <https://doi.org/10.1007/s12272-017-0958-2>.
- Leibbrandt, A., Meier, C., König-Schuster, M., Weimüller, R., Kalthoff, D., Pflugfelder, B., Graf, P., Frank-Gehrke, B., Beer, M., Fazekas, T., Unger, H., Prieschl-Grassauer, E., Grassauer, A., 2010. Iota-carrageenan is a potent inhibitor of influenza A virus infection. *PLoS One* 5 (12), e14320. <https://doi.org/10.1371/journal.pone.0014320>.
- Li, H., Mao, W., Hou, Y., Gao, Y., Qi, X., Zhao, C., Chen, Y., Chen, Y., Li, N., Wang, C., 2012. Preparation, structure and anticoagulant activity of a low molecular weight fraction produced by mild acid hydrolysis of sulfated rhamnan from *Monostroma latissimum*. *Bioresour. Technol.* 114, 414–418. <https://doi.org/10.1016/j.biortech.2012.03.025>.
- Liu, Z., Gao, T., Yang, Y., Meng, F., Zhan, F., Jiang, Q., Sun, X., 2019. Anti-cancer activity of porphyrin and carrageenan from red seaweeds. *Molecules* 24 (23), 4286. <https://doi.org/10.3390/molecules24234286>.
- Lopes, N., Ray, S., Espada, S.F., Bomfim, W.A., Ray, B., Faccin-Galhardi, L.C., Linhares, R. E.C., Nozawa, C., 2017. Green seaweed *Enteromorpha compressa* (Chlorophyta, Ulvaceae) derived sulphated polysaccharides inhibit herpes simplex virus. *Int. J. Biol. Macromol.* 102, 605–612. <https://doi.org/10.1016/j.ijbiomac.2017.04.043>.
- Makarenkova, I.D., Deriabin, P.G., L'vov, D.K., Zviagintseva, T.N., Besednova, N.N., 2010. Antiviral activity of sulfated polysaccharide from the brown alga *Laminaria japonica* against avian influenza A (H5N1) virus infection in the cultured cells. *Vopr. Virusol.* 55, 41–45.
- Malagoli, B.G., Cardozo, F.T., Gomes, J.H.S., Ferraz, V.P., Simões, C.M., Braga, F.C., 2014. Chemical characterization and antihpetic activity of sulfated polysaccharides from *Lithothamnion muelleri*. *Int. J. Biol. Macromol.* 66, 332–337. <https://doi.org/10.1016/j.ijbiomac.2014.02.053>.
- Mandal, P., Mateu, C.G., Chattopadhyay, K., Pujol, C.A., Damonte, E.B., Ray, B., 2007. Structural features and antiviral activity of sulphated fucans from the brown seaweed *Cystoseira indica*. *Antivir. Chem. Chemother.* 18, 153–162. <https://doi.org/10.1177/095632020701800305>.
- Mandal, P., Pujol, C.A., Carlucci, M.J., Chattopadhyay, K., Damonte, E.B., Ray, B., 2008. Anti-herpetic activity of a sulfated xylomannan from *Scinaia hatei*. *Phytochemistry* 69, 2193–2199. <https://doi.org/10.1016/j.phytochem.2008.05.004>.
- Mazumber, S., Ghosal, P.K., Pujol, C.A., Carlucci, M.J., Damonte, E.B., Ray, B., 2002. Isolation, chemical investigation and antiviral activity of polysaccharides from *Gracilaria corticata* (Gracilariaceae, Rhodophyta). *Int. J. Biol. Macromol.* 31, 87–95. [https://doi.org/10.1016/S0141-8130\(02\)00070-3](https://doi.org/10.1016/S0141-8130(02)00070-3).
- Mercur, J., Schelhaas, M., Helenius, A., 2010. Virus entry by endocytosis. *Annu. Rev. Biochem.* 79, 803–833. <https://doi.org/10.1146/annurev-biochem-060208-104626>.
- Morán-Santibáñez, K., Cruz-Suárez, L.E., Riquelme-Marie, D., Robledo, D., Freile-Pelegrín, Y., Peña-Hernández, M.A., Rodríguez-Padilla, C., Trejo-Avila, L.M., 2016. Synergistic effects of sulfated polysaccharides from Mexican seaweeds against measles virus. *Biomed. Res. Int.* 2016. <https://doi.org/10.1155/2016/8502123>.
- Morokutti-Kurz, M., Graf, C., Prieschl-Grassauer, E., 2017. Amylmetacresol/2, 4-dichlorobenzyl alcohol, hexylresorcinol, or carrageenan lozenges as active treatments for sore throat. *Int. J. Gen. Med.* 10, 53. <https://doi.org/10.2147/IJGM.S120665>.
- Muthukumar, J., Chidambaram, R., Sukumaran, S., 2020. Sulfated polysaccharides and its commercial applications in food industries—a review. *J. Food Sci. Technol.* 1–14. <https://doi.org/10.1007/s13197-020-04837-0>.
- Ortiz-Tena, J.G., Schieder, D., Sieber, V., 2017. Carrageenan and more: biorefinery approaches with special reference to the processing of *Kappaphycus*. In: Hurtado, A., Critchley, A., Neish, I. (Eds.), *Tropical Seaweed Farming Trends, Problems and Opportunities*, Developments in Applied Phycology, vol. 9. Springer, Cham, pp. 155–164. http://doi-org-443.webvpn.fjmu.edu.cn/10.1007/978-3-319-6349-8-2_10.
- Ponce, N.M., Pujol, C.A., Damonte, E.B., Flores, M.L., Stortz, C.A., 2003. Fucoidans from the brown seaweed *Adenocystis utricularis*: extraction methods, antiviral activity and structural studies. *Carbohydr. Res.* 338, 153–165. [https://doi.org/10.1016/S0008-6215\(02\)00403-2](https://doi.org/10.1016/S0008-6215(02)00403-2).
- Pujol, C.A., Estevez, J.M., Carlucci, M.J., Ciancia, M., Cerezo, A.S., Damonte, E.B., 2002. Novel DL-galactan hybrids from the red seaweed *Gymnogongrus torulosus* are potent inhibitors of herpes simplex virus and dengue virus. *Antivir. Chem. Chemother.* 13, 83–89. <https://doi.org/10.1177/095632020201300202>.
- Queiroz, K.C.S., Medeiros, V.P., Queiroz, L.S., Abreu, L.R.D., Rocha, H.A.O., Ferreira, C. V., Jucá, M.B., Aoyama, H., Leite, E.L., 2008. Inhibition of reverse transcriptase activity of HIV by polysaccharides of brown algae. *Biomed. Pharmacother.* 62, 303–307. <https://doi.org/10.1016/j.biopha.2008.03.006>.
- Rabanal, M., Ponce, N.M., Navarro, D.A., Gómez, R.M., Stortz, C.A., 2014. The system of fucoidans from the brown seaweed *Dictyota dichotoma*: chemical analysis and antiviral activity. *Carbohydr. Polym.* 101, 804–811. <https://doi.org/10.1016/j.carbpol.2013.10.019>.
- Ray, B., 2006. Polysaccharides from *Enteromorpha compressa*: isolation, purification and structural features. *Carbohydr. Polym.* 66, 408–416. <https://doi.org/10.1016/j.carbpol.2006.03.027>.
- Rhein-Knudsen, N., Ale, M.T., Meyer, A.S., 2015. Seaweed hydrocolloid production: an update on enzyme assisted extraction and modification technologies. *Mar. Drugs* 13, 3340–3359. <https://doi.org/10.3390/md13063340>.
- Roy, S., Bhattacharyya, P., 2020. Possible role of traditional medicinal plant *Neem* (*Azadirachta indica*) for the management of COVID-19 infection. *Int. J. Res. Pharm. Sci.* 11, 122–125. <https://doi.org/10.26452/ijrps.v11i1SPL1.2256>.
- Rudke, A.R., de Andrade, C.J., Ferreira, S.R.S., 2020. *Kappaphycus alvarezii* macroalgae: an unexplored and valuable biomass for green biorefinery conversion. *Trends Food Sci. Technol.* 103, 214–224. <https://doi.org/10.1016/j.tifs.2020.07.018>.
- Ruocco, N., Costantini, S., Guariniello, S., Costantini, M., 2016. Polysaccharides from the marine environment with pharmacological, cosmeceutical and nutraceutical potential. *Molecules* 21 (5), 551. <https://doi.org/10.3390/molecules21050551>.
- Sepúlveda-Crespo, D., Ceña-Díez, R., Jiménez, J.L., Ángeles Muñoz-Fernández, M., 2017. Mechanistic studies of viral entry: an overview of dendrimer-based microbicides as entry inhibitors against both HIV and HSV-2 overlapped infections. *Med. Res. Rev.* 37 (1), 149–179. <https://doi.org/10.1002/med.21405>.
- Shree, P., Mishra, P., Selvaraj, C., Singh, S.K., Chaube, R., Garg, N., Tripathi, Y.B., 2020. Targeting COVID-19 (SARS-CoV-2) main protease through active phytochemicals of ayurvedic medicinal plants—*Withania somnifera* (Ashwagandha), *Tinospora cordifolia* (Giloy) and *Ocimum sanctum* (Tulsi)—a molecular docking study. *J. Biomol. Struct. Dyn.* 1–14. <https://doi.org/10.1080/07391102.2020.1810778>.
- Singh, P.K., Kushwaha, A., Hans, N., Gautam, A., Rani, R., 2019. Evaluation of the cytotoxicity and interaction of lead with lead resistant bacterium *Acinetobacter junii* Pb1. *Braz. J. Microbiol.* 50 (1), 223–230. <https://doi.org/10.1007/s42770-019-00041-1>.
- Song, S., Peng, H., Wang, Q., Liu, Z., Dong, X., Wen, C., Ai, C., Zhang, Y., Wang, Z., Zhu, B., 2020. Inhibitory activities of marine sulfated polysaccharides against SARS-CoV-2. *Food Funct.* 11 (9), 7415–7420. <https://doi.org/10.1039/D0FO02017F>.
- Sozzani, S., Bosio, D., Scarsi, M., Tinca, A., 2010. Type I interferons in systemic autoimmunity. *J. Autoimmun.* 43, 196–203. <https://doi.org/10.3109/08916930903510872>.
- Talarico, L.B., Damonte, E.B., 2007. Interference in dengue virus adsorption and uncoating by carrageenans. *Virology* 363, 473–485. <https://doi.org/10.1016/j.virol.2007.01.043>.
- Talarico, L.B., Zibetti, R.G., Faria, P.C., Scolaro, L.A., Duarte, M.E., Noseda, M.D., Pujol, C.A., Damonte, E.B., 2004. Anti-herpes simplex virus activity of sulfated galactans from the red seaweeds *Gymnogongrus griffithsiae* and *Cryptomenia crenulata*. *Int. J. Biol. Macromol.* 34, 63–71. <https://doi.org/10.1016/j.ijbiomac.2004.03.002>.
- Tang, F., Chen, F., Li, F., 2013. Preparation and potential in vivo anti-influenza virus activity of low molecular-weight κ-carrageenans and their derivatives. *J. Appl. Polym. Sci.* 127, 2110–2115. <https://doi.org/10.1002/app.37502>.
- Thuy, T.T.T., Ly, B.M., Van, T.T.T., Quang, N.V., Tu, H.C., Zheng, Y., Seguin-Devaux, C., Mi, B., Ai, U., 2015. Anti-HIV activity of fucoidans from three brown seaweed species. *Carbohydr. Polym.* 115, 122–128. <https://doi.org/10.1016/j.carbpol.2014.08.068>.
- Trinchero, J., Ponce, N.M., Córdoba, O.L., Flores, M.L., Pampuro, S., Stortz, C.A., Salomón, H., Turk, G., 2009. Antiretroviral activity of fucoidans extracted from the brown seaweed *Adenocystis utricularis*. *Phytother. Res.* 23, 707–712. <https://doi.org/10.1002/ptr.2723>.
- Usov, A.I., 2011. Polysaccharides of the red algae. In: Horton, D. (Ed.), *Advances in Carbohydrate Chemistry and Biochemistry*. Academic Press, pp. 115–217. <https://doi.org/10.1016/B978-0-12-385520-6.00004-2>.
- Wahedi, H.M., Ahmad, S., Abbasi, S.W., 2020. Stilbene-based natural compounds as promising drug candidates against COVID-19. *J. Biomol. Struct. Dyn.* 1–10. <https://doi.org/10.1080/07391102.2020.1762743>.
- Wang, S., Wang, W., Hao, C., Yunjia, Y., Qin, L., He, M., Mao, W., 2018. Antiviral activity against enterovirus 71 of sulfated rhamnan isolated from the green alga *Monostroma latissimum*. *Carbohydr. Polym.* 200, 43–53. <https://doi.org/10.1016/j.carbpol.2018.07.067>.
- Wang, W., Zhang, P., Hao, C., Zhang, X.E., Cui, Z.Q., Guan, H.S., 2011. In vitro inhibitory effect of carrageenan oligosaccharide on influenza A H1N1 virus. *Antivir. Res.* 92, 237–246. <https://doi.org/10.1016/j.antiviral.2011.08.010>.
- Wang, W., Wang, S.X., Guan, H.S., 2012. The antiviral activities and mechanisms of marine polysaccharides: an overview. *Mar. Drugs* 10, 2795–2816. <https://doi.org/10.3390/md10122795>.
- Wang, Y., Di, Y., Ye, J., Wei, W., 2020. Study on the public psychological states and its related factors during the outbreak of coronavirus disease 2019 (COVID-19) in some regions of China. *Psychol. Health Med.* 1–10. <https://doi.org/10.1080/13548506.2020.1746817>.
- Wijesekara, I., Pangestuti, R., Kim, S.K., 2011. Biological activities and potential health benefits of sulfated polysaccharides derived from marine algae. *Carbohydr. Polym.* 84, 14–21. <https://doi.org/10.1016/j.carbpol.2010.10.062>.
- Worldometer [Internet]. COVID-19 Coronavirus Pandemic, 2020. Available from: <http://www.worldometers.info/coronavirus/> (Accessed 20 November 2020).
- Xian, Y., Zhang, J., Bian, Z., Zhou, H., Zhang, Z., Lin, Z., Xu, H., 2020. Bioactive natural compounds against human coronaviruses: a review and perspective. *Acta Pharm. Sin.* B 10, 1163–1174. <https://doi.org/10.1016/j.apsb.2020.06.002>.
- Xianliang, X., Hua, D., Meiyu, G., Pingfang, L., Yingxia, L., Huashi, G., 2000. Studies of the anti-AIDS effects of marine polysaccharide drug 911 and its related mechanisms of action. *Chin. J. Mar. Drugs* 19 (6), 4–8.
- Xu, X., Chen, P., Wang, J., Feng, J., Zhou, H., Li, X., Zhong, W., Hao, P., 2020. Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci. China Life Sci.* 63, 457–460. <https://doi.org/10.1007/s11427-020-1637-5>.

- Yamada, T., Ogamo, A., Saito, T., Uchiyama, H., Nakagawa, Y., 2000. Preparation of O-acylated low-molecular-weight carrageenans with potent anti-HIV activity and low anticoagulant effect. *Carbohydr. Polym.* 41, 115–120. [https://doi.org/10.1016/S0144-8617\(99\)00083-1](https://doi.org/10.1016/S0144-8617(99)00083-1).
- Yu, G., Hu, Y., Yang, B., Zhao, X., Wang, P., Ji, G., Wu, J., Guan, H., 2010. Extraction, isolation and structural characterization of polysaccharides from a red alga *Gloiopeltis furcata*. *J. Ocean Univ. China* 9, 193–197. <https://doi.org/10.1007/s11802-010-0193-7>.
- Yuan, H., Song, J., Li, X., Li, N., Dai, J., 2006. Immunomodulation and antitumor activity of κ-carrageenan oligosaccharides. *Cancer Lett.* 243, 228–234. <https://doi.org/10.1016/j.canlet.2005.11.032>.
- Yuguchi, Y., Bui, L.M., Takebe, S., Suzuki, S., Nakajima, N., Kitamura, S., Thanh, T.T.T., 2016. Primary structure, conformation in aqueous solution, and intestinal immunomodulating activity of fucoidan from two brown seaweed species *Sargassum crassifolium* and *Padina australis*. *Carbohydr. Polym.* 147, 69–78. <https://doi.org/10.1016/j.carbpol.2016.03.101>.
- Zhang, W., Oda, T., Yu, Q., Jin, J.O., 2015. Fucoidan from *Macrocystis pyrifera* has powerful immune-modulatory effects compared to three other fucoidans. *Mar. Drugs* 13, 1084–1104. <https://doi.org/10.3390/md13031084>.
- Zhu, B., Ni, F., Sun, Y., Zhu, X., Yin, H., Yao, Z., Du, Y., 2018. Insight into carrageenases: major review of sources, category, property, purification method, structure, and applications. *Crit. Rev. Biotechnol.* 38 (8), 1261–1276. <https://doi.org/10.1080/07388551.2018.1472550>.
- Zhu, N., Zhang, D., Wang, W., Li, X., Yang, B., Song, J., Zhao, X., Huang, B., Shi, W., Lu, R., Niu, P., Zhan, F., Ma, X., Wang, D., Xu, W., Wu, G., Gao, G.F., Tan, W., 2020. A novel coronavirus from patients with pneumonia in China, 2019. *N. Engl. J. Med.* 382, 727–733. <https://doi.org/10.1056/NEJMoa2001017>.