

RESEARCH ARTICLE

Open Access



The parameters of gait analysis related to ambulatory and balance functions in hemiplegic stroke patients: a gait analysis study

Min Cheol Chang^{1†}, Byung Joo Lee^{2†}, Na-Young Joo³ and Donghwi Park^{3*}

Abstract

Background: Ambulatory and balance functions are important for maintaining general health in humans. Gait analysis allows clinicians and researchers to identify the parameters to be focused on when assessing balance and ambulatory functions. In this study, we performed gait analysis with pressure sensors to identify the gait-analysis parameters related to balance and ambulatory functions in hemiplegic stroke patients.

Methods: We retrospectively reviewed the medical records of 102 patients with hemiplegic stroke who underwent gait analysis. Correlations between various temporospatial parameters in the gait analysis and the motor and balance functions assessed using functional ambulation category, modified Barthel index, and Berg balance scale were analyzed.

Results: Gait speed/height and the lower-limb stance-phase time/height were the only temporal and spatial parameters, respectively, that showed a statistical correlation with motor and balance functions.

Conclusions: Measurements of walking speed and stance-phase time of the unaffected lower limb can allow clinicians to easily assess the ambulatory and balance functions of hemiplegic stroke patients. Rehabilitative treatment focusing on increasing gait speed and shortening the stance-phase time of the unaffected side may improve the ambulatory and balance functions in these patients.

Keywords: Gait analysis, Gait speed, Hemiplegia, Stroke, Gait disturbance

Background

Ambulatory and balance functions, which are basic functions necessary for independence in daily activities, are also important for mobility and maintenance of general health in humans [1]. Stroke, one of the most common causes of impaired ambulatory and balance functions

[2], affects patients' activities of daily living and ultimately limits their participation in community-based activities [3, 4]. Moreover, the stroke-induced impairments in ambulatory and balance functions frequently result in falls and fall-related injuries [5]. Thus, improvement of ambulatory and balance functions is one of the primary goals of stroke rehabilitation programs.

Gait analysis has been widely used in studies assessing ambulatory and balance function during gait.[2] Gait analyses can yield quantitative data for several temporospatial gait-related parameters, including gait speed, stride or step length, stance time, and the angle of each

* Correspondence: bdome@hanmail.net

[†]Min Cheol Chang and Byung Joo Lee was contributed equally as first authors

³Department of Physical Medicine and Rehabilitation, Ulsan University Hospital, University of Ulsan College of Medicine, 877 Bangeojin sunhwando-ro, Dong-gu, 44033 Ulsan, Republic of Korea

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

joint [6]. Knowledge of the elements related to ambulatory and balance functions can help clinicians identify the relevant factors during rehabilitative treatment of stroke patients and assist clinicians and researchers in determining the gait-related parameters that should be focused on when assessing balance and ambulatory functions. Gait dysfunctions in stroke patients are usually attributable to cognitive, motor, and sensory impairments [7–9], and may manifest as decreased walking speed, altered proportions of temporal phases, increased variability in the stride duration, and increased asymmetries in the single-stance phase [10, 11]. However, the multicollinearity of spatiotemporal data has made interpretation of spatiotemporal parameters difficult and led to uncertainty about the relevant gait-analysis parameters for assessment of poststroke gait. Moreover, although gait analysis can be used to simply measure the ambulatory and balance function of patients, the complexity and vastness of gait-analysis data precludes such an approach in actual clinical practice.

Therefore, in the current study, we used a gait-analysis tool with pressure sensors to investigate the parameters related to balance and ambulatory functions in hemiplegic stroke patients.

Methods

Participants

Our study was approved by the Ethics Committee (2020-07-020) and was conducted in accordance with the principles of the Declaration of Helsinki for human experiments. All methods followed the relevant guidelines and regulations. We retrospectively reviewed the medical records of patients with stroke who were admitted to our hospital between January 2017 and August 2020.

The inclusion criteria were as follows: (1) adult patients (age ≥ 20 years), (2) hemiplegia due to stroke, (3) ability to walk independently, (4) completed both gait analysis and clinical assessment of motor and gait functions, including modified Barthel index (MBI) evaluation, manual muscle test (MMT) of both lower extremities, Berg balance scale (BBS) measurements, and functional ambulation category (FAC) assessments. The exclusion criteria were as follows: (1) > 1 -week interval between assessment of gait and motor functions and gait analysis, and (2) a history of other neurologic or musculoskeletal disorders that could affect the results of this study.

Clinical assessment

Patients' clinical records, including demographic data, clinical diagnosis, disease duration, Mini Mental Status Examination (MMSE) score, [12] MBI sub-scores for ambulation and stair-climbing [13], MMT score [14, 15], FAC [16], and BBS score [17], were evaluated. We also evaluated the spasticity of the hemi-side ankle plantar

flexor muscle by using the modified Ashworth scale [18]. Using the MMT scores for both lower extremities, we calculated the total score (both hip flexor MMT + hip extensor MMT + knee extensor MMT + knee flexor MMT + ankle dorsiflexor MMT + ankle plantar flexor MMT; total score, 60), and the total score for the anti-gravity muscles in the lower extremities (both hip extensor MMT + knee extensor MMT + ankle plantar flexor MMT; total score, 30) [14, 15].

All clinical assessments were performed by the patient's main physical therapist within 7 days of gait analysis. All therapists who performed the assessments were blinded to the results of gait analysis.

Gait analysis

A computerized gait-analysis system (Walkway MG-1000; Anima, Japan) was used for gait analysis [19]. This system measures the temporospatial parameters of gait by analyzing on/off signals between the patient's foot and the surface of the sensors at a sampling frequency of 100 Hz [19]. The length and width of the system's walkway are 4.8 m and 0.82 m, respectively. During gait, data were obtained and processed using software embedded in the system [19].

Experiment procedure

Participants wearing short pants were asked to get on a walkway and walk barefoot along a 12-m straight line, including 3.5 m in the front and 3.5 m beyond the end of walking path. Each participant performed one trial at a subjectively determined comfortable speed. A physical therapist with more than 10 years of experience performed gait analysis for all patients included in this study.

Parameters of gait analysis

Measurements for the temporal parameters stance phase, swing phase, double stance phase, and stride were obtained in seconds. In addition, the duration of the stance phase during the total gait cycle, duration of the swing phase during the total gait cycle, and the duration of the double stance phase during the total gait cycle in both lower limbs were expressed as percentages. Measurements for the spatial parameters stride length, step length, and step width were obtained in centimeters. We

Table 1 Clinical and demographic characteristics of included patients with hemiplegic stroke

	Patients
Age (years)	59.44 \pm 13.72
Sex (M:F)	68:34
Etiology (hemorrhagic stroke : infarction stroke)	9:93
Side (Left:Right)	43:59
M:F Male:female	

Table 2 The temporo-spatial parameters of gait in patients with hemiplegic stroke

	Patients	
Height (cm)	165.04±7.61	
Weight (kg)	65.16±11.92	
Disease duration (days)	29.20±99.45	
MMSE	25.22±6.11	
Hemi-side ankle PF spasticity (MAS)	0.39±0.54	
Total BBS score	44.69±11.62	
Total MBI score	80.36±17.32	
MBI ambulation	10.86±5.14	
MBI stair climbing	3.31±4.19	
FAC	2.61±1.35	
Gait speed (cm/s)	48.62±16.75	
Cadence (step/min)	79.11±15.28	
Double stance time (s)	0.31±0.18	
	Affected side	Unaffected side
MMT hip flexor	4.35±0.56	5.0±0.0
MMT hip extensor	4.37±0.53	5.0±0.0
MMT Knee flexor	4.28±0.62	5.0±0.0
MMT knee extensor	4.29±0.61	5.0±0.0
MMT ankle DF	4.32±0.77	5.0±0.0
MMT ankle PF	4.33±0.75	5.0±0.0
MMT sum	25.96±3.56	25.0±0.0
Antigravity MMT sum	13.0±1.76	15.0±0.0
GSR	3.03±1.43	3.29±2.46
Stride length (cm)	72.37±17.85	72.28±19.39
Step length (cm)	36.38±9.34	36.17±10.39
Step width (cm)	12.67±4.61	12.87±4.40
Gait angle (°)	21.06±10.38	21.88±11.51
Toe angle (°)	9.38±7.79	10.50±6.90
Stance phase time (s)	2.23±11.48	1.13±0.40
Swing phase time (s)	0.47±0.13	0.42±0.17

MBI Modified Barthel's Index, MAS Modified Ashworth scale, MMSE Mini Mental Status Examination, LE Lower Extremity, MMT Manual Muscle Test, sum Summation, GSR Gait stability ratio, DF Dorsiflexor, PF Plantar flexor, FAC Functional ambulation classification, BBS Berg balance scale

also calculated the gait stability ratio (i.e., steps/m) for both lower limbs [20]. Additionally, gait angle (degrees), toe out angle (degrees), gait speed (cm/s), and cadence (step) of each lower limb were measured. The gait angle is the angle between the line of progression and the foot axis. It is zero when the foot axis is parallel to the line of progression, and positive when the foot axis points lateral to the line of progression. All measurement values were represented as mean ± standard deviation.

To compensate for variations in patient height, we calculated gait speed/height, gait speed/height², stride length/height, stride length/height², step length/height, step length/height², step width/height, step width/height², stride time/height, and stride time/height².

Statistical analysis

To identify the correlations between gait-analysis parameters and the results indicating motor function (FAC, ambulation sub-score of MBI, and BBS score), multiple linear regression analysis was used. To identify the variables affected by multicollinearity and the strengths of the correlations, testing for multicollinearity with variance inflation factors (VIFs) was performed. Multicollinearity was considered to be present when VIF was higher than 5 to 10. Multiple linear regression tests were performed after discarding parameters showing multicollinearity. Statistical analyses were performed using Statistical Package for the Social Sciences for Windows and R package for Windows (version 2.15.2; R Foundation for Statistical Computing, Vienna, Austria).

Results

Patient characteristics

A total of 102 patients with stroke (68 males and 34 females, 18–91 years of age, 148–185 cm in height, and 43–98 kg in weight) were investigated in this study. The results of the clinical assessments and gait analysis are presented in Tables 1 and 2. Mean height (cm), body weight (kg), disease duration (days), total MBI score, and MMSE were 165.04 ± 7.61 cm, 65.16 ± 11.92 kg, 29.20 ± 99.45 days, 80.36 ± 17.32, and 25.22 ± 6.11, respectively (Table 3).

Table 3 Multiple linear regression analysis for assessing temporo-spatial parameters of gait in predicting of balance and gait function

Dependent variable	Independent variables	R ²	Beta coefficient	Standard error	Odd ratio (95% CI)	P value
BBS	Gait speed/height	0.227	0.476	0.130	0.308 ~ 0.826	<0.001
	Unaffected stance time/height	0.220	-0.470	7.832	-49.22~-17.93	<0.001
FAC	Gait speed/height	0.354	0.595	0.011	0.058 ~ 0.100	<0.001
	Unaffected stance time/height	0.199	-0.446	0.517	-3.597~-1.547	<0.001
Ambulation MBI	Gait speed/height	0.153	0.391	0.047	0.105 ~ 0.290	<0.001
	Unaffected stance time/height	0.136	-0.369	2.042	-12.153~-4.052	<0.001
Stair climbing MBI	Gait speed/height	0.181	0.426	0.037	0.101 ~ 0.249	<0.001
	Unaffected stance time/height	0.096	-0.310	1.700	-8.924~-2.178	0.001

BBS Berg Balance Scale, FAC Functional Ambulation Category, MBI Modified Barthel Index, MMT Manual Muscle Test, Av Average




Fig. 1 (See legend on next page.)

(See figure on previous page.)

Fig. 1 a-b Mean of the "actual vs. predicted plot" and the residual plot between the Berg balance scale (BBS) score and spatial parameters of gait (gait speed/height). **c-d** Mean of the "actual vs. predicted plot" and the residual plot between the ambulation sub-score of the modified Barthel index (MBI) and spatial parameters of gait (gait speed/height). **e-f** Mean of the "actual vs. predicted plot" and the residual plot between the stair-climbing sub-score of the MBI and spatial parameters of gait (gait speed/height). **g-h** Mean of the "actual vs predicted plot" and the residual plot between the functional ambulation category (FAC) and spatial parameters of gait (gait speed/height)

Among them, 93 patients had ischemic stroke lesions, while the remaining 9 had hemorrhagic stroke lesions. The average total MMT score for the lower-extremity muscles was 55.72 ± 4.36 , while that for the antigravity muscles of both lower extremities was 27.44 ± 2.76 .

Correlation between Berg balance scale and gait-analysis parameters

Among the spatial and temporal parameters of gait analysis, only gait speed/height and the unaffected-side stance-phase time/height, respectively, showed a statistically significant correlation with BBS scores ($R^2 = 0.227$, $p < 0.001$; and $R^2 = 0.220$, $p < 0.001$, respectively) (Figs. 1 and 2).

Correlation between the ambulation sub-score of the modified Barthel index and gait-analysis parameters

Similar to the findings for BSS scores, gait speed/height and the unaffected-side stance-phase time/height were the only spatial and temporal parameters of gait analysis, respectively, that showed statistically significant correlations with the MBI ambulation sub-score ($R^2 = 0.153$, $p < 0.001$; and $R^2 = 0.136$, $p < 0.001$, respectively) (Figs. 1 and 2).

Correlation between the stair climbing sub-score of modified Barthel index and gait-analysis parameters

In the analyses based on the stair climbing sub-score of MBI, gait speed/height and the unaffected-side stance-phase time/height were again the only spatial and temporal gait-analysis parameters, respectively, that showed statistically significant correlations with the stair climbing sub-score ($R^2 = 0.181$, $p < 0.001$; and $R^2 = 0.096$, $p < 0.001$, respectively) (Figs. 1 and 2).

Correlation between functional ambulation category and gait-analysis parameters

Consistent with the findings of the previous analyses, gait speed/height and the unaffected-side stance-phase time/height were the only spatial and temporal gait-analysis parameters, respectively, that showed statistically significant correlations with FAC ($R^2 = 0.354$, $p < 0.001$; and $R^2 = 0.199$, $p < 0.001$, respectively) (Figs. 1 and 2).

Discussion

In this study, we analyzed various temporospatial parameters of gait analysis in patients with stroke and investigated the relationships between these parameters and parameters indicating ambulatory and balance functions.

Our results showed that only gait speed/height and the unaffected-side lower-limb stance phase time/height were significantly correlated with the FAC and BBS and MBI scores.

Since gait speed is a reliable measure of ambulatory ability in patients with stroke, it is commonly used by clinicians and researchers [21]. Perry et al. [22] reported that gait speed differed significantly depending on the degree of impairment in ambulatory function: indoor-level ambulators (0.4 m/s), limited outdoor-level ambulators (0.40–0.80 m/s), and unlimited outdoor-level ambulators (> 0.8 m/s). Likewise, in our study, gait speed was significantly correlated with ambulatory and balance functions in all our patients. In addition to reflecting ambulatory and balance functions, gait speed evaluations offer the advantage of quick and easy measurements that can be performed without any equipment. Since the outcome of gait speed is related to mortality, poor quality of life, and physical and cognitive functional decline, clinicians and researchers can obtain information regarding the patient's status and predict the functional prognosis by measuring gait speed [23, 24].

In our study, the unaffected-side stance-phase time was inversely correlated with the patients' ambulatory and balance functions. A short stance-phase time of the lower limb on the unaffected side indicates a relatively longer swing-phase time [25]. In such patients, the lower limb on the affected side has to endure a longer stance phase, which is correlated with the limb's motor function. However, the MMT scores for the affected side did not show correlations with any of the gait-analysis parameters. This finding can be interpreted as follows: Although muscle strength is an important factor influencing the support time of the lower limb on the affected side, other factors, such as proprioception or a subjective fear of falling, might also have a significant impact on ambulatory and balance functions [25, 26].

The results of our study can help identify the components that should be focused on during training to improve ambulatory and balance functions in hemiplegic stroke patients. To achieve this improvement, clinicians should formulate a strategy that enhances the walking speed and shortens the stance-phase time of the unaffected-side lower limb, thereby elongating the stance-phase time of the affected-side lower limb. Thus, a high walking speed and short stance-phase time of the unaffected-side lower limb can indicate good ambulatory




Fig. 2 (See legend on next page.)

(See figure on previous page.)

Fig. 2 a-b Mean of the "actual vs. predicted plot" and the residual plot between the Berg balance scale (BBS) score and temporal parameters of gait (unaffected stance-phase time/height). **c-d** Mean of the "actual vs. predicted plot" and the residual plot between the ambulation sub-score of the modified Barthel index (MBI) and temporal parameters of gait (unaffected stance phase time/height). **e-f** Mean of the "actual vs. predicted plot" and the residual plot between the stair-climbing sub-score of the MBI and temporal parameters of gait (unaffected stance phase time/height). **g-h** Mean of the "actual vs. predicted plot" and the residual plot between the functional ambulation category (FAC) and temporal parameters of gait (unaffected stance phase time/height)

and balance functions in hemiplegic stroke patients, and measurements of the walking speed and the stance-phase time of the unaffected-side lower limb can facilitate assessments of the ambulatory and balance functions in these patients. Rehabilitative treatment that focused on increasing the gait speed and shortening the stance-phase time of the unaffected side may help improve patients' ambulatory and balance functions.

To our knowledge, this study is the first to evaluate the gait-analysis parameters related to ambulatory and balance functions in stroke patients. However, the study had some limitations. First, severely impaired patients were excluded from this study. Since the participants were requested to complete gait analysis, those with severe motor or balance impairment could not participate. Second, the patients underwent gait analysis without using any assistance, such as an ankle foot orthosis or a cane. Thus, if the patients' gait function improved with such assistance, our results may not fully reflect the actual balance or gait functions. Future studies that address this limitation are warranted. Third, only the ambulation and stair-climbing sub-scores of MBI and the FAC were used in this study. However, for better evaluation, additional studies using various parameters that can adequately represent the patient's ambulatory or balance function are needed. Fourth, we used only pressure sensor-based gait analysis without obtaining anthropometric measures. More accurate analysis of gait function would have been possible by using auxiliary data such as anthropometric measures along with pressure sensor-based measurements. However, our gait-analysis system based on pressure sensors allowed easy assessments, and the gait-analysis parameters assessed using pressure sensors adequately represented the ambulatory and balance functions of stroke patients with hemiplegia. The purpose of this study was to use pressure sensor-based gait analysis, which is easy to perform, to identify parameters that could more easily represent the patient's ambulatory and balance function. However, more accurate data can be obtained if additional data such as anthropometric measures are used in future studies. Lastly, while the results showed significant correlations, the correlation coefficients were relatively low ($r^2 = 0.096 \sim 0.354$), which may be attributed to the use of pressure sensor-based gait analysis. However, multiple previous studies using gait analysis provided insufficient

explanation for this phenomenon. To overcome this limitation, further studies with larger patient populations and more varied biometric data may be necessary.

Conclusions

Measurements of walking speed and stance-phase time of the unaffected lower limb can allow clinicians to easily assess the ambulatory and balance functions of hemiplegic stroke patients. Rehabilitative treatment focusing on increasing gait speed and shortening the stance-phase time of the unaffected side may improve the ambulatory and balance functions in these patients.

Abbreviations

MBI: Modified Barthel index (MBI); MMT: Manual muscle test; BBS: Berg balance scale; FAC: Functional ambulation category; MMSE: Mini Mental Status Examination; VIFs: Variance inflation factors

Acknowledgements

None.

Authors' contributions

MCC: Writing – have substantial contribution to the design of the work, and conception, and have drafted the work, substantively revised it and editing. BJL: Writing – have drafted the work, substantively revised it and editing, NJ: have made substantial contributions to the acquisition and analysis of data. DP: have drafted the work, substantively revised it and editing, and have substantial contributions to the conception. The authors read and approved the final manuscript.

Funding

This research was supported by Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Science, ICT, and Future Planning (NRF- 2020R1F1A1075199).

Availability of data and materials

The data sets in this study are available from the corresponding author on reasonable request.

Ethics approval and consent to participate

The study protocols were approved by the Institutional Review Board of Yeungnam University Medical Center. Written informed consent was obtained from all patients participating in the study.

Consent for publication

Each of the authors has read and concurs with the content in the final manuscript.

Competing interests

All authors declare that they have no competing interests.

Author details

¹Department of Rehabilitation Medicine, College of Medicine, Yeungnam University, Daegu, Republic of Korea. ²Department of Rehabilitation Medicine, Daegu Fatima Hospital, Daegu, Republic of Korea. ³Department of Physical Medicine and Rehabilitation, Ulsan University Hospital, University of Ulsan

College of Medicine, 877 Bangeojin sunhwando-ro, Dong-gu, 44033 Ulsan, Republic of Korea.

Received: 23 September 2020 Accepted: 21 January 2021

Published online: 27 January 2021

References

- McPhee JS, French DP, Jackson D, Nazroo J, Pendleton N, Degens H. Physical activity in older age: perspectives for healthy ageing and frailty. *Biogerontology*. 2016;17(3):567–80. <https://doi.org/10.1007/s10522-016-9641-0> PubMed PMID: 26936444; PubMed Central PMCID: PMC6488962 Epub 2016/03/05.
- Eng JJ, Tang PF. Gait training strategies to optimize walking ability in people with stroke: a synthesis of the evidence. *Expert Rev Neurother*. 2007;7(10):1417–36. doi:<https://doi.org/10.1586/14737175.7.10.1417>. PubMed PMID: 17939776; PubMed Central PMCID: PMC63196659. Epub 2007/10/18.
- Dehghani A, Khoramkish M, Shahsavari Isfahani S. Challenges in the daily living activities of patients with multiple sclerosis: a qualitative content analysis. *Int J Community Based Nurs Midwifery*. 2019;7(3):201–10. <https://doi.org/10.30476/IJCBNM.2019.44995>. Epub 2019/07/26. PubMed PMID: 31341919; PubMed Central PMCID: PMC6614347.
- Park YJ, Lee CY. Effects of community-based rehabilitation program on activities of daily living and cognition in elderly chronic stroke survivors. *J Phys Ther Sci*. 2016;28(11):3264–6. doi:<https://doi.org/10.1589/jpts.28.3264>. PubMed PMID: 27942164; PubMed Central PMCID: PMC65140844. Epub 2016/12/13.
- Gardner AW, Montgomery PS. Impaired balance and higher prevalence of falls in subjects with intermittent claudication. *J Gerontol A Biol Sci Med Sci*. 2001;56(7):M454–8. Epub 2001/07/11. doi:<https://doi.org/10.1093/gerona/56.7.m454>. PubMed PMID: 11445605.
- Abid M, Mezghani N, Mitiche A. Knee joint biomechanical gait data classification for knee pathology assessment: a literature review. *Appl Bionics Biomech*. 2019;2019:7472039. Epub 2019/06/21. doi:<https://doi.org/10.1155/2019/7472039>. PubMed PMID: 31217817; PubMed Central PMCID: PMC6536985.
- Verghese J, Wang C, Lipton RB, Holtzer R, Xue X. Quantitative gait dysfunction and risk of cognitive decline and dementia. *J Neurol Neurosurg Psychiatry*. 2007;78(9):929–35. doi:<https://doi.org/10.1136/jnnp.2006.106914>. PubMed PMID: 17237140; PubMed Central PMCID: PMC631995159. Epub 2007/01/24.
- Hollman JH, McDade EM, Petersen RC. Normative spatiotemporal gait parameters in older adults. *Gait Posture*. 2011;34(1):111–8. doi:<https://doi.org/10.1016/j.gaitpost.2011.03.024>. PubMed PMID: 21531139; PubMed Central PMCID: PMC63104090. Epub 2011/05/03.
- Lord S, Galna B, Verghese J, Coleman S, Burn D, Rochester L. Independent domains of gait in older adults and associated motor and nonmotor attributes: validation of a factor analysis approach. *J Gerontol A Biol Sci Med Sci*. 2013;68(7):820–7. doi:<https://doi.org/10.1093/gerona/gls255>. PubMed PMID: 23250001. Epub 2012/12/20.
- Olney SJ, Griffin MP, McBride ID. Multivariate examination of data from gait analysis of persons with stroke. *Phys Ther*. 1998;78(8):814–28. doi:<https://doi.org/10.1093/ptj/78.8.814>. PubMed PMID: 9711207. Epub 1998/08/26.
- Patterson KK, Parafianowicz I, Danells CJ, Closson V, Verrier MC, Staines WR, et al. Gait asymmetry in community-ambulating stroke survivors. *Arch Phys Med Rehabil*. 2008;89(2):304–10. doi:<https://doi.org/10.1016/j.apmr.2007.08.142>. PubMed PMID: 18226655. Epub 2008/01/30.
- Han C, Jo SA, Jo I, Kim E, Park MH, Kang Y. An adaptation of the Korean mini-mental state examination (K-MMSE) in elderly Koreans: demographic influence and population-based norms (the AGE study). *Arch Gerontol Geriatr*. 2008;47(3):302–10. doi:<https://doi.org/10.1016/j.archger.2007.08.012>. PubMed PMID: 17936377. Epub 2007/10/16.
- Yang CM, Wang YC, Lee CH, Chen MH, Hsieh CL. A comparison of test-retest reliability and random measurement error of the Barthel Index and modified Barthel Index in patients with chronic stroke. *Disabil Rehabil*. 2020;1–5. Epub 2020/09/10. doi: <https://doi.org/10.1080/09638288.2020.1814429>. PubMed PMID: 32903114.
- Park D, Lee SH, Shin JH, Park JS. Lower limb muscle magnetic resonance imaging in myotonic dystrophy type 1 correlates with the six-minute walk test and CTG repeats. *Neuromuscul Disord*. 2018;28(1):29–37. doi: 10.1016/j.nmd.2017.08.005. PubMed PMID: 29066035.
- Park D, Park JS. Quantitative assessment of trunk muscles involvement in patients with myotonic dystrophy type 1 using a whole body muscle magnetic resonance imaging. *Eur Neurol*. 2017;77(5–6):238–45. <https://doi.org/10.1159/000460291> Epub 2017/03/14. PubMed PMID: 28288466.
- van Hedel HJ, Group ES. Gait speed in relation to categories of functional ambulation after spinal cord injury. *Neurorehabil Neural Repair*. 2009;23(4):343–50. <https://doi.org/10.1177/1545968308324224>. Epub 2008/11/28. PubMed PMID: 19036717.
- Mackintosh SF, Hill KD, Dodd KJ, Goldie PA, Culham EG. Balance score and a history of falls in hospital predict recurrent falls in the 6 months following stroke rehabilitation. *Arch Phys Med Rehabil*. 2006;87(12):1583–9. doi:<https://doi.org/10.1016/j.apmr.2006.09.004>. PubMed PMID: 17141637. Epub 2006/12/05.
- Park JH, Kim Y, Lee KJ, Yoon YS, Kang SH, Kim H, et al. Artificial neural network learns clinical assessment of spasticity in modified Ashworth scale. *Arch Phys Med Rehabil*. 2019;100(10):1907–15. <https://doi.org/10.1016/j.apmr.2019.03.016> PubMed PMID: 31009599 Epub 2019/04/23.
- Iwai M, Koyama S, Tanabe S, Osawa S, Takeda K, Motoya I, et al. The validity of spatiotemporal gait analysis using dual laser range sensors: a cross-sectional study. *Arch Physiother*. 2019;9:3. doi:<https://doi.org/10.1186/s40945-019-0055-6>. PubMed PMID: 30820352; PubMed Central PMCID: PMC6381722. Epub 2019/03/02.
- Cromwell RL, Newton RA. Relationship between balance and gait stability in healthy older adults. *J Aging Phys Act*. 2004;12(1):90–100. doi:<https://doi.org/10.1123/japa.12.1.90>. PubMed PMID: 15211023. Epub 2004/06/24.
- Fulk GD, Echtennach JL. Test-retest reliability and minimal detectable change of gait speed in individuals undergoing rehabilitation after stroke. *J Neurol Phys Ther*. 2008;32(1):8–13. doi:<https://doi.org/10.1097/NPT0b013e31816593c0>. PubMed PMID: 18463550. Epub 2008/05/09.
- Perry J, Garrett M, Gronley JK, Mulroy SJ. Classification of walking handicap in the stroke population. *Stroke*. 1995;26(6):982–9. doi:<https://doi.org/10.1161/01.str.26.6.982>. PubMed PMID: 7762050. Epub 1995/06/01.
- Abellan van Kan G, Rolland Y, Andrieu S, Bauer J, Beauchet O, Bonnefoy M, et al. Gait speed at usual pace as a predictor of adverse outcomes in community-dwelling older people: an International Academy on Nutrition and Aging (IANA) Task Force. *J Nutr Health Aging*. 2009;13(10):881–9. Epub 2009/11/20. doi: <https://doi.org/10.1007/s12603-009-0246-z>. PubMed PMID: 19924348.
- Bland MD, Sturmoski A, Whitson M, Connor LT, Fucetola R, Huskey T, et al. Prediction of discharge walking ability from initial assessment in a stroke inpatient rehabilitation facility population. *Arch Phys Med Rehabil*. 2012; 93(8):1441–7. doi:<https://doi.org/10.1016/j.apmr.2012.02.029>. PubMed PMID: 22446516; PubMed Central PMCID: PMC635707123. Epub 2012/03/27.
- Pirker W, Katzenschlager R. Gait disorders in adults and the elderly: A clinical guide. *Wien Klin Wochenschr*. 2017;129(3–4):81–95. doi:<https://doi.org/10.1007/s00508-016-1096-4>. PubMed PMID: 27770207; PubMed Central PMCID: PMC635318488. Epub 2016/10/23.
- Reelick MF, van Iersel MB, Kessels RP, Rikkert MG. The influence of fear of falling on gait and balance in older people. *Age Ageing*. 2009;38(4):435–40. <https://doi.org/10.1093/ageing/afp066> PubMed PMID: 19451658 Epub 2009/05/20.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

