

# Depression and its associated factors among pregnant women in central Vietnam

Health Psychology Open  
January-June 2021: 1–10  
© The Author(s) 2021  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/2055102920988445  
journals.sagepub.com/home/hpo  


Bao-Yen Luong-Thanh<sup>1</sup>,  Lan Hoang Nguyen<sup>1</sup>,   
Linda Murray<sup>2</sup>, Manuel Eisner<sup>3</sup>, Sara Valdebenito<sup>3</sup>,  
Tuyen Dinh Hoang<sup>1</sup>, Huyen Phuc Do<sup>4</sup>  and Thang Van Vo<sup>1</sup> 

## Abstract

To date, little attention has been given to prenatal depression, especially in low and middle-income countries. The aim of this research was to assess the prevalence of depression and its associated factors amongst pregnant women in a central Vietnamese city. This cross-sectional study included 150 pregnant women from 29 to 40 weeks of gestation, from eight wards of Hue city, via quota sampling from February to May 2019. We employed the Patient Health Questionnaire (PHQ-9) to assess depression. Findings suggest the need to provide routine screening of pregnant women in primary care for depressive symptoms and other mental health problems.

## Keywords

depression, mental illness, pregnancy, wellbeing, women's health

## Introduction

Depression is a common mental health concern that affects women globally, especially during the perinatal period (Howard et al., 2014). Its prevalence in pregnancy has been estimated to range from 6.5% to 26.7% (Gavin et al., 2005; Okagbue et al., 2019). The prevalence rates of antenatal depression differ within and between countries. Perinatal mental disorders are more prevalent in low- and middle-income countries (LMICs), compared to those from high income countries. In high-income countries, the prevalence of depression in pregnancy is estimated to range from 7.4% to 12.8% (Bennett et al., 2004), compared to 15.6% (95% CI: 15.4–15.9%) in LMICs (Fisher et al., 2012).

Evidence suggests that if not treated and diagnosed early, depression during pregnancy can have more serious consequences in the postpartum period (Falah-Hassani et al., 2017). Moreover, a number of short-term and long-term adverse outcomes for the children of women who suffered from antenatal depression have been identified (Dunkel Schetter and Tanner, 2012; Gentile, 2017; Lefkovic et al., 2014). Such outcomes include increased risk of low birth weight, infants that are small for gestational age, preterm birth (Accortt et al., 2015; Grigoriadis et al., 2013; Staneva

et al., 2015a; Szegda et al., 2014); and increased vulnerability to behavioral and emotional problems such as depression and other mental health issues in later childhood and adolescence (Plant et al., 2016; Van den Bergh et al., 2005).

In Vietnam, most research on depression in the perinatal period has focused on postnatal depression only (Do et al., 2018; Murray et al., 2015; Tran et al., 2018b, 2019; Upadhyay et al., 2019; Vo et al., 2017; Wesselhoeft et al., 2020), with a paucity of research focusing on antenatal depression. The two existing studies of antenatal depression amongst Vietnamese pregnant women report prevalence of 4.9% and 28.2% (Fisher et al., 2013; Ngo et al.,

<sup>1</sup>Faculty of Public Health and Institute for Community Health Research, University of Medicine and Pharmacy, Hue University, Vietnam

<sup>2</sup>College of Health Sciences, Massey University, New Zealand

<sup>3</sup>Institute of Criminology, University of Cambridge, United Kingdom

<sup>4</sup>School of Public Health and Social Work, Queensland University of Technology (QUT), Australia

## Corresponding author:

Thang Van Vo, Faculty of Public Health and Institute for Community Health Research, University of Medicine and Pharmacy, Hue University, 06 Ngo Quyen Street, Hue city, Vietnam.

Email: vovanthang147@hueuni.edu.vn



2018). Both of these studies were conducted in the North of Vietnam, and there has been limited research conducted throughout the rest of the country. Additionally, although a recent systematic review suggested that common mental disorders (including depression) are significantly associated with intimate partner violence during pregnancy (OR: 3.69, 95%CI: 2.51–5.42), the associated factors linking this to maternal depression remain unclear due to the complexity of this relationship and a lack of screening during pregnancy (Kingston et al., 2015). This study aimed to estimate the prevalence of depressive symptoms during pregnancy in 2019, and identify factors affecting depressive symptoms among pregnant women in a central region of Vietnam.

## Methods

### Study design

This research formed part of a pilot baseline study conducted as part of the Evidence for Better Lives Foundational Study (EBLS-FS)—among eight medium-sized cities from low- and middle-income countries, including Ghana, Jamaica, Pakistan, the Philippines, Romania, South Africa, Sri Lanka, and Vietnam. Details of the protocol have been published elsewhere (Valdebenito et al., 2020). In Vietnam, a cross-sectional study was conducted from February to May 2019 to estimate the prevalence of maternal health in Hue city.

### Study setting

The study was conducted in Hue city, the ancient imperial capital of Vietnam and the capital of Thua Thien Hue province. Hue city is located in the central region of Vietnam, covering an area of 70.67 km<sup>2</sup>. Its population in 2018 was approximately 358,754 people and it is divided into 27 wards (Portal of Thua Thien Hue Province, 2020). The official estimate of the poverty rate in 2019 is approximately 1.51% (Thua Thien Hue Provincial People's Committee, 2020).

### Study participants

We included pregnant women with a gestational age ranging from 29 to 40 weeks, aged 18 years and above, residing in Hue city for at least 6 months. Exclusion criteria included women who refused to participate, or who planned to migrate from Hue city within 3 months after giving birth.

### Sample size and sampling

A quota sampling method was employed to select 150 participants from a list of pregnant women who were recruited via ward health centers during their standard antenatal care visits in each ward selected for this study. The geographical

stratification was based on the list of wards in the two Southern and Northern parts of Hue city. From the two parts of Hue City, eight out of 27 wards were randomly selected and then, from each selected ward, the midwife from the ward health center contacted eligible pregnant women for recruitment. As a result, a total of 195 participants were approached, and 150 pregnant women were successfully recruited, achieving a response rate of 76.9%. The 45 women who refused to participate cited reasons such as inadequate time for participating in the interview or imminent due date of delivery.

### Measurements

*The dependent variable. Depression:* The Patient Health Questionnaire (PHQ-9) was used to measure the severity of depressive symptoms experienced in the last 2 weeks (Kroenke et al., 2001). The short module scores the nine DSM-IV criteria for depression, including anhedonia, dysphoria, sleep disturbances, fatigue, changes in eating, low self-esteem, concentration difficulties, hypo-or-hyperactive behaviors, and thoughts of suicide or homicide. Each item was rated on a 4-point Likert scale: Not at all=0; Several days=1; More than half the days=2; Nearly every day=3. The possible range was 0–27. A total score greater or equal to 10 was the cut-off for moderate to severe depression (Kroenke et al., 2001).

*Independent variables. Maternal sociodemographic characteristics:* We included maternal age, maternal education, employment status, type of relationship with the father of the baby, and perceived social status. Perceived social status was measured by using the MacArthur scale of subjective social status—a single-item on a 10-point Likert-type scale that assesses a person's perceived rank relative to others in their group (Adler et al., 2000). Those who scored themselves as three or less were considered as having perceived low social status. Items on reproductive health and prenatal characteristics included unwanted pregnancy, parity, and frequency of prenatal check-ups.

*Adverse Childhood Experiences (ACEs):* The adapted 19 out of 31 items of the ACE-International questionnaire (WHO, 2019) were employed to measure any adverse experience before age 18 years. The domains included in this study were: emotional abuse; physical abuse; sexual abuse; violence against household members; living with household members who were substance abusers; living with household members who were mentally ill or suicidal; living with household members who were imprisoned; growing up with one or no parents, parental separation or divorce; emotional neglect; physical neglect. The number of ACE events were later classified as two groups: those who experienced at least four ACEs, and those who experienced less than four ACEs across the nine domains.

*Exposure to intimate partner violence (IPV) during pregnancy* was recorded if perpetrated by a current husband/ partner during the past 6 months. We used the adapted version from the WHO Multi-country Study on Women's Health and Domestic Violence Against Women to measure prenatal IPV (Garcia-Moreno et al., 2006). The WHO questionnaire comprises four questions on emotional IPV, six questions on physical IPV and three questions on sexual IPV. Emotional violence was defined as being insulted, humiliated, intimidated or threatened to hurt her or someone she cared for. Physical violence referred to being slapped, pushed, hit, kicked, choked or threatened. Sexual violence was defined as being physically forced to have sexual intercourse, consenting due to fear of partner's reaction, or being forced to do a sexual act which she found degrading or humiliating. Pregnant women who experienced any type of prenatal IPV in her lifetime were classified as having experienced at least one type of lifetime IPV. If they experienced at least one type of IPV in the last 6 months, they were considered as IPV exposed during pregnancy.

*Self-report stress:* The Perceived Stress Scale (PSS) (Cohen et al., 1983) was applied to measure the level of maternal stress. This scale was validated in the Vietnamese population (Dao-Tran et al., 2017). It consists of ten items that measures how stressful certain life situations were rated by respondents during the last month via a 4-point Likert scale: Not at all=1; Several days=2; More than half the days=3; Nearly every day=4. A mean score of greater than one was defined as perceived stress.

*Partner's characteristics:* We recorded age, educational attainment, and employment status based on information provided by the women.

*Partner supportiveness:* The quality of the mother-father relationship was measured by the partner supportiveness scales (Goldberg and Carlson, 2014). This was a 5-item scale and each item was rated on a 5-point Likert scale: Never=1, Rarely=2, Sometimes=3, Often=4, Always=5. A mean score of less than two was defined as low support from partner.

*Social support:* The Multidimensional Scale of Perceived Social Support (Zimet et al., 1988) was applied to investigate social support. Twelve items measured perceptions of support from three sources: family, friends, and a significant other. The original scale uses a 7-point Likert-type scale, while our questionnaire reduced it to 5-point Likert-type scale: Strongly disagree=1, disagree=2; Neutral=3; agree=4; Strongly agree=5. A mean score of less than three was defined as low social support.

### Data collection

Before data collection, eight midwives were trained intensively for 3 days regarding the study protocol. One midwife from each ward health center conducted interviews via a

tablet assisted face-to-face interview in either a private room in a ward health center, or at a participants' house if appropriate. Information on IPV during pregnancy were collected using Computer-Assisted Self-Interviewing. The interview data were then submitted directly into a secure web application for building and managing online surveys and databases called "Qualtrics," with data stored at the University of Cambridge (United Kingdom).

### Data analysis and statistical method

Participant characteristics were described using frequency and percentages for categorical data, and mean (SD), for continuous variables. To investigate the factors associated with depression among pregnant women, odds ratios (OR) and 95% confidence intervals (95%CI) were estimated using multivariate logistics models. All analyses were performed using the R program (R Core Team, 2014). Statistical significance was defined as two-tailed p values of less than 0.05.

### Ethical considerations

The study was approved by the Ethical Review Committee of Hue University of Medicine and Pharmacy (No H2018/430 dated November 15, 2018). Permission to conduct the study was also obtained from provincial and wards authorities in Thua Thien Hue province, Vietnam. Participants were informed about the study and provided written informed consent prior to the interviews.

## Results

### General characteristics of the study population

Table 1 describes participant characteristics. The mean age was 29.9 years (SD=5.0 years), with a range between 19 and 47 years. Most women were between 25 and 35 years old (71.3%); finished high school or above (62.0%), had paid work in the last 12 months (71.3%), and lived with the father of the baby (97.3%). Among those participants, 12.7% had perceived low social status. Regarding prenatal characteristics, most of the study participants were multiparous (70.7%), wanted to be pregnant (69.3%) and received at least four antenatal checkups (95.3%). About 30% of pregnant women experienced at least four types of ACEs before 18 years old and 28% perceived stress during the previous month. Twenty-two women (14.7%) reported experiencing at least one type of IPV during the last 6 months. The prevalence of emotional violence was highest (14.0%), followed by physical violence (2.0%) and sexual violence (1.3%). The mean age for the participants' partners was 33.1 (SD=6.0). Most of the women's husbands had paid work in the last 12 months (98%) and more than half of them completed high school or above (57.3%).

**Table 1.** Characteristics of study population ( $n = 150$ ).

Characteristics		Number ( $n$ )	Percentage (%)
<b>Maternal socio-demographic characteristics</b>			
Age (years)	Mean (SD)	29.9 (5.0)	
	Median [Min, Max]	30 [19, 47]	
	18 to <25	20	13.3
	25 to <35	107	71.3
	$\geq 35$	23	15.3
Education	Primary school or below	28	18.7
	Secondary school	29	19.3
	High school or above	93	62.0
Paid work in the last 12 months	No	43	28.7
	Yes	107	71.3
Perceived low social status	No	131	87.3
	Yes	19	12.7
Lives with the father of the baby	No	4	2.7
	Yes	146	97.3
<b>Partner's characteristics</b>			
Age	Mean (SD)	33.1 (6.0)	
	Median [Min, Max]	32 [22, 54]	
Partner same age or younger	Yes	35	23.3
	No	115	76.7
Education	Primary school or below	31	20.7
	Secondary school	33	22.0
	High school or above	86	57.3
Having paid work in the last 12 months	No	3	2.0
	Yes	147	98.0
<b>Maternal obstetrics characteristics</b>			
Parity	Nulliparous	44	29.3
	Multiparous	106	70.7
Unwanted pregnancy	No	104	69.3
	Yes	46	30.7
Number of antenatal check-up	<4	7	4.7
	$\geq 4$	143	95.3
Type of adverse childhood events before 18 years old	<4	105	70.0
	$\geq 4$	45	30.0
Lifetime intimate partner violence	<b>At least one type</b>	15	15.3
	Emotional IPV	20	13.3
	Physical IPV	7	4.6
	Sexual IPV	1	0.6
<b>Intimate partner violence during pregnancy</b>			
Number of types of IPV	No	128	85.3
	One type	18	12.0
	Two types	4	2.7
Emotional IPV	No	129	86.0
	Yes	21	14.0
Physical IPV	No	147	98.0
	Yes	3	2.0
Sexual IPV	No	148	98.7
	Yes	2	1.3
Perceived stress	No	108	72.0
	Yes	42	28.0
Perceived low support from partner	No	146	97.3
	Yes	4	2.7
Perceived low social support	No	79	52.7
	Yes	71	47.3

**Table 2.** Prevalence of antenatal depression among pregnant women ( $n = 150$ ).

	Number ( $n$ )	Percentage (%)
PHQ-9		
Mean (SD)	5.2 (3.7)	
Median [Min, Max]	5.0 [0.0, 20.0]	
Moderate to severe depressive symptoms	19	12.7 [7.3–18.1]

Overall, 2.7% of women perceived low support from their husband and nearly half of the participants perceived themselves as having low social support (47.3%).

### Main findings

Table 2 indicates that the prevalence of moderate to severe depression among study participants was 12.7% (95%CI: 7.3%–18.1%). The mean score of the PHQ-9 was 5.2 (SD=3.7).

Table 3 presents the factors associated with depressive symptoms amongst pregnant women. Pregnant women who self-reported as being stressed (aOR=7.62; 95%CI: 2.66–21.81) and had husbands with education attainment of primary school or below (aOR=4.28; 95%CI: 1.02–17.85) were more likely to experience moderate to severe depression. Perceived low support from partners was significantly associated with the odds of depressive symptoms, but only in univariate analysis (Crude OR=24.37; 95% CI: 2.39–248.52) not multivariable regression (aOR=23.62; 95% CI: 0.83–672.53). Our findings did not show any association between depressive symptoms and other factors, such as maternal age, unwanted pregnancy, antenatal check-ups, ACEs, and prenatal IPV, partner's age, and support from their partner or society.

## Discussion

### The strength of this study

This study was one of the first studies on depression among pregnant women in the central region of Vietnam. The main strength of this study was that we used a standardized, validated instrument to measure self reported depressive symptoms (Collier et al., 2020; Dreher et al., 2017; Hinton et al., 2018; Nguyen et al., 2016; Vo et al., 2019). Furthermore, data collectors were well-trained midwives which ensured the quality of data collection. In addition, the application of Qualtrics also reduced missing information compared to use of a paper-based questionnaire.

### Main findings

Our study revealed that more than one in ten women experienced moderate to severe depression during the antenatal stages of pregnancy (12.7%). In Vietnam, studies that used

the Edinburgh Postnatal Depression Scale (EPDS) to measure antenatal depression reported various findings. A higher prevalence was demonstrated in a study conducted in Hanoi in 2007 (14.7%) (Fisher et al., 2007), which involved 61 pregnant women attending an antenatal clinic. A larger study in Dong Anh, in the North of Vietnam which involved 1274 women in the first and second trimester of pregnancy yielded a lower prevalence of 5.0% (Ngo et al., 2018; Tran et al., 2019). Compared to other countries, our findings reported a lower prevalence of depression during the third trimester than others studies which employed the PHQ-9 scale to measure, which were 26.8% in Ghana and 32.7% in Côte d'Ivoire (Bindt et al., 2012), 27.6% in Peru (Yang et al., 2016), 24.9% in Ethiopia (Bitew et al., 2016), and 31.1% in Italy (Venanzio, 2017). Inconsistent prevalence estimates were seen in other studies that used different tools to measure depression besides the PHQ-9 (Bennett et al., 2004; Sawyer et al., 2010; Underwood et al., 2016). There is a disparity in the proportion of women who experienced antenatal depression across countries and within countries. This disparity is likely attributable to macro-level variation in the living conditions of pregnant women, such as the stability of partnership arrangements, access to public health services, and stressful life events, but also methodological differences between studies. Moreover, the high prevalence of antenatal depression in this study suggests the needs for screening for depression during pregnancy at clinical settings in order to reduce the risk of pregnancy outcomes. The effectiveness of screening interventions followed by interventions has been confirmed in other studies (O'Connor et al., 2016; Ponting et al., 2020).

Our study demonstrated that one factor that might be link to antenatal depression was having a partner with a low level of education. Therefore, further interventions should pay attention to families with husbands of low education backgrounds. In this study, lack of support from partners was found to be associated with the odds of experiencing antenatal depression in bivariate analysis but not in the multivariate analysis. The small sample size might be one explanation for this difference. The high correlation between inadequate partner support and a higher risk of experiencing antenatal depression may be explained by gendered role assignment within families in Hue city. In Hue city, it is common for women to take responsibility for child rearing whilst men assume a the role of “breadwinner” and household leader (Dao et al., 2012). This finding is supported by the literature which suggested that having a supportive husband is associated with good maternal mental health during pregnancy and after childbirth (Murray et al., 2015; Vo et al., 2017). Partner supportiveness is protective for pregnant women and aids coping with stressful life events, and is therefore likely to have a significant impact on maternal and infant well-being postpartum (Dayan et al., 2010; Zegeye et al., 2018). A current synthesis of qualitative research reconfirmed partner

**Table 3.** Factors associated with moderate to severe depressive symptoms among pregnant women ( $n = 150$ ).

Characteristics		Univariate analysis	Multivariable regression model
		Crude OR (95%CI)	Adjusted OR (95%CI)
Maternal age	<35		
	≥35	2.24 (0.72, 6.98)	4.65 (0.9, 24.02)
Unwanted pregnancy	No		
	Yes	2.29 (0.86, 6.08)	1.27 (0.33, 4.98)
Number of antenatal check-ups	<4		
	≥4	2.96 (0.53, 16.49)	2.54 (0.25, 25.94)
Partner same age or younger	No		
	Yes	2.15 (0.77, 5.96)	2.6 (0.68, 9.95)
Father's education attainment	High school or above		
	Below secondary school	6.35 (2.07, 19.47)*	4.28 (1.02, 17.85)*
	Secondary school	1.33 (0.31, 5.67)	0.98 (0.17, 5.77)
Number of adverse childhood events before 18 years old	<4		
	≥4	2.37 (0.89, 6.32)	1.2 (0.28, 5.16)
Experience at least one type of IPV in the last 6 months	No		
	Yes	2.39 (0.77, 7.5)	1.05 (0.2, 5.54)
Perceived low support from partner	No		
	Yes	24.37 (2.39, 248.52)*	23.62 (0.83, 672.53)
Perceived low social support	Yes		
	No	1.59 (0.32, 8.01)	0.28 (0.03, 2.4)
Perceived stress	No		
	Yes	7.62 (2.66, 21.81)*	9.77 (2.58, 36.97)*

\* $p < 0.05$ .

supportiveness as an important resource for maternal well-being during pregnancy (Staneva et al., 2015b).

This study also identified that stress was associated with higher odds of experiencing depression during pregnancy. This finding was consistent with other studies (Ahmed et al., 2019; Bayrampour et al., 2016). Previous studies have revealed that stress can alter hormonal responses, including increasing the activity of the hypothalamic–pituitary–adrenal axis, and reducing levels of norepinephrine which can trigger maternal depressive symptoms (Moret and Briley, 2011; Pariante, 2014; Seth et al., 2016). This finding suggests that the contents of screening for prenatal mental disorders should cover not only antenatal depression, but also stress, anxiety and other mental disorders if possible, which might enhance a better control for prenatal mental problems.

Findings from our study indicated that gender-based risk factors such as experiencing IPV were not significantly associated with prenatal depression. This might be due to small sample size, which prevents measurement of this association. However, our study indicated that 12% of pregnant women experienced prenatal IPV, which was in line with findings of a systematic review reporting a range of IPV prevalence from 5.9% to 32.5% among 4598 pregnant women in Vietnam (Do et al., 2019). These important findings raise a critical question about how to screen for psychological abuse and violent tactics, and how to remove social barriers and encourage women to disclose experiences of violence. Though domestic violence

has become less acceptable among Vietnamese women (Trinh et al., 2016), there is still a perceived social norm that emphasizes women's responsibility to maintain the harmony and reputation of the family, and to build the "cultural family" and "cultural village" (Schuler et al., 2006) which can prevent disclosures of IPV (Kwiatkowski, 2011; Schuler et al., 2016; Tran et al., 2018a; Vu et al., 2014). Moreover, the influence of Confucian social norms may also inhibit women from disclosing family issues (Tran et al., 2016). In addition, this disparity may be due to the willingness of women to disclose IPV exposure, which is likely to be affected by culture and social norms. A recent review also reported an association between IPV and depression (Fisher et al., 2012; Howard et al., 2014; Tsai et al., 2016). Another study in Vietnam also confirmed this relationship (Fisher et al., 2013). Therefore, screening for IPV during antenatal care remains strongly recommended.

Our study revealed that ACEs were not associated with depression among the study participants. This finding was different from a systematic review which identified a strong effect of multiple ACEs on mental illness, especially antenatal depression (Howell et al., 2017; Hughes et al., 2017; Wajid et al., 2019). However, this finding was supported by a previous study in Hue city among 608 adults. The authors of this study did not find a significant association between ACEs' exposure and mental disorders (Do, 2019). This unexpected result may be explained by the cultural context in Hue city which is strongly influenced by Confucianism,

and where parents may apply corporal punishment and harsh discipline to manage their child's behavior (Hoang, 2014). In this context, parental physical punishment may be perceived as normative and necessary to educate children (Rudy and Grusec, 2006) rather than being conceptualized as an adverse experience. Another explanation may be recall bias, as some emotional experiences from childhood might not be recalled clearly in adulthood (Fergusson et al., 2000), or respondents were less likely to report ACEs.

### Limitations of the study

This study had several limitations. First, the sample size of this pilot study was small and participants were recruited using a non-probabilistic sampling method. The response rate of this study was 76.9% which is considered moderate quality based on EPHPP checklist (Effective Public Health Practice Project, 1998). However, with a small sample size and moderate response rate, caution must be applied, as the findings might not be generalized to the whole Vietnamese population. Data were also collected in the third trimester of pregnancy; thus, our findings may not be generalizable to the first and second trimesters of pregnancy. Second, recall bias and reporting bias may have affected the internal validity of this study. For example, the unwillingness of participants to disclose stigmatizing experiences such as IPV and depression, might have led to an underestimation of their true prevalence. To address this issue, we collected data using Qualtrics, and interviews were administered by well-trained midwives who had close rapport with study participants. Third, some items of the PHQ-9 related to symptoms of pregnancy (i.e. fatigue, poor appetite, difficulty sleeping) may mirror somatic symptoms of depression, which may lead to an overestimation of the proportion or severity of depression. Moreover, the PHQ-9 is a self-reported scale rather than clinical assessment and also not specific to assess depression for perinatal periods like other tools (i.e. the EPDS), which may have led to an under- or overestimation of antenatal depression.

### Conclusion

The prevalence of moderate to severe depressive symptoms experienced antenatally was 12.7%. Experiencing stress, having a husband with a low level of education and perceived low levels of support from their husband were found to be significantly associated with higher odds of antenatal depression amongst the study population. However, in the multivariate analysis, husband's support was no longer a significant factor. These findings suggest the need to screen for depression during pregnancy in central Vietnam. A collaborative care approach between antenatal care services and mental health care practitioners will ensure that women who have symptoms of antenatal depression are managed appropriately. In addition, routine screening of pregnant women in

primary care for common perinatal mental disorders could be added to existing antenatal assessments. With appropriate referral systems in place, this could reduce women's vulnerability to its adverse consequences and help to achieve optimal postnatal outcomes. Health promotion programs should encompass psychosocial health, gender equity and women's right to be free from violence, especially during pregnancy. Further studies with larger sample size should be considered.

### Acknowledgements

We are indebted to the mothers who participated in this study and to Thua Thien Hue provincial People Committee, provincial Department of Labor, Invalids, and Social Affairs, provincial Department of Health and especially staffs from Institute for Community Health Research and Faculty of Public health of Hue University of Medicine and Pharmacy for data collection in field-work. Our sincere thank also goes to Hue University for the partial support under the core research program.

### Author contributions

BYLT, TVV, ME conceptualized the research question and participated in the analysis plan. BYLT performed the data analysis and drafted the manuscript. TVV, ME, LM, SVM, LNH, HPD, and THD advised about study design, data collection tools and manuscript contents. All authors read and approved the final draft of the manuscript.

### Declaration of conflicting interests


The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

### Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This paper is an output of the Evidence for Better Lives Foundational Study (EBLS-FS). More information on this study can be found at [www.vrc.crim.cam.ac.uk/vrcresearch/EBLS](http://www.vrc.crim.cam.ac.uk/vrcresearch/EBLS). We are grateful for support for the EBLS-FS from: the UK ESRC Impact Acceleration Fund [grant number ES/M500409/1]; the Fondation Botnar, Switzerland [grant number RG92422, research project 6132]; the Jacobs Foundation, Switzerland; the UBS Optimus Foundation, Switzerland; the University of Cambridge School of Social Sciences Research Support Fund, UK; the British Academy, UK; the Consuelo Foundation, the Philippines; and Queensland University of Technology, Australia.

### ORCID iDs

Bao-Yen Luong-Thanh  <https://orcid.org/0000-0003-3511-6698>

Lan Hoang Nguyen  <https://orcid.org/0000-0003-1710-9478>

Huyen Phuc Do  <https://orcid.org/0000-0003-0461-7529>

Thang Van Vo  <https://orcid.org/0000-0003-2018-0371>

### References

Accortt EE, Cheadle ACD and Dunkel Schetter C (2015) Prenatal depression and adverse birth outcomes: An

- updated systematic review. *Maternal and Child Health Journal* 19(6): 1306–1337. DOI: 10.1007/s10995-014-1637-2.
- Adler NE, Epel ES, Castellazzo G, et al. (2000) Relationship of subjective and objective social status with psychological and physiological functioning: Preliminary data in healthy white women. *Health Psychology: Official Journal of the Division of Health Psychology, American Psychological Association* 19(6): 586–592.
- Ahmed A, Bowen A, Feng CX, et al. (2019) Trajectories of maternal depressive and anxiety symptoms from pregnancy to five years postpartum and their prenatal predictors. *BMC Pregnancy and Childbirth* 19(1): 26. DOI: 10.1186/s12884-019-2177-y.
- Bayrampour H, Ali E, McNeil DA, et al. (2016) Pregnancy-related anxiety: A concept analysis. *International Journal of Nursing Studies* 55: 115–130. DOI: 10.1016/j.ijnurstu.2015.10.023.
- Bennett HA, Einarson A, Taddio A, et al. (2004) Prevalence of depression during pregnancy: Systematic review. *Obstetrics and Gynecology* 103(4): 698–709. DOI: 10.1097/01.AOG.0000116689.75396.5f.
- Bindt C, Appiah-Poku J, Bonle MT, et al. (2012) Antepartum depression and anxiety associated with disability in African women: Cross-sectional results from the CDS study in Ghana and Côte d'Ivoire. *PLoS One* 7(10): e48396. DOI: 10.1371/journal.pone.0048396.
- Bitew T, Hanlon C, Kebede E, et al. (2016) Antenatal depressive symptoms and maternal health care utilisation: A population-based study of pregnant women in Ethiopia. *BMC Pregnancy and Childbirth* 16(1): 301. DOI: 10.1186/s12884-016-1099-1.
- Cohen S, Kamarck T and Mermelstein R (1983) A global measure of perceived stress. *Journal of Health and Social Behavior* 24(4): 385–396. DOI: 10.2307/2136404.
- Collier KM, Weiss B, Pollack A, et al. (2020) Explanatory variables for women's increased risk for mental health problems in Vietnam. *Social Psychiatry and Psychiatric Epidemiology* 55(3): 359–369. DOI: 10.1007/s00127-019-01761-3.
- Dao TD, Hoang C, Le HT, et al. (2012) 'Teach the wife when she first arrives' Trajectories and Pathways into Violent and Non-violent Masculinities in Hue City and Phu Xuyen District, Viet Nam. Hanoi.
- Dao-Tran T-H, Anderson D and Seib C (2017) The Vietnamese version of the Perceived Stress Scale (PSS-10): Translation equivalence and psychometric properties among older women. *BMC Psychiatry* 17(1): 53. DOI: 10.1186/s12888-017-1221-6.
- Dayan J, Creveuil C, Dreyfus M, et al. (2010) Developmental model of depression applied to prenatal depression: Role of present and past life events, past emotional disorders and pregnancy stress. *PLoS One* 5(9): e12942. DOI: 10.1371/journal.pone.0012942.
- Do HP, Tran BX, Nguyen CT, et al. (2019) Inter-partner violence during pregnancy, maternal mental health and birth outcomes in Vietnam: A systematic review. *Children and Youth Services Review* 96: 255–265. DOI: 10.1016/j.childyouth.2018.11.039.
- Do TKL, Nguyen TTH and Pham TTH (2018) Postpartum depression and risk factors among Vietnamese women. *BioMed Research International* 2018: 4028913. DOI: 10.1155/2018/4028913.
- Do TTH (2019) *Depression, anxiety and post traumatic stress disorder and their correlates among adults in Central Vietnam*. PhD Thesis, Queensland University of Technology. DOI: 10.5204/thesis.eprints.131606.
- Dreher A, Hahn E, Diefenbacher A, et al. (2017) Cultural differences in symptom representation for depression and somatization measured by the PHQ between Vietnamese and German psychiatric outpatients. *Journal of Psychosomatic Research* 102: 71–77. DOI: 10.1016/j.jpsychores.2017.09.010.
- Dunkel Schetter C and Tanner L (2012) Anxiety, depression and stress in pregnancy: Implications for mothers, children, research, and practice. *Current Opinion in Psychiatry* 25(2): 141–148. DOI: 10.1097/YCO.0b013e3283503680.
- Effective Public Health Practice Project (1998) *Quality Assessment Tool for Quantitative Studies*. Available at: [https://merst.ca/wp-content/uploads/2018/02/quality-assessment-tool\\_2010.pdf](https://merst.ca/wp-content/uploads/2018/02/quality-assessment-tool_2010.pdf) (accessed 13 December 2020).
- Falah-Hassani K, Shiri R and Dennis C-L (2017) The prevalence of antenatal and postnatal co-morbid anxiety and depression: A meta-analysis. *Psychological Medicine* 47(12): 2041–2053. DOI: 10.1017/S0033291717000617.
- Fergusson DM, Horwood LJ and Woodward LJ (2000) The stability of child abuse reports: A longitudinal study of their reporting behaviour of young adults. *Psychological Medicine* 30(3): 529–544. DOI: 10.1017/s003329179900211.
- Fisher J, Cabral de Mello M, Patel V, et al. (2012) Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: A systematic review. *Bulletin of the World Health Organization* 90(2): 139G–149G. DOI: 10.2471/BLT.11.091850.
- Fisher J, Tran TD, Biggs B, et al. (2013) Intimate partner violence and perinatal common mental disorders among women in rural Vietnam. *International Health* 5(1): 29–37. DOI: 10.1093/inthealth/ihs012.
- Fisher JR, Tran H and Tran T (2007) Relative socioeconomic advantage and mood during advanced pregnancy in women in Vietnam. *International Journal of Mental Health Systems* 1(1): 3. DOI: 10.1186/1752-4458-1-3.
- Garcia-Moreno C, Jansen HAFM, Ellsberg M, et al. (2006) Prevalence of intimate partner violence: Findings from the WHO multi-country study on women's health and domestic violence. *Lancet (London, England)* 368(9543): 1260–1269. DOI: 10.1016/S0140-6736(06)69523-8.
- Gavin NI, Gaynes BN, Lohr KN, et al. (2005) Perinatal depression: A systematic review of prevalence and incidence. *Obstetrics and Gynecology* 106(5 Pt 1): 1071–1083. DOI: 10.1097/01.AOG.0000183597.31630.db.
- Gentile S (2017) Untreated depression during pregnancy: Short- and long-term effects in offspring. A systematic review. *Neuroscience* 342: 154–166. DOI: 10.1016/j.neurosci.2015.09.001.
- Goldberg JS and Carlson MJ (2014) Parents' relationship quality and children's behavior in stable married and cohabiting families. *Journal of Marriage and Family* 76(4): 762–777. DOI: 10.1111/jomf.12120.
- Grigoriadis S, VonderPorten EH, Mamisashvili L, et al. (2013) The impact of maternal depression during pregnancy on perinatal outcomes: A systematic review and meta-analysis. *The Journal of Clinical Psychiatry* 74(4): e321–341. DOI: 10.4088/JCP.12r07968.



- Hinton DE, Pollack AA, Weiss B, et al. (2018) Culturally sensitive assessment of anxious-depressive distress in Vietnam: Avoiding category truncation. *Transcultural Psychiatry* 55(3): 384–404. DOI: 10.1177/1363461518764500.
- Hoang NM (2014) *Borderline features in Vietnamese adolescence: The roles of childhood trauma, parental bonding, and family functioning*. PhD Thesis, University of Minnesota.
- Howard LM, Molyneux E, Dennis C-L, et al. (2014) Non-psychotic mental disorders in the perinatal period. *Lancet (London, England)* 384(9956): 1775–1788. DOI: 10.1016/S0140-6736(14)61276-9.
- Howell KH, Miller-Graff LE, Schaefer LM, et al. (2017) Relational resilience as a potential mediator between adverse childhood experiences and prenatal depression. *Journal of Health Psychology* 25(4): 545–557. DOI: 10.1177/1359105317723450.
- Hughes K, Bellis MA, Hardcastle KA, et al. (2017) The effect of multiple adverse childhood experiences on health: A systematic review and meta-analysis. *The Lancet. Public Health* 2(8): e356–e366. DOI: 10.1016/S2468-2667(17)30118-4.
- Kingston D, Austin M-P, Heaman M, et al. (2015) Barriers and facilitators of mental health screening in pregnancy. *Journal of Affective Disorders* 186: 350–357. DOI: 10.1016/j.jad.2015.06.029.
- Kroenke K, Spitzer RL and Williams JB (2001) The PHQ-9: Validity of a brief depression severity measure. *Journal of General Internal Medicine* 16(9): 606–613. DOI: 10.1046/j.1525-1497.2001.016009606.x.
- Kwiatkowski L (2011) Domestic violence and the “happy family” in Northern Vietnam. *Anthropology Now* 3(3): 20–28. DOI: 10.1080/19492901.2011.11728331.
- Lefkovic E, Baji I and Rigó J (2014) Impact of maternal depression on pregnancies and on early attachment. *Infant Mental Health Journal* 35(4): 354–365. DOI: 10.1002/imhj.21450.
- Moret C and Briley M (2011) The importance of norepinephrine in depression. *Neuropsychiatric Disease and Treatment* 7(Suppl. 1): 9–13. DOI: 10.2147/NDT.S19619.
- Murray L, Dunne MP, Van Vo T, et al. (2015) Postnatal depressive symptoms amongst women in Central Vietnam: A cross-sectional study investigating prevalence and associations with social, cultural and infant factors. *BMC Pregnancy and Childbirth* 15: 234. DOI: 10.1186/s12884-015-0662-5.
- Ngo VT, Gammeltoft T, Nguyen HTT, et al. (2018) Antenatal depressive symptoms and adverse birth outcomes in Hanoi, Vietnam. *PLoS One* 13(11): e0206650. DOI: 10.1371/journal.pone.0206650.
- Nguyen TQ, Bandeen-Roche K, Bass JK, et al. (2016) A tool for sexual minority mental health research: The Patient Health Questionnaire (PHQ-9) as a depressive symptom severity measure for sexual minority women in Viet Nam. *Journal of Gay & Lesbian Mental Health* 20(2): 173–191. DOI: 10.1080/19359705.2015.1080204.
- O'Connor E, Rossom RC, Henninger M, et al. (2016) Primary care screening for and treatment of depression in pregnant and postpartum women: Evidence report and systematic review for the US Preventive Services Task Force. *JAMA* 315(4): 388–406. DOI: 10.1001/jama.2015.18948.
- Okagbue HI, Adamu PI, Bishop SA, et al. (2019) Systematic review of prevalence of antepartum depression during the trimesters of pregnancy. *Open Access Macedonian Journal of Medical Sciences* 7(9): 1555–1560. DOI: 10.3889/oam-jms.2019.270.
- Pariante CM (2014) Depression during pregnancy: Molecular regulations of mothers' and children's behaviour. *Biochemical Society Transactions* 42(2): 582–586. DOI: 10.1042/BST20130246.
- Plant DT, Pawlby S, Sharp D, et al. (2016) Prenatal maternal depression is associated with offspring inflammation at 25 years: A prospective longitudinal cohort study. *Translational Psychiatry* 6(11): e936. DOI: 10.1038/tp.2015.155.
- Ponting C, Mahrer NE, Zelcer H, et al. (2020) Psychological interventions for depression and anxiety in pregnant Latina and Black women in the United States: A systematic review. *Clinical Psychology & Psychotherapy* 27(2): 249–265. DOI: 10.1002/cpp.2424.
- Portal of Thua Thien Hue Province (2020) Thua Thien Hue—General Information. Available at: <https://thuathienhue.gov.vn/en-us/tabid/136/language/en-US/Default.aspx/tid/General-Information/newsid/9D47356C-4CCE-48DC-9B71-A731012AEB3/cid/0529563A-7ED5-42B4-B310-A73100EEB0D7> (accessed 15 March 2020).
- Rasch V, Van TN, Nguyen HTT, et al. (2018) Intimate partner violence (IPV): The validity of an IPV screening instrument utilized among pregnant women in Tanzania and Vietnam. *PLoS One* 13(2): e0190856. DOI: 10.1371/journal.pone.0190856.
- R Core Team (2014) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rudy D and Grusec JE (2006) Authoritarian parenting in individualist and collectivist groups: Associations with maternal emotion and cognition and children's self-esteem. *Journal of Family Psychology* 20(1): 68–78. DOI: 10.1037/0893-3200.20.1.68.
- Sawyer A, Ayers S and Smith H (2010) Pre- and postnatal psychological wellbeing in Africa: A systematic review. *Journal of Affective Disorders* 123(1–3): 17–29. DOI: 10.1016/j.jad.2009.06.027.
- Schuler SR, Hoang TA, Vu SH, et al. (2006) Constructions of gender in Vietnam: In pursuit of the ‘Three Criteria’. *Culture, Health & Sexuality* 8(5): 383–394. DOI: 10.1080/13691050600887949.
- Schuler SR, Lenzi R, Hoang T-A, et al. (2016) Recourse seeking and intervention in the context of intimate partner violence in Vietnam: A qualitative study. *Journal of Family Issues* 37(8): 1151–1173. DOI: 10.1177/0192513X14539155.
- Seth S, Lewis AJ and Galbally M (2016) Perinatal maternal depression and cortisol function in pregnancy and the postpartum period: A systematic literature review. *BMC Pregnancy and Childbirth* 16(1): 124. DOI: 10.1186/s12884-016-0915-y.
- Staneva A, Bogossian F, Pritchard M, et al. (2015a) The effects of maternal depression, anxiety, and perceived stress during pregnancy on preterm birth: A systematic review. *Women and Birth: Journal of the Australian College of Midwives* 28(3): 179–193. DOI: 10.1016/j.wombi.2015.02.003.
- Staneva AA, Bogossian F and Wittkowski A (2015b) The experience of psychological distress, depression, and anxiety during pregnancy: A meta-synthesis of qualitative research. *Midwifery* 31(6): 563–573. DOI: 10.1016/j.midw.2015.03.015.

- Szegda K, Markenson G, Bertone-Johnson ER, et al. (2014) Depression during pregnancy: A risk factor for adverse neonatal outcomes? A critical review of the literature. *The Journal of Maternal-Fetal & Neonatal Medicine* 27(9): 960–967. DOI: 10.3109/14767058.2013.845157.
- Thua Thien Hue Provincial People's Committee (2020) Plan for implementing national targeted program on sustainable poverty reduction in 2020—Thua Thien Hue province No 43/KH-UBND.
- Tran TD, Nguyen H and Fisher J (2016) Attitudes towards intimate partner violence against women among women and men in 39 low- and middle-income countries. *PLoS One* 11(11): e0167438. DOI: 10.1371/journal.pone.0167438.
- Tran TN, Hanh NTT, Hinh ND, et al. (2019) Intimate partner violence among pregnant women and postpartum depression in Vietnam: A longitudinal study. *BioMed Research International* 2019: 4717485. DOI: 10.1155/2019/4717485.
- Tran TN, Nguyen HTT, Nguyen HD, et al. (2018b) Emotional violence exerted by intimate partners and postnatal depressive symptoms among women in Vietnam: A prospective cohort study. *PLoS One* 13(11): e0207108. DOI: 10.1371/journal.pone.0207108.
- Tran TN, Nguyen TTH and Gammeltoft TM (2018a) Emotional violence and maternal mental health: A qualitative study among women in northern Vietnam. *BMC Women's Health* 18(1): 58. DOI: 10.1186/s12905-018-0553-9.
- Trinh OTH, Oh J, Choi S, et al. (2016) Changes and socioeconomic factors associated with attitudes towards domestic violence among Vietnamese women aged 15–49: Findings from the Multiple Indicator Cluster Surveys, 2006–2011. *Global Health Action* 9: 29577. DOI: 10.3402/gha.v9.29577.
- Tsai AC, Tomlinson M, Comulada WS, et al. (2016) Intimate partner violence and depression symptom severity among South African women during pregnancy and postpartum: Population-based prospective cohort study. *PLoS Medicine* 13(1): e1001943. DOI: 10.1371/journal.pmed.1001943.
- Underwood L, Waldie K, D'Souza S, et al. (2016) A review of longitudinal studies on antenatal and postnatal depression. *Archives of Women's Mental Health* 19(5): 711–720. DOI: 10.1007/s00737-016-0629-1.
- Upadhyay AK, Singh A and Singh A (2019) Association between unintended births and risk of postpartum depression: Evidence from Ethiopia, India, Peru and Vietnam. *SSM—Population Health* 9: 100495. DOI: 10.1016/j.ssmph.2019.100495.
- Valdebenito S, Murray A, Hughes C, et al. (2020) Evidence for better lives study: A comparative birth-cohort study on child exposure to violence and other adversities in eight low- and middle-income countries—foundational research (study protocol). *BMJ Open* 10(10): e034986. DOI: 10.1136/bmjopen-2019-034986.
- Van den Bergh BRH, Mulder EJM, Mennes M, et al. (2005) Antenatal maternal anxiety and stress and the neurobehavioural development of the fetus and child: Links and possible mechanisms. A review. *Neuroscience and Biobehavioral Reviews* 29(2): 237–258. DOI: 10.1016/j.neubiorev.2004.10.007.
- Venanzio CD (2017) Perinatal depression screening and early treatment. *Official Journal of the Italian Society of Psychopathology* 23: 99–104.
- Vo TM, Tran QT, Le CV, et al. (2019) Depression and associated factors among infertile women at Tu Du hospital, Vietnam: A cross-sectional study. *International Journal of Women's Health* 11: 343–351. DOI: 10.2147/IJWH.S205231.
- Vo VT, Duong TKH and Hoang TD (2017) Postpartum depressive symptoms and associated factors in married women: A cross-sectional study in Danang City, Vietnam. *Frontiers in Public Health* 5: 93. DOI: 10.3389/fpubh.2017.00093.
- Vu HS, Schuler S, Hoang TA, et al. (2014) Divorce in the context of domestic violence against women in Vietnam. *Culture, Health & Sexuality* 16(6): 634–647. DOI: 10.1080/13691058.2014.896948.
- Wajid A, van Zanten SV, Mughal MK, et al. (2019) Adversity in childhood and depression in pregnancy. *Archives of Women's Mental Health*. DOI: 10.1007/s00737-019-00966-4.
- Wesselhoeft R, Madsen FK, Lichtenstein MB, et al. (2020) Postnatal depressive symptoms display marked similarities across continents. *Journal of Affective Disorders* 261: 58–66. DOI: 10.1016/j.jad.2019.09.075.
- WHO (2019) WHO | Adverse Childhood Experiences International Questionnaire (ACE-IQ). Available at: [http://www.who.int/violence\\_injury\\_prevention/violence/activities/adverse\\_childhood\\_experiences/en/](http://www.who.int/violence_injury_prevention/violence/activities/adverse_childhood_experiences/en/) (accessed 15 October 2019).
- Yang N, Gelaye B, Zhong Q, et al. (2016) Serum brain-derived neurotrophic factor (BDNF) concentrations in pregnant women with post-traumatic stress disorder and comorbid depression. *Archives of Women's Mental Health* 19(6): 979–986. DOI: 10.1007/s00737-016-0638-0.
- Zegeye A, Alebel A, Gebrie A, et al. (2018) Prevalence and determinants of antenatal depression among pregnant women in Ethiopia: A systematic review and meta-analysis. *BMC Pregnancy and Childbirth* 18(1): 462. DOI: 10.1186/s12884-018-2101-x.
- Zimet GD, Dahlem NW, Zimet SG, et al. (1988) The multidimensional scale of perceived social support. *Journal of Personality Assessment* 52(1): 30–41. DOI: 10.1207/s15327752jpa5201\_2.