



# Indian Traditional medicinal plants as a source of potent Anti-diabetic agents: A Review

Vishakha Parab Gaonkar<sup>1</sup> · Kirankumar Hullatti<sup>1</sup>

Received: 27 May 2020 / Accepted: 3 September 2020 / Published online: 9 September 2020  
© Springer Nature Switzerland AG 2020

## Abstract

**Objective** The present review aims to provide an overview of traditional medicinal plants known to be of anti-diabetic potential. **Methods** A literature search was conducted using the scientific databases including PubMed, EMBASE and google scholar and a total of fifty herbs have been described and their possible mechanism of anti-diabetic action has been mentioned. Among them, in-depth discussion on five most potent anti-diabetic herbs has been provided with respect to their mechanism of action, in-vivo studies and clinical efficacies.

**Results** The present review has highlighted the usefulness of the herbal source for the treatment and management of diabetes mellitus. With the help of previous literature published on *In-vivo* animal studies and human clinical studies; the effectiveness of *Gymnema sylvestre*, *Momordica charantia*, *Trigonella foenum graecum*, *Tinospora cordifolia* and *Curcuma longa* in the treatment and management of Diabetes has been proved.

**Conclusion** Based on this review it can be concluded that herbs can serve as more efficient, safer, and cost-effective adjuvant therapy in the management and treatment of diabetes. Further investigations mainly focusing on the isolation of phytochemicals from these herbs can lead to the discovery of newer antidiabetic agents.

**Keywords** Anti-diabetic · Diabetes mellitus · Herbs · Hyperglycaemia · Medicinal plants · Phytochemicals

## Introduction

Diabetes mellitus is one of the most prevalent diseases found in all parts of the world and is becoming a serious threat to mankind's health [1]. It is a complex heterogeneous group of metabolic disorders including hyperglycemia and is associated with the imbalance in carbohydrate, protein, and lipid metabolism [2]. According to WHO, "Diabetes mellitus is a chronic disease caused by inherited and/or acquired deficiency in production of insulin by the pancreas, or by the ineffectiveness of the insulin produced. Such a deficiency results in increased concentrations of glucose in the blood, which in turn damage many of the body's systems, in particular the blood vessels and nerves" [3]. According to the recent data by International Diabetes Federation (IDF) Atlas claims that

around 463 million adults are currently living with diabetes and estimates that there will be 578 million adults with diabetes by 2030, and 700 million by 2045 [4].

The management of diabetes mellitus is considered a global problem. In current allopathic therapy the oral hypoglycaemic agents and insulin, are subsequently used to control the diabetic conditions, however, complications associated with them, limited tolerability, cost, and other side effects reduce its wide acceptance. This could be the main reason for the shift of common people to Ayurveda form allopathic system nowadays [5].

Since ancient times traditional herbal drugs with multiple phytoconstituents and properties have been used as medicines for the treatment of a wide range of diseases [6]. Herbal medicines have been considered to be intrinsically safe, due to their natural occurrence, efficacy, and fewer side effects [7]. India has a long history of use of medicinal plants for the management of diabetes. World ethnobotanical information has reported the usage of about 800 plants for the control of diabetes mellitus, amongst them only 410 are experimentally proven for having anti-diabetic properties but the complete mechanism of action is available only for about 109 plants

✉ Kirankumar Hullatti  
kkhullatti@gmail.com

<sup>1</sup> Department of Pharmacognosy and Phytochemistry, KLE College of Pharmacy Belagavi, KLE Academy of Higher Education and Research (KAHER), Belagavi, India

[8]. The treatment of diabetes using herbs has more advantageous effects and does not cause much side effects. These herbal drugs act by different mechanisms and consequently protect the  $\beta$ -cells during the diabetic condition and reduce the amount of glucose level in the blood [9].

This review aims to provide an overview of the use of medicinal plants in the management of diabetes, focusing on their mechanism of action. Furthermore, an emphasis on the five most commonly available and potent anti-diabetic herbs has been given. These include *Gymnema sylvestre*, *Momordica charantia*, *Trigonella foenum graecum*, *Tinospora cardifolia*, and *Curcuma longa*.

## Methods

A literature search was conducted using the scientific databases including PubMed, EMBASE and google scholar. The aim was to identify published data on traditionally used medicinal plant for the treatment and management of Diabetes mellitus. The search terms used were “diabetes and plants”, “traditional plants”, “medicinal plants and diabetes”, “anti-hyperglycemic plants”, and “mechanism of anti-diabetic action”. Based on the above criteria extensive literature search was carried out and a total of fifty herbs have been described with their possible mechanism of anti-diabetic action. Amongst the fifty herbs in-depth discussion on five most potent and easily available anti-diabetic herbs has been provided with respect to their mechanism of action, in-vivo studies and clinical efficacies.

## Traditional anti-diabetic plants

Since the time of Charaka and Sushruta, traditional medicines have been used for the management of diabetes mellitus [10]. Medicinal plants have always been a valuable source of drugs and many of the currently available drugs such as aspirin, quinine, vincristine, vinblastine, and digitalis have been derived directly or indirectly from them [11]. Most of the anti-diabetic drugs derived from plants are from the phytochemical class of polyphenols, terpenoids, tannins, and steroids. These affect various metabolic cascades, which further affect the level of glucose in the human body [12].

A list of medicinal plants used traditionally for diabetes with proven anti-diabetic and related beneficial effects are compiled along with their family, active principles responsible for diabetes, mechanism of action and use (Table 1).

Amongst the fifty herbs described in Table 1 we have identified five most potent and easily available anti-diabetic herbs namely, *Gymnema sylvestre*, *Momordica charantia*, *Trigonella foenum graecum*, *Tinospora cordifolia* and *Curcuma longa*. In-depth discussion on these herbs has been

provided with respect to their mechanism of action, in vivo studies and clinical efficacies. The rationale behind selection of these five herbs owes to their traditional usage, easy availability, effectiveness and most importantly to their proven clinical significance. The selected plants are a choice of herbal medicine in the treatment of diabetes. They are also utilized by multiple pharmaceutical herbal industries and most of the herbal anti-diabetic preparation consist of these herbs. Although there are other plants which are utilized in the diabetes management as folklore medicine but their commercial utilization is less as compare to the chosen five plants. Hence a need has been felt for understanding their various mechanism of actions and efficacy in management of diabetes.

## *Gymnema sylvestre*

*Gymnema sylvestre* is an indigenous herb, belonging to the family Asclepiadaceae. It is popularly known as “gurmar” for its distinct property as sugar destroyer, it is a reputed herb in the Ayurvedic system of medicine. The plant is indigenous to western and central India, Australia, and tropical Africa [94].

## Phytochemistry of *G. sylvestre*

*G. sylvestre* is a good source of a large number of bioactive substances. The leaves contain Triterpene saponins like gymnemic acids, gymnemasaponins, and gymnemasides. Apart from this, other phytoconstituents include flavones, anthraquinones, pentatriacontane, hentriacontane,  $\alpha$  and  $\beta$ -chlorophylls, phytin, stigmasterol, dquercitol, resins, etc. The major secondary metabolites present in *Gymnema* includes Gymnemic acid. The Gymnemic acids consist of numerous members termed as gymnemic acids I–VII, gymnemasaponins, and gymnemosides A–F. Gurmarin is another essential phytoconstituent isolated from *G. sylvestre* [95].

## Mechanism of Action

Antidiabetic activity of Gymnemic acids appears to be due to a combination of mechanisms. It acts through stimulation in insulin secretion from the pancreas. It also shows a similar effect by delaying the glucose absorption in the blood. In the intestine it attaches to the receptor present in the external layer of the intestine, thereby preventing the absorption of sugar molecules by the intestine, resulting in low blood glucose levels [95]. In a study extract of the plant has showed its effectiveness in regeneration of pancreatic  $\beta$  cells [66]. Gymnemic acid the major phytochemicals present in the plant is reported to interact with glyceraldehyde-3-phosphate dehydrogenase (GAPDH), a key enzyme in glycolysis pathway [96]. Moreover, *G. sylvestre* has been reported to exhibit significant inhibitory activity against  $\alpha$ -glucosidase; Fig. 1.

**Table 1** List of Traditional plants used in management and treatment of Diabetes

Plant name	Family	Parts used	Active Principles	Mechanism of action	Uses	Reference
<i>Acacia arabica</i>	Leguminosaceae	Bark	Gallic acid, pyrocatechol, (+)-catechin, (-) epigallocatechin-7-gallate, (-) epicatechin, quercetin, (+) catechin-5-gallate.	Act as secretagogue to release insulin	Hypoglycemic activity	[13, 14]
<i>Achyranthes aspera</i>	Amaranthaceae	Leaves, seeds.	Betaine, achyranthine, $\beta$ ecdysone	Carbohydrate digestion and absorption	Hypoglycemic effect	[15, 16]
<i>Adhatoda vasica</i>	Acanthaceae	leaves	Vasicine Vasicinol	$\alpha$ -Glucosidase-inhibiting activity	Antidiabetic	[17, 18]
<i>Aegle marmelos</i>	Rutaceae	leaves	Aegelin, marmesin and marmelosin	Regeneration of pancreatic $\beta$ cells and insulin secretion	Hypoglycaemic effect	[16, 19]
<i>Ageratum conyzoides</i>	Asteraceae	leaves	Mono- and sesquiterpenes	Increase peripheral utilization of glucose	Hypoglycaemic effect	[20, 21]
<i>Allium cepa</i>	Amaryllidaceae	bulb	S-methyl cysteine sulfoxide, S-allyl cysteine sulfoxide	Stimulates pancreatic $\beta$ -cells	Hypoglycaemic effect	[22, 23]
<i>Allium sativum</i>	Amaryllidaceae	bulb	Allicin, apigenin, alliin	Stimulates pancreatic $\beta$ -cells	Antidiabetic and anti-oxidant	[16, 23, 24]
<i>Aloe barbadensis</i>	Asphodelaceae	leaves	Aloin, barbaloin, isobarbaloin, aloetic acid.	Insulin secretion and synthesis	Hypoglycemic effect.	[16, 25]
<i>Andrographis paniculata</i>	Acanthaceae	Whole plant	Andrographolide,	Regeneration of pancreatic $\beta$ cells, insulin secretion	Antidiabetic & hepatoprotective.	[26, 27]
<i>Ammonia squamosa</i>	Annonaceae	leaves	Acetogenin	Enhances insulin level from pancreatic islets, increased utilization of glucose in muscle.	Hypoglycemic and antihyperglycemic activities	[28, 29]
<i>Areca catechu</i>	Palmitaceae	Leaves, flowers, seeds	Nitrosamines, arecoline, arecaidine	Carbohydrate digestion and absorption	Hypoglycemic	[16, 30]
<i>Azadirachta indica</i>	Meliaceae	leaves	Azadirachtin, nimbolinin, nimbim, nimbidin, quercetin.	Improves the insulin signaling molecules and glucose utilization in the skeletal muscle.	Antidiabetic, Antibacterial, antioxidant	[31, 32]
<i>Bacopa monnari</i>	Serophulariaceae	Aerial part	Bacosine, brahmine, bacopaside I, II, III, IV and V.	Increase in peripheral glucose consumption	Antihyperglycemic agent	[33, 34]
<i>Bauhinia forficata</i>	Fabaceae	leaves	Kaempferitin	Glycolysis, insulinomimetic activity.	Hypoglycemic effect, antioxidant.	[35, 36]
<i>Berberis aristata</i>	Berberidaceae	Stem bark, roots, leaves	Barberin,	Glucose transport, carbohydrate digestion and absorption, DPP-IV inhibition	Hypoglycemic effect	[37, 38]
<i>Boerhavia diffusa</i>	Nyctaginaceae	leaves	Punamavine, Boeravinone A-F	Increase in hexokinase activity, increase plasma insulin level, antioxidant	Antidiabetic	[39, 40]
<i>Camellia sinensis</i>	Theaceae	leaves	Epigallocatechin-gallate, gallicolocatechin, epicatechin, (+) 1-deoxyxynojirimycin,	Free radical scavenging activity, insulinomimetic activity	Antihyperglycemic activity, antioxidant	[41, 42]
<i>Casaria esculenta</i>	Salicaceae	roots	Leucopelargonidin, Dulcitol, Beta sitosterole.	Insulin secretion	Antihyperglycemic activity	[43]
<i>Cassia auriculata</i>	Fabaceae	roots	Bis (2-ethyl hexyl) phthalate	$\alpha$ -Glucosidase-inhibiting activity	Antihyperglycemic effect	[44, 45]
<i>Centella asiatica</i>	Apiaceae	Whole plant	asiaticoside	Initiate insulin secretion, carbohydrate digestion and absorption.	Antihyperglycemic activity	[46, 47]
<i>Coccinia indica</i>	Cucurbitaceae	Aerial parts	- $\beta$ -Amyrin Acetate, Lupcol, Cucurbitacin B, Taraxerone, Taraxerol, $\beta$ -carotene, Lycopene, 1-deoxyxynojirimycin,	Inhibition of $\alpha$ -glucosidase	Hypoglycemic effect.	[48, 49]
<i>Commelina communis</i>	Commelinaceae	Leaves, stem	(2R,3R,4R,5R)2,5-bis(hydroxymethyl)-3,4-dihydropyrrrolidine		Antihyperglycemic agent.	[50, 51]
<i>Curcuma longa</i>	Zingiberaceae	rhizomes	Curcumin, termerone, germaerone, zingiberene	Inhibition of $\alpha$ -glucosidase, inhibition of GSK-3 $\beta$	Antidiabetic, Antihyperlipidemic, antioxidant	[52, 53]
<i>Cyperus rotundus</i>	Cyperaceae	Whole plant	$\alpha$ cyperone, cyperene, cyperol.	Inhibits intestinal glucose absorption and promoting glucose consumption.	Hypoglycemic agent	[54, 55]
<i>Embllica officinalis</i>	Euphorbiaceae	fruits	Gallic acid, ellagic acid, vitamin c.	Hypoglycemic, Decreases lipid peroxidation, antioxidant.	Hypoglycemic and antioxidant.	[56, 57]
<i>Encostema littorale</i>	Gentianaceae	Whole plant	Swertiamarin, apigenin, isovitexin, swertisin, saponarin, 5-o-glucosylswertisin	Glucose-induced insulin release through K(+)-ATP channel.	Hypoglycemic effect.	[58, 59]

Table 1 (continued)

Plant name	Family	Parts used	Active Principles	Mechanism of action	Uses	Reference
<i>Ficus benghalensis</i> <i>Ficus racemosa</i>	Moraceae Moraceae	Bark, leaves Bark, leaves	Leucocyanidin, pelarogonidin β-sitosterol, racemose acid, Bergenin.	Insulin secretion, glycogen synthesis Glycogenolysis and gluconeogenesis	Antidiabetic Hypoglycemic activity	[60, 61] [62, 63]
<i>Glycyrrhiza glabra</i>	Leguminosaceae	roots	Glycyrrhizin, glycyrrhizic acid liquiritin, isoliquiritin.	Potent PPAR-γ ligand binding activity thus, reduces the blood glucose level	Hypoglycemic agent.	[64, 65]
<i>Gymnema sylvestre</i>	Aselepidaceae	leaves	Gynemic acid, Stigmastrol, Gurmarin, betaine, gymnemosides.	Regeneration of pancreatic β cells, α-glucosidase inhibitor, insulin secretion	Antidiabetic agent.	[66–68]
<i>Ginkgo biloba</i>	Gimkgoceae	leaves	Kaempferol, isorhamnetin	Inhibition of α-amylase and α-glucosidase activity	Hypoglycemic agent.	[69]
<i>Mangifera indica</i> <i>Momordica charantia</i> <i>Morus indica</i> <i>Ocimum sanctum</i>	Anacardiaceae Cucurbitaceae Moraceae Lamiaceae	leaves fruits leaves leaves	Mangiferin Momordin, momordicine, charantin Chrysin, isoquercitrin Eugenol, trans-β ocimene, Carvacrol, linalool.	α-Glucosidase-inhibiting activity Insulin secretion, glycogen synthesis Insulin secretion Insulin secretion, carbohydrate digestion and absorption	Hypoglycemic agent Hypoglycemic agent Hypoglycemic agent Hypoglycemic agent.	[70] [71, 72] [73] [74, 75]
<i>Panax ginseng</i>	Araliaceae	roots	Ginsenosides Rg2, panaxan A, B, C, D, E	Regeneration of pancreatic β cells, free radical scavenging	Antihyperglycemic activity	[76, 77]
<i>Phyllanthus amarus</i>	Phyllanthaceae	leaves	Brevifolin carboxylic acid, ethyl brevifolin carboxylate	α-Amylase inhibitory activity	Hypoglycemic, Anti-oxidant activity.	[78]
<i>Pterocarpus marsupium</i> <i>Sweria chirata</i> <i>Syzygium aromaticum</i>	Leguminosaceae Gentianaceae Myrtaceae	Stem wood Whole plant Flower buds	Marsupin, pterospin, pterostilbene Amatogentin, swerchirin, chirantin Eugenol, Caryophylline	Insulinomimetic activity Stimulates insulin release from islets Insulin secretion, carbohydrate digestion and absorption	Antidiabetic Antihyperglycemic agent Hypoglycemic agent	[79] [80] [81]
<i>Syzygium cumini</i> <i>Terminalia arjuna</i>	Myrtaceae Comberetaceae	Bark, seeds Stem bark	Jambosine, jambolin, anthocyanins. Arjunic acid, arjunolic acid, gallic acid.	α-Glucosidase-inhibiting activity Stimulates insulin release from islets	Anti-hyperglycemic Hypoglycemic activity	[82] [83]
<i>Terminalia chebula</i>	Comberetaceae	fruits	Gallic acid, chebulic acid, chebulannin, ellagic acid, chebulagic acid, chebulinic acid	Secretion of insulin from the β-cells.	Hypoglycemic activity	[84]
<i>Terminalia belerica</i>	Comberetaceae	fruits	β-sitosterol, gallic acid, ellagic acid, ethyl gallate, chebulagic acid.	Insulin secretion, carbohydrate digestion and absorption	Hypoglycemic activity	[85]
<i>Tinospora cardifolia</i>	Memispermaceae	Leaves and stem	Tinosporine, cordifolide, tinosporide, Barberin.	α-Glucosidase-inhibiting activity, glycolysis	Antidiabetic agent.	[86, 87]
<i>Trigonella foenum graecum</i>	Fabaceae	seeds	Trigonellin, Fenugreekine.	Regeneration of pancreatic β cells, insulin secretion	Antidiabetic activity.	[88, 89]
<i>Vinca rosea</i>	Apocynaceae	Whole plant	Catharanthine, vindoline, vindolinene vinblastine, vincristine	Regeneration of pancreatic β cells, insulin release	Hypoglycemic activity	[90]
<i>Vitis vinifera</i>	Vitaceae	Leaves, stem	E-resveratrol, E-ε-viniferin, anthocyanins.	Insulinomimetic activity	Anti-hyperglycemic activity	[91]
<i>Withania somnifera</i> <i>Zingiber officinalis</i>	Solanaceae Zingiberaceae	Leaves, roots rhizomes	Withaferin A, withanolides Gingerol, shoagol, zingerone.	Insulin release from pancreatic β cells Increase insulin level & decrease fasting glucose level	Hypoglycemic activity Hypoglycemic activity	[92] [93]

## Antidiabetic effects of *Gymnema sylvest*

Various in vivo animal studies and clinical experiments have repeatedly shown hypoglycemic effects of leaves of *G. sylvest*. In a study, the methanol extract of *G. sylvest* leaf and callus showed pronounced anti-diabetic activity through the regeneration of  $\beta$ -cells [97]. Sugihara Y., et al. reported the antihyperglycemic effect of gymnemic acid IV, isolated from the leaves of *G. sylvest*. The study reported that gymnemic acid IV decreased blood glucose levels by 13.5–60.0% within 6 hours of administration in comparison with glibenclamide, also increased plasma insulin levels was observed in STZ-diabetic mice at a concentration of 13.4 mg/kg due to gymnemic acid IV [66].

## *Momordica charantia*

*Momordica charantia* also known as karela, the bitter melon is a flowering vine, belonging to family Cucurbitaceae. The herb is commonly used by Ayurvedic and other traditional systems of medicine as an anti-diabetic agent. The plant is widely cultivated in Asia, India, East Africa, and South America [72].

## Phytochemistry of *M. charantia*

Bitter melon is rich in constituents such as glycosides, saponins, alkaloids, reducing sugars, resins, phenolic constituents, fixed oil, and free acids [98]. The main phytoconstituents present in *M. charantia* are charantine, charine, momordin, cucurbitin, cucurbitacin, momordin, etc. [72]. Most of the anti-diabetic potential of *M. charantia* is ascribed to Charantin. The hypoglycemic activity of the compound is similar to that of insulin [99].

*M. charantia* exerts its hypoglycemic effects via multiple mechanisms. The possible mechanism of the hypoglycemic action of *M. charantia* is mainly due to insulin secretion and glycogen synthesis [71, 72]. Some studies indicate that bitter melon may stimulate glucose utilization by peripheral and skeletal muscle, inhibit intestinal glucose uptake, and increase hepatic glycogen synthesis [72]. Hsin-Yi Lo et al., reported that seed extract of *M. charantia* regulates glucose metabolism mainly via the insulin signaling pathway [100]. In an experimental study using cell-based screening assay Hsueh-ling Cheng identified and reported that triterpenoids are the potential hypoglycaemic agents responsible for anti-diabetic action of the plant, also the underlying mechanism to bring about the action was attributed to AMP-activated protein kinase [101]; Fig. 2.

## Mechanism of action

*M. charantia* exerts its hypoglycemic effects via multiple mechanisms. The possible mechanism of the hypoglycemic action of *M. charantia* is mainly due to insulin secretion and glycogen synthesis [71, 72]. Some studies indicate that bitter melon may stimulate glucose utilization by peripheral and skeletal muscle, inhibit intestinal glucose uptake, and increase hepatic glycogen synthesis [72]. Hsin-Yi Lo et al., reported that seed extract of *M. charantia* regulates glucose metabolism mainly via the insulin signaling pathway [100]. In an experimental study using cell-based screening assay Hsueh-ling Cheng identified and reported that triterpenoids are the potential hypoglycaemic agents responsible for anti-diabetic action of the plant, also the underlying mechanism to bring about the action was attributed to AMP-activated protein kinase [101]; Fig. 2.

## Antidiabetic effects of *M. charantia*

Experimental studies have confirmed the hypoglycaemic effect of *M. charantia* on various animal models. A study demonstrated the dose-dependent hypoglycaemic activity exhibited by methanolic fruit extract of *M. charantia* in alloxan-induced diabetic rats [102]. Mahmoud MF et al., studied the





Fig. 1 Schematic representation of Probable molecular mechanism for anti-diabetic effect of *G. sylvest*



**Fig. 2** Schematic representation of Probable molecular mechanism for anti-diabetic effect of *M. charantia*

antidiabetic activity of *M. charantia* fruit juice in streptozotocin-induced diabetic rats [103]. In an investigation study Joo-Hui Han et al., isolated four new cucurbitane-type triterpenoids (C1–C4) from the ethanol extract of *M. charantia* and investigated whether the compounds affect insulin sensitivity both in vitro and in vivo models. The results reported significant decreases in blood glucose level and enhanced glycogen storage by compound C2 in STZ-injected mice [104].

### *Trigonella foenum graecum*

*Trigonella foenum graecum* also known as fenugreek is used primarily as an alternative therapy for diabetes. A member of the Fabaceae family, the plant cultivated in India and North African countries. The herb has a long history of usage as a potent anti-diabetic agent in Ayurvedic and folklore medicine [94, 105].

#### Phytochemistry of *T. foenum graecum*


The phytochemical studies have largely been focused on seeds. The main chemical constituents of seeds are alkaloids approximately 36%, steroidal saponins, mucilage, fibers [105]. Among the alkaloid content of fenugreek seed, Trigonelline is major phytoconstituents which is responsible for most of the activity of the herb. The mucilage (25–30%) is mostly a galactomannan. Steroidal saponins such as diosgenin and yamogenin constitute about 0.1–2.2%. Fenugreekine a saponin peptide ester is also present in the seeds. The free amino acids in the seeds are present as 4-hydroxyisoleucine, which is reported to have directly stimulated insulin [94, 106].

#### Mechanism of action

The mode of action of the herb is through Regeneration of pancreatic  $\beta$  cells and insulin secretion [88, 89]. A study reported the inhibitory role of Trigonelline on the activity of glycogen synthase kinase isoforms in the regulation of glycogen metabolism, to bring about the hypoglycaemic action [106]. Trigonelline has also been found to enhance glucose and lipid hemostasis via the improvement of the insulin signaling pathway [107]. Another possible mode of action of *T. foenum graecum*, are the Inhibition of glucose uptake, GLUT-4 translocation, and improved insulin resistance [108, 109]; Fig. 3.

#### Antidiabetic effects of *T. foenum graecum*

The pharmacological studies have proven the anti-diabetic potential of various extracts of *T. foenum graecum*. In a study, the ethanolic extract of fenugreek seeds was investigated for anti-diabetic action on streptozotocin-induced diabetic rats. The results demonstrated a significant decrease in serum glucose, total cholesterol, triacylglycerol, while an increase in serum insulin in diabetic rats was observed [110]. Shah et al., studied the hypoglycaemic effect of Trigonelline in alloxan-induced diabetic mice. They reported a reduction in blood glucose level and identified the presence of islet cells in the pancreatic duct suggesting its beneficial effect on  $\beta$  cells [111]. Multiple studies on seed extracts, raw powder, and active constituents by investigators have demonstrated the hypoglycaemic action of the herb, confirming its use as a potent herbal remedy for the management of diabetes.



**Fig. 3** Schematic representation of Probable molecular mechanism for anti-diabetic effect of *T. foenum-graecum*

### *Tinospora cordifolia*

*Tinospora cordifolia* (Willd.) Miers, belonging to family Menispermaceae, is a potent herb used to combat diabetes. The herb reported to possess anti-diabetic activity in Ayurvedic literature and is present in many Ayurvedic formulations. The herb is commonly known as Guduchi and is indigenous to the tropical areas of India, Myanmar, and Sri Lanka [94].

#### Phytochemistry of *T. cordifolia*

*T. cordifolia* consists of a variety of phytoconstituents belonging to different classes such as alkaloids, glycosides, steroids diterpenoid, sesquiterpenoid, phenolics, proteins, etc. The major active constituents responsible for the anti-diabetic effect belongs to a class of alkaloids; these consist of Berberine, Palmatine, Tembetarine, Magnoflorine, Tinosporin. The other constituents present are Tinocordiside, Tinocordifolioside, Cordioside, Cordifolioside, Tinosporon, Tinosporides, etc. [94, 112].

#### Mechanism of Action

*T. cordifolia* is reported to act by a different mechanism of action. The possible mechanism to bring about hypoglycemic action is due to inhibition of  $\alpha$ -glucosidase activity and glycolysis. In a study Chougale et al. reported the inhibitory effect of *T. cordifolia* on the  $\alpha$ -glucosidase enzyme [86]. Joladarashi et al., proved the Glucose uptake-stimulatory

effect of stem extracts of *T. cordifolia* by conducting experiments with Ehrlich ascites tumor cell model. The study reported the amelioratory effect of *T. cordifolia* on GLUT 1 and GLUT 3 transporters involved in basal glucose uptake; suggesting it as a possible mechanism to exert the hypoglycemic effect [113]. Another study reported the insulin-releasing, insulin-sensitizing, and inhibition of gluconeogenesis as the possible mechanism exhibited by an alkaloidal fraction of *T. cordifolia* [114]; Fig. 4.

#### Antidiabetic effects of *T. cordifolia*

The stem has been the maximum investigated part of the plant for its anti-diabetic activity. It has been reported that methanol extract of *T. cordifolia* significantly reduces the fasting blood glucose levels in streptozotocin-induced diabetic rats. Improvement in the insulin and C-peptide levels were also reported which indicated the regeneration of  $\beta$  cells in the pancreas [115]. Manikkam et al. isolated a polysaccharide from methanolic extract of *T. cordifolia* stem and demonstrated the  $\beta$ -cell regenerative property of the isolated polysaccharide in streptozotocin-induced diabetic rats suggesting its usage as a potent hypoglycemic agent [116].

### *Curcuma longa*

*Curcuma longa* Linn, belonging to family Zingiberaceae, is reported as a potent herb in Ayurveda system of medicine to combat diabetes. It is commonly known as Turmeric, Haldi,




Fig. 4 Schematic representation of Probable molecular mechanism for anti-diabetic effect of *T. cordifolia*

Haridra, etc. The herb is native to India and is widely cultivated particularly in West Bengal, Tamil Nadu, and Maharashtra [94].

**Phytochemistry of *C. longa***

The rhizomes of *C. longa* consist of a large number of phenolic compounds. Curcuminoids are the major active constituents present in the rhizomes. Curcuminoids are the mixture of three related compounds namely Curcumin,

Demethoxycurcumin, and Bisdemethoxycurcumin among this curcumin constitute about 60% of total curcuminoids. Curcumin is a major active principle responsible for most of the biological activity of *C. longa* [94].

**Mechanism of action**

*C. longa* is known to exert the hypoglycemic action via different mechanisms, of which the most common being the inhibition of  $\alpha$ -glucosidase and  $\alpha$ -amylase enzyme [52, 117,




Fig. 5 Schematic representation of Probable molecular mechanism for anti-diabetic effect of *C. longa*



**Table 2** Clinical studies on Anti-diabetic efficacy of selected five plants

Author (year)	Plant name	Study	Duration	Dose	Number (Cases/ Control)	Outcome	References
Gaytan Martinez L. et al., (2020)	<i>G. sylvestris</i>	RCT	12 weeks	600 mg/day (GS capsule)	15/15	↓ 2-h OGTT (p = 0.003) and A1C (p = 0.025), ↑ insulin sensitivity	[122]
Al-Romaiyan A. et al., (2010)	<i>G. sylvestris</i>	Cohort study	2 months	1 gm/day (Novel GS extract)	11	↓ blood glucose	[123]
Nanda Kumar S. et al., (2010)	<i>G. sylvestris</i>	quasi-experimental design	3 months	500 mg/day (GS capsule)	39/19	↑ plasma insulin and C-peptide levels ↓ blood glucose (fasting and post-prandial), and glycated hemoglobin	[124]
Trakoon-osot W. et al., (2013)	<i>M. charantia</i>	RCT	16 weeks	6 g/day of MC dried-fruit pulp	19/19	↓ A1C from baseline (p = 0.042, ↓ of total advanced glycation end products (AGEs) in serum (p = 0.028)	[125]
Fuangchan A. et al., (2011)	<i>M. charantia</i>	Multicentric double-blind RCT	4 weeks	G1-500 mg/day, G2-1000 mg/day, G3-2000 mg/day, dry fruit pulp (G4-1000 mg/day metformin)	G1 = 33 G2 = 32 G3 = 31 G4 = 33	↓ fructosamine levels in G3 and G4.	[126]
Lim ST. et al., (2010)	<i>M. charantia</i>	RCT		G1- 60 mg/kg/day G2-80 mg/kg/day G3-100 mg/kg/day G4 = Placebo	G1 = 10 G2 = 10 G3 = 10 G4 = 10	G3 showed a more rapid (15 minutes) stimulation of insulin secretion than placebo	[127]
Najdi RA. et al. (2019)	<i>T. foenum graecum</i>	RCT	12 weeks	2 gm/day TGF	6/6	↑ fasting insulin level (P = 0.04).	[128]
Verma N. et al., (2016)	<i>T. foenum graecum</i>	Multicentric double-blind RCT	3 month	1000 mg/day Fenfuro (TGF seed extract) capsule	154	↓ fasting plasma, post-prandial blood sugar levels and HbA1c levels. ↑ fasting and post-prandial C-peptide levels.	[129]
Kumar V. et al., (2015)	<i>T. cordifolia</i>	RCT	15 days	50 mg/ kg body weight/ day TC stem powder	90	↓ fasting blood sugar	[130]
Roy K. (2015)	<i>T. cordifolia</i>	RCT	2 Months	500 mg/day encapsulated stem of TC	29/30	↓ HbA1c levels	[131]
Rahimi HR. et al., (2016)	<i>C. longa</i>	RCT	3 months	80 mg/day Nano-cureumin (as nano-micelle)	39/41	↓ fasting blood sugar (p = 0.004) ↓ HbA1c levels (p = 0.02)	[132]
Chuengsamarn S. et al. (2012)	<i>C. longa</i>	RCT	9 months	1500 mg/day curcuminoids capsule	119/116	↓ fasting plasma glucose and HbA1c levels (p = < 0.01), better β cell functions. ↑HOMA-β (p = < 0.01)	[133]
Na LX. et al. (2012)	<i>C. longa</i>	RCT	3 months	300 mg/day curcuminoids	50/50	↓ fasting blood glucose (p < 0.01), HbA1c (p = 0.031), and insulin resistance index (HOMA-IR) (p < 0.01)	[134]

[118]. Gutierrez et al., reported that increased levels of AKT phosphorylation and GLUT4 translocation in skeletal muscles could be the possible mechanism responsible for the antidiabetic activity of curcumin [53]. Kuroda et al., reported that the hypoglycemic effect exerted by curcumin, demethoxycurcumin, bisdemethoxycurcumin, and ar-turmerone is mainly attributed to PPAR- $\gamma$  ligand-binding activity of the compound [119]. In a molecular docking study Yasser et al., reported that the hypoglycemic effects of curcumin may be due to the inhibition of glycogen synthase kinase 3 $\beta$  [120]; Fig. 5.

### Antidiabetic effect of *C. longa*

Various studies have shown the hypoglycemic activity of rhizomes of *C. longa*. Seo et al. investigated the glucose-lowering potential of curcumin in diabetic db/db mice. A significant decrease in blood glucose and HbA1c levels were observed in animals treated with curcumin. A further study reported the improvement in glucose homeostasis, glucose tolerance, and elevated plasma insulin levels by the administration of curcumin [121]. A study reported the suppression of increased blood glucose levels in Genetically Diabetic KK-A<sup>y</sup> Mice by ethanolic extract of *C. longa* [119].

### Clinical studies

Clinical trials play an important role in assessing the safety and efficacy of a particular medication in humans. Based on the literature obtained on the clinical efficacy of the selected five herbs, only recent publications from the year 2010 onwards have been identified from the database search. The details of clinical experiments for the investigation of antidiabetic effects of the five herbs has been summarized in Table 2.

### Conclusion

Diabetes mellitus is the most common endocrine disorder marked by persistent hyperglycaemia resulting from impaired insulin production or insulin resistance. Regardless of all the developments in therapeutics, diabetes still remains a major cause of morbidity and mortality in the world. Allopathic therapies available currently for the treatment of diabetes have a number of serious side effects; consequently, there is a need for investigation of more effective and safer hypoglycaemic agents. Traditional medicine and ethno-botany have an ever-emerging role to play in the treatment and management of diabetes mellitus.

The present review has highlighted the usefulness of the herbal source for the treatment and management of diabetes mellitus. Around 50 herbs known for their

usefulness in diabetes have been reviewed and a possible mechanism of the action exerted by them to bring about the anti-diabetic action has been highlighted. Among them, light has been shed upon 5 most potent antidiabetic herbs with respect to their phytochemistry, underlying mechanism of action, and anti-diabetic effect exerted by them.

From the evidences gathered from literature it is noticeable that the herbs act by various mechanisms to bring about the anti-diabetic effect. As evident from the literature a single herb displays multiple mechanisms of action for example Regeneration of pancreatic  $\beta$  cells, inhibition  $\alpha$ -glucosidase enzyme, insulin secretion, PPAR- $\gamma$  ligand binding activity, etc. this may be due to the presence of a variety of phytoconstituents in a herb. This can in turn bring about synergistic effects leading to reduction in hyperglycaemic action. As described in the review, most of the reported mechanisms of actions exerted by the herbs have been described. In addition to that, more information on *G. sylvestre*, *M. charantia*, *T. foenum graecum*, *T. cordifolia* and *C. longa* has been provided owing to their extensive utilization by herbal industries for development of anti-diabetic products. By looking at the diverse phytoconstituents, their mechanism of action, and clinical evidences it is clear that these herbs possess anti-diabetic effect.

Based on this comprehensive overview we can conclude that herbal medicines can play a pivotal role in the management and treatment of diabetes with fewer side effects. More investigations mainly focusing on the isolation of phytochemicals from these herbs can lead to the discovery of newer anti-diabetic agents.

**Acknowledgements** The author is thankful to principal KLE College of Pharmacy, Belagavi for his encouragement to write this review.

### Compliance with ethical standards

**Conflict of interest** The authors declare that they have no conflict of interest.

### References

1. Onal S, Timur S, Okutucu B, Zihnioglu F. Inhibition of  $\alpha$ -Glucosidase by Aqueous Extracts of Some Potent Antidiabetic Medicinal Herbs. *Prep Biochem Biotech*. 2005;35(1):29–36.
2. Lin Y, Sun Z. Current views on type 2 diabetes. *J Endocrinol*. 2009;204(1):1–11.
3. WHO | Diabetes mellitus. <https://www.who.int/mediacentre/factsheets/fs138/en/>. Accessed 7 Aug 2020.
4. Worldwide toll of diabetes. *Diabetesatlas.org*. 2020. <https://www.diabetesatlas.org/en/sections/worldwide-toll-of-diabetes.html>. Accessed 7 Aug 2020.
5. Parasuraman S, Thing G, Dhanaraj S. Polyherbal formulation: concept of Ayurveda. *Pharmacogn Rev*. 2014;8(16):73.

6. Calixto JB. Efficacy, safety, quality control, marketing and regulatory guidelines for herbal medicines (phytotherapeutic agents). *Braz J Med Biol Res.* 2000;33(2):179–89.
7. Prabhakar P, Kumar A, Doble M. Combination therapy: A new strategy to manage diabetes and its complications. *Phytomedicine.* 2014;21(2):123–30.
8. Prabhakar P, Doble MA. Target based therapeutic approach towards diabetes mellitus using medicinal plants. *Curr Diabetes Rev.* 2008;4(4):291–308.
9. Jeeva S, Sheebha A. A review of antidiabetic potential of ethnomedicinal plants. *Med Aromat Plants.* 2014;3(4):1–8.
10. Kashikar VS, Kotkar T. Indigenous remedies for Diabetes mellitus. *Int J Pharm Pharm Sci.* 2011;3(3):22–9.
11. Jasmin R, Ganesh kumar R, Rajaram R. Probing the mechanism of the anti-diabetic potential of terpenoids from *Elephantopus scaber* L., an Indian ethnomedical plant in STZ diabetic rats In-vivo and In-silico analysis. *Indian J Biochem Biophys.* 2018;55:384–88.
12. Zhang L, Reddy N. Bioactive molecules from medicinal herbs for life threatening diseases. *J Mol Sci.* 2018;2(4):1–11.
13. Rajvaidhya S, Nagori BP, Singh GK, Dubey BK, Desai P, Alok S, Jain SA. Review on *Acacia arabica* - An Indian medicinal plant. *Int J Pharm Sci Res.* 2012;3(7):1995–2005.
14. Hegazy GA, Alnoury AM, Gad HG. The role of *Acacia Arabica* extract as an antidiabetic, antihyperlipidemic, and antioxidant in streptozotocin-induced diabetic rats. *Saudi Med J.* 2013;34(7):727–723.
15. Vijayaraj R, Sri Kumaran N. Evaluation of in vivo antidiabetic and antioxidant activity of *Achyranthes aspera* Linn. seeds by streptozotocin induced diabetic rats. *Int J Green Pharm.* 2018;12(1):29–37.
16. Bharti S, Krishnan S, Kumar A, Kumar A. Antidiabetic phytoconstituents and their mode of action on metabolic pathways. *Ther Adv Endocrinol Metab.* 2018;9(3):81–100.
17. Gao H, Huang Y, Gao B, Li P, Inagaki C, Kawabata J. Inhibitory effect on  $\alpha$ -glucosidase by *Adhatoda vasica* Nees. *Food Chem.* 2008;108(3):965–72.
18. Kumar V, Prakash O, Kumar S, Narwal S.  $\alpha$ -glucosidase inhibitors from plants: A natural approach to treat diabetes. *Pharmacogn Rev.* 2011;5(9):19.
19. Arawwawala M, Jayaratne M. *Aegle marmelos* (L.) Correa as a potential candidate for treatment of diabetes mellitus: A review. *J Herbmed Pharmacol.* 2017;6(4):141–7.
20. Rafe MR. A review of five traditionally used anti-diabetic plants of Bangladesh and their pharmacological activities. *Asian Pac J Trop Med.* 2017;10(10):933–9.
21. Nyunai N, Njikam N, Abdennebi EH, Mbafor JT, Lamnaouer D. Hypoglycaemic and antihyperglycaemic activity of *Ageratum conyzoides* L. in rats. *Afr J Tradit Complement Altern Med.* 2009;6(2):123–30.
22. Vu NK, Kim CS, Ha MT, Mai Ngo TQ, Park SE, Kwon H, et al. Antioxidant and antidiabetic activities of flavonoid derivatives from the outer skins of *Allium cepa* L. *J Agric Food Chem.* 2020. <https://doi.org/10.1021/acs.jafc.0c02122>.
23. Sabiu S, Madende M, Ajao AA, Aladodo RA, Nurain IO, Ahmad JB. The Genus *Allium* (Amaryllidaceae: Alloioideae): Features, phytoconstituents, and mechanisms of antidiabetic potential of *Allium cepa* and *Allium sativum*. In: Watson RR, Preedy VR, eds. *Bioactive food as dietary interventions for diabetes.* Cambridge: Academic; 2019. pp. 137–154.
24. Mamun MA, Hasan N, Shirin F, Belal MH, Khan MAJ, Tasnin MN, et al. Antihyperglycemic and antihyperlipidemic activity of ethanol extract of garlic (*Allium sativum*) in streptozotocin-induced diabetic mice International. *J Med Health Res.* 2017;3(2):63–6.
25. Yagi A, Hegazy S, Kabbash A, Wahab EA. Possible hypoglycemic effect of *Aloe vera* L. high molecular weight fractions on type 2 diabetic patients. *Saudi Pharm J.* 2009;17:209–15.
26. Komalasari T, Harimurti S. A Review of The Anti-diabetic Activity of *Andrographis paniculata* (Burm. f.) Nees based in-vivo Study. *Int J Med Sci Public Health.* 2015;4(4):265–263.
27. Akhtar M, Bin Mohd Sarib M, Ismail I, Abas F, Ismail A, Lajis N. Anti-diabetic activity and metabolic changes induced by *Andrographis paniculata* plant extract in obese diabetic rats. *Molecules.* 2016;21(8):1026.
28. Tomar RS, Sisodia SS. Antidiabetic activity of *Annona squamosa* Linn. In alloxan – induced diabetic rats. *Int J Green Pharm.* 2014;8(4):237–41.
29. Davis JA, Sharma S, Mitra S, Sujatha S, Kanaujia A, Shukla G, et al. Antihyperglycemic effect of *Annona squamosa* hexane extract in type 2 diabetes animal model: PTP1B inhibition, a possible mechanism of action. *Indian J Pharmacol.* 2012;44(3):326–32.
30. Mannan N, Boucher BJ, Evans SJW. Increased waist size and weight in relation to consumption of *Areca catechu* (betel-nut); a risk factor for increased glycaemia in Asians in East London. *Br J Nutr.* 2000;83:267–75.
31. Satyanarayana K, Sravanthi K, Shaker IA, Ponnulakshmi R. Molecular approach to identify antidiabetic potential of *Azadirachta indica*. *J Ayurveda Integr Med.* 2015;6(3):165–74.
32. Omnia M. Molecular and biochemical effect of neem extract on experimental Diabetes. *IOSR J Appl Chem.* 2014;7(7):24–9.
33. Ghosh T, Maity T, Singh J. Antihyperglycemic activity of bacosine, a triterpene from *Bacopa monnieri*, in alloxan-induced diabetic rats. *Planta Med.* 2011;77:804–8.
34. Kishore L, Kaur N, Singh R. Renoprotective effect of *Bacopa monnieri* via inhibition of advanced glycation end products and oxidative stress in STZ-nicotinamide-induced diabetic nephropathy. *Ren Fail.* 2016. <https://doi.org/10.1080/0886022X.2016.1227920>.
35. Franco RR, Alves VH, Zabisky LF, Justino AB, Martins MM, Saraiva AL, et al. Antidiabetic potential of *Bauhinia forficata* Link leaves: a non-cytotoxic source of lipase and glycoside hydrolases inhibitors and molecules with antioxidant and antiglycation properties. *Biomed Pharmacother.* 2020;123:1–8.
36. Jorge AP, Horst H, de Sousa E, Pizzolatti MG, Silva FR. Insulinomimetic effects of kaempferitrin on glycaemia and on 14C-glucose uptake in rat soleus muscle. *Chem Biol Interact.* 2004;149(2–3):89–96.
37. Al masri IM, Mohammad MK, Tahaa MO. Inhibition of dipeptidyl peptidase IV (DPP-IV) is one of the mechanisms explaining the hypoglycemic effect of berberine. *J Enzyme Inhib Med Chem.* 2009;24:1061–6.
38. Cicero AFG, Tartagni E. Antidiabetic Properties of Berberine: From Cellular Pharmacology to Clinical Effects. *Hosp Pract.* 2012;40(2):57–63.
39. Alam P, Shahzad N, Gupta AK, Mahfoz AM, Bamagous GA, Al-Ghamdi SS, et al. Anti-diabetic Effect of *Boerhavia diffusa* L. root extract via free radical scavenging and antioxidant mechanism. *Toxicol Environ Health Sci.* 2018;10:220–7.
40. Brahima M, Eugene A, Justin B, Madjid A, Machioud S, Rodrigue A, et al. The Effect of Methanolic Leaf Extract of *Boerhavia diffusa* Linn. (Nictaginaceae) on the Activities of Antidiabetic, Anti-inflammatory and Antioxidant Enzymes in Experimental Diabetes. *J Pharm Res Int.* 2018;24(5):1–25.
41. Wang H, Shi S, Bao B, Li X, Wang S. Structure characterization of an arabinogalactan from green tea and its anti-diabetic effect. *Carbohydr Polym.* 2015;124:98–108.
42. Hininger-Favier I, Benaraba R, Coves S, Anderson RA, Roussel AM. Green Tea Extract Decreases Oxidative Stress and Improves Insulin Sensitivity in an Animal Model of Insulin Resistance, the

- Fructose-Fed Rat. *J Am Coll Nutr.* 2009. <https://doi.org/10.1080/07315724.2009.10718097>.
43. Prakasam A, Sethupathy S, Pugalendia KV. Antiperoxidative and antioxidant effects of *Casearia esculenta* root extract in streptozotocin induced diabetic rats. *Yale J Biol Med.* 2005;78:15–23.
  44. Govindasamy C, Al-Numair KS, Alsaif MA, Viswanathan KP. Influence of 3-hydroxymethyl xylitol, a novel antidiabetic compound isolated from *Casearia esculenta* (Roxb.) root, on glyco-protein components in streptozotocin-diabetic rats. *J Asian Nat Prod Res.* 2011;13(8):700–6.
  45. Benalla W, Bellahcen S, Bnouham M. Antidiabetic medicinal plants as a source of alpha glucosidase inhibitors. *Curr Diabetes Rev.* 2010;6(4):247–54.
  46. Kabir AU, Samad MB, D'Costa NM, Akhter F, Ahmed A, Hannan JM. Anti-hyperglycemic activity of *Centella asiatica* is partly mediated by carbohydrase inhibition and glucose-fiber binding. *BMC Complement Altern Med.* 2014;18:14–31.
  47. Nugroho AE, Lindawati NY, Herlyanti K, Widyastuti L, Pramono S. Anti-diabetic effect of a combination of andrographolide-enriched extract of *Andrographis paniculata* (Burm f.) Nees and asiaticoside-enriched extract of *Centella asiatica* L. in high fructose-fat fed rats. *Indian J Exp Biol.* 2013;51(12):1101–8.
  48. Jamwal A, Kumar S. Antidiabetic activity of isolated compound from *Coccinia indica*. *Indian J Pharm Educ.* 2019;53(1):151–9.
  49. Kohli S, Kumar PN. Combined effect of *Coccinia indica* leaf extract with acarbose in type II diabetes induced neuropathy in rats. *J Innov Pharm Biol Sci.* 2014;1(2):77–87.
  50. Youn JY, Park HY, Cho KH. Anti-hyperglycemic activity of *Commelina communis* L.: inhibition of alpha-glucosidase. *Diabetes Res Clin Pract.* 2004;66:149–55.
  51. Shibano M, Kakutani K, Taniguchi M, Yasuda M, Baba K. Antioxidant constituents in the dayflower (*Commelina communis* L.) and their alpha-glucosidase-inhibitory activity. *J Nat Med.* 2008;62(3):349–53.
  52. Riyaphan J, Jhong C, Lin S, Chang C, Tsai M, Lee D, et al. Hypoglycemic efficacy of docking selected natural compounds against  $\alpha$ -Glucosidase and  $\alpha$ -Amylase. *Molecules.* 2018;23:2260.
  53. Gutierrez V, Campos M, Arcaro C, Assis R, Baldan-Cimatti H, Peccinini R, et al. Curcumin Pharmacokinetic and Pharmacodynamic Evidences in Streptozotocin-Diabetic Rats Support the Antidiabetic Activity to Be via Metabolite(s). *Evid Based Complement Alternat Med.* 2015;1–13.
  54. Singh P, Khosa RL, Mishra G, Jha KK. Antidiabetic activity of ethanolic extract of *Cyperus rotundus* rhizomes in streptozotocin-induced diabetic mice. *J Pharm Bioallied Sci.* 2015;7(4):289–92.
  55. Tran H, Nguyen MC, Le HT, Nguyen TL, Pham TB, Chau VM, et al. Inhibitors of  $\alpha$ -glucosidase and  $\alpha$ -amylase from *Cyperus rotundus*. *Pharm Biol.* 2013. <https://doi.org/10.3109/13880209.2013.814692>.
  56. Fatima N, Hafizur RM, Hameed A, Ahmed S, Nisar M, Kabir N. Ellagic acid in *Embllica officinalis* exerts anti-diabetic activity through the action on  $\beta$ -cell of pancreas. *Eur J Nutr.* 2015. <https://doi.org/10.1007/s00394-015-1103-y>.
  57. Nain P, Saini V, Sharma S, Nain J. Antidiabetic and antioxidant potential of *Embllica officinalis* Gaertn. leaves extract in streptozotocin-induced type-2 diabetes mellitus (T2DM) rats. *J Ethnopharmacol.* 2012;142(1):65–71.
  58. Murali B, Upadhyaya UM, Goyal RK. Effect of chronic treatment with *Enicostemma littorale* in non-insulin-dependent diabetic (NIDDM) rats. *J Ethnopharmacol.* 2002;81(2):199–204.
  59. Maroo J, Vasu VT, Aalinkeel R, Gupta S. Glucose lowering effect of aqueous extract of *Enicostemma littorale* Blume in diabetes: a possible mechanism of action. *J Ethnopharmacol.* 2002;81(3):317–20.
  60. Chikaraddy A, Maniyar Y. Evaluation and comparison of hypoglycemic activity of bark and leaf of *Ficus bengalensis* linn. in alloxan induced diabetes in albino rats. *Indian J Pharm Pharmacol.* 2017;4(3):138–42.
  61. Khanal P, Patil BM. Gene set enrichment analysis of alpha-glucosidase inhibitors from *Ficus benghalensis*. *Asian Pac J Trop Biomed.* 2019;9(6):263–70.
  62. Veerapur VP, Prabhakar KR, Thippeswamy BS, Bansal P, Srinivasan KK, Unnikrishnan MK. Antidiabetic effect of *Ficus racemosa* Linn. stem bark in high-fat diet and low-dose streptozotocin-induced type 2 diabetic rats: a mechanistic study. *Food Chem.* 2012;132(1):186–93.
  63. Amin MM, Bhakta S, Das SK. Anti-diabetic potential of *Ficus racemosa*: current state and prospect especially in the developing countries. *J Biosci Agric Res.* 2015;05(02):65–72.
  64. Karthikeson PS, Laxmi T. Anti-Diabetic Activity of *Glycyrrhiza glabra* - An In vitro Study. *Int J Pharm Sci Rev Res.* 2017;44(1):80–1.
  65. Harwansh RK, Parea KC, Pareta SK, Singh J. Pharmacological studies of *Glycyrrhiza glabra* - a review. *Pharmacologyonline.* 2011;2:1032–8.
  66. Sugihara Y, Nojima H, Matsuda H, Murakami T, Yoshikawa M, Kimura I. Antihyperglycemic Effects of Gymnemic Acid IV, a Compound Derived from *Gymnema sylvestre* Leaves in Streptozotocin-Diabetic Mice. *J Asian Nat Prod Res.* 2000;2:321–7.
  67. Chen G, Guo M. Rapid Screening for  $\alpha$ -Glucosidase Inhibitors from *Gymnema sylvestre* by Affinity Ultrafiltration-HPLC-MS. *Front Pharmacol.* 2017;8:1–8.
  68. Dholi SK, Raparla R, Kannappan. In vivo anti diabetic evaluation of gymnemic acid in streptozotocin induced rats. *Pharma Innov J.* 2014;3(7):82–6.
  69. Tanaka S, Han LK, Zheng YN, Okuda H. Effects of the flavonoid fraction from *Ginkgo biloba* extract on the postprandial blood glucose elevation in rats. *Yakugaku Zasshi.* 2004;124(9):605–11.
  70. Ngo DH, Ngo DN, Vo TT, Vo TS. Mechanism of Action of *Mangifera indica* Leaves for Anti-Diabetic Activity. *Sci Pharm.* 2019;87(13):1–12.
  71. Miura T, Itoh C, Iwamoto N, Kato M, Kawai M, Park SR, Suzuki I. Hypoglycemic activity of the fruit of the *Momordica charantia* in type 2 diabetic mice. *J Nutr Sci Vitaminol.* 2001;47(5):340–4.
  72. Joseph B, Jini D. Antidiabetic effects of *Momordica charantia* (bitter melon) and its medicinal potency. *Asian Pac J Trop Dis.* 2013;3(2):93–102.
  73. Pradeep Kumar R, Sujatha D, Mohamed Saleem TS, Madhusudhana Chetty C, Ranganayakulu D. Potential antidiabetic and antioxidant activities of *Morus indica* and *Asystasia gangetica* in alloxan-induced diabetes mellitus. *J Exp Pharmacol.* 2010;2:29–36.
  74. Raja TAR, Reddy RV, Priyadarshini, Buchineni M. An evaluation of anti-hyperglycemic activity of *Ocimum sanctum* Linn (leaves) in Wister rats. *Pharma Innov.* 2016;5(1):1–3.
  75. Patil R, Patil R, Ahirwar B, Ahirwar D. Isolation and characterization of anti-diabetic component (bioactivity-guided fractionation) from *Ocimum sanctum* L. (Lamiaceae) aerial part. *Asian Pac J Trop Med.* 2011;4(4):278–82.
  76. Qiu S, Yang WH, Yao CL, Shi XJ, Li JY, Lou Y, et al. Malonylginsenosides with Potential Antidiabetic Activities from the Flower Buds of *Panax ginseng*. *J Nat Prod.* 2017;80:899–908.
  77. Kang OH, Shon MY, Kong R, Seo YS, Zhou T, Kim DY. Antidiabetic effect of black ginseng extract by augmentation of AMPK protein activity and upregulation of GLUT2 and GLUT4 expression in db/db mice. *BMC Complement Alter M.* 2017;17(341):1–11.
  78. Adedapo AA, Ofuegbe SO. The evaluation of the hypoglycemic effect of soft drink leaf extract of *Phyllanthus amarus*

- (Euphorbiaceae) in rats. *J Basic Clin Physiol Pharmacol.* 2014;25(1):47–57.
79. Pant DR, Pant ND, Saru DB, Yadav UN, Khanal DP. Phytochemical screening and study of antioxidant, antimicrobial, antidiabetic, anti-inflammatory and analgesic activities of extracts from stem wood of *Pterocarpus marsupium* Roxburgh. *J Intercol Ethnopharmacol.* 2017;6(2):170–6.
  80. Kavitha KN, Dattatri AN. Experimental Evaluation of antidiabetic activity of *Swertia Chirata* – Aqueous Extract. *J Pub Health Med Res.* 2013;1(2):71–5.
  81. Kuroda M, Mimaki Y, Ohtomo T, Yamada J, Nishiyama T, Mae T, et al. Hypoglycemic effects of clove (*Syzygium aromaticum* flower buds) on genetically diabetic KK-Ay mice and identification of the active ingredients. *J Nat Med.* 2012;66(2):394–9.
  82. Kumar A, Ilavarasan R, Jayachandran T, Deecaraman M, Aravindan P, Padmanabhan N, et al. Anti-diabetic activity of *Syzygium cumini* and its isolated compound against streptozotocin-induced diabetic rats. *J Med Plant Res.* 2008;2(9):246–9.
  83. Barman S, Das S. Hypoglycemic effect of ethanolic extract of bark of *Terminalia arjuna* Linn. In normal and alloxan-induced noninsulin-dependent diabetes mellitus albino rats. *Int J Green Pharm.* 2012;6(4):279–284.
  84. Akram N, Srivastava M, Mishra MK. Antidiabetic Activity of root of *Terminalia chebula* on alloxan induced diabetic rat. *World J Pharm Med Res.* 2019;5(3):108–12.
  85. Sabu MC, Kuttan R. Antidiabetic and antioxidant activity of *Terminalia bellerica* Roxb. *Indian J Exp Biol.* 2009;47(4):270–5.
  86. Chougale AD, Ghadyale VA, Panaskar SN, Arvindekar AU. Alpha glucosidase inhibition by stem extract of *Tinospora cordifolia*. *J Enzyme Inhib Med Chem.* 2009;24(4):998–1001.
  87. Sharma R, Amin H, Galib, Prajapati PK. Antidiabetic claims of *Tinospora cordifolia* (Willd.) Miers: critical appraisal and role in therapy. *Asian Pac J Trop Biomed.* 2015;5(1):68–78.
  88. Yoshinari O, Igarashi K. Anti-Diabetic Effect of Trigonelline and Nicotinic Acid, on KK-A<sup>y</sup> Mice. *Curr Med Chem.* 2010;17:2196–202.
  89. Zhou J, Zhou S, Zeng S. Experimental diabetes treated with trigonelline: effect on  $\beta$  cell and pancreatic oxidative parameters. *Fundam Clin Pharmacol.* 2011;27(3):279–87.
  90. Ahmed MF, Kazim SM, Ghoris S, Mehjabeen SS, Ahmed SR, Ali SM, Ibrahim M. Antidiabetic activity of *Vinca rosea* extracts in alloxan-induced diabetic rats. *Int J Endocrinol.* 2010;2010:1–6.
  91. Orhan N, Aslan M, Orhan DD, Ergun F, Yeşilada E. In-vivo assessment of antidiabetic and antioxidant activities of grapevine leaves (*Vitis vinifera*) in diabetic rats. *J Ethnopharmacol.* 2006;108(2):280–6.
  92. Kumar V, Dey A, Chatterjee SS. Phytopharmacology of Ashwagandha as an Anti-Diabetic Herb, In: Kaul S., Wadhwa R, editors. *Science of Ashwagandha: Preventive and Therapeutic Potentials.* Berlin: Springer. 2017.
  93. Otunola GA, Afolayan AJ. Antidiabetic effect of combined spices of *Allium sativum*, *Zingiber officinale* and *Capsicum frutescens* in alloxan-induced diabetic rats. *Front Life Sci.* 2015. <https://doi.org/10.1080/21553769.2015.1053628>.
  94. Khare CP. *Indian Medicinal Plants An Illustrated Dictionary.* Berlin: Springer. 2008.
  95. Kanetkar P, Singhal R, Kamat M. *Gymnema sylvestre: A Memoir.* *J Clin Biochem Nutr.* 2007;41:77–81.
  96. Ishijima S, Takashima T, Ikemura T, Izutani Y. Gymnemic acid interacts with mammalian glycerol-3-phosphate dehydrogenase. *Mol Cell Biochem.* 2008;310:203–8.
  97. Ahmed A, Rao AS, Rao M. In vitro callus and in vivo leaf extract of *Gymnema sylvestre* stimulate  $\beta$ -cells regeneration and anti-diabetic activity in Wistar rats. *Phytomedicine.* 2010;17(13):1033–9.
  98. Saeed MK, Shahzadi I, Ahmad I, Ahmad R, Shahzad K, Ashraf M, et al. Nutritional analysis and antioxidant activity of bitter gourd (*Momordica charantia*) from Pakistan. *Pharmacologyonline.* 2010;1:252–60.
  99. Budrat P, Shotipruk A. Extraction of phenolic compounds from fruits of bitter melon (*Momordica charantia*) with subcritical water extraction and antioxidant activities of these extracts. *Chiang Mai J Sci.* 2008;35(1):123–30.
  100. Lo H, Ho T, Lin C, Li C, Hsiang C. *Momordica charantia* and Its Novel Polypeptide Regulate Glucose Homeostasis in Mice via Binding to Insulin Receptor. *J Agric Food Chem.* 2013;61:2461–8.
  101. Hsueh-Ling C, Hsin-Kai H, Chi -IC, Chung-Pao T, Chang-Hung C. A Cell-Based Screening Identifies Compounds from the Stem of *Momordica charantia* that Overcome Insulin Resistance and Activate AMP-Activated Protein Kinase. *J Agric Food Chem.* 2008;56:6835–43.
  102. Nkambo W, Anyama NG, Onegi B. In vivo hypoglycemic effect of methanolic fruit extract of *Momordica charantia* L. *Afr Health Sci.* 2013;13(4):933–9.
  103. Mahmoud MF, El Ashry FE, El Maraghy NN, Fahmy A. Studies on the antidiabetic activities of *Momordica charantia* fruit juice in streptozotocin-induced diabetic rats. *Pharm Biol.* 2017;55(1):758–65.
  104. Han JH, Tuan NQ, Park MH, Quan K, Oh J, Heo KS, et al. Cucurbitane Triterpenoids from the Fruits of *Momordica Charantia* Improve Insulin Sensitivity and Glucose Homeostasis in Streptozotocin-Induced Diabetic Mice. *Mol Nutr Food Res.* 2018;62(7):1700–769.
  105. Bahmani M, Shirzad H, Mirhosseini M, Mesripour A, Rafieian-Kopaei M. A Review on Ethnobotanical and Therapeutic Uses of Fenugreek (*Trigonella foenum-graecum* L.). *Evid Based Complement Alternat Med.* 2016;21(1):53–62.
  106. Roshana Devi V, Sharmila C, Subramanian S. Molecular Docking Studies Involving the Inhibitory Effect of Gymnemic Acid, Trigonelline and Ferulic Acid, the Phytochemicals with Antidiabetic Properties, on Glycogen Synthase Kinase 3 ( $\alpha$  and  $\beta$ ). *J Appl Pharm Sci.* 2018;8(04):150–60.
  107. Amat-Alrazaq A, Aldakinah, Muhammad Y, Al-Shorbagy, Dalaal M, Abdallah, et al. Trigonelline and vildagliptin antidiabetic effect: improvement of insulin signalling pathway. *J Pharm Pharmacol.* 2017:1–9.
  108. Jaiswal N, Maurya CK, Venkateswarlu K, Sukanya P, Srivastava AK, Narender T, Tamrakar AK. 4-Hydroxyisoleucine stimulates glucose uptake by increasing surface GLUT4 level in skeletal muscle cells via phosphatidylinositol-3-kinase-dependent pathway. *Eur J Nutr.* 2012;51(7):893–8.
  109. Maurya CK, Singh R, Jaiswal N, Venkateswarlu K, Narender T, Tamrakar AK. 4-Hydroxyisoleucine ameliorates fatty acid-induced insulin resistance and inflammatory response in skeletal muscle cells. *Mol Cell Endocrinol.* 2014;395(1–2):51–60.
  110. Eidia A, Eidib M, Sokhteha M. Effect of fenugreek (*Trigonella foenum-graecum* L) seeds on serum parameters in normal and streptozotocin-induced diabetic rats. *Nutr Res.* 2007;27:728–33.
  111. Shah SN, Bodhankar SL, Badole SL, Kamble HV, Mohan V. Effect of trigonelline: An active compound from *Trigonella foenum graecum* Linn. In alloxan induced diabetes in mice. *Cell Tissue Res.* 2006;6(1):585–90.
  112. Singh SS, Pandey SC, Srivastava S, Gupta VS, Patro B, Ghosh AC. Chemistry and medicinal properties of *Tinospora cordifolia* (Guduchi). *Indian J Pharmacol.* 2003;35:83–91.
  113. Joladarashi D, Chilkunda ND, Salimath PV. Glucose uptake-stimulatory activity of *Tinospora cordifolia* stem extracts in Ehrlich ascites tumor cell model system. *J Food Sci Technol.* 2014;51(1):178–82.

114. Patel MB, Mishra S. Hypoglycemic activity of alkaloidal fraction of *Tinospora cordifolia*. *Phytomed*. 2011;18:1045–52.
115. Rajalakshmi M, Eliza J, Cecilia E, Nimala A, Daisy P. Anti-diabetic properties of *Tinospora cordifolia* stem extracts on streptozotocin-induced diabetic rats. *Afr J Pharm Pharmacol*. 2009;3(5):171–80.
116. Rajalakshmi R, Roy A. b-cell regenerative efficacy of a polysaccharide isolated from methanolic extract of *Tinospora cordifolia* stem on streptozotocin-induced diabetic Wistar rats. *Chem-Biol Interact*. 2016;243:45–53.
117. Ghani U. Re-exploring promising a-glucosidase inhibitors for potential development into oral anti-diabetic drugs: Finding needle in the haystack. *Eur J Med Chem*. 2015;103:133–62.
118. Jhong C, Riyaphan J, Lin J, Chia Y, Weng C. Screening alpha-glucosidase and alpha-amylase inhibitors from natural compounds by molecular docking in silico. *BioFactors*. 2015;41(4):242–51.
119. Kuroda M, Mimaki Y, Nishiyama T, Mae T, Kishida H, Tsukagawa M, et al. Hypoglycemic Effects of Turmeric (*Curcuma longa* L. Rhizomes) on Genetically Diabetic KK-Ay Mice. *Biol Pharm Bull*. 2005;28(5):937–9.
120. Bustanji Y, Taha MO, Almasri IM, Al-Ghoussein M, Mohammad MK, Alkhatib HS. Inhibition of glycogen synthase kinase by curcumin: Investigation by simulated molecular docking and subsequent in vitro/in vivo evaluation. *J Enzyme Inhib Med Chem*. 2009;24(3):771–8.
121. Seo K, Choi M, Jung U, Kim H, Yeo J, Jeon S, et al. Effect of curcumin supplementation on blood glucose, plasma insulin, and glucose homeostasis related enzyme activities in diabetic db/db mice. *Mol Nutr Food Res*. 2008;52:995–1004.
122. Gaytan Martinez L, Sanchez-Ruiz L, Zuniga L, Gonzalez-Ortiz M, Martinez-Abundis E. Effect of *Gymnema sylvestre* Administration on Glycemic Control, Insulin Secretion, and Insulin Sensitivity in Patients with Impaired Glucose Tolerance. *J Med Food*. 2020. <https://doi.org/10.1089/jmf.2020.0024>.
123. Al-Romaiyan A, Liu B, Asare-Anane H, Maity C, Chatterjee S, Koley N, et al. A novel *Gymnema sylvestre* extract stimulates insulin secretion from human islets in vivo and in vitro. *Phytother Res*. 2010;24(9):1370–6.
124. Nanda Kumar S, Mani UV, Mani I. An Open Label Study on the Supplementation of *Gymnema sylvestre* in Type 2 Diabetics. *J Diet Suppl*. 2010;7(3):273–82.
125. Trakoon-osot W, Sotaphun U, Phanachet P, Porasuphatana S, Udomsubpayakul U, Komindr S. Pilot study: Hypoglycemic and antiglycation activities of bitter melon (*Momordica charantia* L.) in type 2 diabetic patients. *J Pharm Res*. 2013;6:859–64.
126. Fuangchana A, Sonthisombata P, Seubnukarnb T, Chanouanc R, Chotchaisuwatd P, Sirigulsatien V, et al. Hypoglycemic effect of bitter melon compared with metformin in newly diagnosed type 2 diabetes patients. *J Ethnopharmacol*. 2011;134:422–8.
127. Lim ST, Jimeno CA, Razon-Gonzales EB, Velasquez ME. The MOCHA DM study: The Effect of *Momordica Charantia* tablets on glucose and insulin levels during the postprandial state among patients with Type 2 Diabetes Mellitus. *Philipp J Intern Med*. 2010;48(2):19–25.
128. Najdi RA, Hagrass MM, Kamel FO, Magadmi RM. A randomized controlled clinical trial evaluating the effect of *Trigonella foenum-graecum* (fenugreek) versus glibenclamide in patients with diabetes. *Afr Health Sci*. 2019;19(1):1594–601.
129. Verma N, Usman K, Patel N, Jain A, Dhakre S, Swaroop A, et al. A multicenter clinical study to determine the efficacy of a novel fenugreek seed (*Trigonella foenum-graecum*) extract (Fenfurol™) in patients with type 2 diabetes. *Food Nutr Res*. 2016;60(1):32382.
130. Kumar V, Mahdi F, Singh R, Mahdi A, Singh RK. A clinical trial to assess the antidiabetic, antidyslipidemic and antioxidant activities of *Tinospora cordifolia* in management of type – 2 Diabetes Mellitus. *Int J Pharm Sci Res*. 2016;7(2):757–64.
131. Roy K, Shah R, Iyer U. *Tinospora cordifolia* stem supplementation in diabetic dyslipidemia: an open labelled randomized controlled trial. *Funct Food Health Dis*. 2015;5(7):265–74.
132. Rahimi HR, Mohammadpour AH, Dastani M, Jaafari MR, Abnous K, Mobarhan MG, et al. The effect of nano-curcumin on HbA1c, fasting blood glucose, and lipid profile in diabetic subjects: a randomized clinical trial. *Avicenna J Phytomed*. 2016;6(5):567–77.
133. Chuengsamarn S, Rattanamongkolgul S, Luechapudiporn R, Phisalaphong C, Jirawatnotai S. Curcumin Extract for Prevention of Type 2 Diabetes. *Diabetes Care*. 2012;35(11):2121–7.
134. Na LX, Li Y, Pan HZ, Zhou XL, Sun DJ, Meng M, et al. Curcuminoids exert glucose-lowering effect in type 2 diabetes by decreasing serum free fatty acids: a double-blind, placebo-controlled trial. *Mol Nutr Food Res*. 2012;56:1–9.

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.