




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Correspondence

Anxiety, depression, and posttraumatic stress in nurses during the COVID-19 outbreak

Marialaura Di Tella ^{*,1}, Agata Benfante ¹, Lorys Castelli, Annunziata Romeo

Department of Psychology, University of Turin, Turin, Italy

Dear Editor,

During the COVID-19 pandemic, nurses and physicians faced exhausting work conditions, making ethically and morally difficult decisions, and taking health risks for themselves and their loved ones (Maben and Bridges, 2020). This situation also affected their mental health, with high levels of posttraumatic stress symptoms (PTSS) that have been reported by nurses and physicians since the early months of pandemic (Benfante et al., 2020).

The present study mainly aimed to assess the psychological impact of the COVID-19 outbreak on nurses and physicians working in Italy.

The responses of 73 nurses and 72 physicians were collected through an anonymised online survey. The current data were acquired as part of a larger project investigating the psychological impact of COVID-19 in Italian population. Participants were asked to indicate sociodemographic, clinical and work-related information, and to complete three self-report questionnaires to assess

Table 1

Sociodemographic, clinical, and work-related characteristics of nurse and physician groups. Mean (SD), percentage, *t*-test, chi-square test, and Cohen's *d* are listed.

	Nurses (N = 73)	Physicians (N = 72)	Test (df)	<i>p</i>	Effect size
Age (years)	44.3 (10.6)	41.5 (11.7)	<i>t</i> (143) = -1.51	0.134	<i>d</i> = 0.25
Sex			$\chi^2(1) = 0.63$	0.427	
Male	18 (24.7%)	22 (30.6%)			
Female	55 (75.3%)	50 (69.4%)			
Marital status			$\chi^2(1) = 0.61$	0.433	
Not in a relationship	31 (42.5%)	26 (36.1%)			
In a relationship	42 (57.5%)	46 (63.9%)			
Type of ward			$\chi^2(1) = 0.01$	0.925	
COVID-19 unit	32 (43.8%)	31 (43.1%)			
Other unit	41 (56.2%)	41 (56.9%)			
Medical condition			$\chi^2(1) = 2.14$	0.144	
Yes	23 (31.5%)	15 (20.8%)			
No	50 (68.5%)	57 (79.2%)			
Psychological aspects					
Health evaluation (VAS)*	7.11 (2.0)	8.4 (1.4)	<i>t</i> (126.64) = 4.62	<0.001	<i>d</i> = 0.77
Health concern (VAS)#	7.04 (2.5)	6.0 (2.6)	<i>t</i> (143) = -2.42	0.017	<i>d</i> = 0.40
STAI Y1	53.8 (14.5)	48.8 (12.9)	<i>t</i> (143) = -2.21	0.029	<i>d</i> = 0.37
Scored above the STAI Y1 cut-off point (≥ 41)	55 (75.3%)	48 (66.7%)	$\chi^2(1) = 1.33$	0.250	
BDI-II	12.3 (10.5)	10.7 (8.4)	<i>t</i> (143) = -0.96	0.340	<i>d</i> = 0.20
Scored above the BDI-II cut-off point (>13)	21 (28.8%)	24 (33.3%)	$\chi^2(1) = 0.35$	0.552	
PCL-5	29.1 (18.6)	20.3 (14.2)	<i>t</i> (134.52) = -3.19	0.002	<i>d</i> = 0.53
Scored above the PCL-5 cut-off point (≥ 33)	24 (32.9%)	14 (19.4%)	$\chi^2(1) = 3.38$	0.066	

SD = Standard Deviation; VAS = Visual Analogue Scale; BDI-II = Beck Depression Inventory; STAI Y1 = State-Trait Anxiety Inventory Form Y1; PCL-5 = PTSD Checklist for DSM-5.

P-values in bold indicate statistically significant differences between nurses and physicians ($p < .05$).

* Health evaluation question = 'How do you currently rate your health?'

Health concern question = 'How concerned are you about contracting COVID-19?'

* Corresponding author.

E-mail address: marialaura.ditella@unito.it (M. Di Tella).

¹ These authors contributed equally to this work.

anxiety (State-Trait Anxiety Inventory-Form Y1, STAI Y1), depressive (Beck Depression Inventory, BDI-II), and posttraumatic (PTSD Checklist for DSM-5, PCL-5) symptoms. The study was approved by the University of Turin ethics committee (Prot. n. 142069) and was conducted in accordance with the Declaration of Helsinki; all the participants gave their informed consent.

In order to investigate the psychological impact of the COVID-19 outbreak on nurses and physicians, independent *t*-tests were performed to evaluate possible differences between those two groups on sociodemographic, clinical, and work-related variables, health-related questions, and psychological variables. Results of comparisons are presented in Table 1. Significant differences between nurses and physicians were found on both Visual Analogue Scales (VAS) we used for the assessment of health-related aspects. Indeed, nurses rated lower their health and reported to be more worried about contracting COVID-19 with respect to physicians. Similarly, significant differences emerged between nurses and physicians on anxiety symptoms and PTSS, with the former reporting higher scores on the STAI Y1 and PCL-5, respectively.

These results are not surprising considering that nurses are usually one of the healthcare professional groups dealing with patients at the frontline across healthcare settings and for this reason they are continually exposed to stressful events. Particularly, during the COVID-19 pandemic, being more in contact with risky patients than physicians makes nurses more vulnerable and at risk to develop psychological distress. In line with our findings, previous studies showed that PTSS (Şahin et al., 2020; Johnson et al., 2020) and anxiety symptoms (Johnson et al., 2020) were significantly higher among nurses than physicians.

Spending much more time with patients may also explain nurses' greater fear of contracting infection. A previous study confirmed our results showing that nurses reported an increased fear of being infected, and consequently of infecting their loved ones, leading to high levels of psychological distress (Sampaio et al., 2020).

Despite this study being limited by its cross-sectional nature and its reduced sample, our results highlight that nurses experi-

ence higher levels of anxiety symptoms and PTSS, and are more worried about contracting COVID-19 compared to physicians. In conclusion, the present findings underline the importance of paying attention to the psychological health of this healthcare professional group.

Funding source

This research received no specific grant from any funding agency in the public, commercial, or not-for-profit sectors.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement

The authors would like to thank the participants involved in the study.

References

- Benfante, A., Di Tella, M., Romeo, A., Castelli, L., 2020. Traumatic stress in healthcare workers during COVID-19 pandemic: a review of the immediate impact. *Front. Psychol.* 11. <https://doi.org/10.3389/fpsyg.2020.569935>.
- Johnson, S.U., Ebrahimi, O.V., Hoffart, A., 2020. PTSD symptoms among health workers and public service providers during the COVID-19 outbreak. *PLoS One* 15 (10), e0241032. <https://doi.org/10.1371/journal.pone.0241032>.
- Maben, J., Bridges, J., 2020. Covid-19: Supporting nurses' psychological and mental health. *J. Clin. Nurs.* 29 (15–16), 2742–2750. <https://doi.org/10.1111/jocn.15307>. Epub 2020 Jun 2.
- Şahin, M.K., Aker, S., Şahin, G., Karabekiroğlu, A., 2020. Prevalence of depression, anxiety, distress and insomnia and related factors in healthcare workers during COVID-19 pandemic in Turkey. *J. Community Health* 45 (6), 1168–1177. <https://doi.org/10.1007/s10900-020-00921-w>.
- Sampaio, F., Sequeira, C., Teixeira, L., 2020. Nurses' mental health during the Covid-19 outbreak: a cross-sectional study. *J. Occup. Environ. Med.* 62 (10), 783–787. <https://doi.org/10.1097/JOM.0000000000001987>.