

# V374A KCND3 Pathogenic Variant Associated With Paroxysmal Ataxia Exacerbations

Martin Paucar, MD, PhD,\* Richard Ågren, MD, PhD,\* Tianyi Li, PhD, Simon Lissmats, Åsa Bergendal, PhD, Jan Weinberg, MD, PhD, Daniel Nilsson, PhD, Irina Savichetva, MD, PhD, Kristoffer Sahlholm, PhD, Johanna Nilsson, MD, PhD, and Per Svenningsson, MD, PhD

## Correspondence

Dr. Paucar  
martin.paucar-arce@karolinska.se

*Neurol Genet* 2021;7:e546. doi:10.1212/NXG.0000000000000546

## Abstract

### Objective

Ataxia channelopathies share common features such as slow motor progression and variable degrees of cognitive dysfunction. Mutations in potassium voltage-gated channel subfamily D member 3 (*KCND3*), encoding the K<sup>+</sup> channel, Kv4.3, are associated with spinocerebellar ataxia (SCA) 19, allelic with SCA22. Mutations in potassium voltage-gated channel subfamily C member 3 (*KCNC3*), encoding another K<sup>+</sup> channel, Kv3.3, cause SCA13. First, a comprehensive phenotype assessment was carried out in a family with autosomal dominant ataxia harboring 2 genetic variants in *KCNC3* and *KCND3*. To evaluate the physiological impact of these variants on channel currents, in vitro studies were performed.

### Methods

Clinical and psychometric evaluations, neuroimaging, and genotyping of a family (mother and son) affected by ataxia were carried out. Heterozygous and homozygous Kv3.3 A671V and Kv4.3 V374A variants were evaluated in *Xenopus laevis* oocytes using 2-electrode voltage-clamp. The influence of Kv4 conductance on neuronal activity was investigated computationally using a Purkinje neuron model.

### Results

The main clinical findings were consistent with adult-onset ataxia with cognitive dysfunction and acetazolamide-responsive paroxysmal motor exacerbations in the index case. Despite cognitive deficits, fluorodeoxyglucose (FDG)-PET displayed hypometabolism mainly in the severely atrophic cerebellum. Genetic analyses revealed the new variant c.1121T>C (V374A) in *KCND3* and c.2012T>C (A671V) in *KCNC3*. In vitro electrophysiology experiments on *Xenopus* oocytes demonstrated that the V374A mutant was nonfunctional when expressed on its own. Upon equal co-expression of wild-type (WT) and V374A channel subunits, Kv4.3 currents were significantly reduced in a dominant negative manner, without alterations of the gating properties of the channel. By contrast, Kv3.3 A671V, when expressed alone, exhibited moderately reduced currents compared with WT, with no effects on channel activation or inactivation. Immunohistochemistry demonstrated adequate cell membrane translocation of the Kv4.3 V374A variant, thus suggesting an impairment of channel function, rather than of expression. Computational modeling predicted an increased Purkinje neuron firing frequency upon reduced Kv4.3 conductance.

### Conclusions

Our findings suggest that Kv4.3 V374A is likely pathogenic and associated with paroxysmal ataxia exacerbations, a new trait for SCA19/22. The present FDG PET findings contrast with a previous study demonstrating widespread brain hypometabolism in SCA19/22.

\*These authors contributed equally to this work.

From the Department of Clinical Neuroscience (M.P., R.Å., T.L., Å.B., J.N., P.S.), Department of Molecular Medicine and Surgery (D.N.), Center for Molecular Medicine (D.N.), and Science for Life Laboratory (D.N.), Karolinska Institutet (S.L., I.S.), Stockholm; Department of Neurology (M.P., J.W., P.S.), Department of Clinical Genetics (D.N.), Department of Nuclear Medicine (I.S.), and Department of Neurophysiology (J.N.), Karolinska University Hospital (R.Å.), Stockholm; Department of Integrative Medical Biology (K.S.), Umeå University; and Department of Medical Sciences (J.N.), Örebro University, Sweden.

Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures. Funding information is provided at the end of the article.

The Article Processing Charge was funded by the authors.

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 (CC BY-NC-ND), which permits downloading and sharing the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

## Glossary

**ACZ** = acetazolamide; **Fc** = fluorescence density of Alexa Fluor 633 cytoplasm; **FDG** = fluorodeoxyglucose; **Fm** = fluorescence density of Alexa Fluor 633 membrane; **GOF** = gain of function; **KCNC3** = potassium voltage-gated channel subfamily C member 3; **KCND3** = potassium voltage-gated channel subfamily D member 3; **KChIP2** = potassium channel interacting protein 2; **LOF** = loss of function; **SCA** = spinocerebellar ataxia; **SUD** = sudden unexpected death; **WT** = wild-type.

Heterozygous mutations in potassium voltage-gated channel subfamily D member 3 (*KCND3*), encoding the voltage-dependent potassium channel Kv4.3, are associated with spinocerebellar ataxia (SCA) 19, allelic with SCA22.<sup>1–3</sup> So far, only 10 articles describing patients with SCA19/22 have been published.<sup>2–10</sup> The phenotype is variable and includes, besides ataxia, other movement disorders, variable degrees of cognitive impairment and epilepsy.<sup>1–10</sup> The only fluorodeoxyglucose (FDG) PET study on SCA19 demonstrated widespread hypometabolism.<sup>9</sup> Genetic variants in *KCND3* are also associated with cardiac arrhythmia and sudden unexpected death (SUD).<sup>4,11,12</sup>

Functional analyses of some of the *KCND3* mutations demonstrated either channel dysfunction or reduced cell membrane trafficking.<sup>2–5,10,13</sup> These abnormalities are often attenuated by co-expression of potassium channel interacting protein 2 (KChIP2).<sup>4,5,10,13</sup> Here, we report 2 members of a family affected by ataxia and intellectual disability associated with the new V374A variant in *KCND3*. In addition, the index case suffered from paroxysmal ataxia exacerbations responsive to acetazolamide (ACZ). Our results demonstrate a downscaling of current upon co-expression of wild-type (WT) Kv4.3 with the V374A mutant compared with WT Kv4.3 expressed alone, suggesting a decreased expression of functional channels in the membrane.

## Methods

Both patients provided written and oral consent to this study approved by the ethical committee in Stockholm. Clinical assessment included examination with Scale for Assessment and Rating of Ataxia, Index of Non-Ataxia Symptoms, psychometric testing, and neuroimaging. Neuroimaging included MRI and 18F-FDG PET, whereas psychometric evaluation included a comprehensive test battery (e-Methods, links.lww.com/NXG/A357). The index case was investigated with massive parallel sequencing and candidate variants verified with Sanger sequencing. Injection of complementary RNA encoding channel subunits into *Xenopus laevis* oocytes and subsequent oocyte voltage-clamp recordings were performed as described previously.<sup>14</sup> Finally, a computational analysis of the mutational effects of Kv4.3 channels on a Purkinje neuron model was performed using the program NEURON7.7.<sup>15,16</sup> Briefly, this model incorporates resurgent and nonresurgent Nav channels, Kv1, Kv3, Kv4 channels, P-type calcium channels, calcium-activated Kv channels, hyperpolarization-activated cation channels, and leak currents (e-Methods).

## Immunocytochemistry

HEK293T cells were transfected according to previous descriptions,<sup>2,3</sup> the mean fluorescence density of Alexa Fluor 633 of whole cell body, membrane (Fm), and cytoplasm (Fc), and the Fm/Fc ratio was calculated (e-Methods, links.lww.com/NXG/A357).

## Data Availability

Anonymized data will be shared by the investigators upon request.

## Results

### Clinical Findings

Two patients, mother and son, in a Swedish family were diagnosed with adult-onset slowly progressive cerebellar ataxia and bradyphrenia (figure e-1, links.lww.com/NXG/A348). Both patients were also diagnosed with learning difficulties (table 1) and have attended special schools. Common features for these patients include normal general intellectual ability despite low results in cognitive screening tests. Low performance in the following cognitive domains was evident in both patients: information processing speed, visuo-spatial episodic memory, spatial/visual construction, verbal concept formation, phonemic word fluency, and executive function (table 2). Despite these deficits, IQ scores were 90 and 95, respectively. The index case (II:1) is a 35-year-old man who suffered from paroxysmal ataxia exacerbations when exposed to accelerations/decelerations, such as when riding an elevator, bus, or train. These episodes, witnessed in 2 occasions by 2 of the authors (J.W. and M.P.), were responsive to ACZ. Patient II:1 was also treated with clonazepam and mesalazine, and the latter was for ulcerative colitis. Patient I:1 (62 years old) was not treated with any psychotropic drugs. Cerebellar atrophy was severe in the index case and milder in patient I:1; in both cases, FDG PET demonstrated moderate-to-severe hypometabolism in the cerebellum (vermis and upper cerebellar hemispheres) and subtle hypometabolism in the prefrontal cortex (figure e-2, links.lww.com/NXG/A349). We did not find evidence of parkinsonism, epilepsy, polyneuropathy, cardiac signs, or arrhythmias.

### Genetic Analyses

Pathologic CAG expansions associated with SCA 1, 2, 3, 6, 7, dentatorubropallidolulsian atrophy as well as copy number variations were ruled out in the index case. Both patients carry the variants c.1121T>C (V374A) in *KCND3* and c.2012T>C (A671V) in potassium voltage-gated channel subfamily C

**Table 1** The Main Features in Both Patients Are Ataxia and Intellectual Disability

Phenotype features	II:1 (index case)	I:1
<b>Axial ataxia</b>	Y	Y
<b>Dysarthria</b>	Y	Y
<b>Age at ataxia onset/current age, y/sex</b>	25/35/M	46/62/F
<b>Reflexes</b>	Normal	Normal
<b>Cognitive impairment</b>	Y	Y
<b>Impaired smooth pursuit/nystagmus</b>	Y/Y	Y/N
<b>Other features</b>	Paroxysmal ataxia exacerbation; increased muscle tone in legs	Mild hand posturing
<b>SARA score at last examination</b>	13	14
<b>EEG</b>	Slow activity in the temporo-parietal regions	Moderate widespread slow activity
<b>ENeG</b>	Normal	Normal
<b>MRI of the brain</b>	Marked cerebellar atrophy (vermis and superior part of the cerebellar hemispheres)	Mild atrophy in the cerebellum (vermis) and superior cerebellar peduncles
<b>FDG-PET</b>	Cerebellar hypometabolism; subtle hypometabolism in prefrontal cortex	Cerebellar hypometabolism; subtle hypometabolism in prefrontal cortex
<b>Electrocardiogram</b>	Normal	Normal
<b>Comorbidities</b>	Ulcerative colitis	Type 2 diabetes <sup>a</sup> ; hypertension

Abbreviations: ENeG = electroneurography; *KCND3* = potassium voltage-gated channel subfamily D member 3; SARA = Scale for the Assessment and Rating of Ataxia.

Range or SARA score is 0–40 points (>3 points indicate ataxia).

This syndrome is associated with the novel c.1121T>C (V374A) in *KCND3*.

<sup>a</sup> Treated with metformin and enalapril.

member 3 (*KCNC3*). The variant c.1121T>C (V374A) in *KCND3* has not been reported before, whereas c.2012T>C (A671V) in *KCNC3* has a minor allele frequency = 0.00016% in the European population (e-Methods, [links.lww.com/NXG/A357](https://links.lww.com/NXG/A357)). Heterozygous variants in *KCNC3* are associated with SCA13; therefore, we hypothesize a synergistic effect of the 2 K<sup>+</sup> channel variants.

## In Vitro Studies

### Kv4.3 V374A Disrupts Potassium Conductance

Oocytes injected with complementary RNA encoding Kv4.3 + KChIP2 (WT:V374A, 1:0) exhibited peaked currents with swift N-type inactivation (figure 1A), consistent with previous descriptions of Kv4.3 currents.<sup>17</sup> Cells expressing Kv4.3 + Kv4.3 V374A + KChIP2 (WT:V374A, 0.5:0.5) showed smaller currents maintaining the peaked character without shifting the voltage dependence of channel activation (figure 1B) or inactivation (figure e-3, [links.lww.com/NXG/A350](https://links.lww.com/NXG/A350)). In cells expressing Kv4.3 V374A + KChIP2 (WT:V374A, 0:1) alone, virtually no voltage-dependent currents were observed (figure 1C). WT:V374A, 0.5:0.5 demonstrated an average peak current reduction of ~80% as compared to WT:V374A, 1:0 (figure 1D). A simple scaling effect suggests a reduction of functional Kv4.3 channels in the membrane, with no effect of

the V374A mutation on conductance or inactivation curves (figure e-3). Conversely, expression of Kv3.3 + A671V (WT: A671V, 0.5:0.5) resulted in currents similar to those of cells expressing Kv3.3 WT (WT:A671V, 1:0) alone, whereas Kv3.3 A671V (WT:A671V, 0:1) alone exhibited slightly reduced currents compared with WT (figure e-4, [links.lww.com/NXG/A351](https://links.lww.com/NXG/A351)). Again, for A671V, no effects were observed on activation or inactivation curves (figure e-5, [links.lww.com/NXG/A352](https://links.lww.com/NXG/A352)). Application of 100 μM ACZ to oocytes expressing Kv4.3 + Kv4.3 V374A, 1:0, 0.5:0.5, or 0:1, respectively, did not modify the currents (data not shown).

### Kv4.3 Channel Modulates Purkinje Neuron Excitability

The effect of reducing the number of functional Kv4.3 channels on neuronal excitability was analyzed using a Purkinje neuron model incorporating Kv1, Kv3, and Kv4 channels.<sup>15</sup> Based on the similar kinetic properties of Kv4.3 + Kv4.3 V374A, 1:0 and 0.5:0.5, the maximum Kv4 conductance (Kv4 bar) was reduced according to the I-V relationships (figure 1D). In contrast to the WT:V374A, 1:0, the WT: V374A, 0.5:0.5 model demonstrated lower action potential amplitudes and higher firing frequency, indicating modified excitability properties<sup>18</sup> (figure e-6A, [links.lww.com/NXG/A353](https://links.lww.com/NXG/A353)). Action potential slope analysis corroborated the

**Table 2** Summary of Cognitive Features in 2 Patients From a Family With SCA19 Associated With the Novel Pathogenic Variant V374A in *KCND3*

Cognitive domain	Neuropsychological test	I:1	II:1
		At age 63 (z score)	At age 35 (z score)
Brief cognitive status examination	MoCA	12/30 (−7.0) <sup>a</sup>	21/30 (−2.91) <sup>a</sup>
General intellectual ability IQ	Ravens progressive matrices	19 (−0.67); IQ = 90	36 (−0.33); IQ = 95
Verbal episodic memory	RAVLT learning	14 (−4.72) <sup>a</sup>	40 (−0.83)
	RAVLT retention	6 (−1.48)	15 (1.67) <sup>a</sup>
Visuo-spatial episodic memory	ROCFT (delayed recall)	2 (<−3) <sup>a</sup>	10.5 (<−3) <sup>a</sup>
Working memory	Digit span/WAIS-III	4 (−2.33) <sup>a</sup>	9 (−1.67) <sup>a</sup>
Spatial/visual construction	ROCFT copy	3.5 (−8.44) <sup>a</sup>	19.5 (−4.30) <sup>a</sup>
	Block Design/WAIS-III	12 (−2.0) <sup>a</sup>	24 (−1.33)
Verbal concept formation	Similarities/WAIS-III	8 (−2.0) <sup>a</sup>	13 (−1.67) <sup>a</sup>
Phonemic word fluency	COWAS/FAS	7 (−1.65) <sup>a</sup>	11 (−2.29) <sup>a</sup>
Information processing speed	SDMT	16 (−3.56) <sup>a</sup>	19 (−5.41) <sup>a</sup>
Executive function	TMT B	200 (<−3.0) <sup>a</sup>	394 (<−14.0) <sup>a</sup>
Motor speed	FT dominant hand	5 (−3.98) <sup>a</sup>	44 (−0.95)
	FT nondominant hand	22 (−1.39)	35 (−2.18) <sup>a</sup>

Abbreviations: COWAT = Controlled Oral Word Association Test; FAS = the letters used in the phonological word fluency test; FT = Finger-tapping test; *KCND3* = potassium voltage-gated channel subfamily D member 3; MoCA = Montreal cognitive assessment; RAVLT = Rey Auditory Verbal Learning Test; ROCFT = Rey-Osterrieth Complex Figure Test; SCA = spinocerebellar ataxia; SDMT = Symbol Digit Modalities Test; TMT = Trail Making Test; WAIS-III = Wechsler Adult Intelligence Scale, third edition.

<sup>a</sup> A z score ≤−1.5 SD is compatible with a significant deficit.

dynamic differences induced by WT:V374A, 0.5:0.5, with a slower increase in voltage for the reduced amplitude action potentials (figure e-6B). The firing frequency was increased 2.9-fold by WT:V374A, 0.5:0.5, compared with 1:0, and detailed analysis of peak amplitudes revealed a 22 mV reduction for 0.5:0.5 (figure e-6, C and D). Also, peak width and hyperpolarization were increased for WT:V374A, 0.5:0.5 (figure e-6, E and F).

### Immunohistochemistry

HEK293T cells transiently transfected to express Kv4.3 WT: V374A, 1:0, 0.5:0.5, and 0:1, separately, displayed a high level of Kv4.3 protein at the cell surface. The Fm/Fc ratio of Kv4.3 WT protein remained at a similar level with or without Kv4.3 V374A (figure e-7, [links.lww.com/NXG/A354](https://links.lww.com/NXG/A354)).


### Discussion

Paroxysmal ataxia exacerbations responsive to ACZ and the *KCND3* variant V374A are new findings for SCA19. Hypometabolism limited to the cerebellum is unexpected, considering the range of cognitive deficits in this family. These findings contrast with the widespread brain hypometabolism in another SCA19 family.<sup>9</sup> In our studies, the failure of the Kv4.3 V374A mutant to produce appreciable Kv4.3 currents when expressed alone, along with the conservation of WT

channel kinetic properties in the face of a downscaling of current amplitudes, suggests a reduction of functional WT Kv4.3 channels in the membrane upon co-expression with the mutant channel. Our findings neither support a synergistic effect between Kv4.3 V374A and Kv3.3 A671V. However, some caution is required since heterologous expression systems such as *Xenopus* oocytes may not completely mirror mammalian expression and/or trafficking of ion channels.

Kv4 channels are made of up 4 alpha subunits (figure e-8, [links.lww.com/NXG/A355](https://links.lww.com/NXG/A355))<sup>19</sup>; thus, incorporation of mutant subunits that are incompatible with channel function can be expected to decrease the number of functional channels by more than 50% (i.e., a dominant-negative effect).<sup>20</sup> The Kv4.3 V374A variant is located in the pore loop between transmembrane helices S5 and S6 near the selectivity filter; the part of the protein forming the ion conduction pathway, crucial to the channel's ability to pass current. Of interest the majority of the 13 reported pathogenic variants in *KCND3* associated with SCA19 are located in the pore loop or in the nearby extracellular ends of S5 and S6 (figure e-6, [links.lww.com/NXG/A353](https://links.lww.com/NXG/A353)). Judging by our immunohistochemistry data, V374A appears to be localized to the cell membrane, contrary to other pathogenic *KCND3* variants.<sup>2,3</sup> Despite the relatively conservative amino acid substitution, the V374A variant reduced Kv4.3 peak currents by ~81% when co-expressed at a

**Figure 1** Kv4.3 WT/V374A/KChIP2 Currents and I-V Relationships



(A) Representative traces of Kv4.3 WT:V374A (1:0). (B) Representative traces of Kv4.3 WT:V374A (0.5:0.5). Inset demonstrates first 40 ms at voltage increments of 20 mV (between  $-100$  and  $+60$  mV). (C) Representative traces of WT:V374A (0:1). Inset demonstrates first 40 ms at voltage increments of 20 mV (between  $-100$  and  $+60$  mV). (D) I-V relationships for (A–C) measured at the peak current. The holding potential was  $-100$  mV. Voltage steps of 10 mV during 1,000 ms. The interpulse interval was 5 seconds. KChIP2 = potassium channel interacting protein 2; WT = wild-type.

0.5:0.5 ratio with WT Kv4.3 subunits, consistent with a dominant-negative effect. In comparison, the SCA19/22-associated Kv4.3 T352P variant, in the presence of 1:1 KChIP2, has been suggested to reduce the peak current by 52%.<sup>13</sup> Of interest the adjacent pathogenic variant Kv4.3 M373I, also in the presence of KChIP2, did not affect the expression level in oocytes as measured by current amplitudes.<sup>13</sup> Given its critical location, it may be that the V374A mutation perturbs the ability of the channel to conduct potassium ions.

KCND3 mutations are also associated with arrhythmia and SUD. We hypothesize that gain-of-function (GOF) variants, reported for SUD, mainly influence the cardiac action potential, whereas loss-of-function (LOF) variants, associated with SCA19/22, mainly modulate neuronal excitability. Purkinje neuron simulations with a 80% reduction of Kv4.3 channel conductance (e.g., 0.5:0.5, WT:V374A) demonstrated rapid firing, whereas corresponding simulations with only a slightly increased Kv4.3 channel conductance (103%; corresponding to GOF mutations) demonstrate reduced firing frequencies (and no firing with further increased Kv4 conductance). The increase in firing frequency following

Kv4.3 inhibition is in line with previous pharmacologic and genetic experiments with dopamine<sup>21</sup> and pyramidal neurons, respectively.<sup>22</sup> By contrast, a ventricular cell model<sup>23</sup> exhibited slightly shortened action potentials following  $\sim 80\%$  reduction in the number of Kv4 channels, whereas a 50%–100% increase in Kv4 channels, corresponding to GOF variants, prolonged the action potentials substantially<sup>11</sup> (figure e-9, links.lww.com/NXG/A356). Hypothetically, KChIP2 may be involved in regulating the Kv4.3 expression in pathologic conditions—for GOF (LOF) variants, reduced (increased) KChIP2 levels may decrease (increase) the number of Kv4.3 channels. Theoretically, such differences may explain why concomitant ataxia and arrhythmia have not been reported. Expression of KChIP2 has recently been demonstrated to modulate the balance between Kv4.3 and voltage-gated sodium channel 5 (Nav1.5).<sup>24</sup> However, the GOF variant Kv4.3 L450F has been reported separately in a patient with Brugada syndrome<sup>25</sup> and a patient with ataxia,<sup>4</sup> illustrating the complexity in elucidating genotype-phenotype correlations for KCND3.

Although ACZ demonstrated clinical efficacy in the index case, no in vitro effects were observed on Kv4.3 V374A.



Instead, carbonic anhydrase inhibition by ACZ and subsequent pH reduction could indirectly modulate ion channels, thereby possibly mediating the therapeutic effect and attenuating paroxysmal ataxia exacerbations. Decreased neuronal excitability in numerous SCAs associated with LOF variants (SCA27—indirectly Nav1.6, SCA13—Kv3.3, and SCA6—Cav2.1) has been described.<sup>26</sup> For Kv4.3 LOF, increased Purkinje neuron excitability was observed in the simulations. Based on this, evaluation of drugs reducing neuronal excitability, e.g., Nav channel blockers, in a Kv4.3 LOF model may provide further insights of therapeutic relevance.

## Acknowledgment

The authors are grateful to the patients for their kind participation in this study. They are also grateful to the Follin Foundation for supporting M. Paucar's research.

## Study Funding

The authors report no targeted funding.

## Disclosure

The authors report no disclosures relevant to the manuscript. Go to [Neurology.org/NG](https://www.neurology.org/NG) for full disclosures.

## Publication History

Received by *Neurology: Genetics* August 27, 2020. Accepted in final form October 27, 2020.

## Appendix Authors

Name	Location	Contribution
<b>Martin Paucar, MD, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Patient care; study concept and design; analysis and interpretation of data; drafting and revising the manuscript
<b>Richard Ågren, MD, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Study concept and design; electrophysiologic experiments; Purkinje neuron firing model and data analyses; drafting and revising the manuscript
<b>Tianyi Li, PhD</b>	Astrid Lindgren's Hospital, Stockholm, Sweden	Immunohistochemistry; revising the manuscript
<b>Simon Lissmats</b>	Karolinska Institutet, Stockholm, Sweden	Electrophysiologic experiments; revising the manuscript
<b>Åsa Bergendal, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Acquisition and interpretation of cognitive evaluations; revising the manuscript
<b>Jan Weinberg, MD, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Acquisition and interpretation of clinical data, patient care; revising the manuscript
<b>Daniel Nilsson, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Obtaining and interpretation of genetic data; revising the manuscript

## Appendix (continued)

Name	Location	Contribution
<b>Irina Savichetva, MD, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Obtaining and interpreting neuroimaging data; revising the manuscript
<b>Kristoffer Sahlholm, PhD</b>	Umeå University, Sweden	Design, analysis and interpretation of electrophysiologic data; revising the manuscript; editing of the final manuscript version
<b>Johanna Nilsson, MD, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Study concept and design; analysis and interpretation of data, drafting and revising the manuscript; editing of the final manuscript version
<b>Per Svenningsson, MD, PhD</b>	Karolinska University Hospital and Karolinska Institutet, Stockholm, Sweden	Study concept and design; analysis and interpretation of data, drafting and revising the manuscript; editing of the final manuscript version

## References

- Schelhaas HJ, Ippel PF, Hageman G, et al. Clinical and genetic analysis of a four-generation family with a distinct autosomal dominant cerebellar ataxia. *J Neurol* 2001; 248:113–120. doi: 10.1007/s004150170245.
- Duarri A, Jezierska J, Fokkens M, et al. Mutations in potassium channel *kcnd3* cause spinocerebellar ataxia type 19. *Ann Neurol* 2012;72:870–880. doi: 10.1002/ana.23700.
- Lee YC, Durr A, Majczenko K, et al. Mutations in *KCND3* cause spinocerebellar ataxia type 22. *Ann Neurol* 2012;72:859–869. doi: 10.1002/ana.23701.
- Duarri A, Nibbeling E, Fokkens MR, et al. The L450F [Corrected] mutation in *KCND3* brings spinocerebellar ataxia and Brugada syndrome closer together. *Neurogenetics* 2013;14:257–258. doi: 10.1007/s10048-013-0370-0.
- Smets K, Duarri A, Deconinck T, et al. First de novo *KCND3* mutation causes severe Kv4.3 channel dysfunction leading to early onset cerebellar ataxia, intellectual disability, oral apraxia and epilepsy. *BMC Med Genet* 2015;16:51. doi: 10.1186/s12881-015-0200-3.
- Coutelier M, Coarelli G, Monin ML, et al. A panel study on patients with dominant cerebellar ataxia highlights the frequency of channelopathies. *Brain* 2017;140:1579–1594. doi: 10.1093/brain/awx081.
- Huin V, Strubi-Vuillaume I, Dujardin K, et al. Expanding the phenotype of SCA19/22: parkinsonism, cognitive impairment and epilepsy. *Parkinsonism Relat Disord* 2017;45:85–89. doi: 10.1016/j.parkreldis.2017.09.014.
- Kurihara M, Ishiura H, Sasaki T, et al. Novel de novo *KCND3* mutation in a Japanese patient with intellectual disability, cerebellar ataxia, myoclonus, and dystonia. *Cerebellum* 2018;17:237–242. doi: 10.1007/s12311-017-0883-4.
- Paucar M, Bergendal Å, Gustavsson P, et al. Novel features and abnormal pattern of cerebral glucose metabolism in spinocerebellar ataxia 19. *Cerebellum* 2018;17:465–476. doi: 10.1007/s12311-018-0927-4.
- Hsiao CT, Fu SJ, Liu YT, et al. Novel SCA19/22-associated *KCND3* mutations disrupt human KV 4.3 protein biosynthesis and channel gating. *Hum Mutat* 2019;40:2088–2107. doi: 10.1002/humu.23865.
- Giudicessi JR, Ye D, Kritzberger CJ, et al. Novel mutations in the *KCND3*-encoded Kv4.3 K<sup>+</sup> channel associated with autopsy-negative sudden unexplained death. *Hum Mutat* 2012;33:989–997. doi: 10.1002/humu.22058.
- Takayama K, Ohno S, Ding WG, et al. A de novo gain-of-function *KCND3* mutation in early repolarization syndrome. *Heart Rhythm* 2019;16:1698–1706. doi: 10.1016/j.hrthm.2019.05.033.
- Duarri A, Lin MC, Fokkens MR, et al. Spinocerebellar ataxia type 19/22 mutations alter heterocomplex Kv4.3 channel function and gating in a dominant manner. *Cell Mol Life Sci* 2015;72:3387–3399. doi: 10.1007/s00018-015-1894-2.
- Papke RL, Stokes C. Working with OpusXpress: methods for high volume oocyte experiments. *Methods* 2010;51:121–133. doi: 10.1016/j.ymeth.2010.01.012.
- Akemann W, Knöpfel T. Interaction of Kv3 potassium channels and resurgent sodium current influences the rate of spontaneous firing of Purkinje neurons. *J Neurosci* 2006; 26:4602–4612. doi: 10.1523/JNEUROSCI.5204-05.2006.
- Carnevale NT, Hines ML. *The NEURON Book*. Cambridge: Cambridge University Press; 2006.
- Patel SP, Campbell DL, Strauss HC. Elucidating KChIP effects on Kv4.3 inactivation and recovery kinetics with a minimal KChIP2 isoform. *J Physiol* 2002;545:5–11. doi: 10.1113/jphysiol.2002.031856.

18. Izhikevich EM. *Dynamical Systems in Neuroscience*. Cambridge: MIT Press; 2007.
19. Parcej DN, Scott VE, Dolly JO. Oligomeric properties of alpha-dendrotoxin-sensitive potassium ion channels purified from bovine brain. *Biochemistry* 1992;31:11084–11088. doi: 10.1021/bi00160a018.
20. Herskowitz I. Functional inactivation of genes by dominant negative mutations. *Nature* 1987;329:219–222. doi: 10.1038/329219a0.
21. Liss B, Franz O, Sewing S, Bruns R, Neuhoff H, Roeper J. Tuning pacemaker frequency of individual dopaminergic neurons by Kv4.3L and KChip3.1 transcription. *EMBO J* 2001;20:5715. doi: 10.1093/emboj/20.20.5715.
22. Carrasquillo Y, Burkhalter A, Nerbonne JM. A-type K<sup>+</sup> channels encoded by Kv4.2, Kv4.3 and Kv1.4 differentially regulate intrinsic excitability of cortical pyramidal neurons. *J Physiol* 2012;590:3877–3890. doi: 10.1113/jphysiol.2012.229013.
23. Hund TJ, Rudy Y. Rate dependence and regulation of action potential and calcium transient in a canine cardiac ventricular cell model. *Circulation* 2004;110:3168–3174. doi: 10.1161/01.CIR.0000147231.69595.D3.
24. Clatot J, Neyroud N, Cox R, et al. Inter-regulation of K(v)4.3 and voltage-gated sodium channels underlies predisposition to cardiac and neuronal channelopathies. *Int J Mol Sci* 2020;21:4908. doi: 10.3390/ijms21144908.
25. Giudicessi JR, Ye D, Tester DJ, et al. Transient outward current (I<sub>(to)</sub>) gain-of-function mutations in the KCND3-encoded Kv4.3 potassium channel and Brugada syndrome. *Heart Rhythm* 2011;8:1024–1032. doi: 10.1016/j.hrthm.2011.02.021.
26. Hoxha E, Balbo J, Miniaci MC, et al. Purkinje cell signaling deficits in animal models of ataxia. *Front Synaptic Neurosci* 2018;10:6. doi: 10.3389/fnsyn.2018.00006.