

## CASE REPORT

# Cryptococemia in a patient with COVID-19: A case report

Mohamad Y. Khatib<sup>1</sup> | Amna A. Ahmed<sup>1</sup> | Said B. Shaat<sup>2</sup> | Ahmed S. Mohamed<sup>1</sup>  | Abdulqadir J. Nashwan<sup>1,3</sup> 

<sup>1</sup>Critical Care Medicine Department, Hazm Mebaireek General Hospital (HMGH), Hamad Medical Corporation (HMC), Doha, Qatar

<sup>2</sup>Critical Care Medicine Department, Hamad General Hospital (HGH), Hamad Medical Corporation (HMC), Doha, Qatar

<sup>3</sup>University of Calgary in Qatar, Doha, Qatar

## Correspondence

Abdulqadir J. Nashwan, Critical Care Medicine Department, Hazm Mebaireek General Hospital (HMGH), Hamad Medical Corporation (HMC), Doha, Qatar.  
Email: anashwan@hamad.qa

## Abstract

The authors urge clinicians to balance the use of Immunosuppressant drugs and tocilizumab for COVID-19 patients to avoid the development of infections like Cryptococemia, leading to death within 30 days.

## KEYWORDS

corticosteroids, COVID-19, cryptococemia, *Cryptococcus neoformans*, tocilizumab

## 1 | BACKGROUND

*Cryptococcus neoformans* infection is a known cause of meningoencephalitis in immunocompromised patients. However, it is less frequent in immunocompetent individuals. Here, we report a case of a 60-year-old COVID-19 patient with a history of several comorbidities who received tocilizumab and corticosteroids, which lead to Cryptococemia.

*Cryptococcus neoformans* are basidiomycetous; encapsulated yeasts are classified into four serotypes according to their capsule.<sup>1</sup> The way of the infection is by inhalation of basidiospore form or small, poorly encapsulated yeasts.<sup>2</sup> *Cryptococcus neoformans* infection typically happens among immunocompromised patients, and meningoencephalitis is the typical manifestation of cryptococcosis among HIV patients.<sup>3,4</sup> In immunocompetent, the infection can be subclinical and a study on postmortem immunocompetent just showed small granulomatous disease in the lung without disseminating the infection.<sup>5,6</sup> We describe a case of a patient

with COVID-19 who developed Cryptococemia while being treated by immunosuppressive treatment.

## 2 | CASE PRESENTATION

A 60-year-old man with a history of hypertension, diabetes mellitus, and ischemic heart disease admitted to the intensive care unit (ICU) with confirmed COVID-19 infection required mechanical ventilation; on the same day of admission, he was proned for many times due to low PaO<sub>2</sub>/FiO<sub>2</sub> (P/F) ratio. He received three doses of tocilizumab (400 mg/once), (600 mg/once), and (600 mg/once), respectively, due to possible cytokine storm; he received several doses of methylprednisolone as part of the COVID-19 management and hydrocortisone.

Few days after receiving tocilizumab, his bronchial alveolar lavage was positive for *Candida glabrata*; he was started on anidulafungin 200 mg q24 hours for 7 days. The patient

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2020 The Authors. *Clinical Case Reports* published by John Wiley & Sons Ltd.

was sedated and intubated for 22 days, so a tracheostomy was done. He developed acute kidney injury (AKI) after 28 days of ICU admission; hence, hemodialysis was started.

Despite that the patient was on antibiotics, his inflammatory marker is still high. Thus, full septic workups were done, and the patient was having candidemia (*Candida parapsilosis*) after one month of the admission. For which he was started on anidulafungin 200 mg q24 hours for more than 7 days. Many episodes of gastrointestinal (GI) tract bleeding complicated his course manifested as melena and per rectal bleeding, but all the episodes were managed conservatively by the gastroenterology team. The candidemia persists after 2 weeks of treatment and there was an increase in C-reactive protein (CRP) level despite that the patient was under antibiotic and antifungal cover. Hence, the ID team recommended giving antifungal (anidulafungin 100 mg) for a total of 14 days to involve an ophthalmologist to rule out fungal ophthalmitis. Still, it was not applicable and to do an echocardiogram to rule out vegetation and it was ruled out by trans-thoracic echocardiogram.

While the patient was on anidulafungin, his blood culture became positive for *Cryptococcus neoformans* after 18 days of receiving anidulafungin for previous candida infection, for which he was started on amphotericin (300 mg, q24 hours) and flucytosine (500 mg q12 hours). And there was a recommendation from the ID team to do trans esophageal echo and lumbar puncture to rule out vegetation and brain involvement, respectively, but both tests were not done due to poor prognosis and risk of bleeding because the patient had thrombocytopenia (platelet 21, RR =  $150\text{--}400 \times 10^9/L$ ). The Cryptococemia persist despite the management and he developed sepsis and die within 10 days of the Cryptococemia.

### 3 | DISCUSSION

Disseminated *Cryptococcus neoformans* infection is a serious infection that can occur in immunocompromised patients.<sup>7</sup> During the outbreak of COVID-19 disease, there were many studies about the overwhelming immune response to COVID-19 infection through the activation of a large number of T lymphocytes and mononuclear macrophages, producing cytokines like interleukin-6 (IL-6), which bind to the receptor and cause cytokine storm and severe inflammation in the lung and other tissues. Tocilizumab, as an antihuman IL-6 receptor monoclonal antibody and immunosuppressant drugs, was recommended by several studies for alleviating the inflammatory response.<sup>8</sup>

Jean and colleagues have noted in their study of 52 patients with *Cryptococcus neoformans* isolated from blood culture; the primary three predisposing for Cryptococemia

are Acquired immunodeficiency syndrome, immunosuppressive therapy, and decompensated liver cirrhosis.<sup>9</sup> And in our case, the *Cryptococcus neoformans* was isolated from the blood culture. And according to Passerini M et al that *Cryptococcus neoformans* can affect different organs as meningoencephalitis, which is the most common, followed by pulmonary and cutaneous disease.<sup>10</sup>

According to Jean et al that patients with positive blood culture for *Cryptococcus neoformans* developed sepsis by 81%, including those developed septic shock, the lumbar puncture (LP) was positive in 61.5% of patients with positive blood culture for *Cryptococcus neoformans* and 84% of those with LP positive showed meningeal involvement, in our case LP was recommended but not done because the patient was critically sick and unstable. Also, it was noted by Jean et al that the fatality within 30 days of Cryptococemia was 37%.<sup>9</sup> Also, in our case, the patient died within 10 days of Cryptococemia. Therefore, it is essential to suspect *Cryptococcus neoformans* infection in an immunocompromised patient and early management, keeping in mind the high morbidity associated with it. Consequently, it is crucial to avoid misdiagnosis and mismanagement, keeping in mind the high morbidity.

### 4 | CONCLUSION

This case highlights the importance of early suspicion of *Cryptococcus neoformans* infection and other opportunistic infections in immunocompromised patients, putting in mind that patients with Cryptococemia have a high risk of mortality within 30 days, which warrants the use of corticosteroid and immunomodulatory drugs in a critically ill patient with COVID-19. In the current scenario, the use of immunosuppressive therapy should be justified and to be alert for opportunistic infection like *Cryptococcus neoformans* infection, which can lead to sepsis and mortality.

### ACKNOWLEDGMENTS

Open Access funding provided by the Qatar National Library. The consent for publication was obtained.

### CONFLICT OF INTEREST

The authors declare that they have no competing interests.

### AUTHORS' CONTRIBUTIONS

MYK, AAA, SBS, ASM, AJN: involved in data collection, literature search, and manuscript preparation. All authors read and approved the final manuscript.

### ETHICAL APPROVAL

The article describes a case report. Therefore, no additional permission from our Ethics Committee was required.

## DATA AVAILABILITY STATEMENT

All data generated or analyzed during this study are included in this published article.

## ORCID

Ahmed S. Mohamed  <https://orcid.org/0000-0002-2107-5357>

Abdulqadir J. Nashwan  <https://orcid.org/0000-0003-4845-4119>

## REFERENCES

1. Franzot SP, Salkin IF, Casadevall A. *Cryptococcus neoformans* var. *grubii*: separate varietal status for *Cryptococcus neoformans* serotype A isolates. *J Clin Microbiol.* 1999;37(3):838-840.
2. Velagapudi R, Hsueh Y-P, Geunes-Boyer S, Wright JR, Heitman J. Spores as infectious propagules of *Cryptococcus neoformans*. *Infect Immun.* 2009;77(10):4345-4355.
3. Cox G, Perfect J. *Cryptococcus neoformans* var. *neoformans* and *gattii* and *Trichosporon* species. In *Topley and Wilson's Microbiology and Microbial Infections*, 9th edn. London: Arnold Press; 1997:461-484.
4. Lee SC, Dickson DW, Casadevall A. Pathology of cryptococcal meningoencephalitis: analysis of 27 patients with pathogenetic implications. *Hum Pathol.* 1996;27(8):839-847.
5. Baker RD. The primary pulmonary lymph node complex of cryptococcosis. *Am J Clin Pathol.* 1976;65(1):83-92.
6. Baker RD, Haugen RK. Tissue changes and tissue diagnosis in cryptococcosis. A study of 26 cases. *Am J Clin Pathol.* 1955;25(1):14-24.
7. Bamba S, Lortholary O, Sawadogo A, Millogo A, Guiguemde RT, Bretagne S. Decreasing incidence of cryptococcal meningitis in West Africa in the era of highly active antiretroviral therapy. *Aids.* 2012;26(8):1039-1041.
8. Xu X, Han M, Li T, et al. Effective treatment of severe COVID-19 patients with tocilizumab. *Proc Natl Acad Sci.* 2020;117(20):10970-10975.
9. Jean SS, Fang CT, Shau WY, et al. Cryptococcaemia: clinical features and prognostic factors. *QJM.* 2002;95(8):511-518.
10. Passerini M, Terzi R, Piscaglia M, Passerini S, Piconi S. Disseminated cryptococcosis in a patient with metastatic prostate cancer who died in the coronavirus disease 2019 (COVID-19) outbreak. *Cureus.* 2020;12(5):1-7.

**How to cite this article:** Khatib MY, Ahmed AA, Shaat S, Mohamed AS, Nashwan AJ. Cryptococccemia in a patient with COVID-19: A case report. *Clin Case Rep.* 2021;9:853-855. <https://doi.org/10.1002/ccr3.3668>