CORRECTION Open Access



Correction to: Elderly patients with atrial fibrillation in routine clinical practice: peri-procedural management of edoxaban oral anticoagulation therapy is associated with a low risk of bleeding and thromboembolic complications: a subset analysis of the prospective, observational, multinational **EMIT-AF study**

M. Unverdorben^{1*}, C. von Heymann², A. Santamaria³, M. Saxena⁴, T. Vanassche⁵, J. Jin¹, P. Laeis⁶, R. Wilkins⁷, C. Chen¹ and P. Colonna⁸

Correction to: BMC Cardiovasc Disord (2020) 20:504

https://doi.org/10.1186/s12872-020-01766-w

After publication of the original article [1], the authors identified an error in Table 1 related to two values on EHRA bleeding risk-low row.

The correct values are listed below:

EHRA bleeding risk-	174	580	457	297	283	252	45
low n, (%)	(62.1)	(49.9)	(55.1)	(48.5)	(51.5)	(49.0)	(45.9)

The original article has been corrected.

Author details

Global Medical Affairs Specialty and Value Products, Daiichi Sankyo Inc., 211 Mt Airy Road, Basking Ridge, NJ 07920, USA. ² Department of Anaesthesia

The original article can be found online at https://doi.org/10.1186/s1287

Full list of author information is available at the end of the article

and Intensive Care Medicine, Emergency Medicine, and Pain Therapy, Vivantes Klinikum Im Friedrichshain, Landsberger Allee 49, 10249 Berlin, Germany. ³ Hematology Department, University Hospital Vilaopó y Torrevieja, Alicante, Spain. ⁴ William Harvey Research Institute, Barts Health NHS Trust, Charterhouse Square, London EC1M 6BQ, UK. 5 Department of Cardiovascular Sciences, University Hospitals (UZ) Leuven, Leuven, Belgium. ⁶ Daiichi Sankyo, Medical Affairs Europe, Munich, Germany. 7 QPS Consulting, LLC, 19884 Naples Lakes Terrace, Ashburn, VA 20147, USA. ⁸ Department of Cardiology, Polyclinic of Bari - Hospital, 70124 Bari, Italy.

Published online: 15 February 2021

Reference

Unverdorben M, et al. Elderly patients with atrial fibrillation in routine clinical practice-peri-procedural management of edoxaban oral anticoagulation therapy is associated with a low risk of bleeding and thromboembolic complications: a subset analysis of the prospective, observational, multinational EMIT-AF study. BMC Cardiovasc Disord. 2020;20:504. https://doi.org/10.1186/s12872-020-01766-w.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



© The Author(s) 2021. Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit http://creativecommons.org/licenses/by/4.0/. The Creative Commons Public Domain Dedication waiver (http://creativeco mmons.org/publicdomain/zero/1.0/) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

^{*}Correspondence: munverdorben@dsi.com

¹ Global Medical Affairs Specialty and Value Products, Daiichi Sankyo Inc., 211 Mt Airy Road, Basking Ridge, NJ 07920, USA