



Published in final edited form as:

Curr Genet. 2021 February ; 67(1): 85–92. doi:10.1007/s00294-020-01116-5.

Dynamic regulation of Pif1 acetylation is crucial to the maintenance of genome stability

Onyekachi E. Ononye¹, Christopher W. Sausen², Matthew L. Bochman^{2,§}, Lata Balakrishnan^{1,§}

¹Department of Biology, School of Science, Indiana University Purdue University Indianapolis, Indianapolis, Indiana, USA

²Molecular and Cellular Biochemistry Department, Indiana University, Bloomington, Indiana, USA

Abstract

PIF1 family helicases are evolutionarily conserved among prokaryotes and eukaryotes. These enzymes function to support genome integrity by participating in multiple DNA transactions that can be broadly grouped into DNA replication, DNA repair, and telomere maintenance roles. However, the levels of PIF1 activity in cells must be carefully controlled, as Pif1 over-expression in *Saccharomyces cerevisiae* is toxic, and knockdown or over-expression of human PIF1 (hPIF1) supports cancer cell growth. This suggests that PIF1 family helicases must be subject to tight regulation *in vivo* to direct their activities to where and when they are needed, as well as to maintain those activities at proper homeostatic levels. Previous work shows that C-terminal phosphorylation of *S. cerevisiae* Pif1 regulates its telomere maintenance activity, and we recently identified that Pif1 is also regulated by lysine acetylation. The over-expression toxicity of Pif1 was exacerbated in cells lacking the Rpd3 lysine deacetylase, but mutation of the NuA4 lysine acetyltransferase subunit Esa1 ameliorated this toxicity. Using recombinant proteins, we found that acetylation stimulated the DNA binding affinity, ATPase activity, and DNA unwinding activities of Pif1. All three domains of the helicase were targets of acetylation *in vitro*, and multiple lines of evidence suggest that acetylation drives a conformational change in the N-terminal domain of Pif1 that impacts this stimulation. It is currently unclear what triggers lysine acetylation of Pif1 and how this modification impacts the many *in vivo* functions of the helicase, but future work promises to shed light on how this protein is tightly regulated within the cell.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <https://www.springer.com/aam-terms-v1>

§To whom correspondence should be addressed: Matthew L. Bochman, bochman@indiana.edu; Lata Balakrishnan, latabala@iupui.edu.

Author contributions: All authors participating in the writing and editing of this manuscript.

Publisher's Disclaimer: This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Conflicts of interest/competing interests: The authors declare no conflicts of interest.

Ethics approval: Not applicable

Consent to participate: Not applicable

Consent for publication: All authors have read and approved the contents of this manuscript.

Availability of data and material: Not applicable

Code availability: Not applicable

Keywords

Pif1 Helicase; Lysine Acetylation; DNA Replication; DNA Repair; G4 Resolvase; NuA4 (Esa1); Rpd3

Introduction

The *Petite Integration Frequency 1 (PIF1)* gene was identified in 1983 during a screen for mutants that disrupt mitochondrial DNA recombination in the budding yeast *Saccharomyces cerevisiae* (Foury and Kolodinsky 1983). A decade later, the Pif1 gene product was purified and shown to have single-stranded DNA (ssDNA)-dependent ATPase and 5′-3′ DNA helicase activities (Lahaye, et al. 1993, Lahaye, et al. 1991). Little did the field know that this yeast enzyme was the founding member of the PIF1 family of multifunctional DNA helicases found in both prokaryotes and eukaryotes (Bochman, et al. 2011, Bochman, et al. 2010) whose metazoan members would eventually be linked to cancer (Chen, et al. 2020, Chisholm, et al. 2012, Gagou, et al. 2014, Gagou, et al. 2011, Luo, et al. 2009), aging (Zhang, et al. 2006), and obesity (Bannwarth, et al. 2016, Belmonte, et al. 2019).

In many respects, the second wave of PIF1 research began with the “rediscovery” of *S. cerevisiae PIF1* as a gene whose mutation affects telomere length (Schulz and Zakian 1994). Since that time, Pif1 has been shown to have roles in multiple DNA transactions *in vivo*, and insights into the function(s) of *S. cerevisiae Pif1* in these processes are included below. *In vitro*, biochemical and biophysical approaches have demonstrated that Pif1 preferentially unwinds DNA fork substrates compared to substrates with a single ssDNA tail (Lahaye, et al. 1993, Ononye, et al. 2020, Ramanagoudr-Bhojappa, et al. 2013), suggesting that Pif1 undergoes secondary interactions with the strand of DNA complementary to the one upon which it translocates. In support of this, RNA:DNA hybrid forks stimulate Pif1 activity above that displayed on DNA forks, when the DNA strand supplies the 5′ ssDNA tail (translocating) and the RNA supplies the 3′ ssDNA tail (non-translocating) of the fork structure (Boule and Zakian 2007, Chib, et al. 2016, Ononye, et al. 2020, Zhou, et al. 2014). Notably, RNA-DNA hybrids adopt an A-form conformation instead of the canonical B-form of duplex DNA (Wong and Shao 2017), and this is not the only non-canonical structure that is preferentially unwound by PIF1 family helicases. For PIF1 enzymes from multiple species, it has been shown that G-quadruplex structures also stimulate helicase activity (Mohammad, et al. 2018, Paeschke, et al. 2013, Sanders 2010).

Lesser-studied biochemical activities of PIF1 helicases include ssDNA annealing and nuclease activity. Strand annealing by *S. cerevisiae Pif1* was first demonstrated by the Raney group, and this activity occurs *in vitro* under unwinding conditions (*i.e.*, in the presence of ATP), even in the presence of single-stranded binding proteins (Ramanagoudr-Bhojappa, et al. 2014). This strand annealing activity is evolutionarily conserved with hPIF1 and intrinsic to its helicase domain (George, et al. 2009), which is the most conserved portion of all PIF1 proteins (Bochman, et al. 2011, Bochman, et al. 2010). Thus, strand annealing may be a general feature of all PIF1 helicases. Another putative conserved PIF1 biochemical function is 3′-5′ exonuclease activity, which has been reported for recombinant *Candida albicans*

Pif1 and other fungal PIF1 proteins, including *S. cerevisiae* Pif1 (Wei, et al. 2017). However, these findings are controversial because high concentrations of *S. cerevisiae* Pif1 were necessary to observe the nuclease activity, and the absence of a contaminating nuclease from the recombinant protein expression host was not rigorously demonstrated.

Crystal structures of the *S. cerevisiae* helicase domain have been solved (Lu, et al. 2018), as well as structures of PIF1 family helicases from *Bacteroides* species (Chen, et al. 2016, Zhou, et al. 2016) and the hPIF1 helicase domain (Dehghani-Tafti, et al. 2019). Although these structures have revealed some commonalities and species-specific differences between PIF1 family helicases, it is unclear how informative they will be to the genome integrity field at large. This is because research has shown that in certain instances, PIF1 helicase domains are generic motor modules whose activity is either regulated by accessory domains or complemented by flanking domains necessary for proper *in vivo* function (Andis, et al. 2018). The N- and C-terminal domains that flank the central helicase domain are also likely why a full-length eukaryotic PIF1 structure has eluded the field. These domains vary in length and sequence between species (Bochman, et al. 2011, Bochman, et al. 2010), but a common characteristic among them is that they are predicted to be natively disordered (Nickens, et al. 2019). Thus, modern structural biology approaches such as cryo-electron microscopy will be needed to solve the structures of these enzymes, which may exist in an ensemble of conformations in solution. The utility of such natively disordered regions in proteins has gained recent attention as they are often sites of protein-protein interaction (Fong, et al. 2009), post-translational modifications (Bah and Forman-Kay 2016), and/or involved in liquid-liquid phase separation (Kato, et al. 2012).

Multifunctional activities of nuclear PIF1

Based on its ability to unwind DNA and RNA/DNA hybrid duplexes, nuclear Pif1 has been implicated in a multitude of nucleic acid transactions. For instance, during Okazaki fragment maturation, Pif1 stimulates the strand displacement activity of DNA polymerase delta (pol δ), creating long 5' primer flaps (Pike, et al. 2009). Lengthening of this flap gives rise to the accumulation of the ssDNA binding protein, replication protein A (RPA), which when bound, prevents illegitimate recombination, formation of secondary structures, and inappropriate degradation of genetic material (Bartos, et al. 2008). RPA-bound flaps prevent primer cleavage by flap endonuclease 1 (FEN1), pushing Okazaki fragment processing toward an alternative pathway, known as the long-flap pathway. This necessitates the activity of the helicase/nuclease Dna2 mediates RPA displacement and flap removal (Pike, et al. 2010, Stewart, et al. 2008). This proposed mechanism succeeded initial genetic data by Budd et al. (2006), which revealed that the presence of Pif1 necessitates the endonuclease activity of Dna2, presumably by functioning together in the lagging strand maturation process (Budd, et al. 2006). Pif1 also plays a role in repetitive ribosomal DNA (rDNA) replication. Movement of the replication fork proceeds in a manner such that the right moving fork is unhindered, while the left moving fork experiences a stop at the replication fork barrier (RFB) (Ivessa, et al. 2000). The presence of the RFB is essential so that replication proceeds in the same direction as transcription, preventing macromolecular collisions (Kobayashi 2003). The protein Fob1 binds to these RFB sequences and serves as a unidirectional block to fork progression (Kobayashi and Horiuchi 1996). Pif1 directly

interacts with Fob1 to maintain this barrier, though the exact mechanism by which this happens has yet to be fully explored (Muellner and Schmidt 2020). Interestingly, the second PIF1 family helicase in *S. cerevisiae*, Rrm3, has the opposite function during rDNA replication, mitigating fork stalling at the RFB (Ivessa, et al. 2000). Additionally, ScPif1 also plays a backup role to Rrm3 in tRNA gene replication (Tran, et al. 2017).

While majority of the studies investigating the role of PIF1 helicases during DNA replication focus on the elongation phase, more recently, Pif1 and Rrm3 have been associated with replication termination. Their unwinding activity is proposed to relieve torsional strain in the final section of the parental DNA in budding yeast (Deegan, et al. 2019, Su, et al. 2019). Failure to unwind this region prevents the convergence of replication forks, thereby giving rise to late termination intermediates (LTIs) that stall replication termination (Steinacher, et al. 2012). Compared to other *S. cerevisiae* helicases that are known to resolve structured intermediates, such as Sgs1 and Chl1, only Pif1 and Rrm3 allow for proper replication termination of preformed LTIs *in vitro* (Deegan, et al. 2019).

Pif1 also acts as a G-quadruplex (G4) resolvase on both strands of the replication fork. In its absence, the lagging strand containing these stable non-canonical DNA secondary structures experience stalling (Dahan, et al. 2018, Sparks, et al. 2019). Therefore, although other helicases have displayed the capability to resolve G4 structures *in vitro*, such as Sgs1 and Dna2, it is now clear that in comparison to the activity of Pif1, such helicases function much less efficiently on intramolecular G4s, which are more commonly formed during lagging strand replication (Lerner and Sale 2019, Liu, et al. 2010, Paeschke, et al. 2013). Similarly, PIF1 family helicases can also efficiently unwind other non-B form DNA structures such as R-loops. This activity was initially implicated by the stimulation of *S. cerevisiae* Pif1 helicase activity by RNA-DNA hybrids (Boule and Zakian 2007) and the finding that Pif1 binding is enriched at highly transcribed genes genome wide (Paeschke, et al. 2011). More recently, it has been demonstrated that both Pif1 and Rrm3 promote genome stability at tRNA genes by removing R-loops (Tran, et al. 2017). In addition to its role in structure resolution, Pif1 can also remove protein barriers from dsDNA (Koc, et al. 2016). Furthermore, along with the various roles of Pif1 during replication, the helicase is also involved in the repair of single-stranded nicks via break-induced replication (BIR) (Malkova and Ira 2013). During this process, Pif1 recruits pol δ to D-loops that are formed following strand invasion and stimulates pol δ synthesis via bubble migration as the helicase interacts with PCNA (Buzovetsky, et al. 2017).

Finally, Pif1's activities also modulate—and are modulated by—protein–protein interactions such as is the case with telomerase. Biochemical studies reveal that Pif1 inhibits Est2, the reverse transcriptase subunit of telomerase, using its ATPase activity to unwind the telomerase RNA and the telomeric DNA hybrid (Boule, et al. 2005, Phillips, et al. 2015). Overexpression and deletion studies of Pif1 *in vivo* reveal a decrease and increase in telomere length, respectively, showing that telomere length is inversely proportional to the amount of Pif1 activity in *S. cerevisiae* (Boule, et al. 2005). Although Pif1 is a known regulator of telomerase at telomeres, it is equally capable of regulating telomerase at sites of DNA double-strand breaks (DSBs). Indeed, to specifically prevent the *de novo* addition of telomeres at DSBs instead of telomeres, Pif1 is phosphorylated in a Rad53-dependent

manner (Makovets and Blackburn 2009). This regulation of Pif1 telomere maintenance activity also occurs via an interaction with the Hrq1 helicase (Bochman, et al. 2014), and the combined activities of these two helicases has been shown to synergistically tune telomerase activity up and down *in vitro* (Nickens, et al. 2018).

The known central functions of Pif1 in various cellular pathways are featured in Figure 1. Given the myriad of pathways mediated by Pif1, the cell must have multiple mechanisms in place to regulate the activity of this low copy number protein within the cell.

Lysine acetylation of nuclear PIF1

Dynamically reversible protein post-translational modifications (PTMs) function to enhance the diversity of the proteome without much energetic cost to the cell. Following identification of Pif1 phosphorylation in *S. cerevisiae* (Makovets and Blackburn 2009), quantitative proteomic analyses further revealed hPIF1 to also be modified by phosphorylation and by ubiquitination (Kettenbach, et al. 2011, Sharma, et al. 2014, Udeshi, et al. 2013). In our recent work, we show that *S. cerevisiae* Pif1 is acetylated *in vivo* and can also be modified *in vitro* (Ononye, et al. 2020). The acetylation signature of Pif1 spans the entire protein, with lysine residues being modified on the N-terminus (Lys 118, Lys 129), within the helicase domain (Lys 525, Lys 639, Lys 725), and the C-terminus (Lys 800). Using lysine acetyltransferase (KAT) and lysine deacetylase (KDAC) mutant strains, we assessed the impact of altering the acetylation status of Pif1 on cell viability. While overexpression of Pif1 in itself is highly toxic to the cell, specific KDAC knockouts accentuated this toxicity phenotype. In contrast, a specific KAT knockout ameliorated the overexpression toxicity. Based on the viability of Pif1-overexpressing cells, we were able to assign the NuA4 (Esa1) KAT and its counteracting KDAC Rpd3 as responsible for dynamically altering the acetylation status of Pif1 within the cell. Because Pif1 acetylation had a profound impact on cell viability, we characterized the biochemical changes to Pif1 function upon lysine acetylation. Helicase assays using preferred substrates of Pif1 showed that the acetylated form of the helicase was much better at unwinding duplex strands (either DNA/DNA or RNA/DNA hybrid) compared to unmodified Pif1. Analyzing the reason for this increased helicase activity revealed that while acetylated Pif1 did not display an increased rate of unwinding, it was simply able to remain bound to the substrate longer, allowing for improved unwinding function. This observation directly correlates with our binding assays, wherein Pif1 displayed higher binding affinity to its cognate substrates compared to unmodified Pif1. Similar to the stimulated helicase activity and increased DNA binding affinity, we also detected higher ATPase efficiency by acetylated Pif1.

Due to being natively disordered, the functional role of the N-terminus of Pif1 has remained somewhat of an enigma. Though acetylation marks were identified across multiple domains of Pif1, we were particularly interested in understanding the role of this modification on the N-terminus of the helicase. Over-expression of a Pif1 mutant lacking its N-terminus (Pif1^N) resulted in decreased cellular toxicity compared to the full-length helicase. The toxicity phenotype of Pif1^N was further rescued in strains lacking *RPD3*. This observation clarified the role of the N-terminus in contributing to the acetylation-based regulation of Pif1's enzymatic activities. Biochemical studies showed that similar to the full-length

protein, the helicase and ATPase functions of Pif1 N were stimulated by acetylation. However, the mutant helicase did not exhibit increased DNA binding affinity when modified.

We were curious about the role of lysine acetylation in driving conformational changes in the Pif1 structure, which could impact its function. Limited proteolysis assays revealed that acetylation afforded Pif1 protection from cleavage, suggesting a change in conformation compared to unmodified Pif1. The unmodified Pif1 N was more resistant to proteolysis, suggesting that the N-terminus directly impacts overall protein structure, and perhaps acetylation of this domain modifies the three-dimensional architecture of Pif1 to stimulate its biochemical activities. Future structural analyses will be required to delineate the role of the acetylated N-terminus in altering the Pif1 structure, thereby regulating the binding, unwinding, and ATPase functions of the protein.

Current perspectives and future research

The activity of Pif1 in the cell needs to be finely tuned to ensure the maintenance of genome stability and cell viability. Too much or too little Pif1 activity creates havoc in the cell, leading to checkpoint activation and ultimately cell death. Our observation that lysine acetylation leads to increased Pif1 enzymatic activities suggests that PTM of the helicase could be another mode by which the cell can dynamically regulate the functions of this protein. Modification of Pif1 could also change the outcome of a specific biological pathway. For example, increased helicase function of acetylated Pif1 could lead to rampant unwinding of Okazaki fragments, causing the accumulation of checkpoint-triggering ssDNA in the cell. Alternatively, this could also push Okazaki fragment maturation (OFM) to the long-flap pathway of processing, wherein the pol α -synthesized primers would be completely displaced, promoting resynthesis by the higher fidelity polymerase, pol δ . Because deletion of *PIF1* rescues the lethal phenotype of *dna2* in *S. cerevisiae*, it has been proposed that both of these helicases work in conjunction during OFM (Budd, et al. 2006), and multiple biochemical studies lend credence to this proposal (Ayyagari, et al. 2003, Pike, et al. 2009, Pike, et al. 2010, Rossi, et al. 2008). However, recent genetic studies also suggest that formation of long flaps is not widespread in the cell (Kahli, et al. 2019). A very recent study suggests that it is not their activity during OFM but the combined activity of these helicases in replication fork restart that make them vital to cell viability (Appanah, et al. 2020, Falquet, et al. 2020). Human Dna2 was previously shown to be acetylated *in vivo*, and biochemical analysis revealed that its nuclease, helicase and binding activities were stimulated *in vitro* (Balakrishnan, et al. 2010), similar to our report of increased activity by acetylated yeast Pif1 (Ononye, et al. 2020). Combined modification of Pif1 and Dna2 may directly impact their role in both flap processing and/or replication fork restart. A multitude of proteins functioning in replication and repair are also known to be modified by lysine acetylation, including the hBLM (Wang 2017) and hWRN helicases (Lozada, et al. 2014, Muftuoglu, et al. 2008). While the alterations to enzymatic activities of these acetylated replication proteins have not all been characterized, we are now starting to gain a better understanding of this mode of protein functional regulation.

Our recent study serves as good first step in defining the role of lysine acetylation in regulating the biochemical functions of the Pif1 helicase, but many questions remain about this mode of protein regulation. Specifically, it will be important to know what fraction of Pif1 is constitutively acetylated within the cell or if there are specific triggers for lysine acetylation of the helicase, such as cell cycle phases or the DNA damage response. Alterations in the acetylation signature of the helicase upon different triggers would help us to identify how specific lysine modifications could impact Pif1's interactions with other proteins in a biological pathway. Further, it would be interesting to know if the same enzyme modifiers that we identified in our study (*i.e.*, the KAT, NuA4 [Esa1], and KDAC, Rpd3) are exclusively responsible for altering Pif1's acetylation status or if other redundant acetylation-specific modifiers also play a role in the PTM of this enzyme. Further, we need to characterize the impact of Pif1 acetylation on the myriad of biochemical processes that the helicase has been implicated in (outlined in Figure 1) and the consequences of this modification on these biological pathways. Finally, while we have preliminary evidence showing that hPIF1 is acetylated *in vivo* (unpublished observation), we are currently unaware of the biochemical consequences of this modification and its impact on human genome integrity. Multiple avenues exist to further understand this regulatory modification on PIF1 family helicases, and future research on this topic will help to shed light on the significance of this protein in genome maintenance and cell viability.

Acknowledgments

Funding: This work was funded by grants from the National Science Foundation (1929346) to L.B. and National Health Institutes (1R35GM133437) and American Cancer Society (RSG-16-180-01-DMC) to M.L.B.

References

- Andis NM, Sausen CW, Alladin A, Bochman ML (2018) The WYL Domain of the PIF1 Helicase from the Thermophilic Bacterium *Thermotoga elfii* is an Accessory Single-Stranded DNA Binding Module. *Biochemistry* 57: 1108–1118 doi: 10.1021/acs.biochem.7b01233 [PubMed: 29341597]
- Appanah R, Jones D, Falquet B, Rass U (2020) Limiting homologous recombination at stalled replication forks is essential for cell viability: DNA2 to the rescue. *Curr Genet* 10.1007/s00294-020-01106-7
- Ayyagari R, Gomes XV, Gordenin DA, Burgers PM (2003) Okazaki fragment maturation in yeast. I. Distribution of functions between FEN1 AND DNA2. *J Biol Chem* 278: 1618–1625 doi: 10.1074/jbc.M209801200 [PubMed: 12424238]
- Bah A, Forman-Kay JD (2016) Modulation of Intrinsically Disordered Protein Function by Post-translational Modifications. *J Biol Chem* 291: 6696–6705 doi: 10.1074/jbc.R115.695056 [PubMed: 26851279]
- Balakrishnan L, Stewart J, Polaczek P, Campbell JL, Bambara RA (2010) Acetylation of Dna2 endonuclease/helicase and flap endonuclease 1 by p300 promotes DNA stability by creating long flap intermediates. *J Biol Chem* 285: 4398–4404 doi: 10.1074/jbc.M109.086397 [PubMed: 20019387]
- Bannwarth S, Berg-Alonso L, Auge G, Fragaki K, Kolesar JE, Lespinasse F, Lacas-Gervais S, Burel-Vandenbos F, Villa E, Belmonte F, Michiels JF, Ricci JE, Gherardi R, Harrington L, Kaufman BA, Paquis-Flucklinger V (2016) Inactivation of Pif1 helicase causes a mitochondrial myopathy in mice. *Mitochondrion* 30: 126–137 doi: 10.1016/j.mito.2016.02.005 [PubMed: 26923168]
- Bartos JD, Willmott LJ, Binz SK, Wold MS, Bambara RA (2008) Catalysis of strand annealing by replication protein A derives from its strand melting properties. *J Biol Chem* 283: 21758–21768 doi: 10.1074/jbc.M800856200 [PubMed: 18522944]

- Belmonte FR, Dedousis N, Sipula I, Desai NA, Singhi AD, Chu Y, Zhang Y, Bannwarth S, Paquis-Flucklinger V, Harrington L, Shiva S, Jurczak MJ, O'Doherty RM, Kaufman BA (2019) Petite Integration Factor 1 (PIF1) helicase deficiency increases weight gain in Western diet-fed female mice without increased inflammatory markers or decreased glucose clearance. *PLoS One* 14: e0203101 doi: 10.1371/journal.pone.0203101 [PubMed: 31136580]
- Bochman ML, Judge CP, Zakian VA (2011) The Pif1 family in prokaryotes: what are our helicases doing in your bacteria? *Molecular biology of the cell* 22: 1955–1959 doi: 10.1091/mbc.E11-01-0045 [PubMed: 21670310]
- Bochman ML, Paeschke K, Chan A, Zakian VA (2014) Hrq1, a homolog of the human RecQ4 helicase, acts catalytically and structurally to promote genome integrity. *Cell Rep* 6: 346–356 doi: 10.1016/j.celrep.2013.12.037 [PubMed: 24440721]
- Bochman ML, Sabouri N, Zakian VA (2010) Unwinding the functions of the Pif1 family helicases. *DNA Repair (Amst)* 9: 237–249 doi: S1568-7864(10)00011-X [pii] 10.1016/j.dnarep.2010.01.008 [PubMed: 20097624]
- Boule JB, Vega LR, Zakian VA (2005) The yeast Pif1p helicase removes telomerase from telomeric DNA. *Nature* 438: 57–61 doi: 10.1038/nature04091 [PubMed: 16121131]
- Boule JB, Zakian VA (2007) The yeast Pif1p DNA helicase preferentially unwinds RNA DNA substrates. *Nucleic Acids Res* 35: 5809–5818 doi: [PubMed: 17720711]
- Budd ME, Reis CC, Smith S, Myung K, Campbell JL (2006) Evidence suggesting that Pif1 helicase functions in DNA replication with the Dna2 helicase/nuclease and DNA polymerase delta. *Mol Cell Biol* 26: 2490–2500 doi: 10.1128/MCB.26.7.2490-2500.2006 [PubMed: 16537895]
- Buzovetsky O, Kwon Y, Pham NT, Kim C, Ira G, Sung P, Xiong Y (2017) Role of the Pif1-PCNA Complex in Pol delta-Dependent Strand Displacement DNA Synthesis and Break-Induced Replication. *Cell Rep* 21: 1707–1714 doi: 10.1016/j.celrep.2017.10.079 [PubMed: 29141206]
- Chen B, Hua Z, Gong B, Tan X, Zhang S, Li Q, Chen Y, Zhang J, Li Z (2020) Downregulation of PIF1, a potential new target of MYCN, induces apoptosis and inhibits cell migration in neuroblastoma cells. *Life Sci* 256: 117820 doi: 10.1016/j.lfs.2020.117820 [PubMed: 32512012]
- Chen WF, Dai YX, Duan XL, Liu NN, Shi W, Li N, Li M, Dou SX, Dong YH, Rety S, Xi XG (2016) Crystal structures of the BsPif1 helicase reveal that a major movement of the 2B SH3 domain is required for DNA unwinding. *Nucleic Acids Res* 44: 2949–2961 doi: 10.1093/nar/gkw033 [PubMed: 26809678]
- Chib S, Byrd AK, Raney KD (2016) Yeast Helicase Pif1 Unwinds RNA:DNA Hybrids with Higher Processivity than DNA:DNA Duplexes. *J Biol Chem* 291: 5889–5901 doi: 10.1074/jbc.M115.688648 [PubMed: 26733194]
- Chisholm KM, Aubert SD, Freese KP, Zakian VA, King MC, Welch PL (2012) A Genomewide Screen for Suppressors of Alu-Mediated Rearrangements Reveals a Role for PIF1. *Plos One* 7: 10 doi: 10.1371/journal.pone.0030748
- Dahan D, Tsirkas I, Dovrat D, Sparks MA, Singh SP, Galletto R, Aharoni A (2018) Pif1 is essential for efficient replisome progression through lagging strand G-quadruplex DNA secondary structures. *Nucleic Acids Res* 46: 11847–11857 doi: 10.1093/nar/gky1065 [PubMed: 30395308]
- Deegan TD, Baxter J, Ortiz Bazan MA, Yeeles JTP, Labib KPM (2019) Pif1-Family Helicases Support Fork Convergence during DNA Replication Termination in Eukaryotes. *Mol Cell* 74: 231–244 e239 doi: 10.1016/j.molcel.2019.01.040 [PubMed: 30850330]
- Dehghani-Tafti S, Levdkov V, Antson AA, Bax B, Sanders CM (2019) Structural and functional analysis of the nucleotide and DNA binding activities of the human PIF1 helicase. *Nucleic Acids Res* 47: 3208–3222 doi: 10.1093/nar/gkz028 [PubMed: 30698796]
- Falquet B, Olmezer G, Enkner F, Klein D, Challa K, Appanah R, Gasser SM, Rass U (2020) Disease-associated DNA2 nuclease-helicase protects cells from lethal chromosome under-replication. *Nucleic Acids Res* 48: 7265–7278 doi: 10.1093/nar/gkaa524 [PubMed: 32544229]
- Fong JH, Shoemaker BA, Garbuzynskiy SO, Lobanov MY, Galzitskaya OV, Panchenko AR (2009) Intrinsic disorder in protein interactions: insights from a comprehensive structural analysis. *PLoS Comput Biol* 5: e1000316 doi: 10.1371/journal.pcbi.1000316 [PubMed: 19282967]

- Foury F, Kolodynski J (1983) pif mutation blocks recombination between mitochondrial rho+ and rho- genomes having tandemly arrayed repeat units in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci* 80: 5345–5349 doi: [PubMed: 6310571]
- Gagou ME, Ganesh A, Phear G, Robinson D, Petermann E, Cox A, Meuth M (2014) Human PIF1 helicase supports DNA replication and cell growth under oncogenic-stress. *Oncotarget* 5: 11381–11398 doi: [PubMed: 25359767]
- Gagou ME, Ganesh A, Thompson R, Phear G, Sanders C, Meuth M (2011) Suppression of Apoptosis by PIF1 Helicase in Human Tumor Cells. *Cancer Research* 71: 4998–5008 doi: 10.1158/0008-5472.can-10-4404 [PubMed: 21616935]
- George T, Wen Q, Griffiths R, Ganesh A, Meuth M, Sanders CM (2009) Human Pif1 helicase unwinds synthetic DNA structures resembling stalled DNA replication forks. *Nucleic Acids Res* 37: 6491–6502 doi: gkp671 [pii] 10.1093/nar/gkp671 [PubMed: 19700773]
- Ivessa AS, Zhou JQ, Zakian VA (2000) The *Saccharomyces* Pif1p DNA helicase and the highly related Rrm3p have opposite effects on replication fork progression in ribosomal DNA. *Cell* 100: 479–489 doi: 10.1016/s0092-8674(00)80683-2 [PubMed: 10693764]
- Kahli M, Osmundson JS, Yeung R, Smith DJ (2019) Processing of eukaryotic Okazaki fragments by redundant nucleases can be uncoupled from ongoing DNA replication in vivo. *Nucleic Acids Res* 47: 1814–1822 doi: 10.1093/nar/gky1242 [PubMed: 30541106]
- Kato M, Han TW, Xie S, Shi K, Du X, Wu LC, Mirzaei H, Goldsmith EJ, Longgood J, Pei J, Grishin NV, Frantz DE, Schneider JW, Chen S, Li L, Sawaya MR, Eisenberg D, Tycko R, McKnight SL (2012) Cell-free formation of RNA granules: low complexity sequence domains form dynamic fibers within hydrogels. *Cell* 149: 753–767 doi: 10.1016/j.cell.2012.04.017 [PubMed: 22579281]
- Kettenbach AN, Schweppe DK, Faherty BK, Pechenick D, Pletnev AA, Gerber SA (2011) Quantitative phosphoproteomics identifies substrates and functional modules of Aurora and Polo-like kinase activities in mitotic cells. *Sci Signal* 4: rs5 doi: 10.1126/scisignal.2001497 [PubMed: 21712546]
- Kobayashi T (2003) The replication fork barrier site forms a unique structure with Fob1p and inhibits the replication fork. *Mol Cell Biol* 23: 9178–9188 doi: 10.1128/mcb.23.24.9178-9188.2003 [PubMed: 14645529]
- Kobayashi T, Horiuchi T (1996) A yeast gene product, Fob1 protein, required for both replication fork blocking and recombinational hotspot activities. *Genes Cells* 1: 465–474 doi: 10.1046/j.1365-2443.1996.d01-256.x [PubMed: 9078378]
- Koc KN, Singh SP, Stodola JL, Burgers PM, Galletto R (2016) Pif1 removes a Rap1-dependent barrier to the strand displacement activity of DNA polymerase delta. *Nucleic Acids Res* 44: 3811–3819 doi: 10.1093/nar/gkw181 [PubMed: 27001517]
- Lahaye A, Leterme S, Foury F (1993) PIF1 DNA helicase from *Saccharomyces cerevisiae*. Biochemical characterization of the enzyme. *J Biol Chem* 268: 26155–26161 doi: [PubMed: 8253734]
- Lahaye A, Stahl H, Thines-Sempoux D, Foury F (1991) PIF1: a DNA helicase in yeast mitochondria. *EMBO J* 10: 997–1007 doi: [PubMed: 1849081]
- Lerner LK, Sale JE (2019) Replication of G Quadruplex DNA. *Genes (Basel)* 1010.3390/genes10020095
- Liu JQ, Chen CY, Xue Y, Hao YH, Tan Z (2010) G-quadruplex hinders translocation of BLM helicase on DNA: a real-time fluorescence spectroscopic unwinding study and comparison with duplex substrates. *J Am Chem Soc* 132: 10521–10527 doi: 10.1021/ja1038165 [PubMed: 20614884]
- Lozada E, Yi J, Luo J, Orren DK (2014) Acetylation of Werner syndrome protein (WRN): relationships with DNA damage, DNA replication and DNA metabolic activities. *Biogerontology* 15: 347–366 doi: 10.1007/s10522-014-9506-3 [PubMed: 24965941]
- Lu KY, Chen WF, Rety S, Liu NN, Wu WQ, Dai YX, Li D, Ma HY, Dou SX, Xi XG (2018) Insights into the structural and mechanistic basis of multifunctional *S. cerevisiae* Pif1p helicase. *Nucleic Acids Res* 46: 1486–1500 doi: 10.1093/nar/gkx1217 [PubMed: 29202194]
- Luo J, Solimini NL, Elledge SJ (2009) Principles of Cancer Therapy: Oncogene and Non-oncogene Addiction. *Cell* 136: 823–837 doi: 10.1016/j.cell.2009.02.024 [PubMed: 19269363]
- Makovets S, Blackburn EH (2009) DNA damage signalling prevents deleterious telomere addition at DNA breaks. *Nat Cell Biol* 11: 1383–1386 doi: 10.1038/ncb1985 [PubMed: 19838171]

- Malkova A, Ira G (2013) Break-induced replication: functions and molecular mechanism. *Curr Opin Genet Dev* 23: 271–279 doi: 10.1016/j.gde.2013.05.007 [PubMed: 23790415]
- Mohammad JB, Wallgren M, Sabouri N (2018) The Pif1 signature motif of Pfh1 is necessary for both protein displacement and helicase unwinding activities, but is dispensable for strand-annealing activity. *Nucleic Acids Res* 46: 8516–8531 doi: 10.1093/nar/gky654 [PubMed: 30053106]
- Muellner J, Schmidt KH (2020) Yeast Genome Maintenance by the Multifunctional PIF1 DNA Helicase Family. *Genes (Basel)* 11 10.3390/genes11020224
- Muftuoglu M, Kusumoto R, Speina E, Beck G, Cheng WH, Bohr VA (2008) Acetylation regulates WRN catalytic activities and affects base excision DNA repair. *PLoS One* 3: e1918 doi: 10.1371/journal.pone.0001918 [PubMed: 18398454]
- Nickens DG, Rogers CM, Bochman ML (2018) The *Saccharomyces cerevisiae* Hrq1 and Pif1 DNA helicases synergistically modulate telomerase activity in vitro. *J Biol Chem* 293: 14481–14496 doi: 10.1074/jbc.RA118.004092 [PubMed: 30068549]
- Nickens DG, Sausen CW, Bochman ML (2019) The Biochemical Activities of the *Saccharomyces cerevisiae* Pif1 Helicase Are Regulated by Its N-Terminal Domain. *Genes (Basel)* 1010.3390/genes10060411
- Ononye OE, Sausen CW, Balakrishnan L, Bochman ML (2020) Lysine Acetylation Regulates the Activity of Nuclear Pif1. *J Biol Chem* 10.1074/jbc.RA120.015164
- Paeschke K, Bochman ML, Garcia PD, Cejka P, Friedman KL, Kowalczykowski SC, Zakian VA (2013) Pif1 family helicases suppress genome instability at G-quadruplex motifs. *Nature* 497: 458–462 doi: 10.1038/nature12149 [PubMed: 23657261]
- Paeschke K, Capra JA, Zakian VA (2011) DNA Replication through G-Quadruplex Motifs Is Promoted by the *Saccharomyces cerevisiae* Pif1 DNA Helicase. *Cell* 145: 678–691 doi: 10.1016/j.cell.2011.04.015 [PubMed: 21620135]
- Phillips JA, Chan A, Paeschke K, Zakian VA (2015) The pif1 helicase, a negative regulator of telomerase, acts preferentially at long telomeres. *PLoS Genet* 11: e1005186 doi: 10.1371/journal.pgen.1005186 [PubMed: 25906395]
- Pike JE, Burgers PM, Campbell JL, Bambara RA (2009) Pif1 helicase lengthens some Okazaki fragment flaps necessitating Dna2 nuclease/helicase action in the two-nuclease processing pathway. *J Biol Chem* 284: 25170–25180 doi: 10.1074/jbc.M109.023325 [PubMed: 19605347]
- Pike JE, Henry RA, Burgers PM, Campbell JL, Bambara RA (2010) An alternative pathway for Okazaki fragment processing: resolution of fold-back flaps by Pif1 helicase. *J Biol Chem* 285: 41712–41723 doi: 10.1074/jbc.M110.146894 [PubMed: 20959454]
- Ramanagoudr-Bhojappa R, Byrd AK, Dahl C, Raney KD (2014) Yeast Pif1 accelerates annealing of complementary DNA strands. *Biochemistry* 53: 7659–7669 doi: 10.1021/bi500746v [PubMed: 25393406]
- Ramanagoudr-Bhojappa R, Chib S, Byrd AK, Aarattuthodiyil S, Pandey M, Patel SS, Raney KD (2013) Yeast Pif1 helicase exhibits a one-base-pair stepping mechanism for unwinding duplex DNA. *J Biol Chem* 288: 16185–16195 doi: 10.1074/jbc.M113.470013 [PubMed: 23596008]
- Rossi ML, Pike JE, Wang W, Burgers PM, Campbell JL, Bambara RA (2008) Pif1 helicase directs eukaryotic Okazaki fragments toward the two-nuclease cleavage pathway for primer removal. *J Biol Chem* 283: 27483–27493 doi: M804550200 [pii] 10.1074/jbc.M804550200 [PubMed: 18689797]
- Sanders CM (2010) Human Pif1 helicase is a G-quadruplex DNA binding protein with G-quadruplex DNA unwinding activity. *Biochem J* BJ20100612 [pii] 10.1042/BJ20100612
- Schulz VP, Zakian VA (1994) The *Saccharomyces PIF1* DNA helicase inhibits telomere elongation and de novo telomere formation. *Cell* 76: 145–155 doi: [PubMed: 8287473]
- Sharma K, D'Souza RC, Tyanova S, Schaab C, Wisniewski JR, Cox J, Mann M (2014) Ultradeep human phosphoproteome reveals a distinct regulatory nature of Tyr and Ser/Thr-based signaling. *Cell Rep* 8: 1583–1594 doi: 10.1016/j.celrep.2014.07.036 [PubMed: 25159151]
- Sparks MA, Singh SP, Burgers PM, Galletto R (2019) Complementary roles of Pif1 helicase and single stranded DNA binding proteins in stimulating DNA replication through G-quadruplexes. *Nucleic Acids Res* 47: 8595–8605 doi: 10.1093/nar/gkz608 [PubMed: 31340040]

- Steinacher R, Osman F, Dalgaard JZ, Lorenz A, Whitby MC (2012) The DNA helicase Pfh1 promotes fork merging at replication termination sites to ensure genome stability. *Genes Dev* 26: 594–602 doi: 10.1101/gad.184663.111 [PubMed: 22426535]
- Stewart JA, Miller AS, Campbell JL, Bambara RA (2008) Dynamic removal of replication protein A by Dna2 facilitates primer cleavage during Okazaki fragment processing in *Saccharomyces cerevisiae*. *J Biol Chem* 283: 31356–31365 doi: 10.1074/jbc.M805965200 [PubMed: 18799459]
- Su N, Byrd AK, Bharath SR, Yang O, Jia Y, Tang X, Ha T, Raney KD, Song H (2019) Structural basis for DNA unwinding at forked dsDNA by two coordinating Pif1 helicases. *Nat Commun* 10: 5375 doi: 10.1038/s41467-019-13414-9 [PubMed: 31772234]
- Tran PLT, Pohl TJ, Chen CF, Chan A, Pott S, Zakian VA (2017) PIF1 family DNA helicases suppress R-loop mediated genome instability at tRNA genes. *Nat Commun* 8: 15025 doi: 10.1038/ncomms15025 [PubMed: 28429714]
- Udeshi ND, Svinkina T, Mertins P, Kuhn E, Mani DR, Qiao JW, Carr SA (2013) Refined preparation and use of anti-diglycine remnant (K-epsilon-GG) antibody enables routine quantification of 10,000s of ubiquitination sites in single proteomics experiments. *Mol Cell Proteomics* 12: 825–831 doi: 10.1074/mcp.O112.027094 [PubMed: 23266961]
- Wang YL, J. (2017) Acetylation of BLM protein regulates its function in response to DNA damage. *RSC Advances* 7: 55301–55308 doi:
- Wei XB, Zhang B, Bazeille N, Yu Y, Liu NN, Rene B, Mauffret O, Xi XG (2017) A 3′-5′ exonuclease activity embedded in the helicase core domain of *Candida albicans* Pif1 helicase. *Sci Rep* 7: 42865 doi: 10.1038/srep42865 [PubMed: 28216645]
- Wong JR, Shao F (2017) Hole Transport in A-form DNA/RNA Hybrid Duplexes. *Sci Rep* 7: 40293 doi: 10.1038/srep40293 [PubMed: 28084308]
- Zhang DH, Zhou B, Huang Y, Xu LX, Zhou JQ (2006) The human Pif1 helicase, a potential *Escherichia coli* RecD homologue, inhibits telomerase activity. *Nucl Acids Res* 34: 1393–1404 doi: [PubMed: 16522649]
- Zhou R, Zhang J, Bochman ML, Zakian VA, Ha T (2014) Periodic DNA patrolling underlies diverse functions of Pif1 on R-loops and G-rich DNA. *Elife* 3: e02190 doi: 10.7554/eLife.02190 [PubMed: 24843019]
- Zhou X, Ren W, Bharath SR, Tang X, He Y, Chen C, Liu Z, Li D, Song H (2016) Structural and Functional Insights into the Unwinding Mechanism of *Bacteroides* sp Pif1. *Cell Rep* 14: 2030–2039 doi: 10.1016/j.celrep.2016.02.008 [PubMed: 26904952]




Figure 1: Role of nuclear Pif1 in different biological DNA transactions

Pif1 functions in (A) flap lengthening during Okazaki fragment processing; (B) modulating fork convergence during replication termination; (C) aiding G4 structure resolution; (D) promoting break-induced replication; (E) unwinding R-loops; and (F) regulating telomere length. Acetylation of Pif1 mediated by the lysine acetyltransferase NuA4 (Esa1) and lysine deacetylase (Rpd3) may play a regulatory role in how Pif1's functions are altered in all of these outlined pathways. Triple arrows indicate directionality of Pif1 activity.