

Neuroprotective Effect of Cannabidiol Against Hydrogen Peroxide in Hippocampal Neuron Culture

Jungham Kim, Ji Yu Choi, Jeongyeon Seo, and Insung S. Choi*

Abstract

Introduction: Reports on the neurotoxic and neuroprotective effects of cannabidiol (CBD) have not been in complete accord, showing different and somewhat contradictory results depending upon the brain cell types and experimental conditions employed. This work systematically examines the neuroprotective capability of CBD against oxidative stress (i.e., hydrogen peroxide [H₂O₂]) as well as its toxicity profile in the *in vitro* culture platform of primary hippocampal neurons.

Materials and Methods: The low cell-density (100 neurons per mm²) culture was used for analyzing the viability and morphology of neurons at a single-cell level with a confocal laser-scanning microscope (CLSM). Primary neurons were obtained from the hippocampal tissues of embryonic day-18 (E18) Sprague-Dawley rat pups and treated with CBD (0.1–100 μM) and/or H₂O₂ (0.1–50 μM) at 1 DIV (days *in vitro*).

Results: The lethal concentration 50 (LC₅₀) value (the concentration causing 50% cell death) of CBD was calculated to be 9.85 μM after 24 h of incubation, and that of H₂O₂ was 2.46 μM under the same conditions. The neuroprotection ratio of CBD against H₂O₂ ([H₂O₂] = 10 μM) was 2.40 with 5 μM of CBD, increasing the cell viability to 57% from 24%. The CLSM analysis suggested that the cell-death mechanisms were different for CBD and H₂O₂, and CBD did not completely rescue the morphological alterations of primary hippocampal neurons caused by H₂O₂, such as neurite degeneration, at least in the *in vitro* neuron culture.

Conclusion: Although CBD showed both neurotoxic and neuroprotective effects on hippocampal neurons in the *in vitro* setting, the use of low-concentrated (i.e., 5 μM) CBD, not causing toxic effects on the neurons, significantly rescued the neurons from the oxidative stress (H₂O₂), confirming its neuroprotection capability.

Keywords: cannabidiol; cell culture; hydrogen peroxide; neuroprotection; primary hippocampal neurons

Introduction

(–)-Cannabidiol (CBD, IUPAC name: 2-[(1R,6R)-3-methyl-6-prop-1-en-2-ylcyclohex-2-en-1-yl]-5-pentylbenzene-1,3-diol, C₂₁H₃₀O₂), the main nonpsychoactive compound among 100+ phytocannabinoids extracted from the genus *Cannabis*, anecdotally—if not clinically—shows potential therapeutic benefits in various health care applications, including epilepsy, inflammation, cancer, and neurodegenerative diseases.^{1–5} The year of 2018 witnessed the first approval of *Cannabis*-derived CBD (Epidiolex[®]) for the treatment of epileptic seizures associated with the Lennox–Gastaut or Dravet syndrome in patients 2 years of age and older by the U.S. Food and Drug Administration (FDA).^{6–8}

In addition to the therapeutic effect on epileptic seizures, previous reports suggest that CBD is neuroprotective. The earliest research involved the neuroprotective action of CBD in the rat cortical neuron culture exposed to toxic levels of glutamate and tert-butyl hydroperoxide.⁹ The bidirectional Ca²⁺ regulation capability of CBD for primary hippocampal cells (neuron:glia = 2:1) was reported, indicating the regulatory role of CBD in Ca²⁺ homeostasis^{10–12}: for example, the intracellular calcium concentration ([Ca²⁺]_i) was reduced by about 25% of basal Ca²⁺ level in both neurons and glia by CBD application to a HEPES-based buffer solution with double K⁺ concentration, while CBD caused the [Ca²⁺]_i elevation, with no significant cell death, by

Department of Chemistry, KAIST, Daejeon, Korea.

*Address correspondence to: Insung S. Choi, PhD, Department of Chemistry, KAIST, Daejeon 34141, Korea, E-mail: ischoi@kaist.ac.kr

© Jungham Kim et al. 2020; Published by Mary Ann Liebert, Inc. This Open Access article is distributed under the terms of the Creative Commons Attribution Noncommercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and the source are cited.

about 45% of basal Ca^{2+} level under physiological conditions. The neuroprotective properties of CBD against oligomycin, an inhibitor of ATP synthetase,¹⁰ and ammonium acetate and ethanol in the hippocampal neuron culture also have been reported.¹³

Oxidative stress is thought to cause pathological conditions in neurodegenerative diseases, such as Alzheimer's disease, Parkinson's disease, Huntington's disease, and amyotrophic lateral sclerosis.^{14,15} The cytoprotective effect of CBD against hydrogen peroxide (H_2O_2), a crucial metabolic molecule in the oxidative stress,¹⁶ was reported for cerebellar granule cells¹⁷ and oligodendrocyte progenitor cells,¹⁸ but there have been no reports on its effect against H_2O_2 in the neurons that are more relevant to the brain function, such as hippocampal neurons.¹⁹ H_2O_2 has been used as a model molecule to study oxidative stress-induced neural death and also to screen neuroprotective agents on the cultured neurons.^{16,20–25} As our on-going research program on neurochemistry and neurophysiology,^{26–31} this study investigates the neuroprotective effect of CBD against H_2O_2 in the hippocampal neuron culture.

In addition to the neuroprotective property, CBD has also been reported to show different toxicity profiles on different brain cell types. For example, even $0.1 \mu\text{M}$ of CBD increased $[\text{Ca}^{2+}]_i$ and induced cell death in the case of optic nerve oligodendrocytes after 24 h of incubation, while $10 \mu\text{M}$ of CBD did not affect the viability of cortical neurons (CBD exposure time: 20–30 min).³² However, no cytotoxicity was observed with $1 \mu\text{M}$ of CBD for oligodendrocyte progenitor cells (CBD exposure time: 24 h).¹⁸ Furthermore, CBD of below $4 \mu\text{M}$ did not significantly increase hypodiploid apoptotic cells for primary microglial cells, but CBD of above $8 \mu\text{M}$ induced apoptosis in a time- and concentration-dependent manner, confirmed by increase in hypodiploid cells and DNA-strand breaks, and activation of capase-8 and -9 (CBD exposure time: 24 h).³³ Considering the previous somewhat contradictory results, we first systematically evaluated the *in vitro* neurotoxic effect of CBD to hippocampal neurons with different CBD concentrations.

Materials and Methods

Chemicals and reagents

CBD (10-mM stock in dimethyl sulfoxide (DMSO), storage at -20°C) was obtained from Cayman Chemical Company. Neurobasal™ medium (Thermo Fisher Scientific), B-27™ serum-free supplement (50×; Thermo Fisher Scientific), GlutaMAX™ supplement (100×; Thermo Fisher Scientific), calcein acetoxymethyl ester

(AM; Thermo Fisher Scientific), ethidium homodimer-1 (Thermo Fisher Scientific), CellTracker™ Red CMTPX Dye (Thermo Fisher Scientific), Hank's Balanced Salt Solution (HBSS; Welgene), phosphate-buffered saline (PBS, 10 mM, pH 7.4; Welgene), penicillin/streptomycin (5000 U/mL of penicillin and 5000 $\mu\text{g}/\text{mL}$ of streptomycin; Welgene), L-glutamic acid (Sigma), poly-D-lysine hydrobromide (Sigma), paraformaldehyde (Sigma), Triton™ X-100 (Sigma), anti- β -tubulin III antibody produced in rabbit (Sigma), Alexa Fluor™ 488 phalloidin (Invitrogen), Alexa Fluor 594 anti-rabbit goat antibody (Invitrogen), DMSO (Junsei), H_2O_2 (Junsei), and Accumax™ (Stemcell Technologies) were used as received. Antifade mounting medium with 4',6-diamidino-2-phenylindole (DAPI) was obtained from Vector Laboratories.

Primary hippocampal neuron culture

Primary hippocampal neurons were dissected from embryonic day-18 (E18) Sprague-Dawley rat pups and treated with Accumax for 10 min to detach the cells from primary tissues. Hippocampi were washed four times with HBSS, suspended in HBSS, centrifuged at 1000 rpm for 3 min, and resuspended with Neurobasal medium containing B-27 serum-free supplement, 2-mM GlutaMAX, 12.5- μM L-glutamic acid, and 1% penicillin/streptomycin. After filtration using a cell strainer, the hippocampal neurons were seeded on glass coverslips coated with poly-D-lysine in a 24-well plate at density of 100 cells/ mm^2 . The cultures were maintained at 37°C under a humidified atmosphere of 5% CO_2 . This study was approved by IACUC (Institutional Animal Care and Use Committee) of KAIST.

Cell viability assay

For neurotoxicity experiments, the culture media were replaced with the Neurobasal medium containing CBD (0.1–100 μM) or H_2O_2 (0.1–50 μM) at 1 DIV (days *in vitro*). In the case of H_2O_2 , heat potentially accelerated the decomposition rate of H_2O_2 , and, therefore, 0.1- or 1-mM H_2O_2 stock solution was prepared in the 4°C cold medium. The fresh Neurobasal medium was poured to the wells and diluted with a predetermined amount of the H_2O_2 stock solution at 4°C to minimize the decomposition of H_2O_2 , instead of diluting the media at ambient temperature. For neuroprotection experiments, the culture media were replaced, at 1 DIV, with 990 μL of fresh Neurobasal medium containing CBD of various concentrations (0, 0.1, 1, 2, 3, or 5 μM), followed by the addition of 10 μL of 0.1-mM H_2O_2 stock solution. For the 0- μM case of CBD 0.1%

DMSO was used. As another set, the neurons were also preincubated with 5 μM of CBD for 1 h at 1 DIV, before H_2O_2 treatment. After incubation for 24 h, the cells were treated with 2- μM calcein AM and 4- μM ethidium homodimer-1 in PBS for 20 min at 37°C. Images of live and dead cells were obtained with an LSM 700 confocal microscope (Carl Zeiss). Cell viability was estimated by using ImageJ software (NIH) and image analysis of ZEN software (Carl Zeiss), and expressed as %viability (mean \pm standard error) after at least three independent cultures with three replicates.

Fluorescent staining

Neurons were treated with the Neurobasal media containing CBD (1, 5, or 10 μM) or 0.1% DMSO at 1 DIV. After incubation for 24 h, the neurons were treated with Celltracker Red CMTPX Dye at 37°C for 30 min, fixed with 4% paraformaldehyde in PBS for 15 min at room temperature, and washed three times with PBS each for 5 min. The fixed samples were permeabilized with 0.1% Triton X-100 solution in PBS for 10 min at room temperature and washed three times with PBS each for 5 min. The samples were then treated with Alexa Fluor 488 phalloidin for visualizing F-actin or rabbit-produced β -tubulin III primary antibody, incubated at 37°C, and washed with PBS three times each for 5 min. For visualizing β -tubulin III, the samples were additionally treated with Alexa Fluor 594 goat anti-rabbit secondary antibody. Finally, the samples were mounted on slide glasses with the DAPI solution to stain nuclei for 30 min. Fluorescence images were obtained with an LSM 700 confocal microscope.

Results and Discussion

The primary hippocampal neurons, after dissociation from the hippocampal tissues of E18 Sprague-Dawley rat pups, were seeded on a poly-D-lysine-coated glass coverslip at density of 100 cells/ mm^2 in the Neurobasal medium, which was replaced with the medium containing CBD (0.1–100 μM) at 1 DIV. In the case of 1% DMSO, which was the DMSO concentration for preparation of 100- μM CBD solution, we did not observe any noticeable difference compared with the nontreated control in cell viability and morphology. Throughout this article, we used the value of %viability, which was defined as % of cell viability with the nontreated control as a reference (i.e., the observed viability divided by the viability of the nontreated cells). After incubation for 24 h (i.e., at 2 DIV), the cells were stained with calcein AM for live cells

(green) and ethidium homodimer-1 for dead cells (red), and the confocal laser-scanning microscope (CLSM) images were taken for the viability analysis.



The CLSM images (Fig. 1a) arguably showed the concentration-dependent neurotoxicity of CBD in the hippocampal neuron culture. For example, while the cell death was not noticeable up to 5 μM of CBD, the red fluorescence in CLSM images indicated that most hippocampal neurons were dead with 30 μM of CBD. The quantitative analysis showed that the %viability decreased to 44.80% with 10 μM of CBD, and the lethal concentration 50 (LC_{50}) value, the concentration at which the %viability was 50%, was calculated to be 9.85 μM (Fig. 1b). It is to be noted that this result is the first quantitative *in vitro* data, affording the LC_{50} value of CBD, for the hippocampal-neuron culture.

The careful analysis of the CLSM images suggested that CBD might disrupt the cell adhesion. Some CLSM images had green-stained vestiges, and some dying cells, stained in both red and green, had partially detached neurites and cell body. We could frequently find the costained neurons with partial detachment as well as live neurons with no or few neurites, especially in the case of 10 μM of CBD.

To get some insights into the death-related cell detachment, the cells that remained attached to the culture plate were stained, at 2 DIV, with phalloidin for F-actin (green), CellTracker Red CMTPX Dye for cytoplasm (red), and DAPI for nuclei (blue). The CLSM images (Fig. 2) clearly showed that the neurons did not develop properly in the CBD-containing medium ([CBD] = 10 μM), although the quantitative analysis was not feasible due to the cell detachment.


Although the developmental rate, judged by the Banker's classification of developmental stages,³⁴ seemed not to be affected by CBD, it was apparent that the neurons presented few, if not no, filopodia and lamellipodia structures. For example, no axonal branches were observed, and the growth cones were shrunken in size noticeably. In addition, we observed some cellular debris, such as only nuclei (top image in Fig. 2), condensed chromatin, and ruptured cells (middle image in Fig. 2). Based on the observation, it was hypothesized that the neuronal death by CBD might be considered to be a kind of detachment-induced apoptosis/anoikis (i.e., apoptosis following the loss of cell anchorage, especially the loss of cell/matrix interaction).^{35,36}

Detailed elucidation on the neurotoxic mechanism of CBD remains to be seen at this moment, but our results implied that the lipophilic CBD might interfere




with the integrity of cell membranes, causing the cell rupture and death, at its high concentration. However, it is also to mention that our neurotoxicity results are not translated directly to the *in vivo* conditions. The brain is composed of neuronal and non-neuronal cells and others, and the *in vivo* tolerance of hippocampal neurons to CBD would be different from the *in vitro* conditions. For example, glial cells, such as astrocytes, play supporting roles in the brain. Relevant reports showed that the *in vitro* neurotoxicity of unnatural monosaccharides to primary hippocampal neurons was mitigated when a hippocampal tissue was used^{26,28} or a neuron/astrocyte coculture was employed.²⁷ In this respect, it could be explained that the coculture system of neuron and glia (2:1) did not show the neurotoxicity of CBD.^{10–12}

It is to note that the H_2O_2 neurotoxicity experiments should be performed with caution under the tightly controlled conditions to ensure the data reproducibility,



because of the rapid decomposition of H_2O_2 at ambient temperature. Specifically, the H_2O_2 stock solution (0.1 or 1 mM) was to be prepared at the low temperature (4°C) and transferred to the culture medium right after preparation (within 5 min). It was also found to be critical



that the dilution of the H_2O_2 stock solution in the culture medium was done at 4°C , not at ambient temperature.

After 24 h of incubation at 37°C , the %viability was obtained by the analysis of the CLSM images (Fig. 3a). There was no significant difference in viability, compared with the control, up to $1\ \mu\text{M}$ of H_2O_2 , but the %viability decreased with H_2O_2 of $3\ \mu\text{M}$ and above. The LC_{50} value of H_2O_2 for hippocampal neurons was calculated to be $2.46\ \mu\text{M}$, and the %viability decreased to 16.49% with $10\ \mu\text{M}$ of H_2O_2 (Fig. 3b). The LC_{50} value of H_2O_2 has previously been reported to be about $40\ \mu\text{M}$ at the cell density of $700\ \text{cells}/\text{mm}^2$ and about $92\ \mu\text{M}$ at that of $2000\ \text{cells}/\text{mm}^2$, after 24 h of incubation, for cortical neurons.²¹ In comparison, we obtained a lower LC_{50} value ($2.46\ \mu\text{M}$), which might be caused by the difference in cell type and/or the lower cell density ($100\ \text{cells}/\text{mm}^2$) used for our system.

H_2O_2 is known as a trigger of oxidative stress-induced apoptosis.^{16,21} Accordingly, we observed the round neuronal morphology of neurons with no or few neurites, one of the major features of apoptosis, after the treatment of H_2O_2 . In addition, the weak and broken degenerative neurites and cellular debris after degeneration in the CLSM images also suggested H_2O_2 -induced apoptosis.^{37,38}

CBD is known to be an antioxidant,³⁹ with the similar oxidation potential to butylhydroxytoluene (BHT).^{9,40} CBD was reported to reduce the toxic levels of glutamate-induced neurotoxicity to about 40%, with its neuroprotective ability that was similar to BHT and more efficient than commercial antioxidants, such as ascorbate and α -tocopherol, for primary cortical neurons.⁹ Phenolic cannabinoids, including CBD, also have been shown to be the potent cannabinoid receptor 1-independent antioxidants, and protected the mouse hippocampal HT22 cell line up to $\sim 60\%$ from oxidative stress induced by H_2O_2 ($[\text{CBD}] = 5\ \mu\text{M}$).¹⁷

In addition, CBD ($10\ \mu\text{M}$) was reported to reduce the death of neuron-like PC12 cells from 39% to 12% against β -amyloid peptides, and decrease the reactive oxygen species production, lipid peroxidation, intracellular calcium, and caspase-3 levels.⁴¹ Several other compounds have also been studied for potential protection of cortical or hippocampal neurons against H_2O_2 .^{42–44} Y-27632, a Rho-associated kinase (ROCK) inhibitor, increased the cell viability to 87% from 66.5% against H_2O_2 at maximum.⁴² *N*-Acetyl cysteine, inhibiting the oxidative stress by downregulating mitogen-activated protein kinase (MAPK) signal transduction and tau phosphorylation, was reported to be effective in the




FIG. 5. High-magnification CLSM images of hippocampal neurons after 24-h treatment of (middle) H_2O_2 ($10\ \mu\text{M}$), and (right) H_2O_2 ($10\ \mu\text{M}$), and CBD ($5\ \mu\text{M}$). The cells were stained in green (F-actin), red (β -tubulin III), and blue (nuclei).

neuroprotection against H_2O_2 .⁴³ In the case of vitamin E, a fat-soluble antioxidant, also exhibited a neuroprotective capability, increasing the cell viability from 32% to 78% in the hippocampal neuron culture.⁴⁴

In this work, we investigated the neuroprotective effect of CBD against H_2O_2 for primary hippocampal neurons. Specifically, we used $10\ \mu\text{M}$ of H_2O_2 based on our result of the neurotoxicity effect of H_2O_2 : this concentration of H_2O_2 , inducing about 80% cell death, gave a suitable condition for neuroprotective studies. We also limited the CBD concentrations to less than $10\ \mu\text{M}$, considering the previous studies on the plasma and brain pharmacokinetic profile of CBD in rats and mice, indicating that the CBD concentrations of $\geq 10\text{--}20\ \mu\text{M}$ in the *in vitro* studies would be supraphysiological.^{45,46}

The CLSM images clearly showed that CBD was effective in the neuroprotection of primary hippocampal neurons against H_2O_2 ($10\ \mu\text{M}$) (Fig. 4a). The quantitative analysis on the neuroprotection was performed by calculating the protection ratio of CBD, $PR_{[\text{CBD}]}$, by dividing the %viability values for [+ H_2O_2][+CBD] cases by that for the [+ H_2O_2][−CBD] case ($[\text{H}_2\text{O}_2] = 10\ \mu\text{M}$).

The bar graph of $PR_{[\text{CBD}]}$ versus [CBD] confirmed the concentration-dependent neuroprotective effects of CBD against H_2O_2 in the *in vitro* culture of primary hippocampal neurons (Fig. 4b). For example, the $PR_{[\text{CBD}]}$ value was 1.88 for $3\ \mu\text{M}$ of CBD, which further increased to 2.40 for $5\ \mu\text{M}$ of CBD. Interestingly, this $PR_{[\text{CBD}]}$ value was similar to the reported maximum PR value for vitamin E, 2.43,⁴⁴ implying that the antioxidant property of CBD played an important role in the neuroprotection against H_2O_2 . The preincubation of the neurons with CBD ($5\ \mu\text{M}$, 1 h) did not lead to any noticeable neuroprotection against H_2O_2 .

Although CBD increased the %viability against H_2O_2 , showing its neuroprotective capability, the analysis of the high-resolution CLSM images indicated that CBD did not completely prevent the neurite degeneration in apoptotic process even at its concentration of $5\ \mu\text{M}$ (Fig. 5). Therefore, we hypothesized that CBD acted as an antioxidant based on its phenolic structure and partially protected the hippocampal neurons from the initiation and progress of apoptosis caused by oxidative stress. It has been reported that CBD protected oligodendrocyte progenitor cells from apoptosis by decreasing the expression of the proapoptotic proteins or apoptotic effectors, such as Bax and caspase-12, and also increasing the antiapoptotic proteins, such as Bcl-2.¹⁸

Another possibility would be the regulation of $[\text{Ca}^{2+}]_i$.^{10–12} In the H_2O_2 -induced apoptosis was reported

the $[\text{Ca}^{2+}]_i$ increase, by the disruption of Ca^{2+} -storing organelles and also calcium influx through activation of several receptors and channels, such as *N*-methyl-D-aspartate (NMDA) subtype receptor and transient receptor potential melastatin 7 (TRPM7) channel.^{21,23–25} Previous reports also showed that the H_2O_2 -induced cell death could be prevented by controlling the Ca^{2+} influx. For example, H_2O_2 -induced neuronal apoptosis was inhibited by using the antagonist of NMDA receptors, such as MK801.²⁵ The cytotoxicity of H_2O_2 was also attenuated by blocking the TRPM7 channel with the antagonist, 2-APB, or knocking down the TRPM7 channel expression.²⁴

Another plausible mechanism on the neuroprotective effects of CBD against H_2O_2 might be activation of selective receptor mechanisms; H_2O_2 is reported to reversibly sulfenylate cysteines C201 and C208 and deactivate monoacylglycerol lipase, which cleaves 2-arachidonoyl-*sn*-glycerol, a ligand for cannabinoid receptor type 1 receptor.⁴⁷

Conclusions

In summary, we used a low-density, neuron culture platform for evaluating both neurotoxic and neuroprotective effects of CBD on primary hippocampal neurons. Our *in vitro* study showed that CBD was neurotoxic, although its LC_{50} value was high (about $10\ \mu\text{M}$), suggesting that the detailed pharmacokinetic studies on the bioavailability of CBD be done with various administration methods, such as ingestion, inhalation, and transdermal or topical delivery. Biological modes of CBD action on neurons have not been elucidated fully, but our work clearly confirmed that CBD was highly effective in neutralizing the H_2O_2 -induced neurotoxicity on primary hippocampal neurons, which is intimately involved in the onset and progress of neurodegenerative diseases.

Author Disclosure Statement

No competing financial interests exist.

Funding Information

This work was supported by Cannabis Medical, Inc.

References

1. Reekie TA, Scott MP, Kassiou M. The evolving science of phytocannabinoids. *Nat Rev Chem*. 2017;2:0101.
2. Koppel BS, Brust JC, Fife T, et al. Systematic review: efficacy and safety of medical marijuana in selected neurological disorders. *Neurology*. 2014; 82:1556–1563.
3. Scuderi C, Filippis DD, Iuvone T, et al. Cannabidiol in medicine: a review of its therapeutic potential in CNS disorders. *Phytother Res*. 2009;23:597–602.
4. Iuvone T, Esposito G, Filippis DD, et al. Cannabidiol: a promising drug for neurodegenerative disorders? *CNS Neurosci Ther*. 2009;15:65–75.
5. Hayakawa K, Mishima K, Fujiwara M. Therapeutic potential of non-psychoactive cannabidiol in ischemic stroke. *Pharmaceuticals*. 2010;3:2197–2212.

6. Devinsky O, Cross JH, Laux L, et al. Trial of cannabidiol for drug-resistant seizures in the Dravet syndrome. *N Engl J Med*. 2017;376:2011–2020.
7. Devinsky O, Patel AD, Cross JH, et al. Effect of cannabidiol on drop seizures in the Lennox-Gastaut syndrome. *N Engl J Med*. 2018;378:1888–1897.
8. GW Pharmaceuticals. EPIDIOLEX[®] (cannabidiol) oral solution – the first FDA-approved plant-derived cannabinoid medicine – now available by prescription in the U.S. <http://ir.gwpharm.com/news-releases/news-release-details/epidiolex-cannabidiol-oral-solution-first-fda-approved-plant>. (accessed March 1, 2019).
9. Hampson AJ, Grimaldi M, Axelrod J, et al. Cannabidiol and (-) Δ^9 -tetrahydrocannabinol are neuroprotective antioxidants. *Proc Natl Acad Sci U S A*. 1998;95:8268–8273.
10. Ryan D, Drysdale AJ, Lagourcade C, et al. Cannabidiol targets mitochondria to regulate intracellular Ca^{2+} levels. *J Neurosci*. 2009;29:2053–2063.
11. Drysdale AJ, Ryan D, Pertwee RG, et al. Cannabidiol-induced Ca^{2+} elevation in hippocampal cells. *Neuropharmacology*. 2006;50:621–631.
12. Ryan D, Drysdale AJ, Pertwee RG, et al. Interactions of cannabidiol with endocannabinoid signaling in hippocampal tissue. *Eur J Neurosci*. 2007;25:2093–2102.
13. Kinney WA, McDonnell ME, Zhong HM, et al. Discovery of KLS-13019, a cannabidiol-derived neuroprotective agent, with improved potency, safety, and permeability. *ACS Med Chem Lett*. 2016;7:424–428.
14. Kim GH, Kim JE, Rhie SJ, et al. The role of oxidative stress in neurodegenerative diseases. *Exp Neurobiol*. 2015;24:325–340.
15. Barnham KJ, Master CL, Bush AI. Neurodegenerative diseases and oxidative stress. *Nat Rev Drug Discov*. 2004;3:205–214.
16. Ricart KC, Fiszman ML. Hydrogen peroxide-induced neurotoxicity in cultured cortical cells grown in serum-free and serum-containing media. *Neurochem Res*. 2001;26:801–808.
17. Marsicano G, Moosmann B, Hermann H, et al. Neuroprotective properties of cannabinoids against oxidative stress: role of the cannabinoid receptor CB1. *J Neurochem*. 2002;80:448–456.
18. Mecha M, Torrao AS, Mestre L, et al. Cannabidiol protects oligodendrocyte progenitor cells from inflammation-induced apoptosis by attenuating endoplasmic reticulum stress. *Cell Death Dis*. 2012;3:e331.
19. Lupia CR, Hu Y, Devinsky O, et al. Cannabinoids as hippocampal network administrators. *Neuropharmacology*. 2017;124:25–37.
20. Whittemore ER, Loo DT, Cotman CW. Exposure to hydrogen peroxide induces cell death via apoptosis in cultured rat cortical neurons. *Neuroreport*. 1994;5:1485–1488.
21. Whittemore ER, Loo DT, Watt JA, et al. A detailed analysis of hydrogen peroxide-induced cell death in primary neuronal culture. *Neuroscience*. 1995;67:921–932.
22. Desagher S, Glowinski J, Premont J. Astrocytes protect neurons from hydrogen peroxide toxicity. *J Neurosci*. 1996;16:2553–2562.
23. Hoty KR, Gallagher AJ, Hastings TG, et al. Characterization of hydrogen peroxide toxicity in cultured rat forebrain neurons. *Neurochem Res*. 1997;22:333–340.
24. Coombes E, Jiang J, Chu XP, et al. Pathophysiologically relevant levels of hydrogen peroxide induce glutamate-independent neurodegeneration that involves activation of transient receptor potential melastatin 7 channels. *Antioxid Redox Signal*. 2011;14:1815–1827.
25. Mailly F, Marin P, Israël M, et al. Increase in external glutamate and NMDA receptor activation contribute to H_2O_2 -induced neuronal apoptosis. *J Neurochem*. 1999;73:1181–1188.
26. Choi JY, Seo J, Park M, et al. Multiplexed metabolic labeling of glycoconjugates in polarized primary cerebral cortical neurons. *Chem Asian J*. 2018;13:3480–3484.
27. Choi JY, Park M, Cho H, et al. Neuro-compatible metabolic glycan labeling of primary hippocampal neurons in non-contact, sandwich-type neuron-astrocyte co-culture. *ACS Chem Neurosci*. 2017;8:2607–2612.
28. Kang K, Joo S, Choi JY, et al. Tissue-based metabolic labeling of polysialic acids in living primary hippocampal neurons. *Proc Natl Acad Sci U S A*. 2015;112:E241–E248.
29. Marcus M, Baranes K, Park M, et al. Interactions of neurons with physical environments. *Adv Healthcare Mater*. 2017;6:1700267.
30. Kang K, Park YS, Park M, et al. Axon-first neurite outgrowth on vertical nanowires. *Nano Lett*. 2016;16:675–680.
31. Cho WK, Kang K, Kang G, et al. Pitch-dependent acceleration of neurite outgrowth on nanostructured anodized aluminum oxide substrates. *Angew Chem Int Ed*. 2010;49:10114–10118.
32. Mato S, Sánchez-Gómez MV, Matute C. Cannabidiol induces intracellular calcium elevation and cytotoxicity in oligodendrocytes. *Glia*. 2010;58:1739–1747.
33. Wu HY, Goble K, Mecha M, et al. Cannabidiol-induced apoptosis in murine microglial cells through lipid raft. *Glia*. 2012;60:1182–1190.
34. Dotti CG, Sullivan CA, Banker GA. The establishment of polarity by hippocampal neurons in culture. *J Neurosci*. 1988;8:1454–1468.
35. Grossmann J. Molecular mechanisms of “detachment-induced apoptosis-Anoikis.” *Apoptosis*. 2002;7:247–260.
36. Bonfoco E, Chen W, Paul R, et al. Beta1 integrin antagonism on adherent, differentiated human neuroblastoma cells triggers an apoptotic signaling pathway. *Neuroscience*. 2000;101:1145–1152.
37. Lesay A, Hickman JA, Gibson RM. Disruption of focal adhesions mediates detachment during neuronal apoptosis. *Neuroreport*. 2001;10:2111–2115.
38. Gibson RM. Caspase activation is downstream of commitment to apoptosis of Ntera-2 neuronal cells. *Exp Cell Res*. 1999;251:203–212.
39. Borges RS, Batista J Jr., Viana RB, et al. Understanding the molecular aspects of tetrahydrocannabinol and cannabidiol as antioxidants. *Molecules*. 2013;18:12663–12674.
40. Babich H. Butylated hydroxytoluene (BHT): a review. *Environ Res*. 1982;29:1–29.
41. Iuvone T, Esposito G, Esposito R, et al. Neuroprotective effect of cannabidiol, a non-psychoactive component from Cannabis sativa, on beta-amyloid-induced toxicity in PC12 cells. *J Neurochem*. 2004;89:134–141.
42. Wang X, Wang B, Li Z, et al. Neuroprotection effect of Y-27632 against H_2O_2 -induced cell apoptosis of primary cultured cortical neurons. *RSC Adv*. 2016;6:49187.
43. Wu W, Liu BH, Xie CL, et al. Neuroprotective effects of N-acetyl cysteine on primary hippocampus neurons against hydrogen peroxide-induced injury are mediated via inhibition of mitogen-activated protein kinases signal transduction and antioxidative action. *Mol Med Rep*. 2018;17:6647–6654.
44. Lee HJ, Lee JH, Cho NS. Study on the effect of vitamin E on cultured hippocampal neurons damaged by hydrogen peroxide. *Kor J Ori Med Physiol Pathol*. 2003;17:447–450.
45. Deiana S, Watanabe A, Yamasaki Y, et al. Plasma and brain pharmacokinetic profile of cannabidiol (CBD), cannabidivarin (CBDV), Δ^9 -tetrahydrocannabinavarin (THCV) and cannabigerol (CBG) in rats and mice following oral and intraperitoneal administration and CBD action on obsessive-compulsive behaviour. *Psychopharmacology*. 2012;219:859–873.
46. Bih CI, Chen T, Nunn AV, et al. Molecular targets of cannabidiol in neurological disorders. *Neurotherapeutics*. 2015;12:699–730.
47. Dotsey EY, Jung KM, Basit A, et al. Peroxide-dependent MGL sulfenylation regulates 2-AG-mediated endocannabinoid signaling in brain neurons. *Chem Biol*. 2015;22:619–628.

Cite this article as: Kim J, Choi JY, Seo J, Choi IS (2021) Neuroprotective effect of cannabidiol against hydrogen peroxide in hippocampal neuron culture, *Cannabis and Cannabinoid Research* 6:1, 40–47, DOI: 10.1089/can.2019.0102.

Abbreviations Used

AM = acetoxymethyl ester
 BHT = butylhydroxytoluene
 CBD = cannabidiol
 CLSM = confocal laser-scanning microscope
 DAPI = 4',6-diamidino-2-phenylindole
 DIV = days in vitro
 DMSO = dimethyl sulfoxide
 E18 = embryonic day 18
 FDA = U.S. Food and Drug Administration
 H_2O_2 = hydrogen peroxide
 HBSS = Hank's Balanced Salt Solution
 LC_{50} = lethal concentration 50
 MAPK = mitogen-activated protein kinase
 NMDA = N-methyl-D-aspartate
 PBS = phosphate-buffered saline
 ROCK = Rho-associated kinase
 SE = standard error
 TRPM7 = transient receptor potential melastatin 7