



# Draft Genome Sequences of Pelagimyophage Mosig EXVC030M and Pelagipodophage Lederberg EXVC029P, Isolated from Devil's Hole, Bermuda

Holger H. Buchholz,<sup>a</sup> Michelle Michelsen,<sup>a</sup> Rachel J. Parsons,<sup>b</sup> Nicholas R. Bates,<sup>b,c</sup>  Ben Temperton<sup>a</sup>

<sup>a</sup>School of Biosciences, University of Exeter, Exeter, United Kingdom

<sup>b</sup>Bermuda Institute of Ocean Sciences, St. George's, Bermuda

<sup>c</sup>School of Ocean and Earth Sciences, Waterfront Campus, University of Southampton, Southampton, United Kingdom

**ABSTRACT** We present the genomes of two isolated bacteriophages infecting *Pelagibacter ubique* HTCC1062. *Pelagibacter* phage Mosig EXVC030M (*Myoviridae*) and *Pelagibacter* phage Lederberg EXVC029P (*Podoviridae*) were isolated by dilution-to-extinction culturing from the oxygen minimum zone at Devil's Hole (Harrington Sound, Bermuda).

Viruses infecting the heterotrophic bacterial clade of *Pelagibacterales* are an important component of marine microbial communities throughout global oceans (1). Since the discovery and first isolation of four pelagiphages in 2013 (2), 38 more have been isolated and sequenced (3–5). Out of the 38 isolated pelagiphages, 36 belong to the *Podoviridae* family, with only one species each of *Myoviridae* and *Siphoviridae*. Here, we report the draft genome sequences of a novel pelagimyophage and a novel pelagipodophage, both isolated on *Pelagibacter ubique* HTCC1062.

A 2-liter water sample was taken (12 July 2019) using a hand-held Niskin bottle, fired at a 20-m depth at Devil's Hole, Bermuda, a seasonal oxygen minimum zone in Bermuda (6) (latitude 32.32421, longitude –64.71849). The water sample was taken to the Bermuda Institute of Ocean Sciences for processing, where planktonic cells were removed with 0.1- $\mu$ m polyethersulfone filters. Viruses were concentrated by tangential flow filtration (50R VivaFlow 100-kDa Hydrosart filter; Sartorius Lab Instruments, Göttingen, Germany). We used previously described dilution-to-extinction-based methods (4) with HTCC1062 as a bait host (grown in artificial seawater medium ASM1 [7]) in 96-well Teflon plates, which does not rely on plaque formation, because the host does not grow on solid medium. The purification process was repeated five times; nonetheless, final sequence data contained two genomes, suggesting an impure culture.

For DNA isolation, a 50-ml HTCC1062 culture ( $10^6$  cells/ml), amended with 5 ml of 0.1- $\mu$ m-filtered lysate, was grown in ASM1 (18°C) until cell death (detected via flow cytometry). Debris was removed using 0.1- $\mu$ m-pore polyvinylidene difluoride (PVDF) filters, and lysate was subjected to PEG8000/NaCl DNA isolation (modified from <https://doi.org/10.17504/protocols.io.c36yrd>, as described previously (4)).

DNA libraries (Nextera XT) were prepared and sequenced by the Exeter Sequencing Service (Illumina paired end [2 × 250 bp], NovaSeq S Prime [SP], targeting 30-fold coverage). Raw reads (13.18 million) were trimmed, quality controlled, and error corrected using tadpole (default settings [8] within BMAP v38.22 [<https://sourceforge.net/projects/bbmap/>]) and assembled with SPAdes v3.13 (7). Viral contigs were confirmed and gene called with VirSorter v1.05 (9) and imported into DNA Master v5.23.3 (10) for manual curation with additional gene calls using GenMark v2.5 (11), GeneMarkS v4.28 (12), GeneMarkS-2 v1.14 (13), GeneMark.hmm v3.25 (14), Glimmer v3.02 (15), and Prodigal v2.6.3 (16). Open reading frames were annotated with NCBI's nonredundant

**Citation** Buchholz HH, Michelsen M, Parsons RJ, Bates NR, Temperton B. 2021. Draft genome sequences of pelagimyophage Mosig EXVC030M and pelagipodophage Lederberg EXVC029P, isolated from Devil's Hole, Bermuda. *Microbiol Resour Announc* 10:e01325-20. <https://doi.org/10.1128/MRA.01325-20>.

**Editor** Catherine Putonti, Loyola University Chicago


**Copyright** © 2021 Buchholz et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ben Temperton, [b.temperton@exeter.ac.uk](mailto:b.temperton@exeter.ac.uk).

**Received** 24 November 2020

**Accepted** 2 February 2021

**Published** 18 February 2021



**FIG 1** Bayesian inference tree from conserved genes found in pelagiphages (2, 3, 5), contigs from Mediterranean metagenomes (uvMed) (24), and putative pelagimyophages from genome-resolved metagenomics (PMP-MAVG) (25). (A) Terminase large subunit, tail sheath protein, and tail tube protein; (B) head-tail connector protein, capsid assembly protein, major capsid protein, tail tubular protein A, and putative acetyltransferase. Branch support values of 1 were omitted for clarity. The scale bar represents the estimated substitution per site.

protein database (17) and phmmer v2.41.1 (18) against the UniProtKB, uniprotrefprot (19), SWISS PROT (20), and Pfam (21) databases (accessed May 2020) and were evaluated using a previously described scoring system (10). Genome completion was verified with CheckV (22). Sequences similar to our isolates were identified with ClusterGenomes v5.1 (<https://github.com/simroux/ClusterGenomes>) and vConTACT 2 v0.9.19 (23) using previously isolated Pelagiphages (2, 3, 5), fosmid-derived contigs from Mediterranean metagenomes (uvMed) (24), and putative pelagimyophages from genome-resolved metagenomics (PMP-MAVG) (25). Conserved genes were identified (GET\_HOMOLOGUES v09072020 [22]), aligned (MUSCLE v3.8.1551 [26]), curated (Gblocks v0.91b [27]), and concatenated manually (all with default settings). Bayesian inference trees were generated via Phylogeny.fr

(28) using MRBAYES v3.2.7 (29) (100,000 generations, sampled every 10 generations, 5,000 tree burn-in) (Fig. 1).

Pelagimyophage Mosig (named after microbiologist Gisela Mosig in recognition of her work on *Escherichia coli* phage T4) was 141,462 bp long (348× coverage; GC content, 30.01%), linear, and 75.73% complete (CheckV [22]). Out of 208 genes, 98 were putative, 3 were tRNAs, 30 were structural, and 77 were associated with DNA replication.

*Pelagibacter* phage Lederberg (named after microbiologist Esther Lederberg in recognition of her work on the *E. coli* phage λ) was 33,623 bp long (5,849× coverage; GC content, 33.13%) and predicted as circularly permuted/complete. Lederberg had a total of 71 genes, out of which 9 were structural, 8 were associated with DNA replication, and 54 were without known function.

**Data availability.** The complete genome sequences were deposited under GenBank accession numbers [MT647605](#) (Lederberg) and [MT647606](#) (Mosig). The corresponding read data were deposited in the Sequence Read Archive (SRA) under BioProject number [PRJNA625644](#) and SRA accession number [SRR12024324](#).

## ACKNOWLEDGMENTS

The efforts of H.H.B. in this work were funded by the Natural Environment Research Council (NERC) GW4+ Doctoral Training Program. R.J.P., N.R.B., and B.T. were funded by the Simons Foundation BIOS-SCOPE program. M.M. and B.T. were part-funded by NERC (NE/R010935/1). This project used equipment funded by the Wellcome Trust Institutional Strategic Support Fund (WT097835MF), Wellcome Trust Multi-User Equipment Award (WT101650MA), and BBSRC LOLA award (BB/K003240/1).

We acknowledge the use of the University of Exeter High-Performance Computing (HPC) facility in carrying out this work.

## REFERENCES

- Weitz JS, Stock CA, Wilhelm SW, Bourouiba L, Coleman ML, Buchan A, Follows MJ, Fuhrman JA, Jover LF, Lennon JT, Middelboe M, Sonderregger DL, Suttle CA, Taylor BP, Frede Thingstad T, Wilson WH, Eric Wommack K. 2015. A multitrophic model to quantify the effects of marine viruses on microbial food webs and ecosystem processes. *ISME J* 9:1352–1364. <https://doi.org/10.1038/ismej.2014.220>.
- Zhao Y, Temperton B, Thrash JC, Schwalbach MS, Vergin KL, Landry ZC, Ellisman M, Deerinck T, Sullivan MB, Giovannoni SJ. 2013. Abundant SAR11 viruses in the ocean. *Nature* 494:357–360. <https://doi.org/10.1038/nature11921>.
- Zhao Y, Qin F, Zhang R, Giovannoni SJ, Zhang Z, Sun J, Du S, Rensing C. 2019. Pelagiphages in the Podoviridae family integrate into host genomes. *Environ Microbiol* 21:1989–2001. <https://doi.org/10.1111/1462-2920.14487>.
- Buchholz HH, Michelsen ML, Bolaños LM, Brown E, Allen MJ, Temperton B. 2021. Efficient dilution-to-extinction isolation of novel virus–host model systems for fastidious heterotrophic bacteria. *ISME J* <https://doi.org/10.1038/s41396-020-00872-z>.
- Zhang Z, Qin F, Chen F, Chu X, Luo H, Zhang R, Du S, Tian Z, Zhao Y. 2020. Culturing novel and abundant pelagiphages in the ocean. *Environ Microbiol* <https://doi.org/10.1111/1462-2920.15272>.
- Parsons RJ, Nelson CE, Carlson CA, Denman CC, Andersson AJ, Kledzik AL, Vergin KL, McNally SP, Treusch AH, Giovannoni SJ. 2015. Marine bacterioplankton community turnover within seasonally hypoxic waters of a subtropical sound: Devil's Hole, Bermuda. *Environ Microbiol* 17:3481–3499. <https://doi.org/10.1111/1462-2920.12445>.
- Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD, Pyshkin AV, Sirotkin AV, Vyahhi N, Tesler G, Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* 19:455–477. <https://doi.org/10.1089/cmb.2012.0021>.
- Bushnell B, Rood J, Singer E. 2017. BBMerge: accurate paired shotgun read merging via overlap. *PLoS One* 12:e0185056. <https://doi.org/10.1371/journal.pone.0185056>.
- Roux S, Enault F, Hurwitz BL, Sullivan MB. 2015. VirSorter: mining viral signal from microbial genomic data. *PeerJ* 3:e985. <https://doi.org/10.7717/peerj.985>.
- Salisbury A, Tsourkas PK. 2019. A method for improving the accuracy and efficiency of bacteriophage genome annotation. *Int J Mol Sci* 20:3391. <https://doi.org/10.3390/ijms20143391>.
- Borodovsky M, McIninch J. 1993. GenMark: parallel gene recognition for both DNA strands. *Comput Chem* 17:123–133. [https://doi.org/10.1016/0097-8485\(93\)85004-V](https://doi.org/10.1016/0097-8485(93)85004-V).
- Besemer J, Lomsadze A, Borodovsky M. 2001. GeneMarkS: a self-training method for prediction of gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions. *Nucleic Acids Res* 29:2607–2618. <https://doi.org/10.1093/nar/29.12.2607>.
- Lomsadze A, Gemayel K, Tang S, Borodovsky M. 2018. Modeling leaderless transcription and atypical genes results in more accurate gene prediction in prokaryotes. *Genome Res* 28:1079–1089. <https://doi.org/10.1101/gr.230615.117>.
- Zhu W, Lomsadze A, Borodovsky M. 2010. Ab initio gene identification in metagenomic sequences. *Nucleic Acids Res* 38:e132. <https://doi.org/10.1093/nar/gkq275>.
- Delcher AL, Bratke KA, Powers EC, Salzberg SL. 2007. Identifying bacterial genes and endosymbiont DNA with Glimmer. *Bioinformatics* 23:673–679. <https://doi.org/10.1093/bioinformatics/btm009>.
- Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
- Pruitt KD, Tatusova T, Maglott DR. 2007. NCBI reference sequences (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* 35:D61–D65. <https://doi.org/10.1093/nar/gkl842>.
- Potter SC, Luciani A, Eddy SR, Park Y, Lopez R, Finn RD. 2018. HMMER Web server: 2018 update. *Nucleic Acids Res* 46:W200–W204. <https://doi.org/10.1093/nar/gky448>.
- UniProt Consortium. 2019. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 47:D506–D515. <https://doi.org/10.1093/nar/gky1049>.
- Bairoch A, Apweiler R. 2000. The SWISS-PROT protein sequence database and its supplement TrEMBL in 2000. *Nucleic Acids Res* 28:45–48. <https://doi.org/10.1093/nar/28.1.45>.
- Finn RD, Bateman A, Clements J, Coghill P, Eberhardt RY, Eddy SR, Heger

- A, Hetherington K, Holm L, Mistry J, Sonnhammer ELL, Tate J, Punta M. 2014. Pfam: the protein families database. *Nucleic Acids Res* 42:D222–D230. <https://doi.org/10.1093/nar/gkt1223>.
22. Contreras-Moreira B, Vinuesa P. 2013. GET\_HOMOLOGUES, a versatile software package for scalable and robust microbial pangenome analysis. *Appl Environ Microbiol* 79:7696–7701. <https://doi.org/10.1128/AEM.02411-13>.
23. Bolduc B, Jang HB, Doucier G, You Z-Q, Roux S, Sullivan MB. 2017. vConTACT: an iVirus tool to classify double-stranded DNA viruses that infect Archaea and Bacteria. *PeerJ* 5:e3243. <https://doi.org/10.7717/peerj.3243>.
24. Mizuno CM, Rodriguez-Valera F, Kimes NE, Ghai R. 2013. Expanding the marine virosphere using metagenomics. *PLoS Genet* 9:e1003987. <https://doi.org/10.1371/journal.pgen.1003987>.
25. Zaragoza-Solas A, Rodriguez-Valera F, López-Pérez M. 2020. Metagenome mining reveals hidden genomic diversity of pelagimyophages in aquatic environments. *mSystems* 5:e00905-19. <https://doi.org/10.1128/mSystems.00905-19>.
26. Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797. <https://doi.org/10.1093/nar/gkh340>.
27. Talavera G, Castresana J. 2007. Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. *Syst Biol* 56:564–577. <https://doi.org/10.1080/10635150701472164>.
28. Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard J-F, Guindon S, Lefort V, Lescot M, Claverie J-M, Gascuel O. 2008. Phylogeny.fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res* 36:W465–W469. <https://doi.org/10.1093/nar/gkn180>.
29. Huelsenbeck JP, Ronquist F. 2001. MRBAYES: Bayesian inference of phylogenetic trees. *Bioinformatics* 17:754–755. <https://doi.org/10.1093/bioinformatics/17.8.754>.