



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Original Research

New cancer cases at the time of SARS-Cov2 pandemic and related public health policies: A persistent and concerning decrease long after the end of the national lockdown



Emmanuelle Kempf^{a,*}, Guillaume Lamé^b, Richard Layese^c,
Sonia Priou^d, Gilles Chatellier^e, Hedi Chaieb^d, Marc-Antoine Benderra^f,
Ali Bellamine^d, Romain Bey^d, Stéphane Bréant^d, Gilles Galula^g,
Namik Taright^h, Xavier Tannierⁱ, Thomas Guyet^j, Elisa Salamanca^d,
Etienne Audureau^c, Christel Daniel^d, Christophe Tournigand^k on behalf
of the Assistance Publique – Hôpitaux de Paris Cancer Group

^a Assistance Publique – Hôpitaux de Paris, Department of Medical Oncology, Henri Mondor and Albert Chenevier Teaching Hospital, Créteil, France

^b Université Paris-Saclay, CentraleSupélec, Laboratoire Génie Industriel, Gif-sur-Yvette, France

^c Université Paris Est Créteil, INSERM, IMRB, CEpiA Team, Assistance Publique – Hôpitaux de Paris, Clinical Research Unit, Henri Mondor and Albert Chenevier Teaching Hospital, Créteil, France

^d Assistance Publique – Hôpitaux de Paris, IT System Department, Paris, France

^e Université de Paris, Department of Medical Informatics, Assistance Publique Hôpitaux de Paris. Centre-Université de Paris (APHP-CUP), Paris, F-75015, France

^f Assistance Publique – Hôpitaux de Paris, Department of Medical Oncology, Tenon Teaching Hospital, Paris, France

^g Assistance Publique – Hôpitaux de Paris, Head Office, Department of Chronic Diseases and Cancer, Paris, France

^h Assistance Publique – Hôpitaux de Paris, Head Office, Department of Medical Information, Paris, France

ⁱ Sorbonne Université, Laboratoire D'informatique Médicale et D'ingénierie des Connaissances en e-Santé, Paris, France

^j Agrocampus-Ouest / IRISA - Inria, Rennes, France

^k Université Paris Est Créteil, INSERM, IMRB, Assistance Publique – Hôpitaux de Paris, Department of Medical Oncology, Henri Mondor and Albert Chenevier Teaching Hospital, Créteil, France

Received 18 January 2021; accepted 15 February 2021

Available online 25 February 2021

KEYWORDS

Incidence;

Abstract Introduction: The dissemination of SARS-Cov2 may have delayed the diagnosis of new cancers. This study aimed at assessing the number of new cancers during and after the

* Corresponding author: Department of medical oncology, Henri Mondor and Albert Chenevier Teaching Hospital, Assistance Publique – Hôpitaux de Paris, 51 avenue du Mehal de Lattre de Tassigny, Créteil, 94000, France. Fax: +1 4981 2576.

E-mail address: emmanuelle.kempf@aphp.fr (E. Kempf).

Early detection of cancer;
Health policy;
COVID-19

lockdown.

Methods: We prospectively collected the clinical data of the 11.4 million patients referred to the Assistance Publique Hôpitaux de Paris Teaching Hospital. We identified new cancer cases between 1st January 2018 and 31st September 2020 and compared indicators for 2018 and 2019 to 2020 with a focus on the French lockdown (17th March to 11th May 2020) across cancer types and patient age classes.

Results: Between January and September, 28,348, 27,272 and 23,734 new cancer cases were identified in 2018, 2019 and 2020, respectively. The monthly median number of new cases reached 3168 (interquartile range, IQR, 3027; 3282), 3054 (IQR 2945; 3127) and 2723 (IQR 2085; 2,863) in 2018, 2019 and 2020, respectively. From March 1st to May 31st, new cancer decreased by 30% in 2020 compared to the 2018–19 average; then by 9% from 1st June to 31st September. This evolution was consistent across all tumour types: –30% and –9% for colon, –27% and –6% for lung, –29% and –14% for breast, –33% and –12% for prostate cancers, respectively. For patients aged <70 years, the decrease of colorectal and breast new cancers in April between 2018 and 2019 average and 2020 reached 41% and 39%, respectively.

Conclusion: The SARS-Cov2 pandemic led to a substantial decrease in new cancer cases. Delays in cancer diagnoses may affect clinical outcomes in the coming years.

© 2021 Elsevier Ltd. All rights reserved.

1. Introduction

The SARS-Cov2 pandemic led to specific health policies aiming at reducing both its dissemination and hospital crowding. Among them, lockdowns occurred worldwide, and from 17th March to 11th May 2020 in France. Health services adapted to the health situation, in order to avoid contaminations. For example, specialised health care centres adopted triage strategies for patients with acute organ dysfunctions [1]. As a result, in the United States, cardiac catheterisation for STEMI dropped by 38% during the outbreak, revealing a loss of opportunity for severe conditions due to pandemic-related health policy [2]. In the Paris area, the activity of interventional cardiology fell by 60% in April 2020 compared to April 2019 (unpublished data of Greater Paris University Hospital). Cancer screening was also affected. Cancer societies, such as the American Society of Clinical Oncology, asked to postpone any cancer screening procedures during the outbreak [3]. On 2nd April, the French National Institute of Cancer (INCa) decided to interrupt the national cancer screening programs [4]. Outpatient visits were also considerably reduced. In France, general practitioner's, as well as specialist's consultations, dropped by 40% and 50%, respectively, during the first month of lockdown [5]. The number of cancer diagnostic procedures dropped significantly during the lockdown in France, with the use of consumables such as iodinated contrasts for CT-scan, Gadolinium contrasts for RMN and laxative liquids for colonoscopy preparation decreasing by 500,000 (up to –72%), 280,000 (up to –72%) and 250,000 (up to –86%), respectively, between 16th March and 13th

September 2020 [6]. Reduced patient encounters and diagnostic procedures, postponed screening procedures may explain a drop in new cancer cases during and after the outbreak. This would mean that patients with a new cancer are seen later and can, therefore, potentially suffer from a delay in the start of their treatment. Therefore, it is important to assess the importance of such drop, as treatment delays can dramatically impact the prognosis of patients with cancer [7,8].

This study has been aimed at assessing the impact of the pandemic and related French public health policies on the number of new cancer cases, across each type of cancer and main class ages, in Paris and suburban area, during and after the national lockdown.

2. Methods

The Greater Paris University Hospital (Assistance Publique Hôpitaux de Paris, AP-HP) is an organisation comprising 39 specialised health care centres in Paris and suburb area. Since 2015, AP-HP deployed a clinical data warehouse (CDW) based on the Informatics for Integrating Biology and the Bedside (i2b2) open-source software. This CDW prospectively collects from the hospital information system structured data (data directly reusable for further analysis, e.g. demographics, diagnosis codes, procedures, drug prescriptions, laboratory test results...) and unstructured data (i.e. data that need to be transformed before secondary use, including, but not limited to data stored in a free-text format such as radiology reports or discharge summaries). These data are refreshed daily [9], and the

CDW presently contains medical data of 11.4 million patients.

In the present study, more specifically, we used the AP-HP administrative data used for the national hospital reporting system (PMSI – Programme de Medicalisation des Systemes d'Information), a system based on diagnosis-related groups (DRGs). A PMSI record contains, for each patient's hospital stay, diagnoses (coded using International Classification of Diseases 10th revision (ICD-10)), procedures (coded using the Classification Commune des Actes Medicaux (CCAM), age, gender and place of residence. The hospital stay is described by the following variables: diagnostic-related group, length of stay, type of admission and discharge (home, referral, death), month and year of discharge and additional information about specific wards (e.g. intensive care unit) attended.

Patients with cancer were selected using a list of the ICD-10 codes related to a cancer diagnosis: C00 to D48, excluding benign tumours (D10 – D36) [10] (Supplementary Table 1). A patient was considered to have a newly diagnosed cancer if having a cancer-related ICD code and no cancer-related ICD code in the discharge summary recorded within the 2 previous years. We, therefore, counted cancer cases rather than patients, i.e. a patient with two primitive cancers during the study period was recorded twice, once for each cancer. We assessed the monthly number of new cancer cases for all types of cancer and for each cancer type, and for three patient age classes: <70 years, 70–84 years, and 85 years and older. We also assessed the number of *in situ* tumour cases: D00 to D09 [10] (Supplementary Table 1).

We compared the monthly number of cancers recorded in 2020 with those recorded in 2018 and in 2019 and with the average number recorded between 2018 and 2019. We performed aggregated comparisons between the 3-month periods corresponding to the French lockdown (1st March to 31st May 2020) and after the lockdown (1st June to 31st September 2020).

Table 1
Characteristics of the patient population, between January and September each year.

Year	2018	2019	2020
Number of new cancer cases	28,348	27,272	23,734
Number of patients with at least one cancer diagnosis	26,583	24,901	21,182
Number of patients with at least two cancer diagnoses	2847	1941	1159
Age, median (IQR) (years)	65 (54–75)	66 (54–75)	66 (54–75)
Female gender, n (%)	13,098 (46%)	12,593 (46%)	10,980 (46%)

IQR: interquartile range.

3. Results

3.1. Characteristics of the patient population

Between January and September, 28,348, 27,272 and 23,734 patients with a new cancer were referred to the AP-HP greater Paris University Hospital in 2018, 2019 and 2020, respectively. Table 1 summarises the characteristics of the patient population between 2018 and 2020.

3.2. Number of new cases for all tumour types over time

The evolution of the monthly number of overall new cancer cases over time is summarised in Fig. 1. The related figures can be found in Table 2. Between January and September, the monthly median number of new cancer cases reached 3168 (interquartile range, IQR, 3027; 3282), 3054 (IQR 2945; 3127) and 2723 (IQR 2085; 2863). During the French national lockdown period (i.e., from March to May 2020), compared to the two previous years, the median number of new cases dropped: by 30% in 2020 compared to the average of 2018–2019. In April 2020, the decrease reached 43% compared to April 2018 and April 2019. During the four months following the lockdown (i.e., from June to September 2020), the median number of new cases remained 9% lower than that recorded during the same period for the average of 2018–2019. In September 2020, the number of new cancer cases remained 3% and 4% lower than that recorded in September 2018 and September 2019, respectively.

3.3. Number of new cases across cancer types

Fig. 2a summarises the evolution of the monthly number of new breast, colorectal, prostate and lung cancer cases between 2018 and 2020. Table 3 displays the figures across all tumour subtypes from March to May (during the lockdown) and from June to September (after the lockdown) for 2018, 2019 and 2020. Full monthly data for each cancer subtype are provided in Supplementary Table 2.

The decrease in new cancer cases was consistent across all tumour types and has continued after the lockdown: –30% and –9% for colon, –27% and –6% for lung, –29% and –14% for breast, –33% and –12% for prostate cancers, respectively, from March to May and from June to September 2020 compared to the 2018–2019 average, respectively. Similar decrease patterns were found for poor prognosis tumours: –34% and –16% for pancreatic, –30% and –17% for bladder, –32% and –7% for the central nervous system and –29% and –9% for liver cancers, respectively.




Fig. 1. Evolution of the monthly number of new cancer cases of any type over time in 2018, 2019 and 2020 in the Assistance Publique Hôpitaux de Paris Teaching hospital Difference decrease (%).

3.4. Number of new cases across age classes

Fig. 2b summarises the consistent evolution of the monthly number of new cancer cases between 2018 and 2020 across patient age classes: < 70 years; 70–84 years; 85 years and over. Full monthly data per age class for each cancer subtype are provided in [Supplementary Table 2](#). In April 2020, the new cases fell by 42%, 45% and 43% in patients aged <70 years, 70–84 years, 85 years and older, respectively, compared to the April 2018–2019 average. Colorectal and breast cancer diagnoses are partially related to national screening programs that were interrupted during the lockdown. For

patients aged less than 70 years, the decrease of colorectal and breast new cancer cases in April between 2018 and 2019 average and 2020 reached 41% and 39%, respectively ([Supplementary Table 2](#)).

4. Discussion

Public health policies applied for the SARS-Cov2 pandemic led to a substantial drop in new cancer cases in the Paris area. New cases were still lower than expected in September, showing that the situation had not returned to normal even after the lockdown was lifted.

Table 2

Number of new cancer cases of any type in 2018, 2019 and 2020 in the AP-HP hospitals.

Including <i>in situ</i> tumours	Number of new cancer cases			Between-year difference (%)	
	2018	2019	2020	2020 vs. 2018	2020 vs. 2019
Median monthly number of new cases (IQR)	3168 (3027–3282)	3054 (2945–3127)	2723 (2085–2863)		
January	3961	3723	3772	–5	1
February	3027	3127	2952	–2	–6
March	3333	3054	2672	–20	–13
April	3168	3055	1776	–44	–42
May	3166	2984	2082	–34	–30
June	3282	2945	2723	–17	–8
July	3205	3248	2863	–11	–12
August	2307	2211	2085	–10	–6
September	2899	2925	2809	–3	–4
<i>In situ</i> tumour cases solely					
Median monthly number of new cases (IQR)	27 (26–29)	41 (28–44)	19 (17–23)		
January	29	45	23	–21	–49
February	26	45	26	0	–42
March	30	44	17	–43	–61
April	29	28	19	–34	–32
May	42	33	17	–60	–48
June	27	41	18	–33	–56
July	27	42	22	–19	–48
August	16	17	16	0	–6
September	25	19	24	–4	26

iQR: interquartile range.




Fig. 2. Difference (%) in the monthly number of new cancer cases between 2020 and the average of 2018–2019 in the Assistance Publique Hopitaux de Paris Teaching Hospital. a: according to main cancer type (breast, colorectal, prostate, lung cancer). b: according to the patient's age class (<70 years; 70–84 years; 85 years and over).

These results suggest a delay in new diagnoses and treatment of cancer cases.

Our results are in line with the published literature. In the Netherlands, the number of cancer diagnoses fell by 25% for cancers other than skin cancers and by 60% for skin cancers (excluding basal cell carcinomas) after 'social distancing' policies were implemented [11]. A cross-sectional American study showed that the weekly incidence of six main types of cancer fell by 46% during the pandemic crisis compared to baseline [12]. This decrease is consistent with other studies and reached 52% and 49% for breast and colorectal cancers, respectively [13]. An English survey showed a 76%-decrease in urgent cancer referrals for early diagnosis across eight tertiary care centres during the crisis [14]. An Italian study had reported a decrease in oral cancer screening during the first lockdown [15]. Similarly, an American multicentric observational study concluded that the incidence of new cancers had been divided into two between April 2019 and April 2020 across a 28-million patient population [16]. In this study, the drop of cancer incidence was higher for breast, prostate cancers, and melanoma. Screening for breast and colorectal cancers had decreased by 89% and 85%, respectively. An Italian survey concluded that the incidence of neuroendocrine neoplasia (NEN) decreased by 77% during the national lockdown [17].

This is the first study to assess the impact of the SARS-Cov2 outbreak on the number of new cancer

cases of all tumour types across the main age classes during and after a national lockdown. Our results are based on administrative databases and should, therefore, be interpreted with caution, as this kind of epidemiologic studies carries a risk of internal biases [18]. However, since filling the medical record is mandatory for financing reasons, the probability of bias on the number of cases recorded is low. We will link our results with the national administrative database in order to neutralise false-positive cases. Further studies are planned to exploit the content of patients' electronic health records and mitigate these risks of bias.

Our study results confirm the *distraction effect* induced by the SARS-Cov2 outbreak on the management of cancer patients [19]. The pandemic may induce subtle unintended consequences such as psychological stress and social isolation, decrease in care provision for life-threatening conditions, economic and logistic disruptions in drug administrations. In a French study performed on 125,000 healthcare professional practices, patients reported that fear of contamination (38%), reluctance to unduly disturb a physician in the middle of a crisis (28%) and closed medical practices (17%) were the main reason for not seeking medical attention [20]. In a context of reduced healthcare resources, efforts should be balanced between the acute threat of virus dissemination and the long-term clinical issues related to severe diseases, such as cancer. Delaying tumour diagnosis might prevent curable tumours from being treated

Table 3

Number of new cancer cases (any type) during and after the French lockdown period in 2018, 2019 and 2020 in the AP-HP hospitals.

	From 1st March to 31st May				From 1st June to 31st September			
	Number of new cancer cases			Difference between 2020 and the 2018–2019 average (%)	Number of new cancer cases			Decrease between 2020 and the 2018–2019 average (%)
	2018	2019	2020		2018	2019	2020	
Overall	9667	9093	6530	–30	11,693	11,329	10,480	–9
Colon	563	539	388	–30	738	648	629	–9
Rectum	210	209	167	–20	317	294	249	–18
Oesophagus	118	98	61	–44	126	127	99	–22
Gastric	185	168	111	–37	192	208	175	–13
Pancreas	375	340	235	–34	517	464	410	–16
Biliary duct	126	127	73	–42	173	173	119	–31
Bowel	74	77	38	–50	72	88	62	–23
Anus	45	31	35	–8	49	48	51	5
Liver	450	398	303	–29	513	522	469	–9
Digestive other	135	115	53	–58	124	183	116	–24
Endometrium	126	111	97	–18	143	140	143	1
Cervix	96	75	68	–20	103	91	96	–1
Ovary	119	130	117	–6	164	169	157	–6
Breast	792	638	507	–29	896	844	752	–14
Gynaecology other	35	29	25	–22	45	48	39	–16
Lung	828	829	602	–27	1047	949	939	–6
Mesothelioma	23	29	25	–4	27	19	26	13
Pneumology other	46	63	36	–34	57	65	59	–3
Head and neck	353	257	221	–28	409	385	321	–19
Eye	6	3	9	100	8	9	7	–18
CNS	500	430	317	–32	598	538	531	–7
PNS	4	15	3	–68	26	12	13	–32
Soft tissue	134	118	107	–15	164	141	131	–14
Osteosarcoma	132	111	94	–23	158	135	157	7
Myeloma	230	231	201	–13	286	315	318	6
Hodgkin lymphoma	110	95	94	–8	118	122	113	–6
Non-Hodgkin lymphoma	510	465	363	–26	556	617	581	–1
Leukaemia	339	338	258	–24	425	409	426	2
Other haematologic malignancies	378	370	217	–42	464	485	417	–12
Prostate	503	498	335	–33	664	576	544	–12
Testis	44	35	29	–27	62	48	45	–18
Kidney	301	291	195	–34	320	375	336	–3
Bladder	399	388	275	–30	493	487	407	–17
Other urothelial	16	16	11	–31	17	14	12	–23
Melanoma	245	261	213	–16	320	310	295	–6
Skin other	568	645	339	–44	715	693	646	–8
Thyroid	322	305	187	–40	287	297	332	14
CUP	74	54	51	–20	86	89	77	–12
Other endocrine	153	161	70	–55	214	192	181	–11

Abbreviations: CNS, central nervous system; CUP, carcinoma of unknown primitive; PNS, peripheral nervous system.

effectively and impair significantly patient clinical outcomes [21,22]. Through a statistical model, Lai *et al.* estimated that this insufficient care could lead to an excess of 6270 and 33,890 deaths at one year in incident cancer English and American patients, respectively [23]. Another English model study evaluated that the 5-year additional cancer-related deaths due to the pandemic would reach 3500 for four main cancer types [24]. International cancer societies urge cancer patients to come back to clinics, while SARS-Cov2 keeps on disseminating through subsequent waves [25]. Anticipating a loss of opportunity for cancer patients, many scientific communities published guidelines related to cancer care during the pandemic [26–28]. General practitioners

have advocated for a timely diagnosis of symptomatic cancer [29]. National cancer screening programs should go on while claiming patient safety regarding the risk of virus contamination in order to address the backlog within usual cancer screening programs [30]. Innovative, non-invasive, resilient to scarce resources screening procedures should be developed and deployed [31]. In an American survey of 3055 cancer patients and survivors, half of them reported some impact of the pandemic on their health care, and 27% of actively-treated patients expressed some delay to their treatment [32].

5. Conclusion

The SARS-Cov2 pandemic and related public health policies led to a significant decrease in the hospital visits of new patients with cancer in France. Stakeholders should pay attention to this issue during the next pandemic waves in order to avoid an excess in cancer-related deaths over the coming years. Further studies are warranted to analyse data from the second wave and the associated lockdown, as well as the assessment of the clinical prognostic impact of cancer diagnosis delays.

Author contributions

Emmanuelle Kempf: Conceptualisation, Methodology, Formal analysis, Data Curation, Writing - Original Draft, Project administration. **Guillaume Lamé:** Conceptualisation, Methodology, Formal analysis, Writing - Original Draft. **Richard Layese:** Conceptualisation, Methodology, Software, Formal analysis, Data Curation, Visualisation. **Sonia Priou:** Conceptualisation, Methodology, Software, Formal analysis, Data Curation, Visualisation. **Gilles Chatellier:** Conceptualisation, Methodology, Formal analysis, Writing - Original Draft, Supervision, Project administration. **Hedi Chaieb:** Conceptualisation, Methodology, Software, Formal analysis, Data Curation. **Marc-Antoine Benderra:** Conceptualisation, Methodology, Writing - Original Draft. **Ali Bellamine:** Conceptualisation, Methodology, Software, Writing - Original Draft. **Romain Bey:** Conceptualisation, Methodology, Resources. **Stéphane Bréant:** Conceptualisation, Methodology, Resources. **Gilles Galula:** Conceptualisation, Methodology, Writing - Original Draft, Supervision. **Namik Taright:** Conceptualisation, Methodology, Formal analysis, Writing - Original Draft, Supervision. **Xavier Tannier:** Conceptualisation, Methodology, Writing - Original Draft. **Thomas Guyet:** Conceptualisation, Writing - Original Draft. **Elisa Salamanca:** Conceptualisation, Methodology, Resources, Supervision. **Etienne Audureau:** Conceptualisation, Methodology, Supervision. **Christel Daniel:** Conceptualisation, Methodology, Formal analysis, Resources, Writing - Original Draft, Supervision, Project administration. **Christophe Tournigand:** Conceptualisation, Methodology, Formal analysis, Writing - Original Draft, Supervision.

Funding

This research was supported by a grant from the AP-HP Foundation and a grant from the ARC Foundation for cancer research.

Conflict of interest statement

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank Mrs. Patricia Serre and Dr. François Hemery for their help in the data access and analysis.

This research was supported by the teams in charge of the Clinical Data Warehouse of Greater Paris University Hospitals (AP-HP).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.ejca.2021.02.015>.

References

- [1] Elanko A, Khan J, Hamady ZZ, Malik H. Cancer surgery sustainability in the light of COVID-19 pandemic. *Eur J Surg Oncol* 2020 Jun 1;46(6):1174–5.
- [2] Garcia S, Albaghdadi MS, Meraj PM, Schmidt C, Garberich R, Jaffer FA, et al. Reduction in ST-segment elevation cardiac catheterization laboratory activations in the United States during COVID-19 pandemic. *J Am Coll Cardiol* 2020;vol. 75:2871–2. Elsevier USA.
- [3] American Society of Clinical Oncology. Cancer screening, diagnosis, staging, and surveillance. Updated June 22, 2020 [Internet]. Available from: <https://www.asco.org/asco-coronavirus-resources/care-individuals-cancer-during-covid-19/cancer-screening-diagnosis-staging-2>. [Accessed 13 July 2020].
- [4] Ministère des Solidarités et de la Santé. Continuité des activités des Centres régionaux de coordination des dépistages des cancers (CRCDC). *Inst Natl du cancer*; 2020.
- [5] Nicolas Revel directeur général de la Caisse nationale de l'assurance maladie (CNAM). Hearing before the Commission des Affaires Sociales, Senat(15 April, 2020). 2020.
- [6] Weill A, Drouin J, Desplas D, Cuenot F, Dray-Spira R, Zureik M. Usage des médicaments de ville en France durant l'épidémie de la Covid-19 – point de situation jusqu'au 13 septembre 2020. EPI-PHARE GIS ANSM - CNAM; 2020 [Internet]. Available from: <https://www.ansm.sante.fr/S-informer/Points-d-information-Points-d-information/Usage-des-medicaments-de-ville-en-France-durant-l-epidemie-de-Covid-19-point-de-situation-a-la-fin-du-confinement-Point-d-Information>.
- [7] Hanna TP, King WD, Thibodeau S, Jalink M, Paulin GA, Harvey-Jones E, et al. Mortality due to cancer treatment delay: systematic review and meta-analysis. *BMJ* 2020 Nov;371:m4087.
- [8] Hartman HE, Sun Y, Devasia TP, Chase EC, Jairath NK, Dess RT, et al. Integrated survival estimates for cancer treatment delay among adults with cancer during the COVID-19 pandemic. *JAMA Oncol* [Internet] 2020;48:109:1–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33119036>.
- [9] Daniel C. La recherche clinique à partir d'entrepôts de données. L'expérience de l'Assistance Publique – Hôpitaux de Paris (AP-HP) à l'épreuve de la pandémie de Covid-19. *Rev Med Interne* [Internet] 2020 May 1;41(5):303–7 [cited 2020 Oct 20]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7164890/>.

- [10] Ouvrage collectif édité par L'INCa. Algorithme de sélection des hospitalisations liées à la prise en charge du cancer dans les bases nationales d'activité hospitalière de court séjour «algorithme cancer». 2013. p. 1–192.
- [11] Dinmohamed AG, Visser O, Verhoeven RHA, Louwman MWJ, van Nederveen FH, Willems SM, et al. Fewer cancer diagnoses during the COVID-19 epidemic in the Netherlands [Internet]. *Lancet Oncol* 2020;21:750–1. Lancet Publishing Group; [cited 2020 Nov 13]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32359403/>.
- [12] Kaufman HW, Chen Z, Niles J, Fesko Y. Changes in the number of US patients with newly identified cancer before and during the coronavirus disease 2019 (COVID-19) pandemic. *JAMA Netw open* [Internet] 2020 Aug 3;3(8):e2017267 [cited 2020 Oct 20]. Available from: <https://jamanetwork.com/>.
- [13] Team TNUCI of S (NCIS) W. A segregated-team model to maintain cancer care during the COVID-19 outbreak at an academic center in Singapore [Internet]. *Ann Oncol* 2020;31:840–3. Elsevier Ltd; [cited 2020 Nov 13]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7174823/>.
- [14] Lai AG, Pasea L, Banerjee A, Denaxas S, Katsoulis M, Chang WH, et al. Estimating excess mortality in people with cancer and multimorbidity in the COVID-19 emergency. *medRxiv* [Internet] 2020;15(11). 2020.05.27.20083287. Available from: <https://www.medrxiv.org/content/10.1101/2020.05.27.20083287v1>.
- [15] Arduino PG, Conrotto D, Broccoletti R. The outbreak of Novel Coronavirus disease (COVID-19) caused a worrying delay in the diagnosis of oral cancer in north-west Italy: the Turin Metropolitan Area experience [Internet]. *Oral Dis* 2020. Blackwell Publishing Ltd; [cited 2020 Nov 13]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32306459/>.
- [16] London JW, Fazio-Eynullayeva E, Palchuk MB, Sankey P, McNair C. Effects of the COVID-19 pandemic on cancer-related patient encounters. *JCO Clin Cancer Informatics* 2020;4(4): 657–65.
- [17] Panzuto F, Maccauro M, Campana D, Faggiano A, Massironi S, Pusceddu S, et al. Impact of the SARS-CoV2 pandemic dissemination on the management of neuroendocrine neoplasia in Italy: a report from the Italian Association for Neuroendocrine Tumors (Itanet). *J Endocrinol Invest* [Internet] 2020 Aug 16 [cited 2020 Oct 20]; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/32803662>.
- [18] Prada-Ramallal G, Takkouche B, Figueiras A. Bias in pharmacoepidemiologic studies using secondary health care databases: a scoping review [Internet]. *BMC Med Res Methodol* 2019;19: 1–14. <https://doi.org/10.1186/s12874-019-0695-y>. BioMed Central Ltd.; [cited 2020 Dec 11]. Available from: .
- [19] Cortiula F, Pettke A, Bartoletti M, Puglisi F, Helleday T. Managing COVID-19 in the oncology clinic and avoiding the distraction effect. *Ann Oncol* 2020;31:553–5. Elsevier Ltd.
- [20] COMMUNIQUE DE PRESSE Covid-19 : Doctolib alerte sur la chute de fréquentation des cabinets et s'engage pour permettre aux patients de retourner consulter [Internet]. [cited 2020 Nov 13]. Available from: www.community.doctolib.com.
- [21] Sud A, Torr B, Jones ME, Broggio J, Scott S, Loveday C, et al. Effect of delays in the 2-week-wait cancer referral pathway during the COVID-19 pandemic on cancer survival in the UK: a modelling study. *Artic Lancet Oncol* [Internet] 2020;21:1035–79 [cited 2020 Dec 11]; Available from: www.gov.uk/guidance/national-.
- [22] Sud A, Jones ME, Broggio J, Loveday C, Torr B, Garrett A, et al. Collateral damage: the impact on outcomes from cancer surgery of the COVID-19 pandemic. *Ann Oncol* 2020 Aug 1;31(8): 1065–74.
- [23] Lai AG, Pasea L, Banerjee A, Hall G, Denaxas S, Chang WH, et al. Estimated impact of the COVID-19 pandemic on cancer services and excess 1-year mortality in people with cancer and multimorbidity: near real-time data on cancer care, cancer deaths and a population-based cohort study. *BMJ Open* [Internet] 2020; 10 [cited 2020 Dec 11]. Available from: <http://bmjopen.bmj.com/>.
- [24] Maringe C, Spicer J, Morris M, Purushotham A, Nolte E, Sullivan R, et al. The impact of the COVID-19 pandemic on cancer deaths due to delays in diagnosis in England, UK: a national, population-based, modelling study. *Lancet Oncol* 2020 Aug 1;21(8):1023–34.
- [25] Tempero M. COVID-19 and cancer: unintended consequences. *J Natl Compr Canc Netw: JNCCN* 2020;18:1147. NLM (Medline).
- [26] van de Haar J, Hoes LR, Coles CE, Seamon K, Fröhling S, Jäger D, et al. Caring for patients with cancer in the COVID-19 era [Internet]. *Nat Med* 2020;26:665–71. Nature Research; [cited 2020 Nov 13]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32405058/>.
- [27] Qadan M, Hong TS, Tanabe KK, Ryan DP, Lillemoe KD. A multidisciplinary team Approach for triage of elective cancer surgery at the Massachusetts general hospital during the novel coronavirus COVID-19 outbreak. *Ann Surg* [Internet] 2020 Jul 1; 272(1) [cited 2020 Nov 13];e20–1. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7188033/>.
- [28] Ueda M, Martins R, Hendrie PC, McDonnell T, Crews JR, Wong TL, et al. Managing cancer care during the COVID-19 pandemic: agility and collaboration toward a common goal. *JNCCN J Natl Compr Cancer Netw* [Internet] 2020 Apr 1;18(4): 366–9 [cited 2020 Nov 13]. Available from: <https://pubmed.ncbi.nlm.nih.gov/32197238/>.
- [29] Jones D, Neal RD, Duffy SRG, Scott SE, Whitaker KL, Brain K. Impact of the COVID-19 pandemic on the symptomatic diagnosis of cancer: the view from primary care [Internet]. *Lancet Oncol* 2020;21:748–50. Lancet Publishing Group; [cited 2020 Nov 13]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7251992/>.
- [30] Dubois RN. Cancer prevention research | commentary. 2020.
- [31] Dockter AG, Angelos GC. Molecular-based alternatives for colorectal cancer screening during the COVID-19 pandemic. *Surg Technol Int* 2020 May 1;36:1–5.
- [32] American Cancer Society Cancer Action Network (ACS). COVID-19 pandemic impact on cancer patients and survivors [Internet]. Available from: <https://www.fightcancer.org/sites/default/files/National Documents/SARS-COV2-Ongoing-Impact-Polling-Memo.pdf>.