



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

- 5 Mahase E. Covid-19: what have we learnt about the new variant in the UK? *BMJ* 2020; **371**: m4944.
- 6 WHO. SARS-CoV-2 variant—United Kingdom of Great Britain and Northern Ireland. *Disease Outbreak News*, Dec 21, 2020. <https://www.who.int/csr/don/21-december-2020-sars-cov2-variant-united-kingdom/en/> (accessed Feb 9, 2021).
- 7 Chand M, Hopkins S, Dabrera G, et al. Investigation of novel SARS-CoV-2 variant. Variant of Concern 202012/01. Technical briefing document on novel SARS-CoV-2 variant. Public Health England, January, 2021. <https://www.gov.uk/government/publications/investigation-of-novel-sars-cov-2-variant-variant-of-concern-20201201> (accessed Feb 9, 2021).
- 8 Tegally H, Wilkinson E, Giovanetti M, et al. Emergence and rapid spread of a new severe acute respiratory syndrome-related coronavirus 2 (SARS-CoV-2) lineage with multiple spike mutations in South Africa. *medRxiv* 2020; published online Dec 22. <https://www.medrxiv.org/content/10.1101/2020.12.21.20248640v1> (preprint).
- 9 County of Santa Clara Emergency Operations Center. COVID-19 variant first found in other countries and states now seen more frequently in California. Jan 17, 2021. <https://www.sccgov.org/sites/covid19/Pages/press-release-01-17-2021-COVID-19-variant-more-frequent-in-CA.aspx> (accessed Feb 9, 2021).
- 10 Zhang W, Davis BD, Chen SS, Martinez JMS, Plummer JT, Vail E. Emergence of a novel SARS-CoV-2 strain in Southern California, USA. *medRxiv* 2021; published online Jan 20. <https://doi.org/10.1101/2021.01.18.21249786> (preprint).
- 11 PAHO, WHO. Occurrence of variants of SARS-CoV-2 in the Americas. Washington, DC: Pan American Health Organization, 2021.
- 12 Franceschi VB, Caldana GD, de Menezes Mayer A, et al. Genomic epidemiology of SARS-CoV-2 in Esteio, Rio Grande do Sul, Brazil. *medRxiv* 2021; published online Jan 26. <https://doi.org/10.1101/2021.01.21.21249906> (preprint).
- 13 Mwai P. Coronavirus: Africa infections have risen sharply in worst-affected countries. *BBC*, Jan 25 2021. <https://www.bbc.co.uk/news/world-africa-53181555> (accessed Feb 9, 2021).
- 14 Callaway E, Mallapaty S. Novavax offers first evidence that COVID vaccines protect people against variants. *Nature* 2021; **590**: 17.
- 15 European Centre for Disease Prevention and Control. Risk related to spread of new SARS-CoV-2 variants of concern in the EU/EEA. Dec 29, 2020. <https://www.ecdc.europa.eu/en/publications-data/covid-19-risk-assessment-spread-new-sars-cov-2-variants-eueea> (accessed Feb 9, 2021).
- 16 US Centers for Disease Control and Prevention. Emerging SARS-CoV-2 variants. Jan 15, 2021. <https://www.cdc.gov/coronavirus/2019-ncov/more/science-and-research/scientific-brief-emerging-variants.html> (accessed Jan 20, 2021).
- 17 Callaway E. Could new COVID variants undermine vaccines? Labs scramble to find out. *Nature* 2021; **589**: 177–78.
- 18 Callaway E. Fast-spreading COVID variant can elude immune responses. *Nature* 2021; **589**: 50–51.
- 19 Studdert DM, Hall MA, Mello MM. Partitioning the curve—interstate travel restrictions during the Covid-19 pandemic. *N Engl J Med* 2020; **383**: e83.
- 20 McNeill K, Jacobs C. Half of the world’s population lack access to essential health services—are we doing enough? World Economic Forum, Sept 20, 2019. <https://www.weforum.org/agenda/2019/09/half-of-the-world-s-population-lack-access-to-essential-health-services-are-we-doing-enough/> (accessed Feb 9, 2021).
- 21 WHO. COVAX Announces additional deals to access promising COVID-19 vaccine candidates: plans global rollout starting Q1 2021. Dec 18, 2020. <https://www.who.int/news/item/18-12-2020-covax-announces-additional-deals-to-access-promising-covid-19-vaccine-candidates-plans-global-rollout-starting-q1-2021> (accessed Feb 9, 2021).
- 22 Ritchie H, Ortiz-Ospina E, Beltekian D, et al. Coronavirus (COVID-19) vaccinations. OurWorld in Data. 2021. <https://ourworldindata.org/covid-vaccinations> (accessed Feb 11, 2021).
- 23 Council for Trade-Related Aspects of Intellectual Property Rights. Waiver from certain provisions of the TRIPS Agreement for the prevention, containment and treatment of Covid-19. Communication from India and South Africa. World Trade Organization, Oct 2, 2020. <https://docs.wto.org/dol2fe/Pages/SS/directdoc.aspx?filename=q:/IP/C/W669.pdf&Open=True> (accessed Feb 9, 2021).
- 24 Eurasia Group. Ending the Covid-19 pandemic: the need for a global approach. New York: Eurasia Group, 2020.
- 25 International Monetary Fund. World economic outlook, October 2020: a long and difficult ascent. October, 2020. <https://www.imf.org/en/Publications/WEO/Issues/2020/09/30/world-economic-outlook-october-2020> (accessed Feb 9, 2021).
- 26 Battersby B, Lam WR, Ture E. Tracking the \$9 trillion global fiscal support to fight COVID-19. *IMF Blog*, May 20, 2020. <https://blogs.imf.org/2020/05/20/tracking-the-9-trillion-global-fiscal-support-to-fight-covid-19/> (accessed Feb 9, 2021).
- 27 Gurara D, Fabrizio S, Wiegand J. COVID-19: without help, low-income developing countries risk a lost decade. *IMF Blog*, Aug 27, 2020. <https://blogs.imf.org/2020/08/27/covid-19-without-help-low-income-developing-countries-risk-a-lost-decade/> (accessed Feb 9, 2021).
- 28 International Monetary Fund. Debt Service Relief from the Catastrophe Containment and Relief Trust (CCRT). 2020. <https://www.imf.org/en/Topics/imf-and-covid19/COVID-Lending-Tracker#ftn> (accessed Feb 10, 2021).



The changing face of medical professionalism and the impact of COVID-19

Published Online
February 23, 2021
[https://doi.org/10.1016/S0140-6736\(21\)00436-0](https://doi.org/10.1016/S0140-6736(21)00436-0)

Medical professionalism is changing with the increasing gap between what doctors have traditionally been trained to do and the realities of modern clinical practice.¹ In high-income countries, the changing demographics of patients with an ageing population, the large proportion of patients with long-term conditions and multiple comorbidities, and rising health-care costs have placed huge pressures on health systems globally.² Advances in technology and science have changed the way health professionals interact with patients, and democratisation of knowledge and increased accountability that come with changing patient and societal expectations have added to the

demands placed on physicians.³ In many countries, inadequate staffing levels aggravate this situation.⁴ Morale among doctors is generally declining—eg, a survey in the UK showed 54% of physicians reported morale as low or very low⁴—and burnout is rising (prevalence about 66–80%).^{5,6} There is a crisis in staff retention in some countries with up to 48% doctors considering leaving the profession.⁷

COVID-19 has exacerbated these tensions between medical professionalism and physician wellbeing. The pandemic has placed substantial demands on already overstretched, understaffed, and under-resourced health systems. COVID-19 has tested doctors and health-care

workers to the limits of their professional competence and taken a considerable toll on their health and wellbeing.⁸ Core principles of medical professionalism—ie, primacy of patient welfare, patient autonomy, and social justice—have been challenged during the pandemic.⁹ Many doctors worldwide have had to change the way they work, having to prioritise patient care and make difficult decisions based on insufficient resources, including withholding and withdrawing potentially life-saving treatments.^{10,11} Doctors have had to balance their personal risk with their duty to care for patients as well as balance professional versus caring responsibilities for household members in high-risk groups.¹² The need to self-isolate if they or their family members have symptoms of COVID-19 take them away from front-line responsibilities. All these factors have caused a sense of guilt, tension, and moral injury.¹³ Moral injury in this context occurs when doctors are forced to make decisions that contradict their professional and moral commitments—the challenge of knowing what care patients need but being unable to provide it due to constraints beyond their control. The moral injury concept helps reframe such challenges from a focus on the individual to a system-wide perspective.¹⁴

The COVID-19 pandemic has changed how health professionals work, how we behave and interact within our teams and organisations, our understandings of personal health and risk, inequalities between doctors with different risk factors, and wellbeing and mental health.⁸ Globally more than 300 000 health-care workers have been infected with COVID-19 in 79 countries, over 7000 have died, and many more have suffered as a result of stress, burnout, and moral injury.^{13,15}

There is an urgent need for a system-level approach to address the issues that COVID-19 has created to better protect and safeguard our medical workforce for the future. Such approaches need to focus on organisational culture and staff wellbeing as integral to professionalism and central to patient care.¹⁶ Physicians' wellbeing must be recognised as a care quality indicator for all health systems.¹⁷ Improving the working lives of clinicians can optimise the performance of health systems, improve patient experience, drive population health, and reduce costs.^{16,18}

Targeted interventions are likely to be less effective if only aimed at the individual.¹⁷ During COVID-19, there have been many wellbeing initiatives for clinicians that



Tom Pilsdon/Panos Pictures

have been well received.¹⁹ However, they need to be combined with organisational interventions including flexible working arrangements, enhanced teamwork, reductions in administrative burdens, and optimal use of technology.²⁰ Health professionals need to be well supported throughout the COVID-19 pandemic. The medical profession, health systems, and society all have a part to play in ensuring this support is provided. Individual doctors need to be empowered to recognise their own limitations as well as their wellbeing and support needs.

The professions must adapt to the changing needs of modern clinical practice and shape how we balance the many competing demands on us. Health professionals must build on the changes that are good for patient care and resist those that are not. COVID-19 has shown that we must move away from a model of medical professionalism that can lead to moral injury and towards one that provides proactive support for professionals in a systematic way and is focused on supporting moral repair.²¹ With the second and subsequent waves of COVID-19 now well established in many countries, we need to ensure that we as a profession support our doctors and promote ways of working that incorporate the doctor, the patient, teams, health-care organisations, workplace environment, and health systems.¹ Over time, this wider system approach will lead to greater cohesiveness within health care and support individual professionals in a safer, more sustainable way.

AFG is the President and MP the Global Vice President of the Royal College of Physicians, UK. We declare no other competing interests.

**Andrew F Goddard, Mumtaz Patel*
andrew.goddard@rcplondon.ac.uk

Royal College of Physicians, London NW1 4LE, UK

- 1 Royal College of Physicians. Advancing medical professionalism. London: Royal College of Physicians, 2018. <https://www.rcplondon.ac.uk/projects/outputs/advancing-medical-professionalism> (accessed Feb 11, 2021).
- 2 British Medical Association. The changing face of medicine and the role of doctors in the future: presidential project 2017. London: British Medical Association, 2017. <https://www.bma.org.uk/media/2067/bma-the-changing-face-of-medicine-june-2017.pdf> (accessed Feb 11, 2021).
- 3 Feiler T, Gaitskell K, Hordern J et al. Personalised medicine: the promise, the hype and the pitfalls. *New Bioethics* 2017; **1**: 1–12.
- 4 Royal College of Physicians. Focus on physicians: 2018–19 census (UK consultants and higher speciality trainees). London: Royal College of Physicians, 2019. <https://www.rcplondon.ac.uk/projects/outputs/focus-physicians-2018-19-census-uk-consultants-and-higher-specialty-trainees> (accessed Feb 11, 2021).
- 5 Rosenstein LS, Torre M, Ramos MA, et al. Prevalence of burnout amongst physicians—a systematic review. *JAMA* 2018; **320**: 1131–50.
- 6 The Lancet. Physician burnout: a global crisis. *Lancet* 2019; **394**: 93.
- 7 General Medical Council. The state of medical education and practice in the UK 2019. London: General Medical Council, 2019. https://www.gmc-uk.org/-/media/documents/somep-2019---full-report_pdf-81131156.pdf (accessed Feb 11, 2021).
- 8 Shaukat N, Ali D, Razzak J. Physical and mental health impacts of COVID-19 on healthcare workers: a scoping review. *Int J Emerg Med* 2020; **13**: 40.
- 9 American Board of Internal Medicine Foundation, American College of Physicians, American Society of Internal Medicine Foundation, European Federation of Internal Medicine. Medical professionalism in the new millennium: a physician charter. *Ann Intern Med* 2002; **136**: 243–46.
- 10 Liu Q, Luo D, Haase J, et al. The experiences of health-care providers during the COVID-19 crisis in China: a qualitative study. *Lancet Glob Health* 2020; **8**: e790–98.
- 11 Rosenbaum L. Facing Covid-19 in Italy—ethics, logistics, and therapeutics on the epidemic’s front line. *N Engl J Med* 2020; **382**: 1873–75.
- 12 Harkin D. Covid-19: balancing personal risk and professional duty. *BMJ* 2020; **369**: m1606.
- 13 Williamson V, Murphy D, Greenberg N. COVID-19 and experiences of moral injury in frontline key workers. *Occupat Med* 2020; **70**: 317–19.
- 14 Dean W, Talbot S, Dean A. Reframing clinician distress: moral injury not burnout. *Fed Pract* 2019; **36**: 447.
- 15 Amnesty International. Global: Amnesty analysis reveals over 7000 health workers have died from COVID-19. Sept 3, 2020. <https://www.amnesty.org/en/latest/news/2020/09/amnesty-analysis-7000-health-workers-have-died-from-covid19/> (accessed Feb 11, 2021).
- 16 Lemaire JB, Wallace JE. Burnout among doctors. *BMJ* 2017; **358**: j3360.
- 17 Wallace JE, Lemaire JB, Ghali WA. Physician wellness: a missing quality indicator. *Lancet* 2009; **374**: 1714–21.
- 18 Bodenheimer T, Sinsky C. From triple to quadruple aim: care of the patient requires care of the provider. *Ann Fam Med* 2014; **12**: 573–76.
- 19 Dzau VJ, Kirsh D, Nasca T. Preventing a parallel pandemic—a national strategy to protect clinicians’ well-being. *N Engl J Med* 2020; **383**: 513–15.
- 20 Garros D, Austin W, Dodek P. How can I survive this: coping during the COVID-19 pandemic. *Chest* 2020; published online Nov 18. <https://doi.org/10.1016/j.chest.2020.11.012>.
- 21 Shale S. Moral injury and the COVID-19 pandemic: reframing what it is, who it affects and how care leaders can manage it. *BMJ Leader* 2020; **4**: 224–27.



SARS-CoV-2 variants and ending the COVID-19 pandemic

Published Online
 February 11, 2021
[https://doi.org/10.1016/S0140-6736\(21\)00370-6](https://doi.org/10.1016/S0140-6736(21)00370-6)

The COVID-19 pandemic has devastated health-care systems, shut down schools and communities, and plunged the world into an economic recession. While 2020 was a challenging year, 2021 looks to be difficult with the emergence of multiple variants of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The race to vaccinate the world will need to respond to the pathogen’s constant evolution to evade immunity. What marks the path to the end of this pandemic?

2020 saw the successful development and testing of COVID-19 vaccines within timeframes not considered possible before. Two mRNA COVID-19 vaccines produced the first results, with impressive efficacy (94–95%).^{1,2} A disadvantage of these two vaccines is their low temperature storage requirements. Although other COVID-19 vaccines developed to date that use viral vectors, subunit particles, or inactivated viruses have comparatively lower levels of efficacy, with estimates of 70% for ChAdOx1 nCoV-19³ and 92% for Gam-COVID-Vac (Sputnik V)⁴ adenovirus

vector vaccines, they do not have the ultracold storage temperature requirements of the mRNA vaccines and are therefore easier to deliver. Data available so far for COVID-19 vaccines have shown protection only against clinical forms of infection, with the exception of recent data showing reduction in the duration of viral shedding and viral load among recipients of the ChAdOx1 nCoV-19 vaccine compared with placebo recipients, suggesting potential impact on viral transmission as well.⁵

In December, 2020, an unexpected rise in reported COVID-19 cases was attributed to the emergence of the new SARS-CoV-2 variants 501Y.V1 (B.1.1.7) in the UK and 501Y.V2 (B.1.351) in South Africa.^{6,7} In South Africa, high transmission in the context of high population immunity⁸ may have favoured the emergence and subsequent spread of the variant. Both variants had a mutation (N501Y) in the receptor-binding domain of the spike protein that is reported to contribute to increased transmission,⁹ with estimates ranging between 40% and 70% for increased transmission.⁶