

CORRECTION

Open Access



# Correction to: Metformin and sulodexide restore cardiac microvascular perfusion capacity in diet-induced obese rats

Judith van Haare<sup>1</sup> , M. Eline Kooi<sup>2</sup>, Jurgen W. G. E. van Teeffelen<sup>1</sup>, Hans Vink<sup>1</sup>, Jos Slenter<sup>2</sup>, Hanneke Cobelens<sup>1</sup>, Gustav J. Strijkers<sup>4</sup>, Dennis Koehn<sup>5</sup>, Mark J. Post<sup>1</sup> and Marc van Bilsen<sup>1,3\*</sup>

**Correction to:** *Cardiovasc Diabetol* (2017) 16:47

<https://doi.org/10.1186/s12933-017-0525-7>

Following publication of the original article [1], the authors regret errors in Figs. 2b–d. In these figures the

images of the representative Akt and phospho-Akt (pAkt) signals should be replaced with the appropriate images. The representative images shown here are correct. The changes do not affect the scientific conclusion and significance of the article.

The original article can be found online at <https://doi.org/10.1186/s12933-017-0525-7>.


\*Correspondence: marc.vanbilsen@maastrichtuniversity.nl

<sup>1</sup> Department of Physiology, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands

Full list of author information is available at the end of the article



© The Author(s) 2021. This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.



**Author details**

<sup>1</sup> Department of Physiology, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands. <sup>2</sup> Department of Radiology and Nuclear Medicine, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands. <sup>3</sup> Department of Cardiology, CARIM School for Cardiovascular Diseases, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands. <sup>4</sup> Biomedical Engineering and Physics, Academic Medical Center, P.O. Box 22700, 1100 DE Amsterdam, The Netherlands. <sup>5</sup> Pie Medical Imaging, P.O. Box 1132, 6201 BC Maastricht, The Netherlands.

Received: 6 February 2021 Accepted: 6 February 2021

Published online: 26 February 2021

**Reference**

1. van Haare J, Kooi ME, van Teeffelen JWGE, Vink H, Slenter J, Cobelens H, Strijkers GJ, Koehn D, Post MJ, van Bilsen M. Metformin and sulodexide restore cardiac microvascular perfusion capacity in diet-induced obese rats. *Cardiovasc Diabetol*. 2017;16:47. <https://doi.org/10.1186/s12933-017-0525-7>.

**Publisher's Note**

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.