

CARDIOVASCULAR FLASHLIGHT

doi:10.1093/eurheartj/ehab065

Online publish-ahead-of-print 10 February 2021

Late-onset endocarditis after coronavirus disease 2019 infection

Tadao Aikawa ^{1,2*}, Jiro Ogino³, Tamaki Kudo⁴, and Yusuke Kashiwagi¹

¹Department of Cardiology, Hokkaido Cardiovascular Hospital, 1-30, Minami-27, Nishi-13, Chuo-ku, Sapporo 064-8622, Japan; ²Department of Radiology, Jichi Medical University Saitama Medical Center, 1-847 Amanuma-cho, Omiya-ku, Saitama 330-8503, Japan; ³Department of Pathology, JR Sapporo Hospital, Kita-3, Higashi-1, Sapporo 060-0033, Japan; and ⁴Department of Radiology, Hokkaido Cardiovascular Hospital, 1-30, Minami-27, Nishi-13, Chuo-ku, Sapporo 064-8622, Japan

*Corresponding author. Tel: +81 11 563 3911, Fax: +81 11 551 3109, Email: tadao.aikawa@jichi.ac.jp

An 18-year-old man with acute dyspnoea presented to our hospital 3 months after hospitalization due to coronavirus disease 2019 (COVID-19) infection confirmed by reverse transcriptase-polymerase chain reaction (RT-PCR) for SARS-CoV-2. On admission, he had no fever and took no medication. Electrocardiography showed an early repolarization pattern in the precordial leads and ST-segment depression in lead III (Panel A). Transthoracic echocardiography showed subtle wall motion abnormality in the left ventricular anterior wall (Supplementary material online, Video S1). His cardiac troponin T levels were elevated at 642 ng/L, prompting urgent coronary computed tomography (CT) angiography, revealed no coronary artery stenosis in all coronary arteries without any manifestation of pneumonia or pulmonary embolism. Ten minutes after contrast injection, CT showed midwall delayed enhancement in the left ventricular anterior wall (Panel B), indicating COVID-19-related myocardial injury, which were consistent with subsequent β -methyl- ρ -[¹²³I]-iodophenyl-pentadecanoic acid (BMIPP) imaging (Panel C, Supplementary material online, Video S2). The RT-PCR for SARS-CoV-2 on admission returned positive, while serological tests for other virus infections were negative. Right ventricular endomyocardial biopsy revealed the thickened endocardium with fibrin and myeloperoxidase-positive neutrophils (Panels D–F) and the moderate interstitial fibrosis without inflammatory cell infiltration in the myocardium (Panels G–I), indicating endocarditis complicating myocardial fibrosis. No vegetation or thrombus formation was detected by repeat echocardiography. Serum C-reactive protein levels were negative throughout the course. He was discharged without any complications on Day 3 and had no symptoms on Day 21.

In this case, non-bacterial inflammatory processes by SARS-CoV-2 may cause endocarditis. Our findings underscore the importance of histological work-up to elucidate the mechanisms underlying COVID-19-related myocardial injury.

Supplementary material is available at *European Heart Journal* online.

Funding: This work was supported in part by JSPS KAKENHI (grant number 17K10349 to Dr. Aikawa).

Conflict of interest: The authors have submitted their declaration which can be found in the article [Supplementary Material](#) online.

© The Author(s) 2021. Published by Oxford University Press on behalf of the European Society of Cardiology.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

