

REVIEW

Open Access



The effect of additional protein on lean body mass preservation in post-bariatric surgery patients: a systematic review

Marleen M. Romeijn^{1,2*} , Daniëlle D. B. Holthuijsen^{1,3}, Aniek M. Kolen^{1,3}, Loes Janssen¹, Goof Schep⁴, François M. H. van Dielen¹ and Wouter K. G. Leclercq¹

Abstract

Background: As result of bariatric surgery, patients are susceptible to protein deficiency which can result in undesirable lean body mass (LBM) loss. Consumption of high-protein diets or supplements could counteract this, but evidence about the effect is scarce. This paper systematically reviewed the literature to determine the effect of additional protein intake (≥ 60 g/day) on LBM preservation in post-bariatric patients.

Methods: An electronic search of PubMed, EMBASE and the Cochrane Library was conducted. Studies were included if patients received a high-protein diet or protein supplements for at least one month, and LBM was assessed. The primary outcome was difference in mean LBM loss between the experimental (protein) and control group. Secondary outcomes were differences in body fat mass, total body water, body mass index and resting metabolic rate.

Results: Two of the five included studies ($n = 223$) showed that consumption of proteins resulted in significant LBM preservation. Only one study reported a significant difference in the reduction of body fat mass and resting metabolic rate in favour of a high-protein diet, but none of the studies showed a significant difference in total body water loss or body mass index change between the two groups.

Conclusions: This paper showed inconclusive evidence for LBM preservation due to protein supplementation or a high-protein diet in post-bariatric patients. This outcome might be subjected to certain limitations, including a lack of blinding and a low compliance rate reported in the included studies. More specific and personalized recommendations regarding protein intake may need to be established by high quality research. Studies investigating the quantity (g/day) and quality (whey, casein or soy) of proteins are also needed.

Keywords: Bariatric surgery, Protein intake, Body composition, Lean body mass, Systematic review

* Correspondence: Bariatrics.resurge@mmc.nl

¹Department of Surgery, Máxima Medical Center, De Run 4600, Veldhoven 5504 DB, The Netherlands

²Research School NUTRIM, Department of Surgery, Maastricht University Medical Center, Maastricht, the Netherlands

Full list of author information is available at the end of the article



© The Author(s). 2021 **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Introduction

Bariatric surgery (BS) is considered the most effective treatment for severe obesity [1–3]. Despite the successful weight loss, patients are prone to develop nutrient deficiencies due to energy restriction, malabsorption and food intolerances [4, 5]. Current guidelines recommend patients to consume 60–80 g proteins a day or 1.2 g/kg of the ideal body weight (IBW) [6–8], but adherence to these guidelines is known to be problematic in 45% of BS patients [4]. There is a substantial prevalence of excessive lean body mass (LBM) loss in BS patients. Within the first year after laparoscopic Roux-en-Y gastric bypass (RYGB), patients lose about 22% of their LBM [9, 10]. LBM plays an important role in resting energy expenditure, functional capacity, muscle strength and cardiovascular health [11–13]. In post-bariatric surgery patients, an excessive loss of LBM can be detrimental as it may slow down weight loss or even trigger weight regain [14–16]. Mechanisms behind this are a reduced resting metabolic rate (RMR) and a direct change in appetite [14, 16–18]. Moreover, an inadequate protein intake (≤ 60 g/day) potentially results in decreased feelings of satiation and decreased diet-induced thermogenesis, which may hinder weight loss [19]. For these reasons, an adequate protein intake and preservation of LBM in BS patients is of significant importance in long-term weight management.

There is a paucity of data that shows the correlation between protein intake and LBM loss after BS. In 2017, a systematic review concluded that two of the four studies with an adequate protein intake (≥ 60 g/day) was associated with significantly less LBM loss one year after RYGB [17–19]. A major criticism of this review is that the protein intake in eight studies was relatively inadequate (< 60 g/day) generating insufficient evidence. Currently, no systematic review has investigated the effect of an adequate protein intake (≥ 60 g/day) achieved by high-protein diets or protein supplements on LBM preservation, while multiple randomized controlled trials (RCTs) have been performed. Therefore, the aim of this systematic review was to evaluate the effect of protein supplementation or a high-protein diet (≥ 1 month) on LBM preservation in post-bariatric surgery patients, compared to patients following standard treatment.

Method

This review complies with the recommendations of the Cochrane Handbook for Systematic Reviews and Interventions [20] and was recorded according to the PRISMA systematic review guidelines [21]. The systematic review protocol was registered in the International Prospective Register of Systematic Reviews (PROSPERO) under registration number CRD42020176839.

Search strategy

The systematic search was performed in February 2020 and was conducted in three electronic databases: MEDLINE (PubMed Legacy), EMBASE (Ovid) and The Cochrane Library. The search included only human studies that were published in English or Dutch, and was not restricted by publication date. Keywords in the search strategy included [dietary protein], [protein supplementation] and [bariatric surgery], and their synonyms. The full search strategies for all databases can be found in supplementary Table 1. References of relevant reviews and included studies were hand searched for potential eligible studies that have been missed.

Eligibility criteria

Studies were considered eligible if they included: 1) patients in the age of 18–65 years with a body mass index (BMI) of ≥ 35 kg/m² who underwent RYGB or sleeve gastrectomy (SG), 2) daily protein supplementation or a high-protein diet for ≥ 1 month (≥ 60 g/day), started within 2 weeks after surgery, compared to standard treatment (control), 3) body composition as outcome measurement determined by either air displacement plethysmography, bioelectrical impedance analysis (BIA), dual-energy X-ray absorptiometry (DXA) or magnetic resonance imaging (MRI), 4) a follow-up of ≥ 2 months, and 5) an experimental or observational study design including a control group. Exclusion criteria were 1) inclusion of pregnant women, 2) protein supplementation or a high-protein diet combined with ≥ 2 times supervised strength training per week, without data about the effect of proteins only, 3) no data about primary outcome (LBM), or 4) reviews, letters, case series, case reports, conference abstracts and editorials.

Study selection

Initial records were screened for relevance on titles and abstract. Full-texts of relevant articles were obtained for checking final inclusion. Endnote X9 software was used to manage all references, including removal of duplicates.

Data extraction

The following data was extracted by one researcher (DH) using a standardized study form: authors' names, publication year, study design, follow-up period, sample size, gender, mean age, mean BMI, baseline LBM, surgery type, intervention protocol, protein intake prior to surgery, actual protein intake, compliance and study outcomes (LBM, body fat mass (BFM), total body water (TBW), BMI and RMR). A second author (MR) cross-checked the information.

Study outcomes

The primary outcome was difference in mean LBM loss between the experimental (protein) and control group. Secondary outcomes were differences in BFM, TBW, BMI and RMR. If no score (in kg or %) of the predefined outcome was provided, a score was calculated based on the available data (pre- and post-surgery). Effect sizes of the individual studies were calculated using Cohen's *d*. An effect size of ≤ 0.2 was considered trivial, 0.2–0.49 was considered small, 0.5–0.79 was considered moderate and ≥ 0.8 was considered high [22].

Quality assessment

Study quality was assessed by the Cochrane Collaboration's Risk of Bias Tool [23]. The Cochrane Collaboration's Risk of Bias tool subdivides studies into "low", "unclear" or "high" risk for various biases (selection bias, performance bias, detection bias, attrition bias, reporting bias and other bias). Two reviewers (DH, MR) judged the quality of each individual study based on a set criteria. Any disagreements were solved by a third reviewer (LJ).

Results

Study selection

In total, 881 articles were identified in three electronic databases and one article was identified in a reference list. After duplicate removal, 743 articles remained. After screening of titles and abstracts, 23 potentially relevant articles were selected for full-text reading. At the end, five studies met the inclusion criteria and were considered eligible for this systematic review [24–28] (Fig. 1).

Study characteristics

The sample sizes of the included studies varied from 20 [28] to 60 [25, 27] patients (Table 1). The follow-up periods ranged from 8 weeks [24] to 6 months [25, 26, 28] and 12 months [27]. Two studies included only SG patients [25, 27], two studies included only RYGB patients [24, 26] and one study included both types of BS [28]. Three of the five studies used protein supplements [25, 26, 28], one study used amino acid supplements [24] and one study used a protein-enriched diet to increase daily protein intake [27]. The dose of protein supplements or protein content in high-protein diets varied from 15 g/day [28] to 2.0 g/kg IBW/day [27]. Two of the included studies reported a high level of patients' compliance [27, 28], whereas two studies reported a low level [25, 26] and one study did not assess compliance [24]. Some authors reported reasoning for the low compliance, proposing that this can be improved by closer follow-up and face-to-face interviews [25]. Three of the five studies assessed body composition by BIA

[25, 27, 28], whereas the other two studies assessed body composition using DXA [24, 26].

Quality of individual studies

Four studies were free from a high risk of bias in all domains [25–28]. One study contained a high risk of bias based on funding [24]. In addition, in four of the five studies [24–27] the risk of bias was unclear concerning blinding. A summary of the risk of bias for the individual studies can be found in Table 2.

Lean body mass


All studies reported that LBM (kg) decreased significantly from pre-surgery to 8 weeks [24], 6 months [25, 26, 28] and 12 months [27] post-surgery. Two studies showed that protein supplementation [25] and a high-protein diet [27] resulted in significantly more preservation of LBM compared to control, respectively 8% vs. -12% and -12% vs. -19% (Table 3). The other three studies demonstrated no differences towards LBM preservation following protein supplementation [24, 26, 28]. The studies that showed a significant difference in the decrease of LBM had an effect size of 0.31 [25] and 0.61 [27], which is considered small and moderate, respectively.

Total body weight, body fat mass, body mass index and resting metabolic rate

All studies reported that BMI [24–26, 28], TBW and BFM [24–28] decreased significantly following BS. None of the studies observed a significant difference in the decrease of TBW and BMI between the control and protein group. Only one study showed a significant difference in the reduction of BFM between the two groups, indicating a higher decrease in BFM following a high-protein diet [27]. Additionally, only one of the two studies that examined the effect of an additional protein intake on RMR demonstrated a significant higher RMR following a high-protein diet compared to a normal protein diet [27] (Table 3).

Discussion

As result of bariatric surgery, patients are susceptible to protein deficiency which can result in an undesirable LBM loss. Evidence about the effect of protein supplementation or a high-protein diet (≥ 60 g/day) on LBM preservation is scarce. Therefore, this systematic review was conducted to evaluate these effects. Two of the five studies supported the hypothesis that protein supplementation or a high-protein diet resulted in significant LBM preservation [25, 27], whereas the other three studies did not support the hypothesis [24, 26, 28]. This discrepancy can be attributed to differences in protein



intake, type of surgery and measurement tools which are discussed below.

The first explanation for why three studies failed to detect a significant LBM preservation is that the actual daily protein intake of these patients may have been too low. The studies that failed to demonstrate significant LBM preservation following protein supplementation reported an actual daily protein intake of 67 and 82 g/day [26, 28], though this amount is considered as adequate according to literature [6–8]. The actual daily protein intake in one of the studies that observed a significant LBM preservation was much higher, namely 143 g/day [27]. The other study

that showed a significant LBM preservation reported a daily protein intake of just 51 g/day, while the protein intake of the control group was unknown [25]. The amount of 51 g/day should be criticized as protein intake was measured the first month after surgery and it is plausible that patients increased their protein intake hereafter, resulting in a higher protein intake at the time of measuring LBM (3 and 6 months). Based on the abovementioned findings (143 g/day resulting in significant LBM preservation [27]; 67 g/day to 82 g/day not resulting in LBS preservation [26, 28]), it could be questioned whether 60–80 g/day is sufficient to maintain LBM.

Table 1 Study characteristics of the included studies

Author, year	Study design, follow-up	Sample size Gender	Age (yr) BMI (kg/m ²) LBM (kg)	Surgery type	Intervention	Pre-surgery protein intake (g/day)	Actual protein intake (g/day)	Compliance	Protein intake analyzed	Outcomes analyzed
Clements et al., 2011 [24]	Unblinded randomized control pilot study, 8 weeks	30 M (n = 1) F (n = 29)	46.0 ± 7.5 43.6 ± 4.2 54.0 ± 8.1	RYGB (n = 30)	CON: usual care, no use of protein supplements PRO: oral supplement containing 24 g leucine metabolite, glutamine and arginine twice daily for 8 weeks	NA	NA	NA	Log sheets	TBW, BMI Body composition using DXA (BFM, LBM) RMR
		60 M (n = 9) F (n = 51)	43.5 ± 8.4 45.9 ± 6.5 61.7 ± 13.3	SG (n = 60)	CON: standard diet for 1 month, no use of protein supplements PRO: standard diet + 1.2 g/kg IBW/day protein (±80 g) whey powder for 1 month	NA	CON: NA PRO: 51	38% ^a	NA	TBW, BMI Body composition using BIA (BFM, LBM)
Opfert et al., 2018 [26]	Randomized controlled trial, 6 months	53 F (n = 53)	43.9 ± 10.7 43.6 ± 6.2 55.6 ± 8.4	RYGB (n = 53)	CON: usual care with general dietary and physical activity counselling PRO: oral supplement containing whey-protein-enriched powder (24 g) twice daily for 6 months	CON: 80 PRO: 82	CON: 60 PRO: 82	55%	Dietary history method. Return of empty cans of protein powder	TBW, BMI Body composition using DXA (BFM, LBM)
		60 M (n = 60)	41.0 ± 6.2 40.7 ± 5.3 75.0 ± 11.9	SG (n = 60)	NPD (normal protein diet): 1.0 g/kg IBW/day protein (23.3%), 15% fat and 61.7% carbohydrates for 12 months PED (protein-enriched diet): 2.0 g/kg ideal IBW/day protein (±160 g)(47.7%), 15% fat and 37.7% carbohydrates for 12 months	NA	CON: 67 PRO: 143	Very high	Questionnaires, 3 day dietary record, 72 h-recall	TBW, BMI Body composition using BIA (BFM, LBM) RMR
Schollenberger et al., 2016 [28]	Randomized controlled double-blind pilot study, 6 months	20 M (n = 3) F (n = 17)	47.0 ± 11.9 49.0 ± 5.1 68.9 ± 13.4	RYGB (n = 5) SG (n = 15)	CON: pure maltodextrin powder, 15 g/day first month, 30–35 g/day second-sixth month PRO: milk protein powder, 15 g/day first month, 30–35 g/day second-sixth month	CON: 93 PRO: 97	CON: 53 PRO: 67	86%	4 day dietary record, interviews	TBW, BMI Body composition using BIA (BFM, LBM)

Values are expressed as mean ± SD

List of abbreviations: CON control group, PRO protein group, RYGB Roux-en-Y gastric bypass, SG sleeve gastrectomy, IBW ideal body weight, BMI body mass index, BFM body fat mass, LBM lean body mass, RMR resting metabolic rate, DXA dual-energy X-ray absorptiometry, BIA bioelectrical impedance analysis

^a indicates the percentage of patients adhering to the guideline regarding protein intake (≥60 g/day)

Table 2 Assessment of risk of bias [24–28]

Author, year	Selection bias		Performance bias	Detection bias	Attrition bias	Reporting bias	Other bias
	Random sequence generation	Allocation concealment	Blinding of participants and personnel	Blinding of outcome assessment	Incomplete outcome data	Selective reporting	
Clements et al., 2011 [24]	+	?	?	+	+	+	–*
Günes et al., 2019 [25]	+	+	?	+	?	+	?
Oppert et al., 2018 [26]	+	?	?	+	+	+	?
Schiavo et al., 2017 [27]	+	?	?	+	?	+	?
Schollenberger et al., 2016 [28]	+	+	+	+	+	+	?

*indicating bias regarding funding

- = high risk of bias
- = low risk on bias
- = unclear risk bias

*indicating bias regarding funding

- = high risk of bias
- = low risk on bias
- = unclear risk bias

A lack of compliance might explain the relatively low actual protein intake within the first months after surgery. In the study of Oppert et al. this may have attributed to insignificant outcomes [26]. Unfortunately, none of the included studies reported clear causes for poor compliance. We speculate that this could be attributed to the occurrence of side effects, food intolerances and a

lack of understanding regarding the need of adequate proteins [29, 30].

Protein intake and subsequent absorption may have been influenced by the type of surgery as both restrictive and malabsorptive procedures were included in this study. The two studies that found a significant effect included patients who underwent a restrictive procedure

Table 3 Results of the individual studies

Author, year	LBM change (kg)(change%)	Effect size (d)	BFM change (kg)(change%)	TBW change (kg)(change%)	BMI change (kg/m ²)(change%)	RMR change (kcal/day)
Clements et al., 2011 [24] ¹	CON: – 7.9 ± 4.5 (14.6%)* PRO: – 7.7 ± 3.5 (14.7%)*	0.23 (– 0.48 to 0.95)	CON: – 8.0 ± 3.5 (14.3%)* PRO: – 9.2 ± 3.2 (15.7%)*	CON: – 15.7 ± 2.5 (13.8%)* PRO: – 15.8 ± 2.6 (13.9%)*	CON: – 6.1 ± 1.1 (14%)* PRO: – 5.9 ± 0.9 (13.8%)*	CON: – 294.1 ± 207.2 (15.9%)* PRO: – 286.6 ± 271.1 (15.9%)*
Günes et al., 2019 [25] ²	CON: – 7.2 (12%)* PRO: 3.8 (8%)*^	0.31 (– 0.24 to 0.78)	CON: – 25.1 (41%)* PRO: – 36.7 (49.5%)*	CON: – 33.0 (26.9%)* PRO: – 33.1 (27.2%)*	CON: – 12.6 (27.4%)* PRO: – 12.2 (26.5%)*	NA
Oppert et al., 2018 [26] ³	CON: – 8.8 (– 10.1 to – 7.5)* (16%) PRO: – 8.2 (– 9.3 to – 7.1)* (15%)	NA	CON: – 19.7 (– 21.5 to – 17.9)* PRO: – 19.8 (– 21.3 to – 18.2)*	CON: – 28 (– 30.6 to – 25.4)* PRO: – 27.2 (– 29.4 to – 25.1)*	CON: – 10.5 (– 11.4 to – 9.6)* PRO: – 10.2 (– 11.0 to – 9.4)*	NA
Schiavo et al., 2017 [27] ¹	CON: – 14.5 (19%)* PRO: – 8.8 (12%)*^	0.61 (0.07 to 1.15)	CON: – 23.7 (50%)* PRO: – 43.2 (84%)*^	CON: – 38.8 (31%)* PRO: – 46.7 (36%)*	NA	CON: – 645.16 (29%)*^ PRO: – 380.18 (17%)*^
Schollenberger et al., 2016 [28] ¹	CON: – 7.8 (11.3%)* PRO: – 7.6 (11.7%)*	0.3 (– 0.59 to 1.18)	CON: – 21.0 (30.8%)* PRO: – 29.1 (37.2%)*	CON: – 28.7 (20.9%)* PRO: – 36.4 (25%)*	CON: – 10.3 (21%)* PRO: – 13.0 (25%)*	NA

¹ expressed as mean ± SD (change%)

² expressed as mean (change%)

³ expressed as mean (CI) (change%)

*denotes significant difference from baseline. ^denotes significant difference from control

Abbreviations: CON control group, PRO protein group, NA not applicable or not assessed, LBM lean body mass, BFM body fat mass, TBW total body weight, BMI body mass index, RMR resting metabolic rate

(SG) [25, 27], in contrast to the studies that included RYGB patients where no effect was found [24, 26, 28]. It is interesting to note that Schollenberger et al. reported, in a separate analysis of only SG patients, that protein supplementation led to significant LBM preservation [28]. An explanation for these findings may be that protein digestion and absorption is higher after restrictive surgery. This proposes that the additional protein intake is less effective in RYGB patients, but results in more pronounced LBM preservation in SG patients.

A third explanation may be the usage of different tools for measuring body composition (BIA versus DXA), both presenting important limitations. The two studies that detected a significant LBM preservation used BIA [25, 27]. However, BIA is known for overestimating LBM in bariatric patients and the validity of BIA is influenced by fatness [31–33]. As a result, the two significant outcomes are potentially more pronounced. Additionally, DXA is limited by the fact that the fat free mass compartment is measured rather than directly muscle mass [32]. Future research is recommended to measure body composition with a four-compartment model to overcome this limitation.

It is conceivable that physical activity might have influenced study outcomes as it is known from the field of sports physiology that physical activity plays an important role in LBM preservation [34–36]. Four of the five studies did not report anything about physical activity of the patients, implying uncertainty on whether and how physical activity influenced LBM preservation. The study that did report about physical activity (i.e., supervised strength training for 18 weeks plus additional protein intake), failed to show a significant preservation in LBM [26]. Contrarily, Muschitz et al. approved the synergistic effect of physical activity on protein supplementation as they observed significant LBM preservation [37]. The discrepancy in this outcome may be explained by the difference in study length, 18 weeks vs. 24 months respectively [26, 37]. Further studies investigating the synergistic effect of physical activity and protein supplementation in bariatric patients are limited, which implies that it is difficult to draw conclusions based on these two studies.

There are some methodological limitations in this systematic review which should be mentioned. Four of the five studies lacked (double) blinding, which could have influenced the study outcomes. Furthermore, only two of the five studies reported high compliance to protein intake and because of this, outcomes are potentially less pronounced than expected. Moreover, the number of the included studies in this systematic review is small, potentially resulting in a relative low power of this systematic review. A further comparison of the included studies was complicated due to heterogeneity of the

study protocols (e.g. supplementation type, dose and timing) and the measurement tools.

New studies investigating the most effective dose of supplements to preserve LBM in post-bariatric surgery patients are warranted as perhaps the dose of 60–80 g/day is insufficient to maintain muscle mass. In addition, it is advised to conduct studies examining the most effective composition of protein supplements (e.g. whey vs. casein vs. soy) in order to enable interstudy comparison. Special attention needs to be paid to the effect of leucine on LBM preservation, given its key role in muscle protein synthesis. On top of that, studies focusing on the synergistic effect of physical activity and protein intake on LBM preservation are warranted.

Conclusion

Although the preservation of LBM in post-bariatric surgery patients is of extreme importance, our systematic review resulted in the inclusion of only five studies. These studies showed inconclusive evidence for LBM preservation due to protein supplementation or a high-protein diet. Notwithstanding, this work offers awareness to current healthcare providers who should prompt an adequate protein intake in post-bariatric surgery patients. More specific and personalized recommendations regarding protein intake may need to be established by high quality research. New studies investigating the quantity (g/day) and quality (whey, casein or soy) of protein supplements or high-protein diets, possibly in combination with resistance training, in larger study populations are needed.

Abbreviations

BIA: Bioelectrical impedance analysis; BFM: Body fat mass; BMI: Body mass index; BS: Bariatric surgery; DXA: Dual-energy X-ray absorptiometry; IBW: Ideal body weight; LBM: Lean body mass; MRI: Magnetic resonance imaging; RCT: Randomized controlled trial; RMR: Resting metabolic rate; SG: Sleeve gastrectomy; RYGB: Roux-en-Y gastric bypass; TBW: Total body water

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12937-021-00688-3>.

Additional file 1. Appendix Table 1. Search strategy MEDLINE, EMBASE and Cochrane.

Acknowledgements

The authors would like to recognize the contribution of Bart de Vries for assistance in conducting the systematic search.

Authors' contributions

Conceptualization, DB, MM and WL; methodology, DB, MM, AK and LJ; formal analysis of study findings, DB and MM; interpretation, DB, MM, AK, LJ; writing—original draft preparation, DB; writing—review and editing, MM, AK, LJ, GS, FvD and WL; supervision, GS, FvD and WL. The author(s) read and approved the final manuscript.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interest.

Author details

¹Department of Surgery, Máxima Medical Center, De Run 4600, Veldhoven 5504 DB, The Netherlands. ²Research School NUTRIM, Department of Surgery, Maastricht University Medical Center, Maastricht, the Netherlands. ³Maastricht University, Faculty of Health, Medicine and Life Sciences, Maastricht, The Netherlands. ⁴Department of Sport Medicine, Máxima Medical Center, Veldhoven, The Netherlands.

Received: 9 September 2020 Accepted: 5 March 2021

Published online: 14 March 2021

References

- Tsigos C, Hainer V, Basdevant A, Finer N, Fried M, Mathus-Vliegen E, et al. Management of obesity in adults: European clinical practice guidelines. *Obesity facts*. 2008;1(2):106–16.
- Buchwald H, Avidor Y, Braunwald E, Jensen MD, Pories W, Fahrenbach K, et al. Bariatric surgery: a systematic review and meta-analysis. *Jama*. 2004;292(14):1724–37.
- Puzziferri N, Roshek TB, Mayo HG, Gallagher R, Belle SH, Livingston EH. Long-term follow-up after bariatric surgery: a systematic review. *Jama*. 2014;312(9):934–42.
- Moize V, Geliebter A, Gluck ME, Yahav E, Lorence M, Colarusso T, et al. Obese patients have inadequate protein intake related to protein intolerance up to 1 year following Roux-en-Y gastric bypass. *Obesity Surg*. 2003;13(1):23–8. <https://doi.org/10.1381/096089203321136548>.
- Verger EO, Aron-Wisniewsky J, Dao MC, Kayser BD, Oppert JM, Bouillot JL, et al. Micronutrient and Protein Deficiencies After Gastric Bypass and Sleeve Gastrectomy: a 1-year Follow-up. *Obesity Surg*. 2016;26(4):785–96. <https://doi.org/10.1007/s11695-015-1803-7>.
- Mechanick JI, Apovian C, Brethauer S, Garvey WT, Joffe AM, Kim J, et al. Clinical practice guidelines for the perioperative nutrition, metabolic, and nonsurgical support of patients undergoing bariatric procedures—2019 update: cosponsored by American Association of Clinical Endocrinologists/American College of Endocrinology, The Obesity Society, American Society for Metabolic & Bariatric Surgery, Obesity Medicine Association, and American Society of Anesthesiologists. *Surg Obesity Relat Dis*. 2019;25(12):1346–59.
- Heber D, Greenway FL, Kaplan LM, Livingston E, Salvador J, Still C. Endocrine and nutritional management of the post-bariatric surgery patient: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2010;95(11):4823–43.
- Sherf Dagan S, Goldenshluger A, Globus I, Schweiger C, Kessler Y, Kowen Sandbank G, et al. Nutritional recommendations for adult bariatric surgery patients: clinical practice. *Adv Nutr*. 2017;8(2):382–94.
- de Paris FGC, Padoin AV, Mottin CC, de Paris MF. Assessment of changes in body composition during the first postoperative year after bariatric surgery. *Obes Surg*. 2019;29(9):3054–61.
- Nuijten MA, Montpellier VM, Eijssvogels TM, Janssen IM, Hazebroek EJ, Hopman MT. Rate and Determinants of Excessive Fat-Free Mass Loss After Bariatric Surgery. *Obesity Surg*. 2020;30(8):3119–26.
- Lawman HG, Troiano RP, Perna FM, Wang C-Y, Fryar CD, Ogden CL. Associations of relative handgrip strength and cardiovascular disease biomarkers in US adults, 2011–2012. *Am J Prev Med*. 2016;50(6):677–83.
- Cooper R, Kuh D, Hardy R. Objectively measured physical capability levels and mortality: systematic review and meta-analysis. *Bmj*. 2010;341:c4467.
- Wolfe RR. The underappreciated role of muscle in health and disease. *Am J Clin Nutr*. 2006;84(3):475–82.
- de Castro CM, de Lima Montebelo MI, Rasesa I, de Oliveira AV, Gonelli PRG, Cardoso GA. Effects of roux-en-Y gastric bypass on resting energy expenditure in women. *Obes Surg*. 2008;18(11):1376–80.
- Flancbaum L, Choban PS, Bradley LR, Burge JC. Changes in measured resting energy expenditure after roux-en-Y gastric bypass for clinically severe obesity. *Surgery*. 1997;122(5):943–9.
- Ravussin E, Lillioja S, Knowler WC, Christin L, Freymond D, Abbott WG, et al. Reduced rate of energy expenditure as a risk factor for body-weight gain. *N Engl J Med*. 1988;318(8):467–72.
- Ito MK, Gonçalves VSS, Faria SLCM, Moizé V, Porporatti AL, Guerra ENS, et al. Effect of protein intake on the protein status and lean mass of post-bariatric surgery patients: a systematic review. *Obes Surg*. 2017;27(2):502–12.
- Moizé V, Andreu A, Rodríguez L, Flores L, Ibarzabal A, Lacy A, et al. Protein intake and lean tissue mass retention following bariatric surgery. *Clin Nutr*. 2013;32(4):550–5.
- Raftopoulos I, Bernstein B, O'Hara K, Ruby JA, Chhatrala R, Carty J. Protein intake compliance of morbidly obese patients undergoing bariatric surgery and its effect on weight loss and biochemical parameters. *Surg Obes Relat Dis*. 2011;7(6):733–42.
- Higgins J, Wells G. *Cochrane handbook for systematic reviews of interventions*; 2011.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med*. 2009;151(4):264–9.
- Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33(1):159–74.
- Higgins JP, Altman DG, Gøtzsche PC, Jüni P, Moher D, Oxman AD, et al. The Cochrane Collaboration's tool for assessing risk of bias in randomised trials. *Bmj*. 2011;343:d5928.
- Clements RH, Saraf N, Kakade M, Yellumhanthi K, White M, Hackett JA. Nutritional effect of oral supplement enriched in beta-hydroxy-beta-methylbutyrate, glutamine and arginine on resting metabolic rate after laparoscopic gastric bypass. *Surg Endosc*. 2011;25(5):1376–82.
- Gunes Y, Karip B, Ergin A, Esen Bulut N, Fersahoglu MM, Memisoglu K. The Impact of Protein Support on Weight Loss, Sarcopenia, and Quality of Life after Sleeve Gastrectomy. *Bariatric Surg Pract Patient Care*. 2019;14(3):131–6. <https://doi.org/10.1089/bari.2018.0057>.
- Oppert JM, Bellicha A, Roda C, Bouillot JL, Torcivia A, Clement K, et al. Resistance Training and Protein Supplementation Increase Strength After Bariatric Surgery: a Randomized Controlled Trial. *Obesity*. 2018;26(11):1709–20. <https://doi.org/10.1002/oby.22317> PubMed PMID: CN-01650609.
- Schiavo L, Scalera G, Pilone V, De Sena G, Quagliarillo V, Iannelli A, et al. A Comparative Study Examining the Impact of a Protein-Enriched Vs Normal Protein Postoperative Diet on Body Composition and Resting Metabolic Rate in Obese Patients after Sleeve Gastrectomy. *Obesity Surg*. 2017;27(4):881–8. <https://doi.org/10.1007/s11695-016-2382-y>.
- Schollenberger AE, Karschin J, Meile T, Kuper MA, Konigsrainer A, Bischoff SC. Impact of protein supplementation after bariatric surgery: a randomized controlled double-blind pilot study. *Nutrition*. 2016;32(2):186–92. <https://doi.org/10.1016/j.nut.2015.08.005> PubMed PMID: CN-01133406.
- Schweiger C, Weiss R, Keidar A. Effect of different bariatric operations on food tolerance and quality of eating. *Obes Surg*. 2010;20(10):1393–9.
- Di Vetta V, Kraytem A, Giusti V. Gastric bypass: management of complications and food tolerance. *Revue Medicale Suisse*. 2008;4(151):836.
- Faria SL, Faria OP, Cardeal MD, Ito MK. Validation study of multi-frequency bioelectrical impedance with dual-energy X-ray absorptiometry among obese patients. *Obes Surg*. 2014;24(9):1476–80.
- Lee SY, Gallagher D. Assessment methods in human body composition. *Curr Opin Clin Nutr Metab Care*. 2008;11(5):566.
- Schiavo L, Pilone V, Tramontano S, Rossetti G, Iannelli A. May bioelectrical impedance analysis method be used in alternative to the dual-energy X-ray absorptiometry in the assessment of fat mass and fat-free mass in patients with obesity? Pros, cons, and perspectives. *Obes Surg*. 2020;30(8):3212–5.

34. Cermak NM, Res PT, de Groot LC, Saris WH, Van Loon LJ. Protein supplementation augments the adaptive response of skeletal muscle to resistance-type exercise training: a meta-analysis. *Am J Clin Nutr.* 2012;96(6):1454–64.
35. Phillips SM, Hartman JW, Wilkinson SB. Dietary protein to support anabolism with resistance exercise in young men. *J Am Coll Nutr.* 2005;24(2):134S–9S.
36. Biolo G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *Am J Physiol Endocrinol Metab.* 1997;273(1):E122–E9.
37. Muschitz C, Kocijan R, Haschka J, Zendeli A, Pirker T, Geiger C, et al. The impact of vitamin D, calcium, protein supplementation, and physical exercise on bone metabolism after bariatric surgery: the BABS study. *J Bone Miner Res.* 2016;31(3):672–82.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

