

The authors regret that in the October 2013 article “Widespread White Matter Alterations in Patients with End-Stage Renal Disease: A Voxelwise Diffusion Tensor Imaging Study” (*AJNR Am J Neuroradiol* 2013;34:1945–51), the MNI coordinates of brain regions shown in Tables 2–6 were incorrect. The MNI coordinates in Tables 2–6 should be corrected by shifting –20 in Y direction and +20 in Z direction. The tables are reproduced below with the corrected segments shown in bold.

Table 2: MNI coordinates of major areas with significant AD differences along with their means and standard deviations

Brain Regions	MNI Coordinate (mm)			AD (Mean \pm SD) $\times 10^{-3}$ mm 2 /s	
	X	Y	Z	ESRD	Healthy
Rt. RLIC	25	–36	8	1.43 \pm 0.17	1.39 \pm 0.14
Lt. RLIC	–27	–34	6	1.66 \pm 0.21	1.57 \pm 0.13
Lt. Fminor	–21	32	8	1.43 \pm 0.07	1.33 \pm 0.07
Pons	1	–34	–36	1.85 \pm 0.18	1.73 \pm 0.11

Table 3: MNI coordinates of major areas with significant RD differences along with their means and standard deviations

Brain Regions	MNI Coordinate (mm)			RD (Mean \pm SD) $\times 10^{-3}$ mm 2 /s	
	X	Y	Z	ESRD	Healthy
GCC	1	19	19	0.94 \pm 0.2	0.86 \pm 0.14
SCC	1	–34	25	0.65 \pm 0.15	0.56 \pm 0.08
Rt. Fmajor	24	–74	12	0.66 \pm 0.11	0.62 \pm 0.09
Lt. Fmajor	–28	–62	12	0.67 \pm 0.09	0.61 \pm 0.11
Rt. PCR	25	–43	27	0.64 \pm 0.09	0.63 \pm 0.04
Lt. PCR	–28	–40	27	0.66 \pm 0.09	0.65 \pm 0.04
Rt. ACR	17	31	14	0.64 \pm 0.06	0.62 \pm 0.05
Lt. ACR	–20	36	12	0.63 \pm 0.06	0.59 \pm 0.05
Pons	3	–33	–34	0.82 \pm 0.19	0.71 \pm 0.12

Table 4: MNI coordinates of major areas with significant MD differences along with their means and standard deviations

Brain Regions	MNI Coordinate (mm)			MD (Mean \pm SD) $\times 10^{-3}$ mm 2 /s	
	X	Y	Z	ESRD	Healthy
GCC	1	19	12	1.16 \pm 0.07	1.08 \pm 0.05
SCC	1	–33	25	1.12 \pm 0.13	1.02 \pm 0.06
Rt. SS	31	–51	11	1.27 \pm 0.22	1.21 \pm 0.15
Lt. SS	–39	–44	5	1.07 \pm 0.12	1.04 \pm 0.08
Rt. Fmajor	25	–62	17	0.96 \pm 0.11	0.89 \pm 0.04
Lt. Fmajor	–30	–54	17	1.15 \pm 0.11	0.99 \pm 0.09
Rt. PCR	27	–45	24	1.07 \pm 0.12	0.92 \pm 0.06
Lt. PCR	–29	–45	21	1.21 \pm 0.18	1.11 \pm 0.15
Pons	2	–34	–36	1.28 \pm 0.13	1.13 \pm 0.09

Table 5: MNI coordinates of major areas with significant FA differences along with their means and standard deviations

Brain Regions	MNI Coordinate (mm)			FA (Mean \pm SD)	
	X	Y	Z	ESRD	Healthy
GCC	3	20	12	0.6 \pm 0.06	0.67 \pm 0.04
SCC	3	–34	25	0.65 \pm 0.05	0.72 \pm 0.05
Rt. SS	31	–48	14	0.50 \pm 0.04	0.56 \pm 0.04
Lt. SS	–27	–59	14	0.46 \pm 0.05	0.56 \pm 0.07
Rt. MCP	13	–38	–38	0.45 \pm 0.03	0.53 \pm 0.04

Table 6: MNI coordinates of areas with significant correlations along with their correlation coefficients

Brain Regions	MNI Coordinate (mm)			Correlation Coefficient	
	X	Y	Z		
RD and duration					
Lt. Pons	–16		–23	–26	0.7093
RD and CASI					
Lt. ACR	–28		33	39	–0.7442
Rt. SCR	31		–13	41	–0.6154
MD and duration					
Rt. RLIC	33		–40	14	0.6009
Lt. FWM	–21		43	15	0.6287
Lt. Pons	–12		–23	–26	0.6675
MD and CASI					
Lt. ACR	–28		13	39	–0.7657
Rt. SCR	31		–13	41	–0.5463
FA and CASI					
Lt. ACR	–17		20	38	0.7067
Lt. PCR	–20		–49	44	0.6117
Rt. SCR	33		–14	46	0.6815